

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 64 (1989)  
  
**Artikel:** On the outradius of the Teichmüller space.  
**Autor:** Nakanishi, T. / Yamamoto, H.-O  
**DOI:** <https://doi.org/10.5169/seals-48947>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 27.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On the outradius of the Teichmüller space

TOSHIHIRO NAKANISHI and HIRO-O YAMAMOTO

Dedicated to Professor Kôtarô Oikawa on his 60th birthday

**Abstract.** Let  $\Gamma$  be a fuchsian group which preserves the unit disc  $\Delta$  and hence also its complement  $\Delta^*$  in the Riemann sphere  $\hat{\mathbb{C}}$ . The Bers embedding represents the Teichmüller space  $T(\Gamma)$  of  $\Gamma$  in the space  $B(\Delta^*, \Gamma)$  of bounded quadratic differentials for  $\Gamma$  in  $\Delta^*$ . Then,  $T(\Gamma)$  is included in the closed ball centred at the origin of radius 6 in  $B(\Delta^*, \Gamma)$  with respect to the norm employed in a paper by Nehari [The Schwarzian derivative and Schlicht functions; Bull. Amer. Math. Soc. 55 (1949), 545–551]. In other words the outradius  $o(\Gamma)$  of  $T(\Gamma)$  is not greater than 6. The purpose of this paper is to give a complete characterization of a fuchsian group  $\Gamma$  for which the outradius  $o(\Gamma)$  of  $T(\Gamma)$  attains this extremal value 6. The main theorem is: Let  $\Gamma$  be a fuchsian group preserving  $\Delta^*$ . Then the outradius  $o(\Gamma)$  of the Teichmüller space  $T(\Gamma)$  equals 6 if and only if for any positive number  $d$ , either (i) there exists a hyperbolic disc of radius  $d$  precisely invariant under the trivial subgroup, or (ii) there exists the collar of width  $d$  about the axis of a hyperbolic element of  $\Gamma$ .

### Introduction

Let  $\Gamma$  be a fuchsian group which preserves the unit disc  $\Delta = \{z \in \mathbb{C} : |z| < 1\}$  and hence also the complement  $\Delta^*$  of the closure  $\Delta$  in the Riemann sphere  $\hat{\mathbb{C}}$ . In this paper we treat the Teichmüller space  $T(\Gamma)$  of  $\Gamma$  represented as a subregion of the Banach space  $B(\Delta^*, \Gamma)$  of bounded quadratic differentials for  $\Gamma$  by means of the Bers embedding ([1]). By a classical theory of Nehari [8], in  $B(\Delta^*, \Gamma)$  the Teichmüller space  $T(\Gamma)$  is included in the closed ball of radius 6 centered at the origin. In other words the outradius  $o(\Gamma)$  of  $T(\Gamma)$  does not exceed 6. The purpose of this paper is to give a complete characterization of those fuchsian groups  $\Gamma$  for which  $o(\Gamma)$  attains the extremal value 6. Our main theorem is:

**THEOREM 1.1.** *Let  $\Gamma$  be a fuchsian group preserving  $\Delta^*$ . Then the outradius  $o(\Gamma)$  of the Teichmüller space  $T(\Gamma)$  equals 6 if and only if  $\Gamma$  satisfies one of the following conditions:*

(O<sub>1</sub>) *For any positive number  $d$ , there exists a hyperbolic disc of radius  $d$  which is precisely invariant under the trivial subgroup  $\{1\}$  of  $\Gamma$ .*

(O<sub>2</sub>) *For any positive number  $d$ , there exists the collar of width  $d$  about the axis of a hyperbolic element of  $\Gamma$ .*

We shall refer to the notations in the theorem in Section 1. Theorem 1.1 means a geometric condition of the fuchsian group  $\Gamma$  reflects the property that  $o(\Gamma)$  equals 6 or not. First we discuss in Section 1 preliminary notions and definitions concerning fuchsian groups and Teichmüller spaces. In Section 2 we give some lemmas needed to prove Theorem 1.1. A proof of Theorem 1.1 is carried out in Sections 3 and 4. The final section, Section 5, is devoted to some examples concerning the conditions  $(O_1)$  and  $(O_2)$  of Theorem 1.1.

We wish to acknowledge many suggestions by Professors M. Masumoto and M. Shiba which were helpful and improved this paper.

## 1. Preliminaries

1.1. Our basic reference for the content of this section is [2]. Let  $\text{Möb}(\hat{\mathbb{C}})$  be the group of Möbius transformations of the Riemann sphere  $\hat{\mathbb{C}}$  onto itself; that is, mappings

$$\gamma(z) = (az + b)/(cz + d), \quad a, b, c, d \in \mathbb{C}, \quad ad - bc = 1.$$

Let  $\Delta^* = \{z \in \mathbb{C} : |z| > 1\} \cup \{\infty\}$ . The hyperbolic metric defined on  $\Delta^*$ ,

$$\rho^*(z) |dz| = (|z|^2 - 1)^{-1} |dz|$$

has constant curvature  $-4$ . Geodesics with respect to this metric or *hyperbolic lines* are circles or straight lines which are orthogonal to the unit circle  $\{|z| = 1\}$ . Denote by  $\text{Möb}(\Delta)$  the subgroup of  $\text{Möb}(\hat{\mathbb{C}})$  which preserves the unit disc  $\Delta$  (as well as  $\Delta^*$ ). Then all transformations of  $\text{Möb}(\Delta)$  are of the form:

$$\gamma(z) = (az + \bar{b})/(bz + \bar{a}), \quad |a|^2 - |b|^2 = 1. \quad (1.1)$$

A Möbius transformation (1.1) is an orientation preserving hyperbolic motion, which means that  $\rho^*(z) = \rho^*(\gamma(z)) |\gamma'(z)|$  for  $z \in \Delta^*$ . The hyperbolic distance between  $z$  and  $w$  in  $\Delta^*$  will be denoted by  $d(z, w)$ .

A *fuchsian group*  $\Gamma$  is a subgroup of  $\text{Möb}(\Delta)$  which acts discontinuously on  $\Delta$  and hence also on  $\Delta^*$ . Let  $G$  be a subgroup of  $\Gamma$  which is also a fuchsian group. A subset  $D$  of  $\Delta^*$  is said to be *precisely invariant* under  $G$  if  $\gamma(D) = D$  for all  $\gamma \in G$  and  $\gamma(D) \cap D = \emptyset$  for all  $\gamma \in \Gamma - G$ . Let  $\gamma$  be a hyperbolic element of  $\Gamma$ . The hyperbolic line in  $\Delta^*$  connecting the fixed points of  $\gamma$  is called the axis of  $\gamma$  and denoted by  $A^*(\gamma)$ . Let  $\text{Stab}(\Gamma, A^*(\gamma))$  be the stabilizer of  $A^*(\gamma)$  in  $\Gamma$ , that is,

$$\text{Stab}(\Gamma, A^*(\gamma)) = \{\eta \in \Gamma : \eta(A^*(\gamma)) = A^*(\gamma)\}.$$

Then  $\text{Stab}(\Gamma, A^*(\gamma))$  is either the infinite cyclic group generated by a hyperbolic transformation or a group generated by two elliptic elements of order 2. Let  $U^*(d) = U^*(d, A^*(\gamma)) = \{z \in \Delta^* : d(z, A^*(\gamma)) < d\}$ . We say that  $U^*(d, A^*(\gamma))$  is the *collar of width  $d$  about  $A^*(\gamma)$*  if  $U^*(d)$  is precisely invariant under  $\text{Stab}(\Gamma, A^*(\gamma))$ .

Let  $\gamma(z) = (az + \bar{b})/(bz + \bar{a})$  be a transformation in  $\text{Möb}(\Delta)$ . If  $b \neq 0$ , then  $\infty$  is not fixed by  $\gamma$ . In this case the *isometric circle*  $I(\gamma)$  of  $\gamma$  is defined to be  $\{z : |bz + \bar{a}| = 1\}$ . We mean by the exterior of  $I(\gamma)$  the complementary region of  $I(\gamma)$  which contains  $\infty$ . Let  $\Gamma$  be a fuchsian group. Suppose that  $\infty$  is not fixed by any element of  $\Gamma - \{1\}$ . The set  $\mathcal{F}(\Gamma)$  of all points in  $\Delta^*$  exterior to the isometric circles of all the transformations of  $\Gamma - \{1\}$  is referred to here the *Ford region* for  $\Gamma$  in  $\Delta^*$ . The Ford region is a fundamental region for the action of  $\Gamma$  in  $\Delta^*$  ([2, Sec. 15]).

1.2. Let  $\Gamma$  be a fuchsian group. A holomorphic function  $\phi$  in  $\Delta^*$  is a *bounded quadratic differential* for  $\Gamma$  if it satisfies

$$\phi(z) = \phi(\gamma(z))\gamma'(z)^2 \quad \text{for all } \gamma \in \Gamma \text{ and } z \in \Delta^*, \quad (1.2)$$

and

$$\|\phi\| = \sup_{z \in \Delta^*} \rho^*(z)^{-2} |\phi(z)| < \infty, \quad (1.3)$$

where  $\rho^*(\infty)^{-2} |\phi(\infty)|$  means  $\lim_{z \rightarrow \infty} \rho^*(z)^{-2} |\phi(z)|$ . The Banach space of bounded quadratic differentials for  $\Gamma$  with the norm,  $\|\cdot\|$ , defined by (1.3) is denoted by  $B(\Delta^*, \Gamma)$ .

A quasiconformal automorphism  $w$  of  $\hat{\mathbb{C}}$  is said to be compatible with a fuchsian group  $\Gamma$  if the correspondence  $\gamma \rightarrow w\gamma w^{-1}$  defines an isomorphism of  $\Gamma$  into  $\text{Möb}(\hat{\mathbb{C}})$ . Let  $Q(\Delta^*, \Gamma)$  be the family of all quasiconformal automorphisms of  $\hat{\mathbb{C}}$  compatible with  $\Gamma$  and conformal in  $\Delta^*$ . The *Teichmüller space*  $T(\Gamma)$  of  $\Gamma$  is the set of the Schwarzian derivatives  $[w] = (w''/w')' - (1/2)(w''/w')^2$  of  $w \in Q(\Delta^*, \Gamma)$  in  $\Delta^*$ . By the well known properties of the Schwarzian derivative,  $[w](z)$ ,  $w \in Q(\Delta^*, \Gamma)$ , satisfies (1.2). Moreover as the Schwarzian derivative of a univalent function in  $\Delta^*$ , it holds that ([8]):

$$\|[w]\| \leq 6. \quad (1.4)$$

Thus  $T(\Gamma) \subset B(\Delta^*, \Gamma)$ . The outradius  $o(\Gamma)$  of  $T(\Gamma)$  is defined to be  $\sup \{\|\phi\| : \phi \in T(\Gamma)\}$ . By (1.4) we have  $o(\Gamma) \leq 6$ .



## 2. Some lemmas

LEMMA 2.1. *Let  $\{\gamma_n\}$  be a sequence in  $\text{Möb}(\Delta)$ . If there exists  $R > 1$  such that  $|\gamma_n(\infty)| > R$  for all  $n$ , then  $\{\gamma_n\}$  contains a subsequence which converges to a transformation in  $\text{Möb}(\Delta)$ .*

*Proof.* Let  $\gamma_n(z) = (a_n z + \bar{b}_n)/(b_n z + \bar{a}_n)$ ,  $|a_n|^2 - |b_n|^2 = 1$ . By assumption  $|\gamma_n(\infty)| = |a_n/b_n| = |a_n|(|a_n|^2 - 1)^{-1/2} > R > 1$ . It follows that  $|a_n| \leq R(R^2 - 1)^{-1/2}$  and  $|b_n| \leq (R^2 - 1)^{-1/2}$ . Choose subsequences  $|a_{n_j}|$  and  $|b_{n_j}|$  in such a way that  $\lim_{j \rightarrow \infty} a_{n_j} = a_0$  and  $\lim_{j \rightarrow \infty} b_{n_j} = b_0$ , for some  $a_0$  and  $b_0 \in \mathbb{C}$ . Then the subsequence  $\{\gamma_{n_j}\}$  converges to  $\gamma_0(z) = (a_0 z + \bar{b}_0)/(b_0 z + \bar{a}_0)$ .

LEMMA 2.2. *Let  $\{\gamma_n\}$  be a sequence in  $\text{Möb}(\Delta)$  converging to the identity transformation 1. Then there exist a subsequence  $\{\gamma_{n_j}\}$  and a sequence of integers  $\{k(j)\}$  such that  $\gamma_{n_j}^{k(j)}$  converges to a transformation  $\gamma$  which is neither 1 nor an elliptic transformation of order 2.*

*Proof.* By replacing  $\{\gamma_n\}$  with a suitable subsequence, if necessary, we may assume that  $\{\gamma_n\}$  contains only elliptic or parabolic or hyperbolic elements. Here we give a proof only for the case where  $\{\gamma_n\}$  contains only elliptic elements, but other cases can be treated in a similar manner.

Assume that  $\gamma_n$  is elliptic of order  $v_n$ . Let  $p_n$  be the fixed point of  $\gamma_n$  in  $\Delta$ . Then  $\gamma_n^k$  corresponds to the matrix in  $SL(2, \mathbb{C})$

$$\frac{1}{p_n - \bar{p}_n^{-1}} \begin{bmatrix} \lambda_n^{-k} p_n - \lambda_n^k \bar{p}_n^{-1} & (\lambda_n^k - \lambda_n^{-k}) p_n \bar{p}_n^{-1} \\ -(\lambda_n^k - \lambda_n^{-k}) & \lambda_n^k p_n - \lambda_n^{-k} \bar{p}_n^{-1} \end{bmatrix},$$

where  $\lambda_n = e^{2\pi i/v_n}$ . Since  $\gamma_n \rightarrow 1$ , we have that  $v_n \rightarrow \infty$  and  $(\lambda_n - \lambda_n^{-1})/(p_n - \bar{p}_n^{-1}) \rightarrow 0$  as  $n \rightarrow \infty$ . Choose a subsequence  $\{\gamma_{n_j}\}$  so that  $p_{n_j}$  converges to a point  $p$ . If  $|p| < 1$ , then let  $k(j)$  be a nearest integer to  $v_{n_j}/3$ . Then  $\gamma_{n_j}^{k(j)}$  converges to the elliptic transformation  $\gamma$  of order 3 represented by the matrix

$$\frac{1}{p - \bar{p}^{-1}} \begin{bmatrix} \lambda^{-1} p - \lambda \bar{p}^{-1} & (\lambda - \lambda^{-1}) p \bar{p}^{-1} \\ -(\lambda - \lambda^{-1}) & \lambda p - \lambda^{-1} \bar{p}^{-1} \end{bmatrix},$$

where  $\lambda = e^{2\pi i/3}$ . If  $|p| = 1$ , then choose  $k(j)$  so that  $|\lambda_{n_j}^{k(j)} - \lambda_{n_j}^{-k(j)}|/|p_{n_j} - \bar{p}_{n_j}^{-1}|$  ( $j = 1, 2, \dots$ ) are bounded and bounded below by a positive number. Then, by replacing  $\{\gamma_{n_j}\}$  with some subsequence, we may assume that  $(\lambda_{n_j}^{k(j)} -$

$\lambda_{n_j}^{-k(j)}/(p_{n_j} - \bar{p}_{n_j}^{-1})$  converges to a number  $\xi \in \mathbb{C} - \{0\}$ . Then  $|\gamma_{n_j}^{k(j)}(\infty)|$  converges to  $|\xi|^{-1}(1 + |\xi|^2)^{1/2} > 1$ . By Lemma 2.1  $\gamma_{n_j}^{k(j)}$  converges to a Möbius transformation  $\gamma$ . Since  $|p_{n_j} - \bar{p}_{n_j}^{-1}| \rightarrow 0$ , we have that  $|\lambda_{n_j}^{k(j)} - \lambda_{n_j}^{-k(j)}| \rightarrow 0$ . Thus it follows that  $|\operatorname{tr} \gamma_{n_j}^{k(j)}|^2 = |(\lambda_{n_j}^{k(j)} - \lambda_{n_j}^{-k(j)})^2 + 4| \rightarrow 4$  and hence that  $|\operatorname{tr} \gamma|^2 = 4$ . Since  $|\gamma(\infty)| = |\xi|^{-1}(1 + |\xi|^2)^{1/2} \neq \infty$ ,  $\gamma$  cannot be the identity transformation. Now we can conclude that  $\gamma$  is a parabolic transformation.

LEMMA 2.3. *Let  $k(z) = z + z^{-1}$  be defined in  $\Delta^*$ . Then,*

- (i)  $[k](z) = 6(z^2 - 1)^{-2}$ ;
- (ii)  $\rho^*(z)^{-2} |[k](z)| \leq 6$ , where the equality holds if and only if  $z$  lies on the segment  $A_0^* = \{z \in \Delta^* : \operatorname{Im} z = 0\} \cup \{\infty\}$ ;
- (iii) *If  $\gamma$  is a transformation in  $\operatorname{Möb}(\Delta)$  for which*

$$[k](z) = [k](\gamma(z))\gamma'(z)^2, \quad (2.1)$$

*then  $\gamma$  is either the identity transformation, or an elliptic transformation of order 2 which preserves  $A_0^*$ , or a hyperbolic transformation with the axis  $A_0^*$ .*

*Proof.* Conclusions (i) and (ii) are verified by a direct calculation. Now let  $\gamma(z) = (az + \bar{b})/(bz + \bar{a})$  be a transformation satisfying (2.1). Then (i) yields

$$(z^2 - 1)^2 = ((a^2 - b^2)z^2 + 2(a\bar{b} - \bar{a}b)z - (\bar{a}^2 - \bar{b}^2))^2.$$

Comparing the coefficients of the above polynomials in  $z$ , it follows that  $a\bar{b} = \bar{a}b$  and  $a^2 + \bar{b}^2 = (-a + \bar{b})^2/(-b + \bar{a})^2 = 1$ . Therefore  $\gamma$  preserves the set of two points  $\{1, -1\}$ . If  $\gamma$  fixes each of 1 and  $-1$ , then  $\gamma$  is either the identity transformation or a hyperbolic transformation with the axis  $A_0^*$ . If  $\gamma$  interchanges 1 and  $-1$ , then  $\gamma$  is elliptic of order 2 and preserves  $A_0^*$ .

### 3. Proof of theorem 1.1.(I)

In this section we shall prove the “only if” part of Theorem 1.1, that is, that a fuchsian group  $\Gamma$  with  $o(\Gamma) = 6$  satisfies one of the conditions  $(O_1)$  and  $(O_2)$ . The proof is rather lengthy, hence we divide it into several steps (3.1–3.5).

3.1. Suppose that  $o(\Gamma) = 6$  holds for a fuchsian group  $\Gamma$ . Let  $\{\varepsilon_n\}$  be a sequence of positive numbers decreasing to 0. By definition there exists a sequence of bounded quadratic differentials  $\{\phi_n\}$  in  $T(\Gamma)$  for which  $6 - \varepsilon_n < \|\phi_n\| \leq 6$ . Choose a point  $z_n$  of  $\Delta^*$  so that  $6 - \varepsilon_n < \rho^*(z_n)^{-2} |\phi_n(z_n)| \leq 6$ . Here by

replacing  $z_n$  with another point nearby if necessary, we may assume that  $z_n$  is not fixed by any element of  $\Gamma - \{1\}$ . Conjugate  $\Gamma$  by a transformation  $h_n$  in  $\text{Möb}(\Delta)$  with  $h_n(\infty) = z_n$  and denote  $h_n\Gamma h_n^{-1}$  by  $\Gamma_n$ . (Here we remark that, by the choice of  $z_n$ , the Ford region for  $\Gamma_n$  can be defined. We use this fact in the next paragraph.) It is easy to see that  $\psi_n(z) = \phi_n(h_n(z))h'_n(z)^2$  belongs to  $T(\Gamma_n)$ , and that  $6 - \varepsilon_n < \rho^*(\infty)^{-2} |\psi_n(\infty)| \leq 6$ . Let  $f_n$  be the solution of the differential equation  $[f_n](z) = \psi_n(z)$ , which has the following normalized form

$$f_n(z) = z + b_{n1}z^{-1} + b_{n2}z^{-2} + \dots$$

Here we may assume that  $b_{n1}$  is real positive, because if otherwise, we need only to replace  $h_n(z)$  with  $h_n(e^{-i\theta}z)$  and  $f_n(z)$  with  $e^{i\theta}f_n(e^{-i\theta}z)$ , where  $\theta = -(1/2) \arg b_{n1}$ . Since  $\psi_n \in T(\Gamma_n)$ ,  $f_n$  is univalent in  $\Delta^*$ . Then the area theorem (cf. [9; p. 19]) yields

$$b_{n1}^2 + \sum_{v=2}^{\infty} v |b_{nv}|^2 \leq 1. \quad (3.1)$$

Since  $6 - \varepsilon_n < \rho^*(\infty)^{-2} |\psi_n(\infty)| = 6b_{n1} \leq 6$ ,  $b_{n1}$  converges to 1. Then the inequality (3.1) implies that  $f_n$  converges to  $k(z) = z + z^{-1}$  locally uniformly in  $\Delta^*$  with respect to the spherical metric of  $\hat{\mathbb{C}}$ . Also,  $\psi_n$  converges to  $[k](z) = 6(z^2 - 1)^{-2}$  locally uniformly in  $\Delta^*$  with respect to the spherical metric.

3.2. For each  $n$  we choose a  $\gamma_n \in \Gamma_n - \{1\}$  in such a way that the radius  $r_n$  of the isometric circle of  $\gamma_n$  is the largest of those of the elements in  $\Gamma_n - \{1\}$ . If  $\{r_n\}$  contains a subsequence  $\{r_{n_j}\}$  which converges to 0, then the Ford region  $\mathcal{F}(\Gamma_{n_j})$  converges to  $\Delta^*$ . This means that  $\Gamma$  satisfies the condition  $(O_1)$ , since  $\Gamma$  and  $\Gamma_{n_j}$  are conjugate in  $\text{Möb}(\Delta)$  and we can find a disc contained in  $\mathcal{F}(\Gamma_{n_j})$  whose hyperbolic radius tends to  $\infty$  as  $j$  tends to  $\infty$ . Therefore we need only to consider the case that  $\{r_n\}$  is bounded below by a positive number  $r_0$ . Then, for  $\gamma_n(z) = (a_n z + \bar{b}_n)/(b_n z + \bar{a}_n)$ ,  $|a_n|^2 - |b_n|^2 = 1$ , we have  $|\gamma_n(\infty)| = |a_n/b_n| \geq (1 + r_0^2)^{1/2}$ . By Lemma 2.1 it follows that a subsequence of  $\{\gamma_n\}$  converges to a Möbius transformation  $\gamma_0$ . For convenience we denote this subsequence again by  $\{\gamma_n\}$ .

Since  $\gamma_n \in \Gamma_n$ , it holds that  $\psi_n(z) = \psi_n(\gamma_n(z))\gamma'_n(z)^2$ . By letting  $n \rightarrow \infty$  in this equation, we obtain that  $[k](z) = [k](\gamma_0(z))\gamma'_0(z)^2$ . By Lemma 2.3,  $\gamma_0$  is either the identity transformation or an elliptic transformation preserving  $A_0^*$  or a hyperbolic transformation with the axis  $A_0^*$ .

3.3. The above sequence  $\{\gamma_n\}$  contains only finitely many parabolic elements and elliptic elements of order  $> 2$ . To see this, assume that  $\{\gamma_n\}$  contains an

infinite subsequence  $\{\eta_n\}$ , where  $\eta_n$  is either parabolic or elliptic of order  $>2$ . As the limit of  $\eta_n$ ,  $\gamma_0$  is the identity transformation. By Lemma 2.2 there exist a subsequence  $\{\eta_{n_j}\}$  and a sequence of integers  $\{k(j)\}$  such that  $\eta_{n_j}^{k(j)}$  converges to a Möbius transformation  $\eta$  which is neither the identity nor elliptic of order 2. As the limit of parabolic or elliptic elements,  $\eta$  is not hyperbolic. By letting  $j \rightarrow \infty$  in the equation  $\psi_{n_j}(z) = \psi_{n_j}(\eta_{n_j}^{k(j)}(z))(\eta_{n_j}^{k(j)})'(z)^2$ , we obtain that  $[k](z) = [k](\eta(z))\eta'(z)^2$ . However this contradicts Lemma 2.3. In a similar manner we can show that for a subsequence  $\{\eta_n\}$  of hyperbolic elements in  $\{\gamma_n\}$ , if any, the axis of  $\eta_n$  converges to  $A_0^*$ . Assume that the axis of  $\eta_n$  does not converge to  $A_0^*$ . Then  $\gamma_0$  is again the identity transformation. Choose a subsequence  $\{\eta_{n_j}\}$  and a sequence of integers  $\{k(j)\}$  so that  $\eta_{n_j}^{k(j)}$  converges to a nontrivial transformation  $\eta$ . As the limit of hyperbolic elements  $\eta$  is either parabolic or hyperbolic. If  $\eta$  is hyperbolic, then by the assumption the axis of  $\eta$  is different from  $A_0^*$ . Then as above we can deduce a contradiction.

Now, by eliminating a finite number of elements and choosing a subsequence, we can assume that one of the following two cases occurs for  $\{\gamma_n\}$ :

Case (A).  $\{\gamma_n\}$  contains only elliptic elements of order 2, and  $\gamma_0$  is an elliptic transformation of order 2 and preserves  $A_0^*$ .

Case (B).  $\{\gamma_n\}$  contains only hyperbolic elements, and  $\gamma_0$  is either the identity transformation or a hyperbolic transformation. Moreover the axis  $A^*(\gamma_n)$  converges to  $A_0^*$ .

We shall consider the two cases (A) and (B) separately.

3.4. Case (A). In this case we shall show that  $\Gamma$  satisfies the condition  $(O_1)$ , or otherwise that we can transfer the argument to Case (B).

We take a  $\delta_n \in \Gamma_n$  in such a way that the radius  $r'_n$  of the isometric circle of  $\delta_n$  is the largest of those of elements in  $\Gamma_n - \{1, \gamma_n\}$ . If  $\{r'_n\}$  contains a subsequence converging to 0, then we can see easily that  $\Gamma$  satisfies the condition  $(O_1)$ . On the other hand if  $\{r'_n\}$  is bounded below by a positive number, then by Lemma 2.1 a subsequence of  $\{\delta_n\}$  converges to a Möbius transformation  $\delta_0$ . Choosing a subsequence and relabelling if necessary, we can assume that Case (A) or (B) occurs for  $\{\delta_n\}$ . If Case (A) occurs, then both  $\gamma_n$  and  $\delta_n$  are elliptic of order 2. Since  $\gamma_n \neq \delta_n$ ,  $\gamma_n\delta_n$  is hyperbolic and converges to  $\gamma\delta$ . By applying the argument in 3.3 we can show that Case (B) occurs for  $\{\gamma_n\delta_n\}$ . Therefore Case (B) occurs for  $\{\delta_n\}$  or  $\{\gamma_n\delta_n\}$ .

3.5. Case (B). In this case we shall show that  $\Gamma$  satisfies the condition  $(O_2)$ .

Let  $R_n$  be the region exterior to the isometric circles of  $\gamma_n$  and  $\gamma_n^{-1}$ . Among the elements of  $\Gamma_n - \text{Stab}(\Gamma_n, A^*(\gamma_n))$  whose isometric circles meet  $R_n$ , choose a  $\delta_n$  which has the isometric circle of the largest radius  $r'_n$ . Assume that  $\{r'_n\}$  is

bounded below by a positive number. By Lemma 2.1 we can find a subsequence, which is denoted again by  $\{\delta_n\}$ , of  $\{\delta_n\}$  converging to a Möbius transformation  $\delta_0$ . As in 3.2 we can show that  $\delta_0$  preserves  $A_0^*$ . For  $n \geq 0$  denote by  $G_n$  the group generated by  $\gamma_n$  and  $\delta_n$ . Since  $\delta_n \in \Gamma_n - \text{Stab}(\Gamma_n, A^*(\gamma_n))$ ,  $G_n$  is non-elementary for  $n \geq 1$ . Let  $\chi_n: G_0 \rightarrow G_n$  be the mapping which is the canonical extension of the correspondings  $\chi_n(\gamma_0) = \gamma_n$  and  $\chi_n(\delta_0) = \delta_n$ . By [4, Proposition 1 and Theorem 2]  $G_n$  is fuchsian and  $\chi_n$  is homomorphism of  $G_0$  onto  $G_n$  for sufficiently large  $n$ . Thus  $G_0$  is a non-elementary fuchsian group. However  $G_0$  is elementary because both  $\gamma_n$  and  $\delta_n$  preserve  $A_0^*$ . This is a contradiction. Therefore  $\{r'_n\}$  contains a subsequence converging to 0. Since  $\Gamma$  is conjugate to  $\Gamma_n$  in  $\text{Möb}(\Delta)$ ,  $\Gamma$  satisfies the condition  $(O_2)$ . The first half of the proof is now completed.

#### 4. Proof of Theorem 1.1.(II)

In this section we shall prove the “if” part of Theorem 1.1 and complete the proof.

4.1. We consider also the action of a fuchsian group  $\Gamma$  on the unit disc  $\Delta$  with the hyperbolic metric  $\rho(z) |dz| = (1 - |z|^2)^{-1} |dz|$ . Let  $R(z) = 1/\bar{z}$  be the reflection with respect to the unit circle. Note that  $R$  is an isometry of  $(\Delta^*, \rho^* |dz|)$  onto  $(\Delta, \rho |dz|)$  and that  $R$  conjugate  $\Gamma$  to itself;  $R^{-1}\Gamma R = \Gamma$ . Let  $A_0$  be the segment  $\{z \in \Delta: \text{Im } z = 0\}$ . If  $\text{Möb}(A_0)$  denotes the subgroup of  $\text{Möb}(\Delta)$  which preserves  $A_0$  and also  $A_0^*$ , then each element  $\gamma$  of  $\text{Möb}(A_0) - \{1\}$  has the following form:

$$\gamma(z) = \frac{(1 + \lambda)z + (1 - \lambda)}{(1 - \lambda)z + (1 + \lambda)}, \quad \lambda > 0 \quad \text{and} \quad \lambda \neq 1, \quad (4.1a)$$

if  $\gamma$  is hyperbolic, and

$$\gamma(z) = \frac{(1 + \lambda)z - (1 - \lambda)}{(1 - \lambda)z - (1 + \lambda)}, \quad \lambda > 0 \quad \text{and} \quad \lambda \neq 1, \quad (4.1b)$$

if  $\gamma$  is elliptic of order 2. Neither a parabolic transformation nor an elliptic transformation of order  $\neq 2$  belongs to  $\text{Möb}(A_0)$ .

Let  $V$  be the segment  $\{\alpha: -3/2 < \alpha < 1/2\}$ . For each  $\alpha$  in  $V$ , define a function  $\mu_\alpha$  in  $\hat{\mathbb{C}}$ :

$$\mu_\alpha(z) = \begin{cases} (\delta(\alpha) - 1)(1 - z^2)/(1 - \bar{z}^2) & \text{for } z \in \Delta, \\ 0 & \text{for } z \in \bar{\Delta}^*, \end{cases} \quad (4.2)$$

where  $\delta(\alpha) = (1 - 2\alpha)^{1/2}$  (with  $\delta(0) = 1$ ). Note that  $\text{ess. sup } |\mu_\alpha| = |\delta(\alpha) - 1| < 1$ . A direct calculation using (4.1a) and (4.1b) yields:

$$\mu_\alpha(z) = \mu_\alpha(\gamma(z)) \overline{\gamma'(z)} / \gamma'(z) \quad \text{for all } \gamma \in \text{Möb}(A_0). \quad (4.3)$$

Let  $h(z) = i(1 - z)/(1 + z)$  be a Möbius transformation sending  $\Delta^*$  onto the lower half plane  $\mathcal{H}^*$ . Then the Beltrami equation

$$w_{\bar{z}} = \mu_\alpha w_z$$

has a solution  $W_\alpha$  expressed by

$$W_\alpha(z) = \begin{cases} -2i\delta(-i)^\delta / [h(z)\overline{h(z)}^{\delta-1} - (-i)^\delta] & \text{for } z \in \Delta \\ -2i\delta(-i)^\delta / [h(z)^\delta - (-i)^\delta] & \text{for } z \in \bar{\Delta}^*, \end{cases}$$

with  $\delta = \delta(\alpha)$  and a fixed branch of  $z^\delta$  in  $\mathcal{H}^*$ . The function  $W_\alpha$  is a quasiconformal automorphism of  $\hat{\mathbb{C}}$ , conformal in  $\Delta^*$  and has the normalized form:  $W_\alpha(z) = z + O(|z|^{-1})$  as  $z \rightarrow \infty$ . Moreover,  $[W_\alpha](z) = 4\alpha(z^2 - 1)^{-2}$  in  $\Delta^*$ .

*Remark.* We learned the above construction of the function  $W_\alpha$  from a paper by Kalme [5].

For the remainder of this section, the notion of convergence of functions defined in  $\hat{\mathbb{C}}$  will be considered in the spherical metric.

4.2. First we shall show that  $o(\Gamma) = 6$  holds for any fuchsian group  $\Gamma$  satisfying the condition  $(O_2)$ . Now let  $\Gamma$  satisfy  $(O_2)$ . Choose a sequence of positive numbers  $\{d_n\}$  increasing to  $\infty$ . Conjugate  $\Gamma$  by a transformation  $h_n$  in  $\text{Möb}(\Delta)$  so that a hyperbolic element  $\gamma_n$  of  $\Gamma_n = h_n \Gamma h_n^{-1}$  has  $A_0$  as its axis and so that the collar  $U_0(d_n)$  of width  $d_n$  about  $A_0$  exists. Let  $G_n = \text{Stab}(\Gamma_n, A_0)$ . By the definition of collar:

$$\begin{aligned} \gamma(U(d_n)) &= U_0(d_n) \quad \text{for } \gamma \in G_n \quad \text{and} \\ \gamma(U_0(d_n)) \cap U_0(d_n) &= \emptyset \quad \text{for } \gamma \in \Gamma_n - G_n. \end{aligned} \quad (4.5)$$

Define functions  $\mu_{\alpha,n}$  ( $n = 1, 2, \dots$ ) in  $\hat{\mathbb{C}}$ :

$$\mu_{\alpha,n}(w) = \begin{cases} \mu_\alpha(z) \gamma'(z) / \overline{\gamma'(z)} & \text{if } w = \gamma(z) \text{ for some } z \in U_0(d_n) \\ & \text{and for some } \gamma \in \Gamma_n. \\ 0 & \text{otherwise} \end{cases} \quad (4.6)$$

Since  $G_n \subset \text{Möb}(A_0)$ , we can see by (4.3) and (4.5) that  $\mu_{\alpha,n}$  is well defined. The support of  $\mu_{\alpha,n}$  is in the closure of  $\bigcup_{\gamma \in \Gamma_n} \gamma(U_0(d_n))$  and in particular  $\mu_{\alpha,n} = 0$  in  $\Delta^*$ . By definition  $\mu_{\alpha,n}$  is a Beltrami coefficient for  $\Gamma_n$ , that is,

$$\text{ess. sup } |\mu_{\alpha,n}| = |\delta(\alpha) - 1| < 1, \quad (4.7)$$

and

$$\mu_{\alpha,n}(z) = \mu_{\alpha,n}(\gamma(z)) \overline{\gamma'(z)} / \gamma'(z) \quad \text{for all } \gamma \in \Gamma_n. \quad (4.8)$$

4.3. Let  $W_{\alpha,n}$  be the solution of the Beltrami equation

$$(W_{\alpha,n})_{\bar{z}} = \mu_{\alpha,n}(W_{\alpha,n})_z$$

which satisfies the normalization condition:  $W_{\alpha,n}(z) = z + O(|z|^{-1})$  as  $z \rightarrow \infty$ . Let  $K(\alpha) = (1 + |\delta(\alpha) - 1|) / (1 - |\delta(\alpha) - 1|)$ . As a consequence of (4.6), (4.7) and (4.8)  $W_{\alpha,n}$  is  $K(\alpha)$ -quasiconformal automorphism of  $\hat{\mathbb{C}}$ , conformal in  $\Delta^*$  and compatible with  $\Gamma_n$ . From the normalization condition  $\{W_{\alpha,n}\}_{n=1}^\infty$  forms a normal family ([6, Chap. II]). On the other hand the support of  $\mu_\alpha - \mu_{\alpha,n}$  is included in  $\Delta - U_0(d_n)$ . Since the Lebesgue measure of  $\Delta - U_0(d_n)$  decreases to 0 as  $n \rightarrow \infty$ , a subsequence of  $\{\mu_{\alpha,n}\}$  converges a.e. to  $\mu_\alpha$  ([10, Chap. 4, Proposition 17]). by replacing  $\{\mu_{\alpha,n}\}$  and  $\{W_{\alpha,n}\}$  with suitable subsequences, and denoting them again by  $\{\mu_{\alpha,n}\}$  and  $\{W_{\alpha,n}\}$ , the following situation arises:

(i)  $W_{\alpha,n}$  converges uniformly to a  $K(\alpha)$ -quasiconformal automorphism  $W_{\alpha,0}$  of  $\hat{\mathbb{C}}$ .

(ii)  $\mu_{\alpha,n}$  converges a.e. to  $\mu_\alpha$ .

Then  $W_{\alpha,n}$  is a good approximation to  $W_{\alpha,0}$  in the sense given in [6, Chap. IV, 5.4]. Therefore, by the normalization condition,  $W_{\alpha,0} = W_\alpha$ . Let  $\psi_{\alpha,n}(z) = [W_{\alpha,n}](z)$  for  $z \in \Delta^*$ . Then  $\psi_{\alpha,n} \in T(\Gamma_n)$  and  $\psi_{\alpha,n}(z)$  converges to  $[W_\alpha](z) = 4\alpha(z^2 - 1)^{-2}$  locally uniformly in  $\Delta^*$ . In particular it follows that  $\|\psi_{\alpha,n}\| \rightarrow 4|\alpha|$ .

Recall that  $\Gamma_n = h_n \Gamma h_n^{-1}$  for some Möbius transformation  $h_n$ . Thus,  $\phi_{\alpha,n}(z) = \psi_{\alpha,n}(h_n(z)) h_n'(z)^2$  belongs to  $T(\Gamma)$  and  $\|\phi_{\alpha,n}\| = \|\psi_{\alpha,n}\| \rightarrow 4|\alpha|$ . Choose a positive number  $\varepsilon$  to be sufficiently small so that  $\alpha = -3/2 + \varepsilon/4 \in V$ . Then for sufficiently large  $n$  we have that  $6 - 2\varepsilon < \|\phi_{\alpha,n}\|$ . Since  $\varepsilon$  can be arbitrarily small and  $o(\Gamma) \leq 6$  always holds, we can now conclude that  $o(\Gamma) = 6$ , whenever  $\Gamma$  satisfies the condition  $(O_2)$ .

4.4. The proof for the case where  $\Gamma$  satisfies the condition  $(O_1)$  proceeds in a way similar to the preceding case. Now let  $\Gamma$  satisfy  $(O_1)$ . For a choice of a sequence of positive numbers  $\{\delta_n\}$  decreasing to 0, let  $D_n = \{z \in \Delta : |z| < 1 - \delta_n\}$ . Then we can conjugate  $\Gamma$  by a transformation  $h_n$  in  $\text{Möb}(\Delta)$  in such a way that



$\Gamma_n = h_n \Gamma h_n^{-1}$  has the following property:

$$\gamma(D_n) \cap D_n = \emptyset \quad \text{for all } \gamma \in \Gamma_n - \{1\}.$$

For an  $\alpha \in V$ , define functions  $\mu_{\alpha,n}$  ( $n = 1, 2, \dots$ ) in  $\hat{\mathbb{C}}$ :

$$\mu_{\alpha,n}(w) = \begin{cases} \mu_{\alpha}(z)\gamma'(z)/\overline{\gamma'(z)} & \text{if } w = \gamma(z) \text{ for some } z \in D_n \\ & \text{and for some } \gamma \in \Gamma_n, \\ 0 & \text{otherwise.} \end{cases}$$

Then each  $\mu_{\alpha,n}$  is a Bertrami coefficient for  $\Gamma_n$  with  $\text{ess. sup } |\mu_{\alpha,n}| = |\delta(\alpha) - 1| < 1$ . Note that the support of  $\mu_{\alpha} - \mu_{\alpha,n}$  is included in  $\Delta - D_n$ , whose Lebesgue measure decreases to 0. With the normalized solution  $W_{\alpha,n}$  of the Beltrami equation  $(W_{\alpha,n})_{\bar{z}} = \mu_{\alpha,n}(W_{\alpha,n})_z$ , now we are in the same situation as in 4.3. Thus by the same argument there we can conclude in this case also that  $o(\Gamma) = 6$ . Now we complete the proof of Theorem 1.1.

## 5. Examples

We list some examples of fuchsian groups concerning the conditions  $(O_1)$  and  $(O_2)$ .

(a) Any fuchsian group  $\Gamma$  of the second kind satisfies the condition  $(O_1)$ . Thus for such a group we have  $o(\Gamma) = 6$ . Theorem 1.1 includes the former result by Sekigawa and Yamamoto [12], [13].

(b) For each integer  $n$ , define two circles  $A_n = \{z: |z - 2n + 1/2| = 1/2\}$  and  $B_n = \{z: |z - 2n - 1/2| = 1/2\}$ . Then  $\gamma_n(z) = ((1 + 4n)z - 8n^2)/(2z + 1 - 4n)$  maps the exterior of  $A_n$  onto the interior of  $B_n$ . The collection  $\{\gamma_n\}$  generates a subgroup  $\Gamma$  of the modular group  $\text{PSL}(2, \mathbb{Z})$ . The region in the upper half plane which is exterior to every  $A_n$  and  $B_n$  is a fundamental region of  $\Gamma$  and it contains  $\{z: \text{Im } z > 1\}$ . Since we can find there a hyperbolic disc of an arbitrary radius,  $\Gamma$  is a fuchsian group of the first kind satisfying the condition  $(O_1)$ .

(c) Let  $R$  be the complex plane with infinitely many punctures  $\mathbb{C} - \{2^{-n^2}, -2^{-n^2}\}_{n=1}^{\infty}$  endowed with the hyperbolic metric  $\rho_R(z)|dz|$ . Then the annulus  $A_n = \{z: 2^{-(n+1)^2} < |z| < 2^{-n^2}\}$  is included in  $R$ . Let  $g_n$  be a simple closed curve in  $A_n$  which separates the two boundary components of  $A_n$ . There exists a unique simple closed geodesic  $g'_n$  freely homotopic to  $g_n$  in  $R$ . Denote its length by  $l(g'_n)$ . Next let  $g''_n$  be the simple closed geodesic freely homotopic to  $g_n$  with respect to the hyperbolic metric of the annulus  $A_n$ . Denote by  $l(g''_n)$  and  $\tilde{l}(g''_n)$  its length with respect to the hyperbolic metric of  $R$  and that of  $A_n$ , respectively. Since  $A_n \subset R$ ,



it holds that  $\rho_R(z) < \rho_{A_n}(z)$  for  $z \in A_n$ , where  $\rho_{A_n}$  is the density of the hyperbolic metric of  $A_n$ . Therefore we have  $l(g'_n) \leq l(g''_n) < \bar{l}(g''_n)$ . Moreover we have that  $\bar{l}(g''_n) \rightarrow 0$  as  $n \rightarrow \infty$  since the module  $\log 2^{(n+1)^2}/2^{n^2}$  of  $A_n$  tends to  $\infty$ . Thus, for any  $\varepsilon > 0$ , there exists a simple closed geodesic on  $R$  whose length is smaller than  $\varepsilon$ . Then the collar lemma ([3]) ensures that a fuchsian group representing  $R$  satisfies the condition  $(O_2)$ .

(d) A cyclic group  $\Gamma$  generated by a hyperbolic transformation in  $\text{Möb}(\Delta)$  and its extension of index 2 satisfy both  $(O_1)$  and  $(O_2)$ .

(e) A finitely generated fuchsian group  $\Gamma$  of the first kind satisfies neither of the conditions, and hence the outradius  $o(\Gamma)$  is strictly less than 6. This result is first proved by Sekigawa [11]. On the other hand, there exist fuchsian groups  $\Gamma_n$  ( $n = 1, 2, \dots$ ) quasiconformally equivalent to  $\Gamma$ , for which  $o(\Gamma_n) \rightarrow 6$ . This is proved in a generalized form in [7].

## REFERENCES

- [1] BERS, L., *On the boundaries of Teichmüller spaces and on kleinian groups. I*, Ann. of Math., 91 (1970), 570–600.
- [2] FORD, L. R., *Automorphic Functions* (Second edition), Chelsea, New York, 1951.
- [3] HALPERN, N., *A proof of the collar lemma*, Bull. London Math. Soc., 13 (1981), 141–144.
- [4] JØRGENSEN, T., *On discrete groups of Möbius transformations*, Amer. J. Math., 98 (1976), 739–749.
- [5] KALME, C. I., *Remark on a paper by Lipman Bers*, Ann. of Math., 91 (1970), 601–606.
- [6] LEHTO, O. and VIRTANEN, K. I., *Quasiconformal mappings in the plane*, Springer, Berlin–Heidelberg–New York, 1973.
- [7] NAKANISHI, T., *A theorem on the outradii of Teichmüller spaces*, J. Math. Soc. Japan, 40 (1988), 1–8.
- [8] NEHARI, Z., *The Schwarzian derivative and schlicht functions*, Bull. Amer. Math. Soc. 55 (1949), 545–551.
- [9] POMMERENKE, Ch., *Univalent Functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.
- [10] ROYDEN, H. L., *Real Analysis* (Second edition), Macmillan, London, 1968.
- [11] SEKIGAWA, H., *The outradius of the Teichmüller space*, Tôhoku Math. J. 30 (1978), 607–612.
- [12] SEKIGAWA, H. and YAMAMOTO, H., *Outradii of Teichmüller spaces of finitely generated Fuchsian groups of the second kind*, J. Math. Kyoto Univ., 26 (1986), 23–30.
- [13] SEKIGAWA, H. and YAMAMOTO, H., *Outradii of Teichmüller spaces of the second kind*, Tôhoku Math. J., 38 (1986), 365–370.

Shizuoka University  
Department of Mathematics  
Shizuoka 422, Japan

and

National Defence Academy  
Department of Mathematics  
Yokosuka 239, Kanagawa  
Japan

Received March 9, 1987