

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 64 (1989)

**Artikel:** Compactness of isospectral conformal metrics on  $S^3$ .  
**Autor:** Chang, Sun-Yung A. / Yang, Paul C.  
**DOI:** <https://doi.org/10.5169/seals-48952>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 26.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Compactness of isospectral conformal metrics on $S^3$

SUN-YUNG A. CHANG\* and PAUL C. YANG\*

Two Riemannian metrics  $g, g'$  on a compact manifold are said to be isospectral if their associated Laplacian operator on functions have identical spectrum. It is a well known problem to study the extent to which the spectrum determines the metric. In dimension two, Osgood, Phillips and Sarnak [OPS] studied this question and proved that the set of isospectral metrics on a compact surface form a compact family in the  $\mathcal{C}^\infty$  topology. In that case there is available a criterion due to Wolpert [W] for compactness of the conformal structures in the Teichmüller space in terms of the determinant of the Laplacian. This reduces the problem to studying the isospectral conformal metrics on a fixed Riemann surface. It turns out that the determinant of the Laplacian played the key role for the compactness questions. In particular when the underlying surface is the two sphere, which is analytically the least transparent case, the compactness question reduces to an inequality of Onofri ([O], [OPS]) which is a sharp version of the Moser–Trudinger inequality on  $S^2$ .

We are interested in the situation in dimension 3. The well known solution of the Yamabe problem ([A], [S]) says that every conformal class of metrics on a compact Riemannian manifold contains a metric of constant scalar curvature. When  $(M^3, g_0)$  has constant negative scalar curvature, an isospectral set of metrics  $g = u^4 g_0$  conformal to  $g_0$  is compact in the  $\mathcal{C}^\infty$  topology [BPY]. This result was proved directly using the heat invariants of the metric. The first step was to find a pointwise bound  $0 < c_1 \leq u \leq c_2$  and a bound  $\|u\|_{2,2} \leq c_3$  where  $c_i$  depend only on the heat invariants of  $g$ . The higher order derivative bounds required for  $\mathcal{C}^\infty$  compactness is a consequence of this bound for  $u$  and the calculation for the coefficients for the terms involving the highest order derivatives of  $u$  in the asymptotic  $a_k$  of the heat kernel for  $g$  due to Gilkey ([G]).

In this paper we study the situation when  $M$  is the standard three sphere  $(S^3, g_0)$ . As in the case of the two sphere, the conformal group  $G$  complicates the analysis.

**DEFINITION.** For a positive function  $u$  on  $S^3$ , and  $\varphi$  a conformal transforma-

---

\* Research supported by NSF grants DMS-88-16321 and DMS-87-02871.

tion, define

$$u_\varphi = (u \circ \varphi) \cdot |d\varphi|^{1/2}$$

where  $|d\varphi|$  is the linear stretch factor of  $d\varphi$  measured with respect to the standard metric  $g_0$ . Thus  $u_\varphi^4 g_0 = \varphi^*(u^4 g_0)$ . We set

$$[u] = \{u_\varphi \mid \varphi \in G, \text{ the conformal group for } S^3\}.$$

The noncompactness of  $G$  shows that the class  $[u]$  is noncompact in  $H_1(S^3)$ , although the metrics associated to  $v \in [u]: \{g = v^4 g_0; v \in [u]\}$  are all isometric. We have the following result.

**THEOREM 1.** *For  $(S^3, g_0)$ , if  $\{g_i = u_i^4 g_0\}$  is a sequence of isospectral metrics, then there exists a subsequence  $g_i$  and conformal transformations  $\varphi_i$  such that  $\varphi_i^* g_i$  converges in the  $\mathcal{C}^\infty$  topology to a metric  $g$  which is also isospectral to  $\{g_i\}$*

On  $S^3 = \{x \in \mathbb{R}^4 \mid |x| = 1\}$  we have the standard metric  $g_0 = \sum_{i=1}^4 dx_i^2$ , with associated Laplacian  $\Delta$ , scalar curvature  $R_0 = 6$ , volume form  $dv_0$ . For a conformal metric  $g = u^4 g_0$  we have

$$dv = u^6 dv_0,$$

and its scalar curvature  $R$  is determined by the equation

$$8\Delta u + Ru^5 = R_0 u.$$

The trace of the heat kernel  $\exp(-t\Delta_u)$ , where  $\Delta_u$  denotes the Laplacian associated to the metric  $g = u^4 g_0$ , has the well known expansion

$$\text{Trace } \exp(-t\Delta_u) \sim (4\pi t)^{-n/2} (a_0 + a_1 t + a_2 t^2 + \cdots)$$

as  $t \rightarrow 0$ , where

$$a_0 = \int dv = \int u^6 dv_0.$$

$$a_1 = \int R dv = \int (8|\nabla u|^2 + 6u^2) dv_0.$$

$$a_2 = \frac{1}{360} \int (3R^2 + 6|\rho|^2) dv = \frac{1}{360} \int (3R^2 + 6|\rho|^2 u^6) dv_0,$$

$|\rho|$  denoting the norm of the Ricci tensor of  $g$ , measured in the metric  $g$ . As the heat invariants  $a_k$  are determined by the spectrum of the Laplacian associated to

$g$ , we get immediately the following bounds for isospectral conformal metrics  $g_i = u_i^4 g_0$ :

$$\int u^6 dv_0 = a_0,$$

$$\int (8 |\nabla u|^2 + 6u^2) dv_0 = a_1.$$

$$\frac{1}{120} \int R^2 u^6 dv_0 = \int (8\Delta u - 6u)^2 u^{-4} dv_0 \leq a_2.$$

In fact we will prove the following preliminary version of Theorem 1:

**THEOREM 1'.** *On  $(S^3, g_0)$  if  $g = u^4 g_0$  is a metric satisfying*

$$a_0(g) = \alpha_0 \tag{1}$$

$$a_1(g) \leq \alpha_1 \tag{2}$$

$$\int R^2 u^6 dv_0 \leq \alpha_2 \tag{3}$$

$$\Lambda \leq \lambda_1(g) \tag{4}$$

*for positive constants  $\alpha_0, \alpha_1, \alpha_2$  and  $\Lambda$ , where  $\lambda_1(g)$  is the first positive eigenvalue of the Laplacian, then there exist constants  $c_1, c_2$  and  $c_3$  (depending only on  $\alpha_0, \alpha_1, \alpha_2$  and  $\lambda$ ), and a conformal transformation  $\varphi$  so that  $v = u_\varphi$  satisfies:*

$$0 < c_1 \leq v(x) \leq c_2, \tag{5}$$

$$\|v\|_{2,2} \leq c_3, \text{ where } \|v\|_{2,2}^2 = \int v^2 + |\nabla v|^2 + |\nabla \nabla v|^2. \tag{6}$$

Then Theorem 1 will be a direct consequence of Theorem 1' and the following proposition proved in [BPY], which provides the required bounds for all the higher order derivatives of  $u$ .

**PROPOSITION.** *For a compact manifold  $(M^3, g_0)$  the set of conformal metrics  $g = u^4 g_0$  with  $u$  satisfying (5) (6) and the conditions*

$$a_k(g) \leq a_k \text{ for } k = 3, 4, 5, \dots$$

*form a compact set in the  $\mathcal{C}^\infty$  topology.*



We break up the proof of Theorem 1' into four lemmas.

**LEMMA 1.** *Assume  $u$  is a positive smooth function on  $S^n$ , for each  $\varepsilon > 0$  sufficiently small, we can find*

*$v \in [u]$  satisfying with  $n^* = (n+2)/(n-2)$*

$$\int_{S^n} v^{n^*+1+\varepsilon} x_j dv_0 = 0$$

*for  $j = 1, \dots, n+1$ , where  $x_j$  are the coordinate functions in  $\mathbb{R}^{n+1}$ .*

*Proof.* Consider for each  $\varphi \in G$ , the function  $(u_\varphi)^{n^*+1+\varepsilon}$  as a mass distribution on  $S^n$  and we define its center of mass to be

$$\text{C.M.}(\varphi) = \int (u_\varphi)^{n^*+1+\varepsilon} x dv_0 / \int (u_\varphi)^{n^*+1+\varepsilon} dv_0$$

It is clear that C.M.  $(\varphi)$  is continuous in  $\varphi$ . Using the stereographic projection  $\pi: S^n \setminus \{Q\} \rightarrow \{y \in \mathbb{R}^{n+1} \mid \langle y, Q \rangle = 0\}$  as coordinates, we define the conformal transformation  $\varphi_{Q,t}$  via its action on the  $y$  coordinates:

$$\varphi_{Q,t}(y) = ty.$$

The collection of conformal maps  $\{\varphi_{Q,t} \mid Q \in S^n, t \geq 1\}$  is naturally identified with  $B^{n+1} = \{x \in \mathbb{R}^{n+1} \mid |x| \leq 1\}$  by  $\varphi_{Q,t} \rightarrow (t-1)t^{-1}Q$ . Restricting the C.M. map to C.M.:  $\{\varphi_{Q,t} \mid Q \in S^n, t \geq 1\} \approx B^{n+1} \rightarrow B^{n+1}$ , we find easily that it extends continuously to  $\partial B$  to the identity map on  $\partial B$ . This implies that there exists  $Q \in S^n$  and  $t \geq 1$  so that C.M.  $(\varphi_{Q,t}) = 0$ . This is the claim of the lemma.

**LEMMA 2.** *Assume  $u$  is a positive function on  $S^3$  which satisfies the hypothesis of Theorem 1', there exists some  $\varepsilon_0 > 0$  and a constant  $c$  depending only on the data  $(\alpha_0, \alpha_1, \alpha_2, \Lambda)$  and some  $v \in [u]$  with  $\int v^{6+\varepsilon_0} dv_0 \leq c$ .*

*Remarks.* The formulation of Lemma 2 is motivated by the proof of the Yamabe problem (c.f. [A], [S]) where a sequence  $u_k$  of solutions of the equation (for some constants  $\mu_k$  which converges to constant  $\mu$  and  $\varepsilon_k \rightarrow 0$ ) satisfying the equation

$$c_n \Delta u_k + \mu_k u_k^{n^*-\varepsilon_k} = R_0 u_k$$

was shown to converge weakly to a nonnegative limit function  $u$ ; then in order to show the weak limit is not identically zero, it was sufficient to find a similar bound for  $\int u^{6+\varepsilon_0} dv_0$  as above. In that case the limit function  $u$  satisfies the Yamabe equation, so that the maximum principle shows  $u > 0$ . In our situation, because there is no a priori limit for the scalar curvatures  $R_k$  in the equations

$$c_n \Delta u_k + R_k u_k^{n^*} = R_0 u_k.$$

we need to proceed differently to establish the following lemmas to complete our argument:

**LEMMA 3.** *Suppose  $u \geq 0$  satisfies the hypothesis (1), (2), (3) and (4) of Theorem 1 and  $\int u^{6+\varepsilon_0} dv_0 \leq c$ , then there exists a positive  $c_1$  (depending on  $\alpha_0, \alpha_1, \alpha_2, \Lambda$  and  $c$ ) so that  $0 < c_1 \leq u$  on  $S^3$ .*

**LEMMA 4.** *Suppose  $u$  is as in Lemma 3, then there exists a positive  $c_2$  (depending only on  $\alpha_1, \alpha_2, \Lambda$  and  $c$ ), so that  $u \leq c_2$  on  $S^3$ .*

*Proof of Lemma 2.* Multiplying the equation

$$8\Delta u + Ru^5 = 6u \text{ on } S^3 \quad (7)$$

by  $u^\beta$  ( $\beta$  to be chosen later) we have: (denote  $\int_{S^3} dv_0$  as  $\int dv_0$ )

$$8 \frac{4\beta}{(\beta+1)^2} \int |\nabla u^{(\beta+1)/2}|^2 dv_0 + 6 \int u^{\beta+1} dv_0 = \int Ru^4 u^{\beta+1} dv_0. \quad (8)$$

Let  $w = u^{(\beta+1)/2}$  then we have

$$8 \frac{4\beta}{(\beta+1)^2} \cdot \int |\nabla w|^2 dv_0 + 6 \int w^2 dv_0 = \int Ru^4 w^2 dv_0. \quad (9)$$

From the Sobolev inequality for  $w$ :

$$Q \left( \int w^6 dv_0 \right)^{1/3} \leq 8 \int |\nabla w|^2 dv_0 + 6 \int w^2 dv_0 \quad (10)$$

where  $Q = Q(S^3) = 6 \cdot (\int dv_0)^{2/3}$ . We have for  $\beta > 0$ , from (9) and (10):

$$Q \frac{4\beta}{(\beta+1)^2} \left( \int w^6 dv_0 \right)^{1/3} \leq \int Ru^4 w^2 dv_0 + 6 \left( \frac{4\beta}{(\beta+1)^2} - 1 \right) \int w^2 dv_0 \quad (11)$$

We proceed to estimate the term  $I = \int Ru^4 w^2 dv_0$ . Taking  $b$  large to be chosen later, on the region  $|R| \geq b$  we have

$$b^2 \int_{|R| \geq b} u^6 dv_0 \leq \int_{|R| \geq b} R^2 u^6 dv_0 \leq A_2 \leq \alpha_2, \text{ where } A_2 = O(a_2).$$

Thus

$$\begin{aligned} \int_{|R| \geq b} Ru^4 w^2 dv_0 &\leq \left( \int R^2 u^6 dv_0 \right)^{1/2} \cdot \left( \int_{|R| \geq b} u^6 dv_0 \right)^{1/6} \left( \int w^6 dv_0 \right)^{1/3} \\ &\leq A_2^{1/2} \left( \frac{A_2}{b^2} \right)^{1/6} \cdot \left( \int w^6 dv_0 \right)^{1/3}. \end{aligned} \quad (12)$$

For the remaining part of  $I$ , write  $\beta = 1 + 2\varepsilon$  and apply Lemma 1 to  $u$  with  $\varepsilon > 0$ , we can find some  $v = u_\varphi \in [u]$  so that  $\int v^6 v^\varepsilon x_j dv_0 = 0$  for  $j = 1, 2, 3, 4$ . We have  $v$  satisfies the equation

$$8\Delta v + (R \circ \varphi)v^5 = 6v \quad (7)'$$

and that the first eigenvalue  $\lambda_1(v^4 g_0) = \lambda_1(u^4 g_0) \geq \Lambda$ . We have as in (12)' for  $\tilde{w} = v^{(1+\beta)/2}$ ,

$$\int_{|R(v)| \geq b} R(v)v^4 \tilde{w}^2 dv_0 \leq \left( \frac{A_2}{b} \right)^{1/3} \left( \int \tilde{w}^6 dv_0 \right)^{1/3}. \quad (12)'$$

For  $dv = v^6 dv_0$  we have from the Raleigh–Ritz characterization for  $\lambda_1$ :

$$\int_{S^3} \psi^2 dv \leq \left( \int_{S^3} \psi dv \right)^2 + \frac{1}{\lambda_1} \int_{S^3} |\nabla_v \psi|^2 dv \quad (13)$$

where  $|\nabla_v \psi|^2 dv = |\nabla \psi|^2 v^2 dv_0$ .

Choose  $\psi = v^\varepsilon x_j$  in (13), then by our assumption on  $v$ , we have

$$\begin{aligned} \int v^6 v^{2\varepsilon} x_j^2 dv_0 &\leq (1/\Lambda) \int |\nabla v^\varepsilon x_j|^2 v^2 dv_0 \\ &\leq \frac{1}{\Lambda} \left[ 2\varepsilon^2 \int |\nabla v|^2 v^{2\varepsilon} x_j^2 dv_0 + 2 \int |\nabla x_j|^2 v^{2+2\varepsilon} dv_0 \right] \\ &\leq \frac{2\varepsilon^2}{\Lambda(1+\varepsilon)^2} \int |\nabla v^{1+\varepsilon}|^2 x_j^2 dv_0 + L_j \end{aligned} \quad (14)$$

where  $L_j = \int |\nabla x_j|^2 v^{2+2\varepsilon} dv_0 = O(\int v^{2+2\varepsilon} dv_0)$ .

Sum up (14) for  $j = 1, 2, 3, 4$ , we find

$$\int v^{6+2\varepsilon} dv_0 \leq \frac{2\varepsilon^2}{\Lambda(1+\varepsilon)^2} \int |\nabla v^{1+\varepsilon}|^2 dv_0 + L, \quad (14)'$$

with  $L = O(\int v^{2+2\varepsilon} dv_0)$ .

Since  $v^4 g$  is isometric to  $u^4 g_0$  estimates (1) (2) (3) and (9) (10) (11) (12)' hold for  $v$  in place of  $u$ . Applying (8) for  $\beta = 1 + 2\varepsilon$  and (14)' we get

$$\int v^6 v^{2\varepsilon} dv_0 \leq \frac{2\varepsilon^2}{\Lambda(1+\varepsilon)^2} \frac{(1+\varepsilon)^2}{8(1+2\varepsilon)} I + L. \quad (15)$$

Combining (15) with (12)' and recall that  $\tilde{w} = v^{1+\varepsilon}$ , we find

$$\begin{aligned} I &= \int R v^4 \tilde{w}^2 \leq \left(\frac{A_2}{b}\right)^{1/3} \left(\int \tilde{w}^6 dv_0\right)^{1/3} + b \int v^4 \tilde{w}^2 dv_0 \\ &\leq \left(\frac{A_2}{b}\right)^{1/3} \left(\int \tilde{w}^6 dv_0\right)^{1/3} + \frac{2b\varepsilon^2}{8 \cdot \Lambda} I + bL \end{aligned} \quad (16)$$

So that

$$\left(1 - \frac{2b\varepsilon^2}{8\Lambda}\right) I \leq \left(\frac{A_2}{b}\right)^{1/3} \left(\int \tilde{w}^6 dv_0\right)^{1/3} + bL. \quad (17)$$

Now choose  $b$  sufficiently large so that  $\left(\frac{A_2}{b}\right)^{1/3} < \frac{1}{2}Q$ , and then choose  $\varepsilon$  sufficiently small so that

$$\left(1 - \frac{2b\varepsilon^2}{8\Lambda}\right) > \frac{3}{4}, \quad \frac{1+2\varepsilon}{(1+\varepsilon)^2} > \frac{3}{4}.$$

For this choice of  $b$  and  $\varepsilon$ , we have from (11) (16) and (17) that

$$\frac{3}{4}Q \left(\int \tilde{w}^6\right)^{1/3} \leq I \leq \frac{2}{3}Q \left(\int \tilde{w}^6 dv_0\right)^{1/3} + \frac{4}{3}bL.$$

Recall  $\tilde{w} = v^{1+\varepsilon}$ , hence

$$\begin{aligned} \left(\int v^{6+6\varepsilon} dv_0\right)^{1/3} &= \left(\int w^6 dv_0\right)^{1/3} < 16bL = 16b \int v^{2+2\varepsilon} dv_0 \\ &< b \left(\int v^6 dv_0\right)^{(2+2\varepsilon)/6} \left(\int dv_0\right)^{(4-2\varepsilon)/6} = c < \infty. \end{aligned}$$

This proves lemma 2 with  $\varepsilon_0 = 6\varepsilon$ .

*Proof of Lemma 3.* Assume  $u$  satisfies the hypothesis of the Lemma, we will prove the following assertions:

- (a) Denoting  $E_\lambda = \{\xi \in S^3, u(\xi) \geq \lambda\}$ ,  $|E_\lambda| = \int_{E_\lambda} dv_0$  then there exists  $\lambda_0 > 0$  and  $l_0 > 0$  so that  $|E_{\lambda_0}| \geq l_0 > 0$ .
- (b) There exists  $c' < \infty$  with  $\int (\log u)^2 dv_0 \leq c'$ .
- (c) There exists  $c'' < \infty$  with  $-\log u \leq c''$  on  $S^3$ .

It follows from (c) that  $u \geq c_1 > 0$  for some fixed  $c_1$ .

To prove (a), we have

$$\int u^{6+\varepsilon} dv_0 \leq c, \text{ with } \varepsilon \leq \varepsilon_0,$$

and

$$a_0^2 = \left( \int u^6 dv_0 \right)^2 \leq \left( \int u^{6-\varepsilon} dv_0 \right) \left( \int u^{6+\varepsilon} dv_0 \right) \leq c \int u^{6-\varepsilon} dv_0.$$

Thus

$$\frac{a_0^2}{c} \leq \int u^{6-\varepsilon} dv_0.$$

On the other hand, for all  $\lambda > 0$ , we have

$$\int u^{6-\varepsilon} dv_0 = \int_{E_\lambda} u^{6-\varepsilon} dv_0 + \int_{E_\lambda^c} u^{6-\varepsilon} dv_0 \leq \left( \int_{E_\lambda} u^6 \right)^{(6-\varepsilon)/6} |E_\lambda|^{\varepsilon/6} + \lambda^{6-\varepsilon} |E_\lambda^c|.$$

So for  $\lambda$  sufficiently small, say  $\lambda^{6-\varepsilon} \text{Vol}(S^3) < \frac{1}{2} \frac{a_0^2}{c}$  we have

$$\frac{1}{2} \frac{a_0^2}{c} \leq a_0^{(6-\varepsilon)/6} |E_\lambda|^{\varepsilon/6}$$

thus

$$|E_\lambda| \geq (a_0/2c)^{6/\varepsilon} a_0 = l_0.$$

To prove (b), choose  $\lambda_0, l_0$  as in (a), and consider the Raleigh–Ritz characterization for  $\lambda_1(D)$ ,  $D$  being the set  $E_{\lambda_0}^c$ , to find

$$\int_D \left| \log \left( \frac{u}{\lambda} \right) \right|^2 dv_0 \leq \frac{1}{\lambda_1(D)} \int_D \left| \nabla \log \frac{u}{\lambda_0} \right|^2 dv_0. \quad (18)$$

Since  $|D| = \text{Volume}(S^3) - |E_\lambda| \leq |S^3| - l_0$ , the well known Faber–Krahn inequality ([C]), says that  $\lambda_1(D) \geq C(l_0) > 0$ . Thus we find

$$\begin{aligned}
 \int_{u \leq \lambda_0} \left( \log \frac{u}{\lambda_0} \right)^2 dv_0 &\leq \frac{1}{c(l_0)} \int \left| \frac{\nabla u}{u} \right|^2 dv_0 \\
 &\leq \frac{1}{c(l_0)} \int \frac{\Delta u}{u} dv_0 \\
 &\leq \frac{1}{c(l_0)} \frac{1}{8} \int (Ru^4 - 6) dv_0 \\
 &\leq \frac{1}{c(l_0)} \frac{1}{8} \left[ \left( \int R^2 u^6 dv_0 \right)^{1/2} \left( \int u^6 dv_0 \right)^{1/6} \left( \int 1 dv_0 \right)^{1/3} + 6 \int dv_0 \right] \\
 &\leq \frac{1}{c(l_0)8} \cdot [\alpha_0^{1/2} \alpha_2^{1/2} \text{Vol}(S^3) + 6 \text{Vol}(S^3)]. \tag{19}
 \end{aligned}$$

We have also

$$\int_{u \geq \lambda_0} \left( \log \frac{u}{\lambda_0} \right)^2 dv_0 \leq \int_{u \geq \lambda_0} \left( \frac{u}{\lambda_0} \right)^2 dv_0 \leq \frac{1}{\lambda_0^2} \int u^2 dv_0 \leq \frac{1}{\lambda_0^2} \alpha_0^{1/3}. \tag{20}$$

Combining (19) and (20) we have a bound for  $\int (\log u)^2$  as claimed.

To prove (c). We use the integral identity:

$$-\psi(\xi) + \bar{\psi} = \int (\Delta \psi)(Q) G(\xi, Q) dv_0(Q) \tag{21}$$

where  $G(\xi, Q)$  is the Green's function for  $\Delta$  on  $S^3$ , and  $\bar{\psi}$  is the average of  $\psi$  on  $S^3$ . We may add a suitable constant to  $G$  to make it positive. Apply this identity to  $\psi = \log u$ :

$$\begin{aligned}
 -\log u(\xi) + \overline{\log u} &= \int (u^{-1} \Delta u - u^{-2} |\nabla u|^2)(Q) G(\xi, Q) dv_0 \\
 &\leq \int (u^{-1} \Delta u)(Q) G(\xi, Q) dv_0 \\
 &\leq \frac{1}{8} \|Ru^4 - 6\|_p \|G(\xi, \cdot)\|_p
 \end{aligned}$$

for  $\frac{1}{p} + \frac{1}{p'} = 1$ . Choose  $p = \frac{3}{2} + \delta$ ,  $\delta = \frac{\varepsilon}{16 + 2\varepsilon}$ , we find  $p' < 3$  so that both  $\|Ru^4 - 6\|_p$  and  $\|G(\xi, \cdot)\|_{p'}$  are bounded. It now follows from (b) that  $-\log u$  is bounded from above.

*Proof of Lemma 4:* Applying equation (21) to the function  $u$  we have

$$\begin{aligned} -u(p) + \bar{u} &= \int \Delta u(Q) G(p, Q) dv_0(Q) \\ &= \frac{1}{8} \int (6u - Ru^5) G(p, Q) dv_0(Q), \end{aligned}$$

where  $G(p, Q) \sim \frac{1}{d(p, Q)} + \text{smooth function}$ . We recall the following estimate

([A] p. 37): For  $h(y) = \int_{\mathbb{R}^3} \frac{f(x)}{\|x - y\|} dx$  we have, when  $\frac{1}{r} = \frac{1}{q} - \frac{2}{3}$ ,  $r > 1$

$$\|h\|_r \leq c(q) \|f\|_q. \quad (22)$$

We will iterate this estimate with a sequence of suitably chosen  $r_j$ ,  $q_j$ . Start with  $q_0 = \frac{2r_0}{4 + r_0}$ ,  $r_0 = 6 + 6\varepsilon$  where  $6\varepsilon \leq \varepsilon_0$ , we have

$$\int (Ru^5)^{q_0} \leq \left( \int R^2 u^6 \right)^{q_0/2} \left( \int u^{r_0} \right)^{1 - (q_0/2)}$$

Thus by (22) we find a bound for  $\|u\|_{r_1}$  with

$$r_1 = \frac{6r_0}{12 - r_0} > r_0.$$

Continuing with

$$r_2 = \frac{6r_1}{12 - r_1}, q_1 = \frac{2r_1}{r + r_1}, \dots, r_k = \frac{6r_{k-1}}{12 - r_{k-1}}, q_k = \frac{2r_k}{4 + r_k}.$$

We find

$$r_{k+1} - r_k = \frac{r_k - 6}{12 - r_k} r_k \geq \varepsilon r_k > 0 \quad \text{if } 6 < r_k < 12.$$

Thus there will be a  $k_0$  with  $r_{k_0} > 12$  and  $r_0 < r_1 < \dots < r_{k_0-1} < 12 < r_{k_0}$  with

$$q_{k_0} = \frac{2r_{k_0}}{4 + r_{k_0}} > \frac{3}{2}.$$

So at the end of the iteration we find a bound for  $\|u\|_{r_{k_0}}$ ,  $\frac{3}{2} < q_{k_0} < 2$ . This implies  $u \in L^\infty$  from the Holder estimate:

$$\|u\|_\infty \leq \|u\|_1 + \|Ru^5\|_{q_{k_0}} \|G\|_{q'}, \quad \text{where} \quad \frac{1}{q'} + \frac{1}{q_{k_0}} = 1, \quad \text{with} \quad q' < 3.$$

This finishes the proof of lemma 4.

### End of the proof of Theorem 1':

From Lemmas 3 and 4 we have

$$0 < c_1 \leq u \leq c_2.$$

From

$$\alpha_2 \geq a_2 \geq \int R^2 u^6 = \int \left( 64 \frac{(\Delta u)^2}{u^4} - 96 \frac{\Delta u}{u^2} + \frac{36}{u^2} \right) dv_0$$

we conclude that

$$\int \frac{(\Delta u)^2}{u^4} \leq \text{constant}.$$

This together with the uniform upper bound for  $u$ , yields a bound for  $\int (\Delta u)^2$ . Thus  $\int (\Delta u)^2 + u^2 dv_0$  is bounded, hence we have a bound for  $\|u\|_{2,2}$ .

*Remark.* In general for  $S^n$  with  $n \geq 4$ , Theorem 1' continues to hold provided that we substitute condition (3) with the following

$$\int R^{(n/2)+\delta} u^{(2n/n-2)} dv_0 \leq \alpha_2 \quad \text{for some } \delta > 0. \quad (3')$$



## REFERENCES

- [A] AUBIN, J. P., *Nonlinear analysis on manifolds, Monge-Ampere equations*, Springer-Verlag.
- [BGM] BERGER, M., GAUDUCHON, P., and MAZET, E., *Le spectre d'une Variété Riemannienne*, Lecture notes in Mathematics, 194, Springer-Verlag.
- [BPY] BROOKS, R., PERRY, P., and YANG, P., *Isospectral sets of conformally equivalent metrics*, to appear.
- [C] CHAVEL, I., *Eigenvalues in Riemannian geometry*, Academic Press.
- [G] GILKEY, P., *Leading terms in the asymptotics of the heat equation*, preprint.
- [O] ONOFRI, E., *On the positivity of the effective action in a theory of random surfaces*, *Comm. Math. Physics* 86 (1982) 321–326.
- [OPS] OSGOOD, G., PHILLIPS, R., and SARNAK, P., *Compact isospectral sets of Riemann surfaces*, to appear in *J. of Functional Analysis*.
- [S] SCHOEN, R., *Conformal deformations of a Riemannian metric to constant scalar curvature*, *Jour. Diff. Geom.*, 20 (1984), 479–496.
- [T] TRUDINGER, N., *Remarks concerning the conformal deformation of Riemannian structure on compact manifolds*, *Ann. Scuola norm. Sup. Pisa*, (4) 3 (1968) 265–274.
- [W] WOLPERT, S., *Asymptotics of the spectrum and the Selberg zeta function on the space of Riemann surfaces*, preprint.

*Sun-Yung A. Chang*  
*Department of Mathematics*  
*University of California*  
*Los Angeles CA 90024*

*Paul C. Yang*  
*Department of Mathematics*  
*University of Southern California*  
*Los Angeles CA 90089*

Received December 7, 1987