

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 64 (1989)

Artikel: On the linearization of actions of linearly reductive groups.
Autor: Jurkiewicz, Jerzy
DOI: <https://doi.org/10.5169/seals-48960>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the linearization of actions of linearly reductive groups

JERZY JURKIEWICZ

Let G be a linearly reductive algebraic group over an algebraically closed field k . Assume G acts on k^n by a morphism $A: G \times k^n \rightarrow k^n$. Then A may be viewed as a polynomial in n variables with coefficients in $\mathcal{O}(G)^n$. We assume

- 1° k^n has a fixed point, say the origin O , under the action A ,
- 2° the action is of degree ≤ 2 with respect to k^n , i.e. the polynomial considered above is of degree ≤ 2 .

Our aim, roughly speaking, is to prove that under the above assumptions the action is linear in some coordinate system (see the Theorem below). The case of $G = k^*$ and $\text{char } k \neq 2, 3$, has been studied in [J] by the Author, who thanks H. Kraft for important suggestions concerning the present paper. A well-known conjecture states that every action of a linearly reductive group on an affine space is linearizable. For other results and references see e.g. [B-B], [K, P], [Ka, R], [Ko, R], [P] and [P, R]). Notice for example that by Lemma 3.2 of [ibid] the assumption 1° is satisfied for all commutative groups of order $\leq 2^2 \cdot 3^2 \cdot 5^2 - 1 = 899$.

Let $\text{End}(k^n)$ denote the set of morphisms $k^n \rightarrow k^n$. A map $f: G \rightarrow \text{End}(k^n)$ is called algebraic if the corresponding map $\bar{f}: G \times k^n \rightarrow k^n$ is a morphism. Then $f \in R[X_1, \dots, X_n]$, where $R = \mathcal{O}(G)^n$. The Reynolds operator, i.e. the canonical G -equivariant projection $\mathcal{O}(G) \rightarrow k$ induces a G -equivariant projection (sending X_i to X_i)

$$\int := \int_G : R[X_1, \dots, X_n] \rightarrow k^n[X_1, \dots, X_n] = \text{End}(k^n),$$

so that $\int_G f$, the *mean value* of f is an endomorphism of k^n . Using the Reynolds operator corresponding to the group $G \times G$ we can consider $\int_{s \in G} \int_{t \in G} f(s, t)$, for an algebraic map $G \times G \rightarrow \text{End}(k^n)$ as well. We will use the following property of the mean value operator.

For $F \in \text{End}(k^n)$ and an algebraic map $f: G \rightarrow \text{End}(k^n)$ we have $\int_{s \in G} (f(s) \circ F) = (\int f(s)) \circ F$, and if F is linear, also $\int (F \circ f(s)) = F \circ (\int f(s))$. (1)

Assume $A : G \times k^n \rightarrow k^n$ is any group action with the point 0 fixed. Let $t \mapsto A(t)$ denote the corresponding homomorphism $G \rightarrow \text{Aut}(k^n) \subseteq \text{End } k^n$. We have $A(t) = L(t) + C(t) + \dots$, where $t \mapsto L(t)$ (resp $C(t)$) is the morphism from G to the space of linear maps (resp. quadratic maps) $k^n \rightarrow k^n$. From

$$A(st) = A(s)A(t) \quad (2)$$

it follows easily that $L(st) = L(s)L(t)$, i.e. L is a linear representation, and

$$L(s)C(t) + C(s)L(t) = C(st). \quad (3)$$

For $F : k^n \rightarrow k^n$ let $(\) \circ F$ and $F \circ (\)$ denote the respective right and left composition operator.

PROPOSITION. *Assume the condition 1° satisfied. Then there exists a unique quadratic map $Q : k^n \rightarrow k^n$ (independent of t) such that*

- a) $C(t) = L(t) \circ Q - Q \circ L(t)$,
- b) $\int L(t^{-1}) \circ Q \circ L(t) = 0$.

Proof. Apply $L(s^{-1}) \circ (\)$ to (3) and rewrite it in the form

$$C(t) = L(t) \circ L((st)^{-1}) \circ C(st) - L(s^{-1}) \circ C(s) \circ L(t). \quad (4)$$

Here $(s, t) \in G \times G$. Set

$$Q := \int L(s^{-1}) \circ C(s). \quad (5)$$

Applying the operator $\int_{s \in G}$ to (4) one gets the identity a). Apply $L(t^{-1}s^{-1}) \circ (\)$ to (3). The result may be written in the form

$$L((st)^{-1}) \circ C(st) - L(t^{-1}) \circ C(t) = L(t^{-1}) \circ (L(s^{-1}) \circ C(s)) \circ L(t)$$

Now apply $\int_{s \in G} \int_{t \in G}$ to both sides to get b). Finally the identity (5) follows from a) and b), hence the uniqueness. ■

REMARK. Suppose 1° and 2° satisfied. Let I stand for the identity on k^n . By (5), $I + Q = \int L(t^{-1}) \circ A(t)$. The expression under the integral may be viewed as the deviation of the action $A(t)$ from its linear part $L(t)$. So $I + Q$ is the mean

value of that deviation. This last morphism turns out to be a conjugating automorphism in case of action of degree two:

THEOREM. *Let G be linearly reductive and assume 1° and 2°. Let Q be the quadratic map defined in the proposition above. Suppose either of the following holds*

- a) $\text{char}(k) \neq 2$
- b) G is commutative (hence diagonalizable).

Then $A(t) = (I - Q) \circ L(t) \circ (I + Q)$ and $I - Q$, $I + Q$ are mutually invers automorphisms of k^n . In particular the action A is linearizable.

Recall, that G is diagonalizable if and only if it is a finite product of multiplicative groups k^* and a finite commutative group of order prime to $\text{char}(k)$ ([B], ch. III, §8).

Proof of the Theorem. Set $S := L(s)$, $T := L(t)$. Then $A(t) = T + TQ - QT$ and the identity (2) reduces to $(SQ - QS) \circ T = (SQ - QS) \circ (T + TQ - QT)$, for all $(s, t) \in G \times G$. Apply $(\) \circ T^{-1}$:

$$(SQ - QS) = (SQ - QS) \circ (I - Q + TQT^{-1}) \quad (6)$$

Then applying $S^{-1} \circ (\)$ we get $Q - S^{-1}QS = (Q - S^{-1}QS) \circ (I - Q + TQT^{-1})$. Further, $\int_{s \in G}$ gives $Q = Q \circ (I - Q + TQT^{-1})$, by Prop., b). Then $SQ = SQ \circ (I - Q + TQT^{-1})$ and subtracting (6) we have also $QS = QS \circ (I - Q + TQT^{-1})$. Apply $(\) \circ S^{-1}$ to get

$$Q = Q \circ (I - SQS^{-1} + (ST) \circ Q \circ (ST)^{-1}). \quad (7)$$

Now assume a). Let $Q' : k^n \times k^n \rightarrow k^n$, be the bilinear symmetric map such that $Q(x) = Q'(x, x)$. We have

$$Q = Q(I - SQS^{-1}) + Q \circ (ST) \circ Q \circ (ST)^{-1} + 2Q'(I - SQS^{-1}, (ST) \circ Q \circ (ST)^{-1}). \quad (8)$$

Now apply $\int_{t \in G}$. We have $\int Q \circ (ST) \circ Q \circ (ST)^{-1} = \int QTQT^{-1}$ and by (1) and Prop., b), $\int_{t \in G}$ of the last summand of (8) vanishes. So

$$Q = Q \circ (I - SQS^{-1}) + \int_{s \in G} QSQS^{-1} \quad (9)$$

Extract now the parts of degree 4 with respect to k^n to get

$$0 = QSQS^{-1} + \int QSQS^{-1}. \quad (10)$$

Applying $\frac{1}{2} \int_{s \in G}$ we get

$$\int_{s \in G} QSQS^{-1} = 0. \quad (11)$$

By (9) we have $Q = Q \circ (I - SQS^{-1})$. Since the part of degree 3 vanishes, $Q = Q \circ (I + SQS^{-1})$, too. Apply $(\) \circ S$ to get $QS = QS \circ (I + Q)$. Now we are ready to conjugate the linear action $(s, x) \mapsto S(x)$:

$(I - Q) \circ S \circ (I + Q) = S + SQ - QS \circ (I + Q) = S + SQ - QS = A(s)$, as required. For $t = 1$ one gets $(I - Q) \circ (I + Q) = I$. Replacing the action $A(t)$ by $A'(t) := (-I) \circ A(t) \circ (-I)$ one gets by an analogous argument that $(I + Q) \circ (I - Q) = I$. This completes the proof in case a).

So we assume $\text{char}(k) = 2$, and G diagonalizable. Then the proof alters as follows. Choose a bilinear map $\hat{Q}: k^n \times k^n \rightarrow k^n$ such that $\hat{Q}(x) = \hat{Q}(x, x)$ for all x . Then (8) holds with $2Q'$ replaced by the map $(x, y) \mapsto \hat{Q}(x, y) + \hat{Q}(y, x)$. To obtain (11) we must replace the Reynolds operator by a more precise tool, available for diagonalizable groups:

Denote by $X = X(G)$ the group of characters $G \rightarrow k^*$, with the additive notation. Let t^i stand for the value of $i \in X$ at $t \in G$. It follows easily from [B], Ch. III, §8, that

any algebraic map $f: G \rightarrow \text{End}(k^n)$ can be written in a unique way as

$$f(t) = \sum_{i \in X} t^i f_i, \text{ for some } f_i \in \text{End}(k^n). \quad (12)$$

Notice that $\int_{t \in G} f$ coincides with f_0 in this case. For $i \in X(G)$ let the morphisms $b_i: k^n \rightarrow k^n$ be defined by the expansion $SQS^{-1} = \sum s^i b_i$ introduced in (12), and set $q_{i,j} := \hat{Q}(b_i, b_j)$. Since $b_0 = 0$ we have $q_{0,j} = q_{i,0} = 0$. Now extract the part of degree 4 in (7) to get

$$0 = Q \circ (SQS^{-1} + (ST)Q(ST)^{-1}) = \sum_{i,j \in X} s^{i+j} (t^i - 1)(t^j - 1) q_{i,j}.$$

Apply $\int_{s \in G}$:

$$0 = \sum_{i \in X} (t^i + 1)(t^{-i} + 1) q_{i,-i} = \sum (t^i + t^{-i}) q_{i,-i}. \quad (13)$$

Let P be a subset of $X \setminus \{0\}$ such that for all $i \in X \setminus \{0\}$ one and only one of the characters i , $-i$ belongs to P . Hence X is the disjoint union of P , $-P := \{i \mid -i \in P\}$ and $\{0\}$ (indeed, $i + i \neq 0$ for $i \neq 0$ by the condition $(\text{char } k, G : G_{\text{conn}}) = 1$). Since $q_{0,0} = 0$ we have by (13) that

$$\sum_{i \in P} (t^i + t^{-i})(q_{i,-i} + q_{-i,i}) = 0.$$

By (12) this implies that

$$\sum_{i \in P} t^i (q_{i,-i} + q_{-i,i}) = 0.$$

For $t = 1$ we obtain

$$0 = \sum_{i \in P} (q_{i,-i} + q_{-i,i}) = \sum_{i \in X} q_{i,-i}.$$

On the other hand

$$\int_{s \in G} QSQS^{-1} = \int_{s \in G} Q \circ \left(\sum_X s^i b_i \right) = \int \left(\sum_{i,j} s^{i+j} q_{i,j} \right) = \sum_{i \in X} q_{i,-i}.$$

So we get again (11), and the rest of the proof follows as in the part a), obviously simplified. ■

REMARK. If $\text{char}(k) \neq 2, 3$ then exactly as in [J] one proves that the conjugating automorphism $I + Q$ of k^n is triangular in some basis of this vector space.

REFERENCES

- [B] BOREL, A. *Linear algebraic groups*, W. A. Benjamin, 1969.
- [B-B] BIAŁYNICKI-BIRULA, A. *Remarks on the action of an algebraic torus on k^n* , Bull. Ac. Pol. Math., 14 No 4 (1966), 177–181.
- [J] JURKIEWICZ, J. *Linearization of the multiplicative group action of degree two and Jordan algebras*, Bull. Ac. Pol. Math., 32 no 7–8 (1984), 387–391.
- [K, P] KRAFT, H. and POPOV, V. L. *Semi-simple group actions on the three-dimensional affine space are linear*, Comm. Math. Helv. 60 (1985), 466–479.
- [Ka, R] KAMBAYASHI, T. and RUSSELL, P. *On linearizing algebraic torus action*, Jour. of Pure Appl. Algebra, 23 (1984), 243–250.
- [Ko, R] KORAS, M. and RUSSELL, P. *G_m -Actions on A_3* , Canadian Mathematical Society, Conference Proceeding, Vol. 6 (1986), 269–277.

- [P] PANYUSHEV, D. J. *Semi-simple automorphisms group of four-dimensional affine space* (in Russian), Izv. Akad. Nauk SSSR, Ser. Math. 47, no 4 (1983).
- [P, R] PETRIE, T. and RANDALL, J. D. *Finite-order algebraic automorphisms of affine varieties*, Comm. Math. Helv. 61, (1986), 203–221.

Jerzy Jurkiewicz
Institute of Mathematics
Warsaw University, PKiN, 9p.
00-901 Warsaw, Poland

Received September 11, 1987/December 21, 1988

Buchanzeigen

JOHANN JAKOB BURCKHARDT, **Die Symmetrie der Kristalle**, von René-Just Haüy zur kristallographischen Schule in Zürich, mit einem Beitrag von Erhard Scholz, Birkhäuser Verlag, Basel–Boston–Berlin, 1988, 195 pages, SFr. 58.—.

Teil 1 Von Haüy zu Laue: A.-M. Legendre – R.-J. Haüy – Weiss, Frankenheim – Die 32 Kristallklassen. Bravis, Hessel, Neumann, Mieler, Gadolin – Sohncke und regelmässigen Punktsysteme – Fedorovos Entdeckung der 230 kristallographischen Raumsymmetriesysteme – Schoenflies' Theorie-Ausblick – M. Laue – Teil 2 Die kristallographische Schule in Zürich; Paul Niggeli – Kreislagerungen und reduzierte quadratische Formen – Nowacki – Lavas – Heesch – Ornamente – Schwarzweiss-Gruppen – Raumgruppen nach L. Weber und H. Heesch – Die 17 Ornamente – Graphen.

JIOMHUA WANG, **The Theory of Games**, Oxford Science Publications, 1988, 162 pages, £ 25.—.

1 Matrix Games: Saddle points – Mixed strategies – minimax theorem – Domination of strategies – Graphical solution of $2 \times n$ and $m \times 2$ matrix games – Matrix games and linear programming – 2 Continous Games: Continous, convex, separable games – 3 N -Person noncooperative games: Imputations – The core – Stable sets – The kernel – The nucleolus – shapley value.

V. I. ARNOLD, S. M. GUSEIN-ZADE and A. N. VARCHENKO, **Singularities of Differentiable Maps. Vol. II**, Birkhäuser Verlag, Boston–Basel–Berlin, 1988, 492 pages, SFr. 198.—.

Part I. The topological of isolated critical points of functions: Theory of Picard-Lefschetz – Non-singular level – Bifurcations sets, monodromy group – Interaction matrices – Part II. Oscillatory integrals: Asymptotics and Newton polyhedra – The singular index – Part III Integrals of holomorphic forms over vanishing cycles: Complex oscillatory integrals – Hodge filtrations and the spectrum of a critical point – The mixed Hodge structure – The period map and the intersection form.

RICHARD BEALS and PETER GREINER, **Calculus on Heisenberg Manifolds**, Princeton University Press, 1988, 194 pages \$ 40.—.

Preface – Introduction – The Model Operators – Inverting the Model Operator – Pseudodifferential operators on Heisenberg Manifolds – Invariance Theorem – \mathcal{V} -operators on Compact Manifolds, Hilbert Space Theory – Applications to the $\bar{\partial}_b$ -complex – Bibliography, Index of Terminology, List of Notation.

MOHAN S. PUTCHA, **Linear Algebraic Monoids**, Cambridge University Press, 1988, 171 pages, \$ 24.95/£ 15.—.

Linear Algebraic Semigroups – Reductive Groups and Regular Semigroups – Diagonal Monoids – Cross-Section Lattices – Renner's Decomposition – Tits Building – Renner's Extension Principle and Classification.

K. W. GRÜNBERG and J. E. ROSEBLAD, **The Collected Works of Philip Hall**, Clarendon Press, Oxford, 1988, 495 pages, \$ 80.—.

Obituary by J. E. ROSEBLAD (Bibliography).

L. LE BRUYN, M. VAN DEN BERGH and F. VAN OYSTAEYEN, **Graded Orders**, Birkhäuser Verlag, Boston-Basel, 208 pages, SFr. 42.—.