Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 65 (1990)

Artikel: Mean curvature functions of codimension-one foliations.

Autor: Oshikiri, Gen-Ichi

DOI: https://doi.org/10.5169/seals-49713

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Mean curvature functions of codimension-one foliations

GEN-ICHI OSHIKIRI*

0. Introduction

Let F be a transversely oriented codimension-one foliation of a closed connected manifold M. If we choose a Riemannian metric g on M, then we have a smooth function H(x) on M. Here H(x) is the mean curvature function at x of the leaf L_x of F through x with respect to the unit vector field N orthogonal to F and its direction coincides with the given transverse orientation. We call H(x) the mean curvature function of F with respect to g. Recently, Walczak [8], [9] studied the following problem for some special foliations:

[Q] Which smooth function on M can be written as a mean curvature function with respect to some Riemannian metric on M?

In this paper, we consider this problem from the view point given by Sullivan [6], and give an answer to this question. As a corollary, we can give another proof of the results in Walczak [8], [9]. These are done in Sections 1 and 2. In Section 3, we apply it for the case of Reeb foliations of S^3 .

The author is grateful to the hospitality of the Mathematical Institute, University of Köln, especially to Professor Dombrowski and the members of his seminar.

1. Preliminary and result

In this paper, we work in the C^{∞} -category. In what follows, we always assume that foliations are of codimension-one and transversely oriented, and that the ambient manifolds are closed, connected, oriented, and of dimension $n+1 \ge 3$, unless otherwise stated.

^{*}The author is partially supported by a grant from the Alexander von Humboldt Foundation.

Let g be a Riemannian metric on M. Then there is a unique vector field orthogonal to F whose direction coincides with the given transverse orientation. We denote this vector field by N. We given an orientation to F as follows: Let $\{E_1, \ldots, E_n\}$ be an oriented local orthonormal frame for TF. Then the orientation of M given by $\{N, E_1, \ldots, E_n\}$ coincides with the given one of M.

We denote the mean curvature of a leaf L at x with respect to N by H(x), that is,

$$H = \sum_{i=1}^{n} \langle \nabla_{E_i} E_i, N \rangle$$

where \langle , \rangle means g(,) and ∇ is the Riemannian connection of (M, g) and $\{E_i\}$ is a local orthonormal frame for TF with dim F = n. We also define an n-form X_F on M by

$$X_F(V_1,\ldots,V_n) = \det(\langle E_i,V_j\rangle)_{i,j=1,\ldots,n}$$
 for $V_i \in TM$,

where $\{E_1, \ldots, E_n\}$ is an oriented local orthonormal frame for TF. The restriction $X_F \mid L$ is the volume element of $(L, g \mid L)$ for $L \in F$. First we have the following.

PROPOSITION 1 (Rummler [5]). $dX_F = -HdV(M, g) = \operatorname{div}_g(N)dV(M, g)$, where dV(M, g) is the volume element of (M, g) and $\operatorname{div}_g(N)$ is the divergence of N with respect to g, i.e.,

$$\operatorname{div}_{g}(N) = \sum_{i=1}^{n} \langle \nabla_{E_{i}} N, E_{i} \rangle$$

Let f be a smooth function on M. We call f admissible if there is a Riemannian metric g on M so that -f coincides with the mean curvature function of F with respect to g, and set

$$C_{Ad} = \{f : admissible\}.$$

Note that if $H \equiv 0$, then F is a minimal foliation. If we can find such a g, then we call F taut. We also set

$$C_{\pm} = \{ f : f(x) > 0 > f(y) \text{ for some } x, y \in M \}.$$

Then, by Proposition 1, we have $C_{Ad} \subset C_{\pm} \cup \{0\}$. From this view point, we have:

THEOREM (Sullivan [7]). $0 \in C_{Ad}$, i.e., F is taut if and only if each compact leaf of F is cut out by a closed transversal.

THEOREM (Walczak [9]). F is taut if and only if $C_{Ad} = C_{\pm} \cup \{0\}$.

Now recall the set-up of Sullivan [6]. Let \mathbf{D}_p be the space of smooth p-forms on M and \mathbf{D}_p^* be the dual space of \mathbf{D}_p , i.e., the space of p-currents. Then we have:

THEOREM (Schwartz [5]). $(\mathbf{D}_p^*)^* = \mathbf{D}_p$.

Let $x \in M$ and $B = \{e_1, \dots, e_n\}$ be an oriented basis of $T_x F$. We define a Dirac current $\delta_{x,B}$ by

$$\delta_{x,B}(\phi) = \phi_x(e_1 \wedge \cdots \wedge e_n) \text{ for } \phi \in \mathbf{D}_n.$$

And set

 C_F = the closed convex cone in \mathbf{D}_n^* spanned by Dirac currents $\delta_{x,B}$ for $x \in M$.

PROPOSITION 2 (Sullivan [6]). C_F is a compact convex cone cone in \mathbf{D}_n^* . Here "compact" means that there is a continuous linear functional $L: \mathbf{D}_n^* \to \mathbb{R}$ so that the set $L^{-1}(1) \cap C_F$ is compact.

We shall call a compact set $L^{-1}(1) \cap C_F$ of the cone C_F the base of C_F and denote it by \mathbf{C} . Let $d: \mathbf{D}_p \to \mathbf{D}_{p+1}$ be the exterior differentiation and $\partial: \mathbf{D}_{p+1}^* \to \mathbf{D}_p^*$ be the dual of d, i.e., $\langle d\phi, c \rangle = \langle \phi, \partial c \rangle$ for $\phi \in \mathbf{D}_p$, $c \in \mathbf{D}_{p+1}^*$, and \langle , \rangle means the natural coupling $\mathbf{D}_p \times \mathbf{D}_p^* \to \mathbb{R}$. Set $B = \partial(\mathbf{D}_{n+1}^*)$ and $Z = \operatorname{Ker} \partial: \mathbf{D}_n^* \to \mathbf{D}_{n-1}^*$.

MAIN THEOREM. For $f \in C^{\infty}(M)$, the following three conditions are equivalent.

- (1) $f \in C_{Ad}$.
- (2) There are an n-form ω and an oriented volume form dV on M so that $d\omega = f \, dV$ and ω is positive on F. Here "positive" means that $\omega_x(e_1 \wedge \cdots \wedge e_n) > 0$ for all oriented basis $\{e_1, \ldots, e_n\}$ of T_xF and $x \in M$.
- (3) There is an oriented volume form dV on M so that
 - (i) $\int_{M} f dV = 0$, and
 - (ii) $\int_c f dV > 0$ for all $c \in \partial^{-1}(\mathbb{C} \cap B)$.

In case F is taut, Sullivan [7] showed that $\mathbb{C} \cap B = \emptyset$. And it is easy to show the existence of dV so that $\int_{\mathcal{M}} f \, dV = 0$ if $f \in C_{\pm}$. Thus we have another proof of the results of Walczak [8], [9].

2. Proof of main theorem

We follow the proofs given in Sullivan [6], [7]. To do this we need a Hahn-Banach theorem of the following type (cf. [1]):

THEOREM OF HAHN-BANACH. Let V be a Frechet space, W be a closed subspace of V, and C be a compact convex cone at the origin $0 \in V$. And let $\rho: W \to \mathbb{R}$ be a continuous linear functional of W with $\rho(v) > 0$ for $v \in C \cap W - \{0\}$. Then there is an extension $\eta: V \to \mathbb{R}$ of ρ so that $\eta(v) > 0$ for $v \in C - \{0\}$.

Proof of Main Theorem. First note that (2) implies (3). Because the condition (ii) of (3) reads as $\langle f dV, c \rangle = \langle d\omega, c \rangle = \langle \omega, \partial c \rangle$, and $\int_c f dV > 0$, for $\partial c \in \mathbb{C}$ and ω is positive on F.

- (1) implies (2). Now fix a Riemannian metric g on M with -f = H, and set dV = dV(M, g) and $\omega = X_F$ (see §1). Then, by Proposition 1, $dX_F = -H dV(M, g) = f dV$, that is, (1) implies (2).
- (3) implies (1). Here we use the Hahn-Banach Theorem quoted above under the following situation: $V = D_n^*$, W = B, $C = C_F$, and ϕ is given as follows: As $\int_M f \, dV = 0$, there is an *n*-form ω so that $d\omega = f \, dV$. By the theorem of Schwartz, we can regard ω as a continuous linear functional $k: \mathbf{D}_n^* \to \mathbb{R}$. If we restrict k on B, then this map $k \mid B$ is independent of the choice of ω . And, by condition (ii), $\phi = k \mid B$ satisfies the hypotheses. Thus, we have a continuous linear functional $L: \mathbf{D}_n^* \to \mathbb{R}$ which satisfies $L(C_F \{0\}) > 0$, and by the theorem of Schwartz, we have an *n*-form X on M which is positive on F. Further, the condition $d\omega = f \, dV$ implies $dX = f \, dV$, because $\omega \mid B = X \mid B$.

Now choose a Riemannian metric g as follows: On each leaf L, $X \mid L$ is the volume form of $(L, g \mid L)$, Ker X is orthogonal to F, and on Ker X the metric is determined by requiring dV(M, g) = dV, where dV is the n-form in the condition (3). By the relation f dV = dX = -H dV(M, g), we have $f \in C_{Ad}$. Q.E.D.

3. Examples and a concluding remark

In this section, we study, as an example, $C_{\rm Ad}$ for Reeb foliations of S^3 and close this paper with a conjecture.

EXAMPLE 1. A Reeb foliation F_R of S^3 . A Reeb component F_{RC} is a codimension-one foliation of $S^1 \times D^2$ with the boundary $\partial(S^1 \times D^2)$ as a unique compact leaf (cf. Lawson [2]). To get a Reeb foliation F_R of S^3 , decompose S^3 as $S^1 \times D^2 \cup D^2 \times S^1$ by identifying $(x, y) \in S^1 \times \partial D^2$ with $(x, y) \in \partial D^2 \times S^1$, and consider a Reeb component F_{RC} on each $S^1 \times D^2$. We give a transverse orientation to F_R and denote by R_+ (resp. R_-) the compact saturated set $S^1 \times D^2$ on that boundary the given transverse orientation is outward (resp. inward).

In this case, by our theorem, $C_{Ad} = \{ f \in C^{\infty}(S^3) : f(x) > 0 \text{ and } f(y) < 0 \text{ for some } x \in R_+ \text{ and } y \in R_- \}.$

Note that $0 \notin C_{Ad}$. This fact also shows that we cannot find any Riemannian metric of S^3 which makes each leaf of F_R a hypersurface of constant mean curvature (cf. Ex. 2).

EXAMPLE 2. A generalized Reeb foliation F_{GR} of S^3 . Here we also use the same notations as in Example 1. We decompose S^3 as $S^1 \times D^2 \cup T^2 \times [0, 1] \cup D^2 \times S^1$ with the canonical identification given in Example 1. A generalized Reeb foliation F_{GR} is given by $F_{RC} \cup \{T^2 \times (t)\}_{t \in [0, 1]} \cup F_{RC}$. Then we also have $C_{Ad} = \{f : f(x) > 0 \text{ and } f(y) < 0 \text{ for some } x \in R_+ \text{ and for some } y \in R_-\}$.

We choose a transverse orientation so that R_+ is the one of $S^1 \times D^2$ whose boundary is identified with $T^2 \times \{0\}$, and set $R_+(a) = R_+ \cup \{T^2 \times (t); 0 \le t \le a\}$. The above condition is equivalent to the following:

For each $f \in C_{Ad}$ there is a volume element dV of S^3 so that $\int_{S^3} f \, dV = 0$ and $\int_{R_+(a)} f \, dV > 0$ for all $a \in [0, 1]$.

Note that $0 \notin C_{Ad}$. But we can find a Riemannian metric of S^3 so that each leaf of F_{GR} is a hypersurface of constant mean curvature. To see this, we simply choose a smooth function h of [0, 1] satisfying $h(t) \equiv 1$ near 0, and $h(t) \equiv -1$ near 1, and lift this function onto $T^2 \times [0, 1]$ naturally, and set f on S^3 to be 1 on R_+ , -1 on R_- , and h on $T^2 \times [0, 1]$. It is clear that $f \in C_{Ad}$ (cf. Oshikiri [3]).

Condition (3) in our theorem is not easy to apply to arbitrary foliations. By considering the above examples, it seems to be plausible to conjecture that condition (3) is equivalent to the following:

- (3*) There is an oriented volume form dV on M so that
 - (i) $\int_M f \, dV = 0$, and
 - (ii) $\int_D f dV > 0$ for any foliated compact domain D with the transverse orientation of F being outward everywhere on ∂D .

REFERENCES

- [1] KELLY, J. L. and NAMIOKA, I., Linear Topological Spaces. D. Van Nostrand Company Inc., Canada, 1963.
- [2] LAWSON, H., Foliations. Bull. Amer. Math. Soc. 80 (1974), 369-418.
- [3] OSHIKIRI, G., On codimension-one foliations of constant mean curvature. Math. Zeit. (to appear).
- [4] RUMMLER, H., Quelques notions simples en geometrie riemannienne et leur applications aux feuilletages compacts. Comment. Math. Helv. 54 (1979), 224-239.
- [5] SCHWARTZ, L., Theorie des distributions. Nouvelle Edition, Hermann, Paris, 1966.
- [6] SULLIVAN, D., Cycles for the dynamical study of foliated manifolds and complex manifolds. Inventiones Math. 36 (1976), 225-255.
- [7] SULLIVAN, D., A homological characterization of foliations consisting of minimal surfaces. Comment. Math. Helv. 54 (1979), 218-223.
- [8] WALCZAK, P., Mean curvature functions for codimension-one foliations with all leaves compact. Czech. Math. J. 34 (1984), 146-155.
- [9] WALCZAK, P., Foliations which admits the most mean curvature functions. (preprint).

Department of Mathematics, College of General Education, Tohoku University, Kawauchi, Sendai 980 JAPAN

Received March 28, 1989