

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 65 (1990)

Artikel: The monodromy of a series of hypersurface singularities.
Autor: Siersma, Dirk
DOI: <https://doi.org/10.5169/seals-49720>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 06.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The monodromy of a series of hypersurface singularities

DIRK SIERSMA

Abstract. Let $\{f = 0\}$ be a hypersurface in \mathbb{C}^{n+1} with a 1-dimensional singular set Σ . We consider the series of hypersurfaces $\{f + \varepsilon x^N = 0\}$ where x is a generic linear form.

We derive a formula, which relates the characteristic polynomials of the monodromies of f and $f + \varepsilon x^N$. Other ingredients in this formula are the horizontal and the vertical monodromies of the transversal (isolated) singularities on each branch of the singular set. We use polar curves and the carousel method in the proof.

The formula is a generalization of the Iomdin formula for the Milnor numbers:

$$\mu(f + \varepsilon x^N) = \mu_n(f) - \mu_{n-1}(f) + Ne_0(\Sigma).$$

§1. Introduction

Let $f : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ be a germ at $\underline{0} \in \mathbb{C}^{n+1}$ of an analytic function. Let f have a 1-dimensional critical locus Σ . We consider for each $N \in \mathbb{N}$ the functions:

$$f_N = f + \varepsilon x^N \quad \varepsilon \in \mathbb{C}$$

where x is an admissible linear form, which means that $f^{-1}(0) \cap \{x = 0\}$ has an isolated singularity. We call the series $\{f_N\}$ a *linear series* or *Iomdin series* of hypersurface singularities.

An important invariant of a singularity is the Milnor fibration [Mi]: For $\varepsilon > 0$ small enough there exists $\eta > 0$ such that

$$f : B_\varepsilon \cap f^{-1}(S_n^1) \rightarrow S_n^1$$

is a locally trivial fibre bundle. $[B_\varepsilon$ is the closed ε -ball in \mathbb{C}^{n+1} , S_n^1 is the circle with radius η in \mathbb{C}]. A typical fibre $F = f^{-1}(\eta) \cap B_\varepsilon$ is called a *Milnor fibre* of f . Let

$$\mu_k(f) = \dim \tilde{H}_k(F).$$

If $\dim \Sigma = 1$ then it is known (cf [KM]) that $\mu_k(f) = 0$ if $k \neq n - 1, n$. If f has an isolated singularity then $\mu_k(f) = 0$ if $k \neq n$, and $\mu(f) = \mu_n(f)$ is called the Milnor

number of f . Iomdin studied the Milnor numbers for the linear series $f_N = f + \varepsilon x^N$ and related them to the non-isolated singularity f .

THEOREM [Iomdin] ([Io], cf. also [Lê-5]). *Let $f : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ have a 1-dimensional critical locus Σ and let x be an admissible linear form. There exists an N_0 such that for all $N > N_0$:*

1. f_N has an isolated singularity for all $\varepsilon \in \mathbb{C}$, $\varepsilon \neq 0$.
2. $\mu(f_N) = \mu_n(f) - \mu_{n-1}(f) + Ne_0(\Sigma)$ where $e_0(\Sigma)$ is the algebraic multiplicity of Σ at $\underline{0}$

In this paper we study the *monodromy* of such a linear series in connection with the monodromy of f .

A geometric monodromy is a diffeomorphism $h : F \rightarrow F$, which is a characteristic map for the Milnor fibration over the circle S_n^1 . It has the property that there exists an diffeotopy $H : F \times [0, 2\pi] \rightarrow B_\varepsilon \cap f^{-1}(S_n^1)$ such that:

$$f(H(x, t)) = \eta e^{it}$$

$$H(x, 0) = \text{identity}$$

$$H(x, 2\pi) = h(x).$$

The induced map:

$$\mathbb{T} : \tilde{H}_*(F) \rightarrow \tilde{H}_*(F)$$

is called the algebraic monodromy.

We now consider the case of a 1-dimensional critical locus in more detail. As we mentioned above: $\tilde{H}_k(F) = 0$ if $k \neq n-1, n$. So the algebraic monodromy can only act non trivially at the levels $n-1$ and n ;

$$\mathbb{T} | H_n(F) : H_n(F) \rightarrow H_n(F)$$

$$\mathbb{T} | H_{n-1}(F) : H_{n-1}(F) \rightarrow H_{n-1}(F)$$

For every irreducible branch Σ_i of Σ we have on $\Sigma_i - \{\underline{0}\}$ a local system of transversal singularities: Take at any $x \in \Sigma_i - \{\underline{0}\}$ the germ of a generic transversal section. This gives an isolated singularity, whose μ -constant class is well defined. We denote a typical Milnor fibre of this transversal singularity by F'_i . The only non-vanishing homology group is $\tilde{H}_{n-1}(F'_i)$. This is a special case of Deligne's

sheaf of vanishing cycles (cf. [De]). On the level of homology we have got in this way a local system, which has two different monodromies:

(a) the *vertical monodromy*

$$A_i : \tilde{H}_{n-1}(F'_i) \rightarrow \tilde{H}_{n-1}(F'_i)$$

which is the characteristic mapping of the local system over the punctured disc $\Sigma_i - \{0\}$

(b) the *horizontal monodromy*

$$T_i : \tilde{H}_{n-1}(F'_i) \rightarrow \tilde{H}_{n-1}(F'_i)$$

which is the Milnor fibration monodromy, when we restrict f to a transversal slice through $x \in \Sigma_i$

Note that the two monodromies A_i and T_i commute, since they are defined on

$$(\Sigma_i - \{0\}) \times S^n$$

which is homotopy equivalent to a torus.

The two monodromies play an important role in the relation between the monodromies of f and f_N .

MAIN THEOREM. *Let $f : \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ have a 1-dimensional critical locus $\Sigma = \Sigma_1 \cup \dots \cup \Sigma_n$ (irreducible components). Let x be an admissible linear form. Let $M[f](\lambda)$ be the alternating product of the characteristic polynomials of the monodromy \mathbb{T} of f in dimensions n and $n - 1$. Let $M[f + \varepsilon x^N](\lambda)$ be the characteristic polynomial of the monodromy of $f + \varepsilon x^N$ in dimension n . Then for all N sufficiently large $N \geq N_0$*

$$M[f + \varepsilon^N](\lambda) = M[f](\lambda) \prod_{i=1}^r \det(\lambda^{Nd_i} I - A_i T_i^{Nd_i}),$$

where:

$$A_i : \tilde{H}_{n-1}(F'_i) \rightarrow \tilde{H}_{n-1}(F'_i) \quad \text{vertical monodromy}$$

$$T_i : \tilde{H}_{n-1}(F'_i) \rightarrow \tilde{H}_{n-1}(F'_i) \quad \text{horizontal monodromy}$$

$$d_i = e_0(\Sigma_i^{red}), \quad \text{the multiplicity of } \Sigma_i^{red}.$$

Remarks

1. $m_i = d_i \mu_i$ where $m_i = e_0(\Sigma_i)$ and $\mu_i = \dim \tilde{H}_{n-1}(F'_i)$
2. $M[f](\lambda) = \det(\lambda I - \mathbb{T} | H_n(F))$. $[\det(\lambda I - \mathbb{T} | H_{n-1}(F))]^{-1}$
3. $M[f]$ is related to ζ_f , the zeta function of the monodromy which is defined as follows: $\zeta_f(\lambda) = \prod_{q>0} \{\det(I - \lambda \mathbb{T} | H^q(F))\}^{(-1)^{q+1}}$ cf. A'Campo [Ac]

COROLLARY. *The eigenvalues of the monodromy satisfy Steenbrink's spectrum conjecture [St] (2.2).*

The spectrum of a singularity, which is defined in [St] is a set of a real numbers $Sp(f)$. A spectrum number $\sigma \in Sp(f)$ is via

$$\lambda = e^{2\pi i \sigma}$$

related to the eigenvalues of the monodromy (including multiplicities). The multi-valuedness of

$$\sigma = \frac{1}{2\pi i} \log \lambda$$

is normalized with the Hodge filtration on the Milnor fibre. The semi simple part of the monodromy respects the Hodge filtration and the level of the $\log \lambda$.

Our main theorem can be written as

$$e^{2\pi i [\text{Spectrum conjecture}]}$$

As a general reference for hypersurface singularities we refer to the book of Arnol'd-Guzein Zade-Varchenko [AGV].

The proof of the main theorem arose from a discussion with Jozef Steenbrink in the train from Nancy to Maastricht.

Almost simultaneously M. Saito [Sa] announced a proof of Steenbrink's spectrum conjecture. His proof appears to be completely different from ours and uses the deep theory of Mixed Hodge Modules.

§2. The polar curve and the carousel

We start with a summary about the polar filtration and the carousel method. For details we refer to [Lê-4].


Let $f: \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ be a germ of an analytic function. Let $x: \mathbb{C}^{n+1} \rightarrow \mathbb{C}$ be an

admissible linear form. After a change of coordinates we can assume that x is the first coordinate in \mathbb{C}^{n+1} . We denote points in \mathbb{C}^{n+1} by $(x, y) \in \mathbb{C} \times \mathbb{C}^n = \mathbb{C}^{n+1}$.

Let $\Phi : \mathbb{C}^{n+1} \xrightarrow{(f, x)} \mathbb{C} \times \mathbb{C}$. We denote the singular locus of Φ by C . The polar curve of f (relative to x) is the closed set:

$$\Gamma = \overline{C \setminus f'(0)}.$$

The image of Γ in \mathbb{C}^2 is called the *Cerf diagram* Δ . Let $\Gamma = \Gamma_1 \cup \dots \cup \Gamma_s$ (decomposition in irreducible components) and $\Delta = \Delta_1 \cup \dots \cup \Delta_s$ with $F(\Gamma_i) = \Delta_i$. According to [Lê-1] the components of Δ are tangent to $z = 0$ in \mathbb{C}^2 . [We use (z, x) as coordinates in the target].



The curves Δ_i have Puiseux expansions of the form

$$x = a_i z^{r_i} + \dots \quad \text{with } a_i \neq 0 \quad r_i \in \mathbb{Q}.$$

The numbers r_1, \dots, r_s are called the *polar ratios* (of f at $\mathbb{0}$). Note that several branches of Δ can have the same polar ratio. Now let ρ_1, \dots, ρ_l be the different polar ratios and assume

$$\rho_1 > \rho_2 > \dots > \rho_l.$$

Let Δ_i^1 be the first approximation of Δ_i , defined by

$$x = a_i z^{r_i}.$$


Δ_i^1 has the same polar ratio as Δ_i . With the help of the polar ratios one defines a filtration by concentric discs in the x -plane

$$0 \subset D_1 \subset \dots \subset D_l$$

such that $\{\eta\} \times (D_j \setminus D_{j-1})$ contains the intersections

$$\Delta_i \cap \{z = \eta\} \text{ and } \Delta_i^1 \cap \{z = \eta\}$$

for all branches with polar ratio ρ_i .



Let $F' = \Phi^{-1}(\eta, 0)$ and $F_j = \Phi^{-1}(\{\eta\} \times D_j)$. The filtration:

$$F' \subset F_1 \subset \cdots \subset F_l$$

is called the *polar filtration* of the Milnor fibre F of f .

Lê constructed a geometric monodromy of f , which keeps the polar filtration invariant. As mentioned in [Lê-4] the geometric monodromy can be constructed as a carousel. Lê used this construction to show that:

- there exists a geometric monodromy without fixed points [Lê-2],
- the algebraic monodromy is quasi-unipotent [Lê-4].

The construction of the geometric monodromy involves a vectorfield v on $S_n^1 \times D$ which lifts the unit vectorfield on S_n^1 .

The x -component integrates in a first approximation to a rotation by an angle $2\pi\rho_j$; on $D_j \setminus D_{j-1}$ and to the identity on ∂D . Moreover the vectorfield is tangent to the intersection of the Cerf diagram Δ with $S_n^1 \times D$. Near the intersection points of the approximated branch $\Delta_i^1 : x = a_i z^{r_i}$ one uses a similar construction, which involves step by step also the next Puiseux pairs.

The geometric monodromy of F now is a lift by Φ of the vectorfield. For more details see the original papers. [Lê-2], [Lê-3] and [Lê-4].

§3. The geometric monodromy

Let now as in the introduction $f: \mathbf{C}^{n+1} \rightarrow \mathbf{C}$ have a 1-dimensional critical locus Σ and x be an admissible linear coordinate.

We want to compare the polar filtrations and the monodromy operators, associated to f and to $f_N = f + \varepsilon x^N$. We studied f already in §2, we now treat f_N . Let

$$\Phi_N = (f + \varepsilon x^N, x) : \mathbf{C}^{n+1} \rightarrow \mathbf{C}^2.$$

Φ_N has the same critical locus as Φ , namely $C = \{\partial f / \partial y = 0\}$. This is no surprise since the diffeomorphism

$$h : \mathbf{C}^2 \rightarrow \mathbf{C}^2$$

defined by $h(z, x) = (z + \varepsilon x^N, x)$ has the property

$$\Phi h = \Phi_N.$$

Next we consider the question when f_N has an isolated singularity. (cf. [Pe], p. 106). Consider the composition

$$\begin{aligned} f_N : \mathbf{C}^{n+1} &\xrightarrow{\Phi_N} \mathbf{C}^2 \xrightarrow{\pi} \mathbf{C} \\ &(w, x) \mapsto w. \end{aligned}$$

Note that π is submersive, so we can restrict ourselves to the critical set C of Φ_N . The condition we have to satisfy is:


$$\text{image}(d\Phi_N) \subset \text{Ker}(d\pi) = \{w = 0\}.$$

Outside the origin image $(d\phi_N)$ is the tangent space to $\Phi_N(C)$. Since $\Phi(C)$ is given by fractional power series $z = f = \alpha(x)$ we need for each branch

$$w = f + \varepsilon^N = \alpha(x) + \varepsilon x^N = 0.$$

This has a non-isolated solution at $x = 0$ only if $\alpha(x)$ is identical to $-\varepsilon x^N$. This is a strong condition! A necessary condition is that the polar ratio of the branch is $1/N$. Moreover if for a given n we have $\alpha(x) \equiv -\varepsilon x^N$ then the number of ε -solutions is finite.

Next assume N and ε are chosen such that f_N has an isolated singularity. Observe that f and f_N have almost the same polar curves Γ and Γ_N , since $\Gamma_N = \Gamma \cup \Sigma$. As a consequence the Cerf diagram Δ_N of f_N has one component more than the Cerf diagram Δ of f . More precisely h maps $\Delta \cup \{z = 0\}$ bijectively onto Δ_N . The extra component can be non reduced.



Remember that the polar ratios of f are $\rho_1 > \dots > \rho_l$. Let now $N_0 = 1/\rho_l$. If $N \geq N_0$ then f_N has polar ratios $\rho_1 > \dots > \rho_l \geq 1/N$. N.B. If $N < N_0$ the polar ratios become $\max\{\rho_i, 1/N\}$. We have the following propositions:

PROPOSITION. *Let $N \geq N_0$. There is a map $g : F \rightarrow F^N$ from a representative of the Milnor fibre F of f into the Milnor fibre F^N of f_N which induces a diffeomorphism on the polar filtrations up to level l :*


$$\begin{array}{ccccccc}
 F' \subset F_1 \subset F_2 \subset \dots \subset F_l = F & & & & & & \\
 \parallel & \downarrow h & \downarrow h & & \downarrow h & & \\
 F' \subset F_1^N \subset F_2^N \subset \dots \subset F_l^N \subset F^N & & & & & &
 \end{array}$$

So f and f_N have the same polar filtration, except f_N has one level more.

Proof. We take a representative of the map germ

$$(f + \varepsilon x^N, x, \varepsilon) : \mathbb{C}^{n+1} \times \mathbb{C} \rightarrow \mathbb{C} \times \mathbb{C} \times \mathbb{C}.$$

We use polydisc neighborhoods; let D be the corresponding disc in the x -coordinate plane. We take an admissible $\varepsilon > 0$ and choose next $\eta > 0$ such that the solutions of $x^N = \eta/\varepsilon$ are contained in D . We put the information for $\varepsilon = 0$ and $\varepsilon \neq 0$ in one picture.



We use the filtration $0 \subset D_1 \subset \dots \subset D_l$ (associated to f) to define the first l steps in the polar filtrations of both f and f_N . For the Milnor fibres we have:

$$F = \Phi^{-1}[\{z = \eta\} \cap x^{-1}(D_l)]$$

$$F^N = \Phi^{-1}[\{z + \epsilon x^N = \eta\} \cap x^{-1}(D)]$$

and for the polar filtrations

$$F_j = \Phi^{-1}[\{z = \eta\} \cap x^{-1}(D_j)] \quad j = 1, \dots, l$$

$$F_j^N = \Phi^{-1}[\{z + \epsilon x^N = \eta\} \cap x^{-1}(D_j)] \quad j = 1, \dots, l.$$

We next use ϵ as a parameter, and we lift the isotopy

$$\{z = \eta\} \cap x^{-1}(D_j) \rightarrow \{z + \epsilon x^N = \eta\} \cap x^{-1}(D_j)$$

with the help of the Thom isotopy-lemma to a diffeomorphism

$$h : F_l \rightarrow F_l^N$$

which induces diffeomorphisms on each level

$$h | F_j : F_j \rightarrow F_j^N.$$

Remark. We can make the construction and the arguments finer such that also the whole carrousel of F_l and F_l^N are diffeomorphic. Since the carrousel in connection with the Puiseux data determine a geometric monodromy it follows that F_l and F_l^N have the same carrousel monodromy.

Remark. From now on we use h to identify $F_l^N = F$.

Next we want to complete the monodromy over $D \setminus D_l$.

For f we make the x -component of the carrousel such that it becomes the identity on $D \setminus D_l$ except near ∂D_l , where we interpolate it with the rotation on $D_l \setminus D_{l-1}$. This has the advantage that over $D \setminus D_l$ the geometric monodromy preserves the x -coordinate.

Moreover this monodromy extends from $S'_n \times (D \setminus D_l)$ over $D_\eta \times (D \setminus D_l)$ and gives rise to a geometric monodromy T of Φ .

For f_N we have one extra level in the polar filtration. The Cerf-diagram has the extra branch

$$w = \varepsilon x^N,$$

which has exactly N intersection points with the Milnor fibre $\{f_N = \eta\}$.

These points are of the form $x_k = x_1 e^{2\pi i k/N}$ ($k = 1, \dots, N$), where x_1 is one intersection point. They all have the same absolute value and are contained in $D \setminus D_l$.

The x -component of the carrousel for f_N is a rotation by $2\pi/N$ on $D \setminus D_l$.

Let S be the diffeomorphism $F^N \setminus F \rightarrow F^N \setminus F$ which integrates a lift of the x -component of the carrousel vector field.

CLAIM. *The geometric local (relative) monodromy on $F \setminus F_l$ is just $T \cdot S$.*

Proof. Let S_t be the integral of the lifted x -component over the interval $[0, t]$, and T_t be similar for the z -component.

$$S_{2\pi} = S, \quad T_{2\pi} = T,$$

S_t preserves the levels $f = c$; T_t preserves the x -coordinate.


So

$$T_t S_t \{f + \varepsilon x^N\} = T_t \{f + \varepsilon x^N e^{it}\} = f e^{it} + \varepsilon x^N e^{it} = (f + \varepsilon x^N) e^{it}$$

which tells us that TS is the monodromy of f_N on $F^N \setminus F$.

§4. The algebraic monodromy

Next we study the algebraic monodromy.



Let x_1, \dots, x_N be the intersection points of $z + \epsilon x^N = \eta$ with $z = 0$. At these points the projection

$$g : \Phi \mid F^N \setminus F \rightarrow D \setminus D_l$$

fails to be a submersion. Around x_1, \dots, x_N we choose small discs A_1, \dots, A_N and points a_1, \dots, a_N on $\partial A_1, \dots, \partial A_N$.

By deformation retraction and excision we have

$$H_*(F^N, F) \cong \bigoplus_{k=1}^n H_*(g^{-1}(A_k), g^{-1}(A_k)).$$

Before doing more excision we remember that the projections

$$\Phi \mid \Sigma_i : \Sigma_i \rightarrow \{z = 0\}$$

are branched coverings of topological degree d_i .

So each a_k has d_i preimages under this map. Let $\{b_{i,1}, \dots, b_{i,Nd_i}\}$ be the preimages of $a_1, \dots, a_k, \dots, a_N$ numbered in such a way that

$$\Phi(b_{i,k}) = a_{k(\text{mod } N)} \text{ where } 1 \leq k(\text{mod } N) \leq N \text{ and } k(\text{mod } N) \equiv k \pmod{N}.$$

Next we choose $E_{i,k}$ as a Milnor ball and $F_{i,k} = g^{-1}(a_j) \cap E_{i,k}$ as Milnor fibre of the isolated singularity $g : (F^N \setminus F_i^N, b_{i,k}) \rightarrow (\mathbb{C}, a_{k(\text{mod } N)})$. Further deformation and excision gives us

$$H_q(F^N, F_i^N) = \bigoplus_{i=1}^r \bigoplus_{k=1}^{Nd_i} H_q(E_{i,k}, F_{i,k}) = \bigoplus_{i=1}^r \bigoplus_{k=1}^{Nd_i} \tilde{H}_{q-1}(F_{i,k}).$$

These local Milnor fibres $F_{i,k}$ are nothing other than the Milnor fibre of the transversal singularity on Σ_i , taken at the point $b_{i,k}$.

They are all isomorphic with a typical fibre F'_i . Later on this isomorphism will be important; for now:

$$H_q(F^N, F) \cong \bigoplus_{i=1}^r [\tilde{H}_{q-1}(F'_i)]^{Nd_i}.$$

Note also that only non vanishing dimension is $q = N$.

Let $T_{\text{rel}}: H_n(F^N, F) \rightarrow H_n(F^N, F)$ denote the induced action of the geometric monodromy on $H_n(F^N, F)$. We call it the *relative monodromy*.

The definitions, the excisions and the deformations from above, can be made compatible with the actions of S and T . We suppose we have done so. The construction of S was related to the cyclic permutation:

$$x_1 \rightarrow x_2 \rightarrow \cdots \rightarrow x_N \rightarrow x_1$$

and we can assume the same permutation on

$$a_1 \rightarrow a_2 \rightarrow \cdots \rightarrow a_N \rightarrow a_1.$$

In the same way S permutes the $g^{-1}(A_k)$. $k = 1, \dots, N$

We split $H_n(F^N, F)$ according to the branches $\Sigma_1, \dots, \Sigma_r$. From the above arguments, it follows that T_{rel} acts diagonal on this decomposition

$$H_n(F^N, F) = \bigoplus_{i=1}^r \left[\bigoplus_{k=1}^{Nd_i} \tilde{H}_{n-1}(F_{i,k}) \right].$$

So we can treat each branch separately. Since Σ_i is irreducible and the branched covering

$$\Sigma_i \rightarrow \{z = 0\}$$

has degree d_i the induced action of S is also a full cyclic permutation of $\{b_{i,1}, \dots, b_{i, Nd_i}\}$. We assume that we numbered our $b_{i,j}$ such that

$$b_{i,1} \rightarrow b_{i,2} \rightarrow \cdots \rightarrow b_{i, Nd_i} \rightarrow b_{i,1}.$$

So S induces permutations:

$$E_{i,1} \rightarrow E_{i,2} \rightarrow \cdots \rightarrow E_{i, Nd_i} \rightarrow E_{i,1}$$

$$F_{i,1} \rightarrow F_{i,2} \rightarrow \cdots \rightarrow F_{i, Nd_i} \rightarrow F_{i,1}.$$

Note also that T preserves the x -coordinate and can be assumed to be the

identity outside a neighborhood of the polar curve, more precisely on:

$$g^{-1}(A_k) \setminus \bigcup_{k'} \bigcup_{i=1, \dots, r} E_{i, k'}$$

with $k' \pmod{N} = k$. We denote the resulting actions on the homology as follows:

$$\begin{aligned} T_{i, k} &: \tilde{H}_{n-1}(F_{i, k}) \rightarrow \tilde{H}_{n-1}(F_{i, k}) \\ S_{i, k} &: \tilde{H}_{n-1}(F_{i, k}) \rightarrow \tilde{H}_{n-1}(F_{i, k+1}). \end{aligned}$$

The description of the monodromy in the block decomposition

$$\bigoplus_{k=1}^{Nd_i} \tilde{H}_{n-1}(F_{i, k})$$

is as follows:

$$\begin{bmatrix} 0 & 0 & 0 & 0 & S_{i, Nd_i} T_{i, Nd_i} \\ S_{i, 1} T_{i, 1} & 0 & 0 & \cdot & \cdot \\ 0 & S_{i, 2} T_{i, 2} & 0 & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & 0 & \cdot \\ 0 & \cdot & 0 & S.T. & 0 \end{bmatrix}.$$

LEMMA.

$$(S_{i, Nd_i} T_{i, Nd_i}) \cdot \dots \cdot (S_{i, 2} T_{i, 2}) \cdot (S_{i, 1} T_{i, 1}) = A_i(T_i)^{Nd_i}$$

where $A_i = \tilde{H}(F_{i, 1}) \rightarrow \tilde{H}(F_{i, 1})$ is the vertical monodromy of the transversal local system and $T_i = T_{i, 1} : \tilde{H} \rightarrow \tilde{H}(F_{i, 1})$ is the horizontal monodromy of a transversal section.

Proof. This is clear since Φ is a submersion over the torus

$$|x| = |a_j|, |z| = \eta.$$

Moreover:

$$A_i = A_{i, 1} = S_{i, Nd_i} \cdot S_{i, 2} \cdot S_{i, 1}.$$

LEMMA. *Let*

$$B = \begin{bmatrix} 0 & 0 & & 0 & B_m \\ B_1 & 0 & & & 0 \\ 0 & B_2 & 0 & & \\ 0 & 0 & \cdot & \cdot & \\ & & & \cdot & \cdot \\ 0 & & & B_{m-1} & 0 \end{bmatrix}$$

a $m \times m$ matrix consisting of blocks of size $v \times v$, then

$$\det(\lambda I - B) = \det(\lambda^m I - B_1 \cdot \dots \cdot B_m).$$

Proof. As shown to me by Rob Schrauwen, the proof is an elementary exercise in block matrix computation.

COROLLARY. *The characteristic polynomial of the monodromy on*

$$\bigoplus_{k=1}^{d_i N} \tilde{H}_{n-1}(F_{i,k})$$

is equal to:

$$\det[\lambda^{Nd_i} I - A_i(T_i)^{Nd_i}].$$

We have shown:

PROPOSITION. *For the relative monodromy we have:*

$$\det(\lambda I - \mathbb{T}_{\text{rel}}) = \prod_{i=1}^r \det[\lambda^{N_i} I - A_i(T_i)^{N_i}].$$

As a last step in the proof of the main theorem, we consider the action of the monodromy \mathbb{T} on the long exact sequence of the pair (F^N, F) :

$$\begin{array}{ccccccc} 0 & \rightarrow & H_n(F) & \rightarrow & H_n(F^N) & \rightarrow & H_N(F^N, F) \rightarrow H_{n-1}(F) \rightarrow 0 \\ & & \downarrow \mathbb{T} & & \downarrow \mathbb{T}_N & & \downarrow \mathbb{T}_{\text{rel}} & & \downarrow \mathbb{T} \\ 0 & \rightarrow & H_n(F) & \rightarrow & H_n(F^N) & \rightarrow & H_N(F^N, F) \rightarrow H_{n-1}(F) \rightarrow 0. \end{array}$$

It follows that:

$$\begin{aligned} M[f_N](\lambda) &= \det(\lambda I - \mathbb{T}_N) = M[f](\lambda) \cdot \det(\lambda I - \mathbb{T}_{\text{rel}}) \\ &= M[f](\lambda) \cdot \prod_{i=1}^r \det[\lambda^{N d_i} I - A_i(T_i)^{N d_i}] \end{aligned}$$

$$\left(\text{since } M[f](\lambda) \text{ is the quotient: } \frac{\det(\lambda - \mathbb{T} \mid H_n(F))}{\det(\lambda I - \mathbb{T} \mid H_{n-1}(F))} \right).$$

EXAMPLE. Consider the case that f is homogeneous of degree d . Notice that:

- $d_i = 1$ for all i .
 - the only polar ratio is $1/d$,
 - the horizontal and vertical algebraic monodromies are related by $A_i = T_i^{-d}$
- (cf. [St])

So if $N \geq d$ the formula of the main theorem reduces to

$$M[f + \varepsilon x^N](\lambda) = M[f](\lambda) \cdot \prod_{i=1}^r \det(\lambda^N I - T_i^{N-d}).$$

If $N = d$ it follows that

$$M[f + \varepsilon x^d](\lambda) = M[f](\lambda) \cdot \prod_{i=1}^r (\lambda^d - 1)^{\mu_i}.$$

Observe that $f + \varepsilon x^d$ is homogeneous of degree d . For almost all ε it has an isolated singularity and its monodromy depends only on the degree d :

$$M[f + \varepsilon x^d](\lambda) = M_d^{\text{reg}}(\lambda),$$

which is described in detail in [Mi], p. 71. We have now an expression for $M[f]$! After substitution we find: ($N \geq d$):

$$M[f + \varepsilon x^N](\lambda) = \frac{M_d^{\text{reg}}(\lambda)}{(\lambda^d - 1)^{\sum \mu_i}} \cdot \prod_{i=1}^r \det(\lambda^N I - T_i^{N-d})$$

which is similar to the formula in [Ste].

So it is possible to compute $M[f + \varepsilon x^N]$ if we know the degree d and the eigenvalues of the vertical algebraic monodromies. For the Milnor numbers it follows that:

$$\mu(f + \varepsilon x^N) = (d - 1)^n + \Sigma(N - d)\mu_i.$$

Remark. Linear series $f + \varepsilon x^N$ as studied by Yomdin, sometimes correspond to the series in Arnol'd's lists of singularities:

$$A_{N-1}: y^2 + \varepsilon x^N$$

$$D_{N+1}: xy^2x^N.$$

But in other cases one does not get the full series; examples are:

$$Y_{r,s}: x^2y^2 + x^{r+4} + y^{s+4}$$

$$\begin{cases} W_{1,2q-1}^\# : (y^2 - x^3)^2 + x^{4+q}y & q \geq 1 \\ W_{1,2q}^\# : (y^2 - x^3)^2 + x^{3+q}y & q \geq 1 \end{cases}$$

This especially occurs when the multiplicity of Σ is different from one.

Schrauwen [Sch] gave in the case of a plane curve a definition of a topological series of singularities and derived [in the case of transversal type A_1] a formula for the monodromy in such a series, which is similar to our main formula.

REFERENCES

- [Ac] N. A'CAMPO, *La fonction zêta d'une monodromie*, Comment Math. Helv. 50(2) (1975), 233–248.
- [AGV] V. I. ARNOL'D, S. M. GUSEIN ZADE and A. N. VARCHENKO, *Singularities of differentiable Maps*, Vols. I and II (Birkhauser 1985 and 1988).
- [D] P. DELGINE, *Le formalisme des cycles évanescents*. SGA VII². Exp XIII. Lectures Notes in Mathematics 340 (1973), 82–115.
- [Io] I. N. IOMDIN, *Complex surfaces with a 1-dimensional set of singularities*. Sibirskii Mat. Z. 15(5) (1974), 1061–1082.
- [KM] M. KATO and Y. MATSOMOTO, *On the connectivity of the Milnor fibre of a holomorphic function at a critical point*. Proc. Tokyo Manifolds Conference, (1973), 131–136.
- [Lê-1] D. T. LÊ, *Calcul des cycles évanouissants des hypersurfaces complexes*. Ann. Inst. Fourier. 23(4) (1973) 261–270.
- [Lê-2] D. T. LÊ, *La monodromie n'a pas de points fixes*. J. of Fac. Sc. Univ. Tokyo, Sec. 1A, 22 (1975), 409–427.
- [Lê-3] D. T. LÊ, *Some remarks on relative monodromy* Proc. of Nordic Summer School on Real and Complex singularities, Oslo (1976), 397–403.
- [Lê-4] D. T. LÊ, *The geometry of the monodromy theorem*. C. P. Ramanujam—A tribute; Studies in Mathematics, No. 8. Tata Institute of Fundamental Research, (1978), 157–173.

- [Lê-5] D. T. LÊ, *Ensembles analytiques complexes avec lieu singulier de dimension un* (d'après I. N. Yomdin). Séminaire sur les singularités. Publ. Math. Univ. Paris VII, (1980), 87–95.
- [Mi] J. MILNOR, *Singular points of complex surfaces*. Ann. Math. Studies (Princeton University Press Princeton NJ, 1968)
- [Pe] G. R. PELLIKAAN, *Hypersurface singularities and resolutions of Jacobi modules*. Thesis Rijksuniversiteit Utrecht, (1985).
- [Sch] R. SCHRAUWEN, *Topological series of isolated plane curve singularities*. To appear in L'Enseignement Mathématique.
- [St] J. H. M. STEENBRINK, *The spectrum of hypersurface singularities*. Preprint 8810 Catholic University Nijmegen April 1988. To appear in the Proceedings of the Luminy Conference on Hodge theory, 1987 (ed. F. Barlet)
- [Ste] J. STEVENS, *On the μ -constant stratum and the V -filtration; An example*. Math. Z. 201 (1989), 139–144.
- [Sa] J. SAITO: *Vanishing cycles and Mixed Hodge Modules*, preprint I.H.E.S. (August 1988).

*Mathematisch Instituut
der Rijksuniversiteit
Postbus 80.010
NL-3508TA Utrecht
Holland*

Received November 1, 1988