

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 65 (1990)

Artikel: Smooth algebras and vanishing of Hochschild homology.
Autor: Rodicio, Antonio G.
DOI: <https://doi.org/10.5169/seals-49738>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Smooth algebras and vanishing of Hochschild homology

ANTONIO G. RODICIO

For a given homomorphism of commutative noetherian rings $A \rightarrow B$, we consider the $B \otimes_A B$ -module structure in B induced by the canonical surjective homomorphism $B \otimes_A B \rightarrow B$, $b \otimes b' \mapsto bb'$. We denote by $fd_{B \otimes_A B}(B)$ the flat dimension of this module.

Let K be a field; then a noetherian K -algebra C is called geometrically regular if for any finite field extension $L | K$, the ring $C \otimes_K L$ is regular. A homomorphism $A \rightarrow B$ of noetherian rings is called regular if it is flat and its fibers are geometrically regular.

In this paper we prove the following result

THEOREM 1. *Let A be a noetherian ring and let B be a flat and noetherian A -algebra. If $fd_{B \otimes_A B}(B) < \infty$, then the homomorphism $A \rightarrow B$ is regular.*

This result has been obtained in [8] with two additional hypotheses: B has characteristic zero and $B \otimes_A B$ is a noetherian ring. Moreover, in [8] has been proved that if $A \rightarrow B$ is regular and $B \otimes_A B$ noetherian, then $fd_{B \otimes_A B}(B) < \infty$. Nevertheless, it is not true for an arbitrary regular homomorphism. Indeed, for a regular homomorphism $A \rightarrow B$, we have isomorphisms [2, Proposition C]

$$\bigwedge^n \Omega_{B|A} \xrightarrow{\sim} \text{Tor}_n^{B \otimes_A B}(B, B),$$

where $\Omega_{B|A}$ is the Kahler differentials module of B over A .

The proof of the version of Theorem 1 given in [8] uses the Hodge decomposition of Hochschild cohomology in characteristic zero, as well as a strong result of L. Avramov [3], which characterizes the local complete intersection rings by the finiteness of flat dimension of its cotangent complex. Here we give a more elementary proof based on the properties of Hochschild homology exposed in [5]. We also use a result of J.-L. Brylinski [4] concerning to the localization of this homology.

Observe that the condition $fd_{B \otimes_A B}(B) < \infty$ is equivalent to the existence of an integer n such that the Hochschild homology

$$H_n(B, M) = \text{Tor}_n^{B \otimes_A B}(B, M) = 0$$

for all $B \otimes_A B$ -modules M .

The results we will be using are the following

R.1. Let K be a commutative ring and let Λ, L be commutative K -algebras such that Λ is K -flat. Then for any $(L \otimes_K \Lambda) \otimes_L (L \otimes_K \Lambda)$ -module M there are isomorphisms $H_n(\Lambda, M) \simeq H_n(L \otimes_K \Lambda, M)$ (on the left hand side we have homology of Λ over K , and on the right one, homology of $L \otimes_K \Lambda$ over L).

R.2. Let K be a field and let Λ be a commutative K -algebra. Then for any Λ -modules M and N we have isomorphisms $H_n(\Lambda, M \otimes_K N) \simeq \text{Tor}_n^{\Lambda}(M, N)$.

R.3. Let $A \rightarrow B$ be a flat homomorphism of commutative rings. Let q be a prime ideal of B and let p be its contraction in A . Then for any $B \otimes_A B$ -module M we have isomorphisms

$$\text{Tor}_n^{B \otimes_A B}(B, M)_q \simeq \text{Tor}_n^{B_q \otimes_{A_p} B_q}(B_q, M_q).$$

R.1 and R.2 are in [5]. R.3 is in [4].

Proof of Theorem 1. Using R.3 it is sufficient to show that if $A \rightarrow B$ is a flat homomorphism of noetherian local rings, K the residue field of A , and $fd_{B \otimes_A B}(B) < \infty$, then $B \otimes_A K$ is a geometrically regular K -algebra. Let $C = B \otimes_A K$. It follows from R.1 that $fd_{C \otimes_K C}(C) < \infty$. Since C is noetherian, R.2 implies that the global dimension of C is finite. Then C is a regular local ring (by the classical result of J. P. Serre [10]). Moreover, a new application of R.1 and R.2, shows that C remains regular after a finite extension of K . Therefore C is geometrically regular over K , and Theorem 1 is proved.

As a particular case of Theorem 1 we obtain that a field extension $L | K$ is separable if $fd_{L \otimes_K L}(L) < \infty$. It is a generalization of Theorem 9 in [8].

COROLLARY 2. *Let A be a noetherian ring and let B be an A -algebra of finite type. The following conditions are equivalent*

- (i) B is a smooth A -algebra
- (ii) B is a flat A -module and $fd_{B \otimes_A B}(B) < \infty$.

Proof. Let us recall the characterizations of regularity and smoothness using the André-Quillen 1-dimensional (co)homology functors $H(A, B, -)$ (see [1]). This

functors can be described as follows. Let $B \simeq R/J$ be a presentation of B where R is a polynomial A -algebra, and consider the canonical homomorphism $J/J^2 \rightarrow \Omega_{R|A} \otimes_R B$. Then, for a B -module M we have

$$H_1(A, B, M) \simeq \text{Ker}(J/J^2 \otimes_B M \rightarrow \Omega_{R|A} \otimes_R M)$$

$$H^1(A, B, M) \simeq \text{Coker}(\text{Hom}_B(\Omega_{R|A} \otimes_R B, M) \rightarrow \text{Hom}_B(J/J^2, M)).$$

The above mentioned characterizations are the following:

- (1) $A \rightarrow B$ is regular if and only if $H_1(A, B, -) = 0$, i.e., $H_1(A, B, B) = 0$ and $\Omega_{B|A}$ is a flat B -module;
- (2) B is a smooth A -algebra if and only if $H^1(A, B, -) = 0$, i.e., $H_1(A, B, B) = 0$ and $\Omega_{B|A}$ is a projective B -module. Since B is noetherian and $\Omega_{B|A}$ is a finitely generated B -module, these two conditions on $\Omega_{B|A}$ are equivalent. Then the result is a consequence of [8] and Theorem 1.

REMARK 3. The proof of $H_1(A, B, -) = 0$ for a regular homomorphism is easy if B is an A -algebra of finite type. In the general case, the proof is difficult since it involves the homology functors of higher dimensions (see [1], pp. 330–331).

The homomorphism of rings $B \otimes_A B \rightarrow B$ enable us to consider a B -module M as a $B \otimes_A B$ -module. If $B \otimes_A B$ is a noetherian ring, then the vanishing of $H_n(B, M)$ for an integer n and all B -modules M , is sufficient for $\text{fd}_{B \otimes_A B}(B) < \infty$. It is an easy consequence of the following well known result: if R is a local noetherian ring with residue field K , and I is an ideal of R , then $\text{fd}_R(R/I) < \infty$ if and only if there exists an integer n such that $\text{Tor}_n^R(R/I, K) = 0$. On the other hand, the calculation of Hochschild homology of complete intersections in [11] shows that there exist locally complete intersection algebras A , of finite type over a field of characteristic zero such that $H_n(A, A) \neq 0$ for infinitely many n . We consider this as an evidence for the following conjecture:

CONJECTURE. *Let K be a field of characteristic zero and let A be a K -algebra of finite type. If $H_n(A, A) = 0$ for n sufficiently large, then A is a smooth K -algebra.*

A conjecture of J. Herzog implies that such an algebra must be at least a locally complete intersection. In fact, since K has characteristic zero, $H_n(K, A, A)$ is a summand of $H_{n+1}(A, A)$ (see [6], where $H_n(K, A, A)$ is called $H_{1,n}(A, A)$). Then $H_n(K, A, A) = 0$ for n sufficiently large. Let R be a polynomial K -algebra of finite type such that there exists a surjective K -homomorphism $R \rightarrow A$. Since the André–Quillen homology of R vanishes in dimensions > 0 , the Jacobi–Zriski sequence

associated to $K \rightarrow R \rightarrow A$ yields $H_n(R, A, A) = 0$ for n sufficiently large. In this situation, Herzog conjectures [7] (p. 62) that A must be a locally complete intersection.

REFERENCES

- [1] ANDRE, M., *Homologie des Algèbres Commutatives*. Springer, Berlin, 1974.
- [2] ANDRE, M., *Algèbres graduées associées et algèbres symétriques plates*. Comm. Math. Helv. **49** (1974), 277–301.
- [3] AVRAMOV, L., *Local rings of finite simplicial dimension*. Bull. AMS **10** (1984), 289–291.
- [4] BRYLINSKI, J.-L., *Central localization in Hochschild homology*. J. Pure and Appl. Alg. **57** (1989), 1–4.
- [5] CARTAN, H., and EILENBERG, S., *Homological Algebra*. Princeton Univ. Press, 1956.
- [6] GERSTENHABER, M., and SCHACK, S. D., *A Hodge-type decomposition for commutative algebra cohomology*. J. Pure and Appl. Alg. **48** (1987), 229–247.
- [7] HERZOG, J., *Homological properties of the module of differentials*. Colecao Atas Soc. Brasileira de Mat. **14** (1981).
- [8] RODICIO, A. G., *Some characterizations of smooth, regular, and complete intersection algebras*. Manuscripta Math. **59** (1987), 491–498.
- [9] ROSENBERG, A., and ZELINSKY, D., *Cohomology of infinite algebras*. Trans. AMS **82** (1956), 85–98.
- [10] SERRE, J. P., *Sur la dimension homologique des anneaux et des modules noethériens*. Proc. Intern. Symp., Tokio, Nikko 1955.
- [11] WOLFFHARDT, K., *The Hochschild homology of complete intersections*. Trans. AMS **171** (1972), 51–66.

*Departamento de Algebra
 Facultad de Matemáticas
 Universidad de Santiago de Compostela
 15771 Santiago do Compostela
 Spain*

Received October 30, 1989