Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 66 (1991)

Erratum: Leafwise hyperbolicity; a correction. **Autor:** Cantwell, John / Conlon, Lawrence

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Leafwise hyperbolicity; a correction

JOHN CANTWELL* AND LAWRENCE CONLON**

In [1], a proof of the following theorem was proposed.

THEOREM 1. Let (M, \mathcal{F}) be a C^2 -foliated manifold of codimension 1, transversely orientable and such that M is compact, every leaf is proper, and \mathcal{F} is tangent to ∂M . If no leaf of \mathcal{F} is a torus or a sphere, then there is a Riemannian metric on M relative to which each leaf of \mathcal{F} has constant curvature -1.

This theorem is correct, but there was an erroneous step in the proof, namely [1, Lemma (2.2)]. We are grateful to S. Matsumoto and N. Tsuchiya for pointing this out to us.

We fix the hypotheses of Theorem 1. A metric g with the property in that theorem will be called leafwise hyperbolic.

Let $M_0 \subseteq M_1 \subseteq \cdots \subseteq M_k \subseteq M_{k+1} \subseteq \cdots$ denote the level filtration [2]. Each M_k is a compact, nonempty, \mathscr{F} -saturated set, the leaves in $M_k \setminus M_{k-1}$ being the leaves of \mathscr{F} at level k. When all leaves are proper, it has become customary to use the term "depth" rather than "level". Since all leaves are proper and the foliation is of class C^2 , every leaf of \mathscr{F} has finite depth, hence $M = \bigcup_{k=0}^{\infty} M_k$.

PROPOSITION 1. Let M_k denote the union of leaves at dephts at most k. Then there is a nest $W_0 \subseteq W_1 \subseteq \cdots \subseteq W_k \subseteq W_{k+1} \subseteq \cdots \subseteq M$, where W_k is an open neighborhood of M_k , and there is a Riemannian metric g_k on M such that $g_k \mid W_k$ is leafwise hyperbolic for $\mathscr{F} \mid W_k, \forall k \geq 0$.

Theorem 1 follows. Indeed, $\{W_k\}_{k=0}^{\infty}$ is an open, nested cover of the compact manifold M, hence passing to a finite subcover yields a value of k for which $W_k = M$. It remains, then, to prove Proposition 1.

We fix a smooth, 1-dimensional foliation \mathcal{F}^{\perp} , everywhere transverse to \mathcal{F} . Projections along the leaves of \mathcal{F}^{\perp} can be used to define local diffeomorphisms between leaves of \mathcal{F} .

^{*} Research partially supported by N.S.F. Contract DMS-8900127

^{**} Research partially supported by N.S.F. Contract DMS-8822462

If $U \subseteq M$ is an open, connected, \mathscr{F} -saturated set, we use the notations $\widehat{U}, \widehat{i}: \widehat{U} \to M$, $\widehat{\mathscr{F}} = \widehat{i}^{-1}(\mathscr{F})$, and $\widehat{\mathscr{F}}^{\perp} = \widehat{i}^{-1}(\mathscr{F}^{\perp})$ from [1], [2], et al., for the completion of U, its natural immersion into M, and the induced foliations of \widehat{U} , respectively. Recall that U and \widehat{U} are called foliated products if \widehat{U} is diffeomorphic to $L \times [0, 1]$ in such a way that the leaves of $\widehat{\mathscr{F}}^{\perp}$ are the [0, 1]-fibers. Recall that, if U is a foliated product, then $\widehat{i}(\partial \widehat{U})$ is either a single leaf or a pair of leaves of \mathscr{F} .

DEFINITION 1. A closed subset $X \subset M$ that is a finite union of leaves of \mathscr{F} will be called a skeleton if each component of $M \setminus X$ is a foliated product. If k is the highest depth of the leaves in X, the skeleton has depth k. We will say that X (of depth k) is a full skeleton if, for each component U of $M \setminus X$, at least one of the following holds.

- (1) Every leaf L of $\widehat{\mathscr{F}}$ has image $\widehat{\imath}(L)$ at the same depth $k_0 \le k$.
- (2) If $L \subset \partial \hat{U}$ is a boundary leaf, then $\hat{i}(L)$ is a leaf at depth k.

If X is a skeleton, it was proven in [1, (1.2)] that there is an open neighborhood $W \supset X$ and a Riemannian metric g on M such that $g \mid W$ is leafwise hyperbolic for $\mathscr{F} \mid W$. Furthermore, projection along the leaves of \mathscr{F}^{\perp} defines local isometries between the leaves of $\mathscr{F} \mid W$. Finally, $\widehat{U} \setminus \widehat{i}^{-1}(W)$ is compact, for each component U of $M \setminus X$.

LEMMA 1. If there is a full skeleton X of depth N, then there is a neighborhood $W_N \supset M_N$ and a Riemannian metric g_N on M which is leafwise hyperbolic on W_N .

Proof. Let U be a component of $M \setminus X$. There are two cases, corresponding to possibilities (1) and (2) of Definition 1.

- (1) In this case, the proof of [1, Lemma (2.1)] shows how to extend the metric smoothly over all of U so as to make the curvature of the leaves of $\mathcal{F} \mid U$ constantly -1. Indeed, the metric was already appropriately defined on all but a compact submanifold $A \times [0, 1] \subset \widehat{U}$ and \mathcal{F} induces the product foliation on this submanifold. A deformation argument, using the Teichmüller space of A, created the extension. (The error in [1] was to claim that, even in the second case, where the foliation of $A \times [0, 1]$ was not a product, the above metric on the product could be "tilted" to give a hyperbolic metric along the leaves.)
- (2) We assume that the situation in (1) does not also occur. In this case, the argument is actually easier. Since M_N is compact [2, (4.6)], $\hat{i}^{-1}(M_N) \cap \hat{U} = L \times C$, where $C \subset [0, 1]$ is a closed subset containing $\{0, 1\}$. Since $U \setminus M_N \neq \emptyset$, $[0, 1] \setminus C$ has at least one component (a, b). Let a < a' < b' < b. The metric g is already defined on $W \cap \hat{i}(\hat{U})$ in such a way that projections along \mathscr{F}^{\perp} are local isometries between leaves. Using the projections $p^+: L \times (b', 1] \to L \times \{1\}$ and $p^-: L \times [0, a') \to 0$

 $L \times \{0\}$, one lifts this metric smoothly to $L \times [0, a') \cup L \times (b', 1]$. This metric agrees with g wherever both are defined.

Finite repetition of this argument, as U ranges over the components of $M \setminus X$, completes the proof.

LEMMA 2. For some integer $N \ge 0$, there exists a full skeleton of depth N.

Proof. As in [1, (1.1)], one constructs a skeleton X. Let N be the depth of X. If X is not full, consider a component U of $M \setminus X$ with boundary component(s) at depth k < N. If every leaf of $\mathcal{F} \mid U$ is at depth k, there is nothing to do. Otherwise, there is a leaf $L \subset U$ at depth $k + 1 \le N$. It is elementary that $X' = X \cup L$ is again a skeleton of depth N. If X' is not full repeat the process for X'. Finite repetition will ultimately produce a full skeleton of depth N.

For $0 \le k \le N$, we set $W_k = W_N$ and $g_k = g_N$. We also set $X = X_N$.

Each component U_i of $M \setminus X_N$ that has not been engulfed by W_N must contain a leaf L_i at depth N+1. Throwing these finitely many leaves in with X_N provides a full skeleton X_{N+1} of depth N+1. An application of Lemma 1 produces W_{N+1} and g_{N+1} as desired. It is not hard to see that W_{N+1} can be chosen to engulf W_N . Proceeding in this way, we construct the nest of open sets and the metrics as in Proposition 1.

REMARK. Projection along the leaves of \mathscr{F}^{\perp} does not always define local isometries between the leaves of \mathscr{F} . In the pieces $A \times [0, 1]$, where the metric is extended by a deformation in Teichmüller space, these projections will not be isometric. If it were possible to avoid introducing these regions, it would follow that the leafwise hyperbolic metric for \mathscr{F} is a bundlelike metric for \mathscr{F}^{\perp} , hence that the leaves of \mathscr{F} are totally geodesic in this metric. But totally geodesic foliations of compact 3-manifolds by surfaces are relatively rare.

REFERENCES

- [1] CANTWELL J. and CONLON L. Leafwise hyperbolicity of proper foliations. Comment. Math. Helv., 64 (1989) 329-337.
- [2] CANTWELL J. and CONLON L. Poincaré-Bendixson theory for leaves of codimension one. Trans. Amer. Math. Soc., 265 (1981) 181-209.

St. Louis University
Department of Mathematics
Ritter Hall
St. Louis, MO 63103, USA

Washington University
Department of Mathematics
Campus Box 1146
St. Louis, MO 63130-4899, USA