

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 66 (1991)

**Artikel:** Note on phantom phenomena and groups of self-homotopy equivalences.  
**Autor:** Roitberg, Joseph  
**DOI:** <https://doi.org/10.5169/seals-50410>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 17.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Note on phantom phenomena and groups of self-homotopy equivalences

JOSEPH ROITBERG

### §1. Introduction

The group  $\text{Aut}(X)$  of (pointed) homotopy classes of self-homotopy equivalences of a (pointed) space  $X$  has been extensively studied by homotopy-theorists. For a summary of results about  $\text{Aut}(X)$ , see the survey article by Arkowitz [A].

Typically, computations of and general results about  $\text{Aut}(X)$  have been given for spaces  $X$  which are either finite-dimensional cell complexes or finite Postnikov spaces. In this note, we focus on the hybrid space  $X = K(\mathbf{Z}, 2) \times S^3$  and obtain a complete computation of  $\text{Aut}(K(\mathbf{Z}, 2) \times S^3)$ .<sup>1</sup>

It is convenient to study  $\text{Aut}(X)$  by placing it in a short exact sequence

$$WI(X) \twoheadrightarrow \text{Aut}(X) \twoheadrightarrow \text{Aut}(X)/WI(X), \quad (1.1)$$

where  $WI(X)$  is the normal subgroup of  $\text{Aut}(X)$  consisting of the weak identities of  $X$  (see [R<sub>1</sub>]). A complete analysis of (1.1) in the case  $X = K(\mathbf{Z}, 2) \times S^3$  is possible thanks to results of Zabrodsky [Z], the author [R<sub>1</sub>] and Hopkins [H]. The analysis is interesting both because of the structure of the “phantom-like” subgroup  $WI(K(\mathbf{Z}, 2) \times S^3)$  and the interaction of this subgroup with the rather pedestrian quotient group

$$\text{Aut}(K(\mathbf{Z}, 2) \times S^3)/WI(K(\mathbf{Z}, 2) \times S^3).$$

It turns out, in fact, that

$$WI(K(\mathbf{Z}, 2) \times S^3) \cong \mathbf{R},$$

$$\text{Aut}(K(\mathbf{Z}, 2) \times S^3)/WI(K(\mathbf{Z}, 2) \times S^3) \cong \mathbf{Z}/2 \times \mathbf{Z}/2$$

---

<sup>1</sup>Partial information on  $\text{Aut}(K(\mathbf{Z}, 2) \times S^3)$  was obtained in [R<sub>1</sub>].

and that  $\text{Aut}(K(\mathbf{Z}, 2) \times S^3)$  is the semi-direct product of  $\mathbf{R}$  and  $\mathbf{Z}/2 \times \mathbf{Z}/2$  with respect to an action – made explicit in §2 – of  $\mathbf{Z}/2 \times \mathbf{Z}/2$  on  $\mathbf{R}$ .

The rest of the note is organized as follows. §2 contains the details of the computation of  $\text{Aut}(K(\mathbf{Z}, 2) \times S^3)$ . Results of Hopkins on phantom maps ([H]), relevant to the computation in §2, suggest a certain direction for studying the group-theoretic structure of  $\text{Aut}(X)$ . §3 consists then of a more or less random walk in this direction, leading to speculations about residual properties of  $\text{Aut}(X)$  and  $\text{Aut}_1(X)$ , the subgroup of  $\text{Aut}(X)$  consisting of those self-equivalences<sup>2</sup> inducing the identity on all homotopy groups, at least when  $X$  is grouplike.

A preliminary version of the main result of this note may be found in [R<sub>3</sub>].

I would like to thank Martin Bendersky for some helpful discussions of the material contained herein and Gilbert Baumslag for some useful group-theoretic information. I am also grateful to the Department of Mathematics of the University of Rochester for providing such a stimulating atmosphere while this note was being completed during the Fall 1990 semester.

## §2. Computation of $\text{Aut}(K(\mathbf{Z}, 2) \times S^3)$

Given (pointed) spaces  $X, Y$ , we follow [R<sub>1</sub>] in writing  $Ph(X, Y)$  for the (pointed) homotopy classes of phantom maps from  $X$  to  $Y$  and  $Ph(X)$  for  $Ph(X, X)$ . As in [R<sub>1</sub>], we require all spaces to be nilpotent of finite type and with finite fundamental group. If  $X = U \times V$  and  $Y$  is grouplike, there is a short exact sequence of groups

$$[U \wedge V, Y] \twoheadrightarrow [U \times V, Y] \twoheadrightarrow [U \vee V, Y]. \tag{2.1}$$

Hopkins observes ([H; Cor. 1.4]) that an element in  $[U \times V, Y]$  lies in the normal subgroup  $Ph(U \times V, Y)$  if and only if its “components” in  $[U \vee V, Y]$ ,  $[U \wedge V, Y]$  lie in  $Ph(U \vee V, Y)$ ,  $Ph(U \wedge V, Y)$  respectively.

Abbreviating

$$K = K(\mathbf{Z}, 2), \quad S = S^3,$$

we begin our study of  $\text{Aut}(K \times S)$  by examining the short exact sequence of groups

$$Ph(K \times S) \twoheadrightarrow [K \times S, K \times S] \twoheadrightarrow [K \times S, K \times S]/Ph(K \times S) \tag{2.2}$$

---

<sup>2</sup>As is customary, we often blur the distinction between a map and its homotopy class.

deriving from the standard grouplike structure on  $K \times S$ , which we write additively. Now,

$$\begin{aligned} [K \vee S, K \times S] &\cong [K, K \times S] \times [S, K \times S] \\ &\cong [K, K] \times [K, S] \times [S, S] \\ &\cong \mathbf{Z} \times [K, S] \times \mathbf{Z}; \end{aligned} \tag{2.3}$$

while

$$\begin{aligned} [K \wedge S, K \times S] &\cong [K \wedge S, K] \times [K \wedge S, S] \\ &\cong [K \wedge S, S]. \end{aligned} \tag{2.4}$$

According to [Z; Th.D],

$$\begin{aligned} [K, S] &\cong Ph(K, S) \cong \text{Ext}(\mathbb{Q}, \mathbf{Z}) \cong \mathbf{R}, \\ [K \wedge S, S] &\cong Ph(K \wedge S, S) = 0. \end{aligned} \tag{2.5}$$

Combining (2.1), (2.3), (2.4) and (2.5), we infer that (2.2) reduces to the split short exact sequence

$$\mathbf{R} \twoheadrightarrow [K \times S, K \times S] \twoheadrightarrow \mathbf{Z} \times \mathbf{Z} \tag{2.6}$$

with trivial action of  $\mathbf{Z} \times \mathbf{Z}$  on  $\mathbf{R}$ . For future use, we call attention to the isomorphism  $Ph(K, S) \cong Ph(K \times S) (\cong \mathbf{R})$  which associates to  $\varphi$  in  $Ph(K, S)$  the element  $\Phi$  in  $Ph(K \times S)$  defined by

$$\Phi : (k, s) \rightarrow (0, \varphi(k)). \tag{2.7}$$

In terms of the splitting (2.6),  $\text{Aut}(K \times S)$  is characterized as the subset of  $[K \times S, K \times S]$  having each of its  $\mathbf{Z}$ -components equal to  $\pm 1$ . Let  $C$  be the subgroup of  $\text{Aut}(K \times S)$  generated by the automorphisms  $\xi, \eta$  defined by

$$\xi : (k, s) \rightarrow (k, -s), \quad \eta : (k, s) \rightarrow (-k, s).$$

In terms of the splitting (2.6), the  $\mathbf{Z}$ -components of  $\xi$  and  $\eta$  are  $(1, -1)$  and  $(-1, 1)$  respectively. Plainly,

$$C \cong \mathbf{Z}/2 \times \mathbf{Z}/2.$$

Our main result is the following:

**THEOREM 2.1.** *Aut*  $(K \times S)$  is the semidirect product of the normal subgroup  $WI(K \times S) \cong \mathbf{R}$  and the subgroup  $C$ , with the (conjugation) action of  $C$  on  $WI(K \times S)$  given by

$$\xi * w = w^{-1} = \eta * w, \quad w \in WI(K \times S).$$

*Proof.* It was proved in [R<sub>1</sub>; Th.3.1] that for any grouplike space  $X$ , the map  $Ph(X) \rightarrow WI(X)$  defined by

$$\Phi \rightarrow \Phi + 1_X$$

is an isomorphism of groups.<sup>3</sup> Thus,  $WI(K \times S) \cong Ph(K \times S) \cong \mathbf{R}$  and any  $w \in WI(K \times S)$  has the form

$$w : (k, s) \rightarrow (k, \varphi(k) + s) \tag{2.8}$$

for an element  $\varphi$  in  $Ph(K, S)$  uniquely determined by  $w$  (see (2.7)). In terms of the splitting (2.6),  $WI(K \times S)$  is therefore characterized as the subset of  $[K \times S, K \times S]$  having each of its  $\mathbf{Z}$ -components equal to  $+1$ . In other words,  $WI(K \times S)$  coincides with  $\text{Aut}_1(K \times S)$  – see the penultimate paragraph of §1 for the definition of the latter.

For  $\alpha$  in  $\text{Aut}(K \times S)$ , there exists  $\gamma$  in  $C$  such that  $\alpha \circ \gamma$  lies in  $\text{Aut}_1(K \times S) = WI(K \times S)$ ; moreover,  $\gamma$  is uniquely determined by  $\alpha$ . We may therefore write

$$\alpha = w \circ \gamma, \quad w \in WI(K \times S), \quad \gamma \in C$$

with both  $w$  and  $\gamma$  uniquely determined by  $\alpha$ , and so  $\text{Aut}(K \times S)$  is, indeed, the semidirect product of  $WI(K \times S)$  and  $C$ . It remains to establish that

$$\xi \circ w \circ \xi = w^{-1} = \eta \circ w \circ \eta, \quad w \in WI(K \times S).$$

---

<sup>3</sup>The question of whether the map  $Ph(X) \rightarrow WI(X)$  defined by  $\Phi \rightarrow 1_X + \Phi$  is an isomorphism of groups when  $X$  is a cogroup was raised in [R<sub>2</sub>]. This question has now been settled in the affirmative by Touhey [T].

First we compute

$$\begin{aligned}\xi \circ w \circ \xi : (k, s) &\rightarrow (k, -s) \\ &\rightarrow (k, \varphi(k) - s) \text{ (by (2.8))} \\ &\rightarrow (k, -(\varphi(k) - s)).\end{aligned}$$

But the map

$$(k, s) \rightarrow (k, -(\varphi(k) - s))$$

is homotopic to the map

$$(k, s) \rightarrow (k, s - \varphi(k))$$

which, in turn, is homotopic to the map

$$w^{-1} : (k, s) \rightarrow (k, -\varphi(k) + s)$$

as  $[K \times S, K \times S]$  is abelian. [It would be sufficient to know that  $Ph(K \times S)$  is a central subgroup of  $[K \times S, K \times S]$ .]

Next we compute

$$\begin{aligned}\eta \circ w \circ \eta : (k, s) &\rightarrow (-k, s) \\ &\rightarrow (-k, \varphi(-k) + s) \\ &\rightarrow (-(-k), \varphi(-k) + s),\end{aligned}$$

which is homotopic to the map

$$(k, s) \rightarrow (k, \varphi(-k) + s).$$

Allowing for a moment that  $\varphi$  is an  $H$ -map, we infer that the latter map is homotopic to the map

$$w^{-1} : (k, s) \rightarrow (k, -\varphi(k) + s).$$

The following lemma then completes the proof of Theorem 2.1.

**LEMMA 2.1.** *All elements of  $Ph(K, S)$  are represented by  $H$ -maps, no matter which  $H$ -space structure is used on  $S$ .*

*Proof.* Given  $\varphi$  in  $Ph(K, S)$ , the unique multiplication  $m_K$  on  $K$  and any multiplication  $m_S$  on  $S$ , we must prove that  $\varphi \circ m_K$  and  $m_S \circ (\varphi \times \varphi)$  – both of which are clearly phantoms – are equal as elements of  $[K \times K, S]$ . To this end, it suffices to show that the components of  $\varphi \circ m_K$  in  $[K \vee K, S]$  and  $[K \wedge K, S]$  coincide with the corresponding components of  $m_S \circ (\varphi \times \varphi)$ , bearing in mind that the components of both  $\varphi \circ m_K$  and  $m_S \circ (\varphi \times \varphi)$  are phantoms (see (2.1) et seq). That the components of  $\varphi \circ m_K$  and  $m_S \circ (\varphi \times \varphi)$  in  $[K \vee K, S]$  coincide is evident since  $m_K$  and  $m_S$  are multiplications. On the other hand,

$$Ph(K \wedge K, S) = 0$$

by [Z; Th.D] since  $H_2(K \wedge K; \mathbb{Q}) = 0$ . Hence the components of  $\varphi \circ m_K$  and  $m_S \circ (\varphi \times \varphi)$  in  $[K \wedge K, S]$  are both 0, and so coincide.

For reference in §3, we record a corollary to Theorem 2.1.

**COROLLARY 2.1.** *Aut  $(K \times S)$ , while solvable, is not residually nilpotent.*

*Proof.* To see that  $\text{Aut}(K \times S)$  is not residually nilpotent, it suffices to show that the intersection of the terms  $\Gamma_N(\text{Aut}(K \times S))$ ,  $N \geq 1$ , in the lower central series of  $\text{Aut}(K \times S)$  is non-trivial. From Theorem 2.1,

$$\Gamma_N(\text{Aut}(K \times S)) = WI(K \times S), N \geq 1$$

and Corollary 2.1 is established.

### §3. Aut $(X)$ and residual properties

The point of departure for the discussion in this section is the short exact sequence of groups

$$Ph(X, Y) \hookrightarrow [X, Y] \rightarrow \varprojlim [X_\alpha, Y], \tag{3.1}$$

where  $X_\alpha$  runs over the finite (connected) subcomplexes of the cell complex  $X$  and  $Y$  is grouplike.<sup>4</sup>

Hopkins remarks ([H; proof of Prop. 1.1]) that since each  $[X_\alpha, Y]$  is nilpotent (according to a classical theorem of G. W. Whitehead), it is clear that  $\varprojlim [X_\alpha, Y]$

---

<sup>4</sup>If  $Y$  is not grouplike, (3.1) is still a short exact sequence of sets.

is residually nilpotent. He then argues that, provided  $Y$  has the homotopy type of a finite cell complex,  $Ph(X, Y)$  intersects the  $N^{\text{th}}$  term of the lower central series of  $[X, Y]$  trivially for sufficiently large  $N$  (and therefore that  $[X, Y]$  is residually nilpotent) but his argument seems to have a flaw. Nevertheless, later in [H], Hopkins conjectures that, again provided  $Y$  has the homotopy type of a finite cell complex,  $[X, Y]$  is nilpotent – not merely residually nilpotent – and establishes this conjecture in many cases, for instance when the integral homology of  $Y$  is torsion-free ([H; Cor. 2.2]).<sup>5</sup>

More generally, if each  $[X_\alpha, Y]$  possesses some group-theoretic property  $\mathcal{P}$ , then  $\varprojlim [X_\alpha, Y]$  is residually  $\mathcal{P}$  and  $Ph(X, Y)$  is the sole possible obstruction to  $[X, Y]$  being residually  $\mathcal{P}$ . As an example, referring to §2,  $Ph(K, S)$  is the (genuine!) obstruction to the residual finiteness of  $[K, S]$ . This example illustrates

**THEOREM 3.1.**  $[X, Y]$  is residually finite  $\Leftrightarrow Ph(X, Y) = 0$ .

*Proof.* We know, by the already-cited theorem of G. W. Whitehead, that  $[X_\alpha, Y]$  is (finitely generated) nilpotent, hence residually finite (see, e.g., [B; Cor. 1.21]). It is then clear that  $\varprojlim [X_\alpha, Y]$  is residually finite.

To complete the proof, it suffices to observe that a non-0 element in the (divisible) group  $Ph(X, Y)$  cannot be detected in a finite homomorphic image of  $[X, Y]$ .

In a similar vein, the following question may be posed.

**QUESTION 3.1.** Is  $[X, Y]$  residually finitely generated (or residually finitely presented)  $\Leftrightarrow Ph(X, Y) = 0$ .

The implications  $\Leftarrow$  are certainly valid. However, finitely presented groups and, a fortiori, finitely generated groups, have complicated subgroup structures – for instance, any countable, abelian, divisible group embeds in some finitely presented group – so it is conceivable that the implications  $\Rightarrow$  fail.

We now seek analogues of the foregoing for the group  $\text{Aut}(X)$  and begin by noting a variant of (3.1). For convenience, we impose the technical assumption that  $X$  has a cell complex structure for which each  $n$ -skeleton  $X_n$  is a finite cell complex.

**PROPOSITION 3.1.** *There is a short exact sequence of groups*

$$Ph(X, Y) \rightarrow [X, Y] \xrightarrow{P_*} \varprojlim [P_n X, P_n Y], \tag{3.1'}$$

---

<sup>5</sup>Added later: The conjecture is false in general. See V. K. Rao,  $SO(n)$  is not homotopy nilpotent for  $n \neq 2^m - 3, 2^m - 2$  (preprint).

where  $P_n X, P_n Y$  are the  $n^{\text{th}}$  Postnikov approximations of  $X, Y$  and  $P_*$  is the epimorphism induced by associating to a map  $f: X \rightarrow Y$  the compatible family of induced maps  $P_n f: P_n X \rightarrow P_n Y$ .

*Proof.* We show that  $Ph(X, Y) = \ker P_*$ . First, let  $f: X \rightarrow Y$  be such that  $P_n f: P_n X \rightarrow P_n Y$  is trivial for all  $n$ . From the commutative diagram

$$\begin{array}{ccc} X_n \hookrightarrow X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ P_m X & \xrightarrow{P_m f} & P_m Y, \end{array}$$

and the fact that  $[X_n, Y] \xrightarrow{\cong} [X_n, P_m Y]$  for  $m \geq n$ , we infer that  $f|_{X_n}$  is trivial for all  $n$ , hence that  $f$  is phantom.

Conversely, let  $f: X \rightarrow Y$  be phantom and consider (3.2) for  $m = n - 1$ . If  $C$  is the cofiber of  $X_n \hookrightarrow P_{n-1} X$ , we have an exact sequence of groups

$$[C, P_{n-1} Y] \rightarrow [P_{n-1} X, P_{n-1} Y] \rightarrow [X_n, P_{n-1} Y].$$

Since the image of  $P_{n-1} f$  in  $[X_n, P_{n-1} Y]$  is trivial and since  $[C, P_{n-1} Y]$  is trivial, we infer that  $P_{n-1} f$  is trivial for all  $n$ , hence that  $P_* f$  is trivial.

Setting  $X = Y$  in Proposition 3.1, we have

**COROLLARY 3.1.** *There is a short exact sequence of groups*

$$Ph(X) \hookrightarrow [X, X] \rightarrow \varprojlim [P_n X, P_n X].$$

An analogue of Corollary 3.1 for the group  $\text{Aut}(X)$  is readily available, namely

**PROPOSITION 3.2.** *There is a short exact sequence of groups*

$$WI(X) \hookrightarrow \text{Aut}(X) \rightarrow \varprojlim \text{Aut}(P_n X).$$

The proof of Proposition 3.2 is similar to that of Proposition 3.1 and is omitted. Note the contrast of the short exact sequence in Proposition 3.2 with that in (1.1).

To what extent does  $WI(X)$  obstruct  $\text{Aut}(X)$  from satisfying a group-theoretic property  $\mathcal{P}$  residually when each  $\text{Aut}(P_n X)$  satisfies  $\mathcal{P}$ ?

The situation for residual finiteness is as follows. Here we need not bother to determine whether  $\text{Aut}(P_n X)$  is residually finite (it is) as we may simply appeal to  $[R_1; \text{Th.3.2}]$  to conclude the following exact analogue of Theorem 3.1.

**THEOREM 3.2.** *Aut (X) is residually finite  $\Leftrightarrow$  WI(X) is trivial.*

With regard to finite presentability, we recall that for any (not necessarily grouplike) X, a theorem of Wilkerson [W] and Sullivan [Su] implies that Aut (P<sub>n</sub>X) is finitely presented. We ask

**QUESTION 3.2.** Is Aut (X) residually finitely presented  $\Leftrightarrow$  WI(X) is trivial?

The same caveat issued following Question 3.1 applies to Question 3.2.

Turning next to nilpotence, there does not seem to be an issue since, ordinarily, Aut (X) is far from being nilpotent even for X a finite cell complex or a finite Postnikov space. Corollary 2.1 shows that Aut (X) need not be residually nilpotent even if it accidentally occurs that  $\varprojlim$  Aut (P<sub>n</sub>X) is nilpotent. Therefore, we shift attention to the subgroup, Aut<sub>1</sub> (X), of Aut (X) consisting of those self-equivalences inducing the identity on all homotopy groups. For any (not necessarily grouplike) X, a theorem of Dror–Zabrodsky [DZ] asserts that Aut<sub>1</sub> (P<sub>n</sub>X) is nilpotent. We ask

**QUESTION 3.3.** Is Aut<sub>1</sub> (X) residually nilpotent?

We point out a (very tenuous) link between the situations in [H] and Question 3.3. Let [X, Y]<sub>0</sub> denote the subgroup of [X, Y] consisting of those f in [X, Y] inducing the zero map on all homotopy groups. The short exact sequences in Proposition 3.1, Corollary 3.1 and Proposition 3.2 plainly all have versions in which [ , ] is replaced by [ , ]<sub>0</sub> and Aut ( ) by Aut<sub>1</sub> ( ). Now the map [X, X]<sub>0</sub> → Aut<sub>1</sub> (X) defined by

$$f \mapsto f + 1_X \tag{3.3}$$

is bijective; indeed there is a commutative diagram

$$\begin{array}{ccccc} Ph(X) & \hookrightarrow & [X, X]_0 & \twoheadrightarrow & \varprojlim [P_n X, P_n X]_0 \\ \downarrow & & \downarrow & & \downarrow \\ WI(X) & \hookrightarrow & \text{Aut}_1(X) & \twoheadrightarrow & \varprojlim \text{Aut}_1(P_n X) \end{array} \tag{3.4}$$

with the three vertical maps induced by (3.3), hence bijective. However, we emphasize that only the leftmost vertical map in (3.4) is asserted to be an isomorphism of groups.

A more conservative version of Question 3.3 would be

**QUESTION 3.4.** Is Aut<sub>1</sub> (X) residually solvable?

Though Question 3.4 seems much easier to handle than Question 3.3 – the class of solvable groups being closed under group extensions (cf. Corollary 2.1) – it must be pointed out that there are examples of non-residually solvable groups  $G$  such that  $G$  admits a residually solvable quotient group with abelian kernel.

FINAL REMARK. I am informed by C. A. McGibbon and J. M. Møller that Proposition 3.2 and the fact that  $WI(X)$  is abelian, divisible follow from results of A. K. Bousfield/D. M. Kan and C. U. Jensen, even without assuming  $X$  grouplike. Thus Theorem 3.2 and Questions 3.2, 3.3 and 3.4 may be formulated without this assumption on  $X$ . However (3.3), and hence (3.4), makes no sense without some sort of structure (grouplike, cogroup) on  $X$ .

## REFERENCES

- [A] M. ARKOWITZ, The group of self-homotopy equivalences – a survey, *Groups of Self-Equivalences and Related Topics, Proceedings, Montreal 1988* (Ed. R. A. Piccinini) 170–203, Lecture Notes in Mathematics 1425, Springer-Verlag (1990).
- [B] G. BAUMSLAG, *Lecture Notes on Nilpotent Groups*, Amer. Math. Soc., Providence, Regional Conf. Ser. 2 (1971).
- [DZ] E. DROR and A. ZABRODSKY, Unipotency and nilpotency in homotopy equivalences, *Topology* 18 (1979), 187–197.
- [H] M. J. HOPKINS, Nilpotence and finite  $H$ -spaces, *Israel Jour. Math.* 66 (1989), 238–246.
- [R<sub>1</sub>] J. ROITBERG, Weak identities, phantom maps and  $H$ -spaces, *Israel Jour. Math.* 66 (1989), 319–329.
- [R<sub>2</sub>] J. ROITBERG, Phantom maps, cogroups and the suspension map, *Quaestiones Math.* 13(3/4) 1990, 335–347.
- [R<sub>3</sub>] J. ROITBERG, On the self-homotopy equivalences of an infinite cell complex, *Abstracts AMS*, April 1990, Issue 69, Volume 11, Number 3, 253.
- [Su] D. SULLIVAN, Infinitesimal computations in topology, *I.H.E.S. Publ. Math.* 47 (1978), 269–331.
- [T] P. TOUHEY, CUNY doctoral dissertation, in preparation.
- [W] C. WILKERSON, Applications of minimal simplicial groups, *Topology* 15 (1976), 115–130.
- [Z] A. ZABRODSKY, On phantom maps and a theorem of H. Miller, *Israel Jour. Math.* 58 (1987), 129–143.

*Department of Mathematics & Statistics  
Hunter College, CUNY  
695 Park Avenue  
New York, NY 10021, USA*

and

*Department of Mathematics  
Graduate Center, CUNY  
33 West 42nd Street  
New York, NY 10036, USA*

Received November 20, 1990