

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 67 (1992)

**Artikel:** On the minimal surfaces of Riemann.  
**Autor:** Toubiana, Eric  
**DOI:** <https://doi.org/10.5169/seals-51110>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 03.05.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## On the minimal surfaces of Riemann

ERIC TOUBIANA

### §1. Introduction

Let  $\gamma_i, i = 1, 2$  be plane Jordan curves in horizontal planes  $P_i, i = 1, 2, P_1 \neq P_2$ , we know that under conditions on  $\gamma_i$ , for example  $\gamma_1$  not too far from  $\gamma_2$ ,  $\gamma_1$  and  $\gamma_2$  bound a least area minimal annulus  $M$  between  $P_1$  and  $P_2$ . Meeks and White [7] were able to prove that when the  $\gamma_i$  are convex there are at most two minimal annuli bounded by  $\gamma_1 \cup \gamma_2$ . Assuming that the  $\gamma_i$  are convex, Shiffman [10] proved that if  $M$  is a minimal annulus bounded by  $\gamma_1 \cup \gamma_2$  then for each horizontal plane  $P$  between  $P_1$  and  $P_2$ , the intersection  $P \cap M$  is again a convex Jordan curve, furthermore if  $\gamma_1$  and  $\gamma_2$  are circles, then  $P \cap M$  is also a circle.

In view of this last result it is natural to ask what happens when two straight lines replace the Jordan curves. Namely, let  $D_i, i = 1, 2$ , be straight lines in horizontal planes  $P_i, i = 1, 2, P_1 \neq P_2$ . Let us assume that  $D_1$  makes an angle  $\theta$  with  $D_2, \theta \in [0, \pi]$ . Now let  $M$  be a minimal annulus between  $P_1$  and  $P_2$  bounded by  $D_1 \cup D_2$ . If  $P$  is a horizontal plane between  $P_1$  and  $P_2$ , what can we say about the intersection  $P \cap M$ ?

For example, if  $\theta = 0$  i.e.  $D_1$  and  $D_2$  are parallel, Riemann [9] has constructed a minimal embedded annulus  $S$  between  $P_1$  and  $P_2$  bounded by  $D_1 \cup D_2$ . Furthermore the intersection with any horizontal plane is a circle, see [3] for a detailed description of Riemann's examples.

In the same paper [3], Hoffman, Karcher and Rosenberg proved that if  $D_1$  and  $D_2$  are parallel, i.e.  $\theta = 0$ , then the only minimal properly embedded annulus between  $P_1$  and  $P_2$ , bounded by  $D_1 \cup D_2$  is precisely Riemann's example.

Here we shall prove that the case  $\theta \neq 0$  does not occur, namely we show the following.

**THEOREM 1.** *Let  $D_i, i = 1, 2$  be straight lines in horizontal planes  $P_i, i = 1, 2, P_1 \neq P_2$ , let  $\theta \in [0, \pi]$  be the angle that  $D_1$  makes with  $D_2$ . Let us assume there exists a minimal properly embedded annulus  $M$  between  $P_1$  and  $P_2$  bounded by  $D_1 \cup D_2$ .*

*Then necessarily  $\theta = 0$  and  $M$  is Riemann's example  $S$ .*

Let us assume now  $\theta = 0$ , i.e.  $D_1$  and  $D_2$  parallel. We can generalize Riemann's examples to yield a family of minimal surfaces  $S_k$  with the following properties:

- (1)  $S_0 = S$  where  $S$  is Riemann's example.
- (2) For every integer  $k, k \geq 0$ ,  $S_k$  is a minimal immersed annulus between  $P_1$  and  $P_2$ , bounded by  $D_1 \cup D_2$  such that after reflection about the lines  $D_1, D_2, \dots$  we get a complete minimal surface in  $R^3$ , which we call again  $S_k$ . Furthermore  $S_k$  is invariant under the translation  $X \rightarrow X + 2u$  where  $u$  is the vector orthogonal to  $D_1$  translating  $D_1$  to  $D_2$ . Also each end of  $S_k$  in  $R^3$  is a flat horizontal end, i.e. an end asymptotic to a horizontal plane, and the projection of every end over any horizontal plane is a  $(4k + 1)$  to 1 map.

The following gives the Weierstrass representation of this family.

**THEOREM 2.** Let  $T$  be a rectangular torus, i.e.  $T^2 = C/\Gamma$  where  $\Gamma = \{2\omega_1 p + 2\omega_3 q, p, q \in \mathbf{Z}, \omega_1 \in \mathbf{R}_*^+, \omega_3 \in i\mathbf{R}_*^+\}$ .

Then for every  $k, k \in \mathbf{N}$  the following data  $(g_k, \eta_k)$  is the Weierstrass representation of the surface  $S_k$  described above.

$$g_k = \lambda_k [P(z) - P(\omega_2)]^{2k+1},$$

$$\eta_k = \frac{dz}{g_k(z)} = \frac{dz}{\lambda_k [P(z) - P(\omega_2)]^{2k+1}},$$

where  $\omega_2 = \omega_1 + \omega_3, z \in T^2 - \{0, \omega_2\}, \lambda_k = i\sqrt{(-2/P''(\omega_2))^{2k+1}}$  and  $P$  is the  $P$ -function of Weierstrass.

We prove Theorems 1 and 2 in §2 and §3 respectively.

### §2. Proof of Theorem 1

We are going to use the Weierstrass representation for a minimal surface, see Lawson [5] p. 113, and the reflection principle, namely if a minimal surface in  $\mathbf{R}^3$  has a piece  $\gamma$  of a straight line in its boundary then we can extend minimally this surface along  $\gamma$  by the reflection about the line defined by  $\gamma$ , see [5] p. 82.

Let us take the notation of Theorem 1. By the reflection principle we can extend  $M$  to a complete properly embedded minimal surface in  $\mathbf{R}^3$ , let us call  $M$  again this surface. As the composition of the two reflections about  $D_1$  and  $D_2$  is a screw-motion  $S_{2\theta}$ , i.e. the  $2\theta$ -rotation with respect to the  $x_3$ -axis composed with a vertical

translation, we get that  $M$  is globally invariant by the screw-motion  $S_{2\theta}$ :

$$S_{2\theta}(M) = M.$$

Then we get a quotient surface  $M/S_{2\theta}$  in  $\mathbf{R}^3/S_{2\theta}$ . By construction this surface is topologically a two-punctured torus properly and minimally embedded in  $\mathbf{R}^3/S_{2\theta}$ . By a theorem of Meeks and Rosenberg [6] we deduce that  $M/S_{2\theta}$  has finite total curvature. From this we get that each end of  $M$  is conformally a punctured disc, see [5] p. 130. At last we deduce that the ends of  $M$  are parallel, flat, embedded and also the Gauss map  $g$  of  $M$  has order two at each end (this last claim comes from the fact that if  $E_1$  is the end of  $M$  passing through the straight line  $D_1$  then, by construction,  $E_1 - P_1$  has two connected components). Subsequently we assume that  $D_1$  is the  $x_1$ -axis in the plane  $P_1 = \{x_3 = 0\}$ .

**LEMMA 1.** *The extended complete minimal surface  $M$  in  $\mathbf{R}^3$  is parameterized by  $C = C - (\Gamma \cup (z_0 + \Gamma))$  where:*

$$\begin{cases} \Gamma = \{2\omega_1 \cdot p + 2\pi i \cdot q, p, q \in \mathbf{Z}, \omega_1 \in \mathbf{R}_*^+\}, \\ z_0 \in \omega_1 + y\pi i, \quad 0 \leq y \leq 1. \end{cases}$$

Furthermore if  $X$  is the minimal immersion of  $C$  onto  $M \subset \mathbf{R}^3$ ,  $X$  must satisfy:

- $X$  sends the vertical lines  $\{\operatorname{Re}(z) = c, c \neq k\omega_1, k \in \mathbf{Z}\}$  to horizontal Jordan curves in  $\mathbf{R}^3$ .
- $X$  sends the vertical lines  $\{\operatorname{Re}(z) = k\omega_1, k \in \mathbf{Z}\}$  to horizontal straight lines in  $\mathbf{R}^3$  and particularly  $X$  sends  $\{\operatorname{Re}(z) = 0\}$  to the  $x_1$ -axis  $D_1$  and  $\{\operatorname{Re}(z) = \omega_1\}$  to  $D_2$ .

At last we have:

$$\forall z \in C, \quad X(z + 2\pi i) = X(z), \tag{1}$$

$$X(2\omega_1 - \bar{z}) = S_{D_2}[X(z)], \tag{2}$$


$$X(z + 2\omega_1) = S_{2\theta}[X(z)], \tag{3}$$

where  $S_{D_2}$  is the reflection about  $D_2$  and  $S_{2\theta} = S_{D_1} \circ S_{D_2}$ .

*Proof of Lemma 1.* Let  $M_1$  be the piece of  $M$  bounded by  $D_1 \cup D_2$  and let  $M_2$  be the reflection of  $M_1$  about  $D_2$ , i.e.  $M_2 = S_{D_2}(M_1)$ . Let  $D_3$  be  $S_{D_2}(D_1)$ , so that  $M_2$  is bounded by  $D_2$  and  $D_3$ .

By construction  $M' = M_1 \cup M_2$  is homeomorphic to a one-punctured planar annulus  $A$  bounded by two Jordan curves  $\gamma_1, \gamma_3$  each one with a point  $p_i, i = 1, 3$  removed.

Let  $Z$  be the minimal immersion of  $A$  onto  $M'$ , let  $A$  have the conformal structure induced by  $Z$ . As each end of  $M$  is conformally a one-punctured disc we deduce that  $A$  is conformally a one-punctured annulus bounded by two Jordan curves  $\gamma_1, \gamma_3$  each one with a point  $p_i, i = 1, 3$  removed.



Assume that  $Z(\gamma_i) = D_i, i = 1, 3$ .  $Z$  sends a neighbourhood of  $p$  to the end passing through the line  $D_2$ . Assume that  $D_2$  belongs to the plane  $P_2 = \{x_3 = c, c > 0\}$ . Then  $D_3$  belongs to the horizontal plane  $P_3 = \{x_3 = 2c\}$  (recall that  $P_1 = \{x_3 = 0\}$ ).

Let  $Z_3$  be the third coordinate function of  $Z$ ,  $Z_3$  is a harmonic function. By construction of  $M$ , the line  $D_2$  is the only part of  $M$  in the plane  $P_2$ , that is:

$$M \cap P_2 = D_2,$$

as  $Z_3(\gamma_1) = 0$  and  $Z_3(\gamma_3) = 2c$ , we deduced that  $\{Z_3^{-1}(c) \cup p\}$  is an embedded closed curve in the interior of  $\bar{A}$ , and by the maximum principle applied to the function  $Z_3$ , this curve must be connected, so that  $\{Z_3^{-1}(c) \cup p\}$  is a Jordan curve  $\gamma_2$  in the interior of  $A$ .

Now let  $\alpha$  be a real number with  $\alpha \neq c$  and  $0 \leq \alpha \leq 2c$ , again by the maximum principle we get that  $Z_3^{-1}(\alpha)$  is a Jordan curve  $\gamma$  of  $\bar{A}$ . Hence we have a foliation  $(\gamma_i)_{1 \leq i \leq 3}$  of  $\bar{A}$  with Jordan curves  $\gamma_i$ , so that  $Z_3$  is constant over each curve  $\gamma_i$ , i.e.:

$$\forall i \in [1, 3], \quad Z_3(\gamma_i) = c_i \quad \text{with: } 0 \leq c_i \leq 2c \quad \text{and} \quad c_1 = 0; c_2 = c; c_3 = 2c.$$

Let  $Z_3^*$  be a harmonic conjugate of  $Z_3$  over  $\bar{A}$ , we have locally:

$$Z_3^*(z) = \int_{\alpha} *dZ_3,$$

where  $\alpha$  is a path between a base point  $z_0$  and  $z$ , and  $*dZ_3$  is defined by:

$$*dZ_3 = \frac{\partial Z_3}{\partial x} dy - \frac{\partial Z_3}{\partial y} dx, \quad z = x + iy.$$

It can happen that  $Z_3^*$  is not globally defined on  $\bar{A}$ , i.e. if  $\gamma$  is any Jordan curve in  $\bar{A}$  generating  $\Pi_1(\bar{A})$ , let  $a$  be the real number defined by:

$$a = \int_{\gamma} *dZ_3.$$

Then  $Z_3^*$  is well defined on  $\bar{A}$  if and only if  $a = 0$ . In case  $a \neq 0$  let us consider the function:


$$F(z) = \exp \left[ \frac{2\pi}{a} (Z_3(z) + iZ_3^*(z)) \right].$$

In case  $a = 0$  we put:

$$F(z) = \exp [Z_3(z) + iZ_3^*(z)],$$

in any case  $F$  is a well defined map of  $\bar{A}$  into  $C$ ,  $F$  maps the Jordan curves  $\{\gamma_i\}_{1 \leq i \leq 3}$  to concentric circles of  $C$ :  $\{|u| = \exp [(2\pi/a)c_i]$  or  $\exp [c_i]$  depending on the expression of  $F\}$ .  $F$  maps  $\bar{A}$  on an annulus  $B$  of  $C$  bounded by two concentric circles  $C_1, C_3$  with

$$F(\gamma_i) = C_i, \quad 1 \leq i \leq 3.$$



It is easy to see that  $F$  is a  $n$ -covering map of  $\bar{A}$  onto  $B$  in  $C$ , so after composing  $F$  with the  $n$ th root map, we can assume that  $F$  is an embedding of  $\bar{A}$  onto  $B$ .

Let us call  $Y$  the composed map  $Z \circ F^{-1}$ ,

$$B \xrightarrow{Y = Z \circ F^{-1}} \mathbf{R}^3$$


we clearly have  $Y(C_i) = D_i, i = 1, 2, 3$  so by the reflection principle if  $I_C$  denotes the reflection in  $\mathbf{C}$  about a circle  $C$  we have:

$$C_3 = I_{C_2}(C_1) \quad (\text{because } Y(C_3) = D_3 = S_{D_2}(D_1))$$

and then  $F(P_3) = I_{C_2}(F(P_1))$ .

Now by reflection principle we extend  $Y$  to a minimal immersion of  $\mathbf{C} - E$  onto  $M$  in  $\mathbf{R}^3$ , where  $E$  is the infinite discrete set of  $\mathbf{C}$  obtained by taking  $F(P_1), F(P)$  and all the successive images of those points by the reflections about the circles  $C_1, C_2, I_{C_1}(C_2), I_{C_2}(C_1) \dots$

Then the exponential map  $z \rightarrow u = \exp(z)$  of  $\mathbf{C}$  sends the vertical lines  $\{\text{Re}(z) = a\}$  of the  $z$ -complex plane on the concentric circles  $\{|u| = e^a\}$  of the  $u$ -complex plane.



Up to a homothety of the  $u$ -plane we can assume that  $C_1$  is the circle of radius one. So that if we call  $X$  the composed map:  $X = Y \circ \exp$ ,  $X$  sends the imaginary axis  $\{\text{Re}(z) = 0\}$  to  $D_1$  in  $\mathbf{R}^3$ . Furthermore up to a rotation in the  $u$ -plane, we can assume that  $F(p_1) = 1$ , so that  $\exp(0) = F(p_1) = 1$ . Let denote  $z_0$  the inverse image, by the exponential map, of  $F(p)$  with  $0 \leq \text{Im } z_0 \leq 2\pi$ , and  $\omega_1$  the real part of  $z_0$ .

$$\omega_1 = \text{Re}(z_0).$$

Then  $X$  sends the line  $\{\text{Re}(z) = \omega_1\}$  to the horizontal line  $D_2$  in  $\mathbf{R}^3$ .

It is clear that the inverse image of  $E$  by the exponential map is just  $\Gamma \cup (z_0 + \Gamma)$ , with

$$\Gamma = \{2\omega_1 p + 2\pi i q, p, q \in \mathbf{Z}\}.$$

Furthermore by construction of  $Y$  and by the geometric properties of  $M$  in  $\mathbf{R}^3$  we must have:

$$\forall z \in C = \mathbf{C} - (\Gamma \cup (z_0 + \Gamma)), \quad X(z + 2\pi i) = X(z), \tag{1}$$

$$X(2\omega_1 - \bar{z}) = S_{D_2}[X(z)], \tag{2}$$

$$X(4\omega_1 - \bar{z}) = S_{D_3}[X(z)]. \tag{2}'$$

(1) comes from the fact that  $X = Y \circ \exp$ , and (2) (resp. (2)') comes from the fact that  $M$  is invariant by the reflection about  $D_2$  (resp.  $D_3$ ).

So that combining (2) and (2)' we have:

$$\forall z \in C, \quad X(z + 2\omega_1) = S_{2\theta}[X(z)]. \quad (3)$$

Finally considering the map  $z \rightarrow -z$  of the  $z$ -plane keeping fixed the point 0, and because of (1) we can assume that:

$$0 \leq \operatorname{Re}(z_0) \quad \text{and} \quad \operatorname{Im}(z_0) = y\pi, \quad 0 \leq y \leq 1.$$

### Remarks

(1) From Lemma 1 we see that  $X$  induces a minimal embedding of  $C/_{2\pi i\mathbf{Z}}$  onto  $M$  in  $\mathbf{R}^3$  and a minimal embedding of  $T^2 - \{0, z_0\} = C/_{\Gamma \cup (z_0 + \Gamma)}$  onto  $M/_{S_{2\theta}}$  in  $\mathbf{R}^3/_{S_{2\theta}}$ .

(2) Until now we don't know if  $\omega_1$  is bigger or not than  $\pi$ , so from now we assume  $\pi$  bigger than  $\omega_1$ , the rest of the proof will not be affected.

Let  $\omega_3 = i\pi$ , so that the lattice  $\Gamma$  is:

$$\Gamma = \{2\omega_1 p + 2\pi i q, p, q \in \mathbf{Z}\}.$$

**LEMMA 2.** *With the hypothesis of Theorem 1 we have  $\theta = 0$  and the Weierstrass representation of the minimal immersion  $X$  of  $C$  in  $\mathbf{R}^3$  in Lemma 1 is:*

$$\forall z \in C, \quad g(z) = \lambda[P(z) - P(\omega_2)],$$

$$\eta = \frac{dz}{g(z)} = \frac{dz}{\lambda[P(z) - P(\omega_2)]},$$

where  $P$  is the  $P$ -function of Weierstrass associated to the torus  $T^2 = C/\Gamma$ ,  $\omega_2 = \omega_1 + \omega_3$  and  $\lambda = i\sqrt{-2/P''(\omega_2)} \in i\mathbf{R}$ .

*Proof of Lemma 2.* We are going to use some basic facts about the  $\sigma$  and  $\zeta$  functions associated with the  $P$ -function of Weierstrass (we recall these facts in the Appendix).

Let us call  $(\Phi_1, \Phi_2, \Phi_3)$  the coordinate forms of  $X$ , i.e.

$$X(z) = (X_1(z), X_2(z), X_3(z)) = \operatorname{Re} \int_{z_0}^z (\Phi_1, \Phi_2, \Phi_3)$$

with  $z_0$  a base point in  $C$  and:

$$\Phi_1 = \frac{1}{2} \eta(1 - g^2); \quad \Phi_2 = \frac{i}{2} \eta(1 + g^2); \quad \Phi_3 = \eta g.$$

As the third coordinate function  $X_3$  is constant on vertical lines  $\{\text{Re}(z) = c\}$  we can assume  $\Phi_3 = 1$  and then  $\eta = dz/g(z)$ .

As  $M/S_{20}$  has two flat horizontal ends parameterized by 0 and  $z_0$  (with  $\text{Re}(z_0) = \omega_1$ , see Lemma 1), we can assume that

$$g(0) = \infty \quad \text{and} \quad g(z_0) = 0,$$

also  $g$  is a 2 to 1 map near each end.

Furthermore on a fundamental domain of  $T^2$  in  $C$  there is no other point where  $g$  is vertical, if not, up to a rotation in  $R^3$ , another such point would be an  $n$ -pole of  $g$ ,  $n \geq 1$ , but this point is a regular point of  $M$ , so it must be a  $2n$ -zero of  $\eta$  and then it is an  $n$ -pole of  $\Phi_3 = g\eta$ , but  $\Phi_3 = 1$  as we saw before.

As  $g$  is the Gauss map of  $M$ , up to the stereographic projection of  $S^2$  to  $C \cup \{\infty\}$ , by the geometric properties of  $M$  in  $R^3$ ,  $g$  must satisfy:

$$\forall z \in C \quad \begin{cases} g(z + 2\omega_3) = g(z), & (1) \\ g(2\omega_1 - \bar{z}) = -e^{2i\theta} \overline{g(z)}, & (2) \\ g(z + 2\omega_1) = e^{2i\theta} g(z). & (3) \end{cases}$$

Those properties (1), (2), (3) of  $g$  come from respectively the properties (1), (2), (3) of  $X$  established in Lemma 1.

From (3) we deduce that the map  $(g'/g)(z)$  is well defined on the torus  $T^2 = C/\Gamma$ . Furthermore  $(g'/g)(z)$  is an elliptic function on  $T^2$  and has a single pole at a point  $z$  in  $T^2$  if and only if  $z$  is a pole or a zero of  $g$ , and  $(g'/g)(z)$  has no other pole on  $T^2$ . From what we saw before we see that  $(g'/g)(z)$  has two single poles on  $T^2$ , so it is an elliptic function of degree two on  $T^2$  with a single pole at 0 and  $z_0$ . The most general form of such a function is:

$$\frac{g'}{g}(z) = \frac{a_1 P(z) + a_0 + bP'(z)}{P(z) - P(z_0)}, \quad a_1, a_0, b \in C$$

with  $a_1 P(z_0) + a_0 = bP'(z_0)$  because  $-z_0$  is not a pole of  $(g'/g)(z)$ .

From (2) we get that

$$\forall z \in \mathbf{C}, \quad 2\omega_1 - \bar{z} = z \Rightarrow \frac{g'}{g}(z) \in i\mathbf{R}.$$

So  $(g'/g)(z)$  must have purely imaginary values on the line  $\{\operatorname{Re}(z) = \omega_1\}$  in  $\mathbf{C}$ . But as  $\operatorname{Re}(z_0) = \omega_1$ ,  $P(z_0) \in \mathbf{R}$  and then  $P(z) - P(z_0)$  has real values on the line  $\{\operatorname{Re}(z) = \omega_1\}$  also  $P'(z)$  has purely imaginary values on this line, so we have:

$$\forall y \in \mathbf{R}, \quad a_1 P(\omega_1 + y\omega_3) + a_0 + bP'(\omega_1 + y\omega_3) \in i\mathbf{R}.$$

For  $y = 0$  and  $y = 1$  we deduce (as  $P'(\omega_1) = 0 = P'(\omega_1 + \omega_3)$ )

$$a_1 P(\omega_1) + a_0 \in i\mathbf{R},$$

$$a_1 P(\omega_2) + a_0 \in i\mathbf{R},$$

as  $P(\omega_1)$  and  $P(\omega_2)$  are two distinct real numbers, we get that  $a_1$  and  $a_0$  are purely imaginary numbers,  $a_1, a_0 \in i\mathbf{R}$ , so that:

$$\forall y \in \mathbf{R}, \quad bP'(\omega_1 + y\omega_3) \in i\mathbf{R}$$

and then  $b$  is a real number:

$$a_1, a_0 \in i\mathbf{R}, \quad b \in \mathbf{R}.$$

Now let us look for  $g$ , we have:

$$\forall z \in \mathbf{C}, \quad \frac{g'}{g}(z) = a_1 + \frac{a_1 P(z_0) + a_0}{P(z) - P(z_0)} + b \frac{P'(z)}{P(z) - P(z_0)}.$$

We want to show that  $z_0 = \omega_2$ . Let us suppose now that  $z_0 \neq \omega_i$ ,  $i = 1, 2$ , that is:

$$z_0 = \omega_1 + y\omega_3, \quad 0 < y < 1.$$

As  $P'(z_0) \neq 0$  we have

$$\frac{1}{P(z) - P(z_0)} = \frac{1}{P'(z_0)} [\zeta(z - z_0) - \zeta(z + z_0) + 2\zeta(z_0)]$$

as  $a_1 P(z_0) + a_0 = bP'(z_0)$  we get:

$$\frac{g'}{g}(z) = a_1 + 2\zeta(z_0) + b \left[ \zeta(z - z_0) - \zeta(z + z_0) + \frac{P'(z)}{P(z) - P(z_0)} \right]$$

and so

$$\forall z \in \mathbf{C}, \quad g(z) = \lambda e^{[a_1 + 2b\zeta(z_0)]z} \cdot \left[ \frac{\sigma(z - z_0)}{\sigma(z + z_0)} (P(z) - P(z_0)) \right]^b$$

with  $\lambda \in \mathbf{C}, a_1 \in i\mathbf{R}, b \in \mathbf{R}$ .

But  $g$  must have a double pole at 0 and a double zero at  $z_0$ , now we remark that the function in brackets  $(\sigma(z - z_0)/\sigma(z + z_0))(P(z) - P(z_0))$ , has the same zeros and poles, so we must have  $b = 1$ , so:

$$\forall z \in \mathbf{C}, \quad g(z) = \lambda e^{Az} \frac{\sigma(z - z_0)}{\sigma(z + z_0)} (P(z) - P(z_0))$$

where,  $\lambda \in \mathbf{C}, A = a_1 + 2\zeta(z_0)$ .

From the property ( $\#$ ) of the  $\sigma$ -function, see the Appendix, we have:

$$\forall z \in \mathbf{C}, \quad g(z + 2\omega_i) = e^{2\omega_i A - 4\eta_i z_0} g(z), \quad i = 1, 2, 3,$$

from conditions (1) and (3) on  $g$  we get:

$$\begin{cases} 2\omega_1 A - 4\eta_1 z_0 = 2i\theta + 2p\pi i, & p \in \mathbf{Z}, \\ 2\omega_3 A - 4\eta_3 z_0 = 2q\pi i, & q \in \mathbf{Z} \end{cases}$$

and then

$$z_0 = z_0(\theta) = -\frac{2\theta}{2\pi} \omega_3 - p\omega_3 + q\omega_1.$$

By considering the real part and the imaginary part of  $z_0$ , as we know that  $\text{Re}(z_0) = \omega_1$ , and  $0 < \text{Im}(z_0) < -i\omega_3$  we have  $q = 1$  and  $p = -1$ , so:

$$z_0(\theta) = \omega_1 + \left( 1 - \frac{\theta}{\pi} \right) \omega_3, \quad \theta \in [0, \pi]$$

and

$$2\omega_1 A = 4\eta_1 z_0(\theta) + 2i\theta - 2\pi i. \quad (*)$$

On the other hand as the map  $X$  is well defined on  $C$  we must have:

$$\forall \gamma \in \Pi_1(C), \quad \operatorname{Re} \int_{\gamma} (\Phi_1, \Phi_2, \Phi_3) = 0.$$

In particular if  $\gamma$  is a small circle around 0, we must have:

$$\operatorname{Re} \int_{\gamma} (\Phi_1, \Phi_2) = 0,$$

that is

$$\int_{\gamma} \eta = \overline{\int_{\gamma} \eta g^2}$$

and as  $\eta = dz/g(z)$  then:

$$-2\pi \overline{\operatorname{Re} s(g, 0)} = 2\pi \operatorname{Re} s\left(\frac{1}{g}, 0\right) = 0 \quad \left(\text{because } 0 \text{ is a zero of } \frac{1}{g}\right)$$

and so we must have

$$\operatorname{Re} s(g, 0) = 0.$$

A computation shows that:

$$\operatorname{Re} s(g, 0) = -\lambda(A - 2\zeta(z_0)).$$

So we get that  $A = 2\zeta(z_0)$ . In view of (\*) we have

$$2\omega_1 \zeta(z_0(\theta)) = 2\eta_1 z_0(\theta) + i\theta - i\pi. \quad (T)$$

Let  $h(\theta)$  be the function on  $[0, \pi]$  defined by:

$$\forall \theta \in [0, \pi], \quad h(\theta) = \frac{1}{i} [2\omega_1 \zeta(z_0(\theta)) - 2\eta_1 z_0(\theta) - i\theta + i\pi].$$

As  $T^2$  is a rectangular torus we have:

$$\forall y \in \mathbf{R}, \quad \operatorname{Re} (\zeta(\omega_1 + y\omega_3)) = \eta_1 = \zeta(\omega_1),$$

and then:

$$\forall \theta \in [0, \pi], \quad \operatorname{Re} [\zeta(z_0(\theta))] = \eta_1;$$

we deduce that  $h(\theta)$  is a real-valued function.

Now let us look for solutions  $\theta$  of  $(T)$ , i.e. zeros of the  $h$ -function, we have:

$$\begin{aligned} h(0) &= \frac{1}{i} [2\omega_1 \zeta(\omega_2) - 2\eta_1 \omega_2 + \pi i] \\ &= \frac{1}{i} [2\omega_1 \eta_2 - 2\eta_1 \omega_2 + \pi i] \\ &= 0 \quad \text{by Legendre relation, (see the Appendix)} \end{aligned}$$

and

$$h(\pi) = \frac{1}{i} [2\omega_1 \zeta(\omega_1) - 2\eta_1 \omega_1] = \frac{1}{i} [2\omega_1 \eta_1 - 2\eta_1 \omega_1] = 0.$$

So 0 and  $\pi$  are solutions of  $(T)$ . Let us show there is no other solution in  $[0, \pi]$ . Suppose there is another solution  $\theta$  in  $]0, \pi[$ , then by Rolle's theorem the function  $h'(\theta)$  would have at least two distinct zeros in  $]0, \pi[$ , but we have

$$h'(\theta) = \frac{1}{i} \left[ \frac{2\omega_1 \omega_3}{\pi} P(z_0(\theta)) + 2 \frac{\eta_1 \omega_3}{\pi} \right].$$

$z_0(\theta)$  is a strictly monotone function of  $\theta$  with values in  $L = \{\omega_1 + y\omega_3, 0 \leq y \leq 1\}$ . Furthermore  $P$  is a strictly monotone and real-valued function on  $L$ . We deduce that  $h'(\theta)$  is a strictly monotone function on  $[0, \pi]$  and then  $h'(\theta)$  cannot have two zeros on  $]0, \pi[$ .

So the only solutions of  $(T)$  are  $\theta = 0$  and  $\theta = \pi$ , but we then get  $z_0 = \omega_2$  or  $z_0 = \omega_1$ . Let us show that  $z_0 = \omega_1$  does not work. For that let us assume that  $z_0 = \omega_1$ . Then  $(g'/g)(z)$  must have a single pole at 0 and  $\omega_1$ , so:

$$\forall z \in \mathbf{C}, \quad \frac{g'}{g}(z) = \frac{a_1 P(z) + a_0 + bP'(z)}{P(z) - P(\omega_1)}, \quad a_1, a_0, b \in \mathbf{C},$$

with:  $a_1 P(\omega_1) + a_0 = 0$ , we deduce that:

$$\frac{g'}{g}(z) = a_1 + b \frac{P'(z)}{P(z) - P(\omega_1)}.$$

As before we must have  $a_1 \in i\mathbf{R}$  and  $b \in \mathbf{R}$ , furthermore:

$$\forall z \in \mathbf{C}, \quad g(z) = \lambda e^{a_1 z} [P(z) - P(\omega_1)]^b.$$

As  $g$  must have a double pole at 0 and a double zero at  $\omega_1$  we deduce  $b = 1$ , and

$$\forall z \in \mathbf{C}, \quad g(z) = \lambda e^{a_1 z} [P(z) - P(\omega_1)], \quad \lambda \in \mathbf{C}, \quad a_1 \in i\mathbf{R}$$

but as we must have  $g(z + 2\omega_3) = g(z)$ , we deduce that  $a_1 \cdot 2\omega_3 = 2\pi i q$ ,  $q \in \mathbf{Z}$  and so  $a_1 = 0$  because  $a_1 \in i\mathbf{R}$ , at last we have:

$$\begin{cases} g(z) = \lambda(P(z) - P(\omega_1)), & \lambda \in \mathbf{C}, \\ \eta = \frac{dz}{g(z)}. \end{cases}$$

As we have (property (1) of Lemma 1 with  $\omega_3 = i\pi$ )

$$\forall z \in \mathbf{C}, \quad X(z + 2\omega_3) = X(z)$$

we deduce that  $\operatorname{Re} \int_{\omega_{1/2}}^{\omega_{1/2} + 2\omega_3} (\Phi_1, \Phi_2) = 0$ , and then

$$\int_{\omega_{1/2}}^{\omega_{1/2} + 2\omega_3} \eta = \overline{\int_{\omega_{1/2}}^{\omega_{1/2} + 2\omega_3} \eta g^2}.$$

So

$$\int_{\omega_{1/2}}^{\omega_{1/2} + 2\omega_3} \frac{dz}{P(z) - P(\omega_1)} = \lambda \overline{\int_{\omega_{1/2}}^{\omega_{1/2} + 2\omega_3} (P(z) - P(\omega_1)) dz} \quad (4)$$

as we have:

$$\frac{1}{P(z) - P(\omega_1)} = \frac{2}{P''(\omega_1)} (P(z - \omega_1) - P(\omega_1)), \quad \text{and} \quad \zeta' = -P$$

we conclude:

$$\Leftrightarrow -\frac{4}{P''(\omega_1)}(\eta_3 + \omega_3 P(\omega_1)) = -2\lambda\bar{\lambda}(\eta_3 + \omega_3 P(\omega_1)) \quad (4)$$

and then: (as  $\eta_3, \omega_3 \in i\mathbf{R}$  and  $P(\omega_1) \in \mathbf{R}$ )

$$\lambda\bar{\lambda} = -\frac{2}{P''(\omega_1)} \quad (5)$$

but as the  $P$ -function satisfies:

$$P'^2(z) = 4(P(z) - e_1)(P(z) - e_2)(P(z) - e_3), \quad e_i = P(\omega_i), \quad i = 1, 2, 3$$

with  $e_1, e_2, e_3$  real numbers,  $e_1 + e_2 + e_3 = 0$ ,  $e_3 < e_2 < e_1$ , we deduce:

$$\begin{aligned} P''(\omega_1) &= 2(3e_1^2 + e_1e_2 + e_1e_3 + e_2e_3) \\ &= 4\left(e_1 + \frac{e_3}{2}\right)(e_1 - e_3) = 2(e_1 - e_2)(e_1 - e_3) \end{aligned}$$

and then  $P''(\omega_1)$  is a positive number (as  $e_1 > e_2$  and  $e_1 > e_3$ ) so (5) cannot be satisfied, and  $z_0 \neq \omega_1$ .

The only case remaining is  $z_0 = \omega_2 = \omega_1 + \omega_3$ , as before we have:

$$\begin{cases} g(z) = \lambda(P(z) - P(\omega_2)), \\ \eta = \frac{dz}{g(z)} \end{cases}$$

and the condition (5) becomes

$$\lambda\bar{\lambda} = \frac{-2}{P''(\omega_2)}$$

but now we have

$$\begin{aligned} P''(\omega_2) &= 2(3e_2^2 + e_1e_2 + e_1e_3 + e_2e_3) \\ &= 2(e_1 + 2e_3)(2e_1 + e_3) = 2(e_3 - e_2)(e_1 - e_2) \end{aligned}$$

and  $P''(\omega_2)$  is a negative number because  $e_1 > e_2$  and  $e_3 < e_2$ .

At last as we assume that  $X$  sends the line  $\{\operatorname{Re}(z) = 0\}$  to the  $x_1$ -axis of  $\mathbf{R}^3$ ,  $g$  must have purely imaginary values on the line  $\{\operatorname{Re}(z) = 0\}$ , so  $\lambda$  must be in  $i\mathbf{R}$  and we conclude:

$$\lambda = \pm i \sqrt{\frac{-2}{P''(\omega_2)}},$$

and without loss of generality we can assume

$$\lambda = i \sqrt{\frac{-2}{P''(\omega_2)}} \in i\mathbf{R}.$$

This concludes the proof of Lemma 2.

*Remark.* A computation shows that the minimal surface given by the data  $(g, \eta)$  in Lemma 2 is invariant by the  $u$ -translation where  $u$  is given by:

$$\begin{aligned} u &= \operatorname{Re} \int_{\omega_3/2}^{\omega_3/2 + 2\omega_1} (\Phi_1, \Phi_2, \Phi_3) \\ &= \left( 0, \sqrt{\frac{-2}{P''(\omega_2)}} (2\eta_1 + e_2), 2\omega_1 \right). \end{aligned}$$

LEMMA 3. *The data  $(g, \eta)$  given in Lemma 2 is the Weierstrass representation of Riemann's example.*

*Proof.* Riemann [9] constructed for every rectangular torus an embedded minimal annulus bounded by two parallel horizontal straight lines such that every intersection with any horizontal plane is a circle. Inversely we just saw in Lemmas 1 and 2 that if such a surface exists then its Weierstrass representation is necessarily the one given in Lemma 2; this concludes the proof of lemma 3 and the proof of Theorem 1.

### Remarks

(1) In [3], Hoffman, Karcher and Rosenberg gave an explicit proof that each horizontal intersection of the surface defined by the data  $(g, \eta)$  of Lemma 2 is a circle.

(2) Darboux [1] gave explicitly the equations of Riemann's examples in terms of a parameter  $k$ ,  $0 \leq k \leq 1$  associated to a rectangular torus  $T^2$  (that is for every rectangular torus we can associate a real number  $k$ ,  $0 \leq k \leq 1$  and conversely for

each such number there is a corresponding rectangular torus) and the Jacobi function associated to  $T^2$ . For more details about these concepts see Gerretsen and Sansone [2] p. 286.

(3) In [11] Appendix C we gave the following Weierstrass representation for Riemann's examples using the equations given by Darboux, (see Remark 2 above). Following the notations of [11], we set  $2K' = \omega_1$ ,  $2iK = \omega_3$ ,  $2K' + 2iK = \omega_2$  and we use the variable  $z$  instead of  $(z - K')$  in [11]. Let  $\Gamma$  denote the lattice generated by  $(2\omega_1, 2\omega_3) = (4K', 4iK)$  and let  $P$  stand for the Weierstrass  $P$ -function on the torus  $C/\Gamma$ . Set

$$\begin{cases} g_1(z) = \frac{kk'}{4} \frac{1}{P(z) - P(\omega_2)}, \\ \eta_1 = \frac{2}{k} (P(z) - P(\omega_2)) dz = \frac{k'}{2} \frac{dz}{g(z)}. \end{cases}$$

We want to show that the data  $(g_1, \eta_1)$  defines the same surface as the data given in Lemma 2, up to a rigid motion of  $\mathbf{R}^3$ . This will give a new proof of Lemma 3.

To see this recall that (see again [11] p. 60):

$$\begin{cases} k^2 + k'^2 = 1; & h^2 = \frac{e_2 - e_3}{e_1 - e_3}; & k, k' \in [0, 1], \\ e_1 - e_3 = \frac{1}{4} \left( \text{since } K' = h\omega_1 \text{ with } h = \frac{1}{2} \right). \end{cases} \tag{*}$$

We have (since  $1/(P(z) - P(\omega_2)) = (2/P''(\omega_2))(P(z - \omega_2) - P(\omega_2))$ ):

$$g_1(z) = \frac{kk'}{2P''(\omega_2)} (P(z - \omega_2) - P(\omega_2)),$$

but again we can make a change of variable and put  $z$  instead of  $(z - \omega_2)$ , so:

$$g_1(z) = \frac{kk'}{2P''(\omega_2)} (P(z) - P(\omega_2)).$$

Using (\*) and since  $P''(\omega_2) = 2(e_3 - e_2)(e_1 - e_2)$  a simple computation shows that:

$$\frac{kk'}{2P''(\omega_2)} = -\sqrt{\frac{-2}{P''(\omega_2)}}.$$

So we have:

$$\begin{cases} g_1(z) = -\sqrt{\frac{-2}{P''(\omega_2)}}(P(z) - P(\omega_2)), \\ \eta_1 = \frac{k'}{2} \frac{dz}{g_1(z)}. \end{cases}$$

Finally the  $3\pi/2$  rotation about the  $x_3$ -axis in  $\mathbf{R}^3$  applied to the surface defined by  $(g_1, \eta_1)$  gives the following data for the rotated surface:

$$\begin{cases} g_2(z) = i\sqrt{\frac{-2}{P''(\omega_2)}}(P(z) - P(\omega_2)), \\ \eta_2 = \frac{k'}{2} \frac{dz}{g_2(z)}, \end{cases}$$

that is:

$$\begin{cases} g_2(z) = g(z), \\ \eta_2 = \frac{k'}{2} \eta, \end{cases}$$

where  $(g, \eta)$  is the data given in Lemma 2. Then  $(g_2, \eta_2)$  defines the same surface as  $(g, \eta)$  up to the  $k'/2$  homothety in  $\mathbf{R}^3$ , and so the data  $(g, \eta)$  given in Lemma 2 defines Riemann's examples.

(4) In fact in Lemmas 1 and 2 we just need  $M/S_{20}$  to be an immersed two-punctured torus in  $\mathbf{R}^3/S_{20}$  with finite total curvature and two embedded planar ends. That is we can remove the hypothesis " $M$  is properly embedded" by " $M/S_{20}$  has finite total curvature and embedded planar ends".

### §3. Proof of Theorem 2

To show that the data  $(g_k, \eta_k)$  defines an immersed minimal surface in  $\mathbf{R}^3/z$  we just need to verify the period conditions, that is if  $\gamma$  and  $\mu$  are the paths on  $T^2$  defined by:

$$\gamma(t) = \frac{\omega_1}{2} + 2t\omega_3, \quad t \in [0, 1],$$

$$\mu(t) = 2t\omega_1 + \frac{\omega_3}{2}, \quad t \in [0, 1],$$

it is enough to show that:

$$\operatorname{Re} \int_{\gamma} (\Phi_1, \Phi_2, \Phi_3) = (0, 0, 0), \tag{1}$$

where:

$$\Phi_1 = \frac{\eta_k}{2} (1 - g_k^2), \quad \Phi_2 = i \frac{\eta_k}{2} (1 + g_k^2), \quad \Phi_3 = \eta_k g_k.$$

This claim holds since the forms  $\Phi_i, i = 1, 2, 3$  have no residue (because the  $P$ -function of Weierstrass is an even function). Let us remark that if the conditions (1) are satisfied then the forms  $(\Phi_1, \Phi_2, \Phi_3)$  must have periods on the path  $\mu$ , that is:

$$\operatorname{Re} \int_{\mu} (\Phi_1, \Phi_2, \Phi_3) \neq (0, 0, 0),$$

otherwise the data  $(g_k, \eta_k)$  defines an immersed minimal surface with finite total curvature and two parallel flat ends in  $\mathbf{R}^3$ , but the ‘‘Half space theorem’’ of Hoffman and Meeks [4] shows this situation is impossible.

Let us assume for a while that (1) is satisfied. Let  $I$  be the map on  $T^2$  defined by:

$$\forall z \in T^2, \quad I(z) = -\bar{z}.$$

Let us remark that, as  $P''(\omega_2)$  is a negative real number (see the proof of Lemma 2 in §2),  $\lambda_k \in i\mathbf{R}$ . We deduce that:

$$\begin{aligned} \forall z \in T^2, \quad g_k(I(z)) &= \lambda_k [P(-\bar{z}) - P(\omega_2)]^{2k+1} \\ &= \overline{-g_k(z)}. \end{aligned}$$

Then:

$$I_*(\eta_k) = -\frac{d\bar{z}}{g_k(I(z))} = \frac{d\bar{z}}{g_k(z)} = \bar{\eta}_k,$$

so that:

$$I_*(\Phi_1) = \bar{\Phi}_1; \quad I_*(\Phi_2) = -\bar{\Phi}_2; \quad I_*(\Phi_3) = -\bar{\Phi}_3.$$

Calling  $X_k$  the minimal immersion defined, up to a translation in  $\mathbf{R}^3$ , by  $(g_k, \eta_k)$ , we deduce that:

$$\forall z \in T^2 - \{0, \omega_2\}, \quad X_k(I(z)) = S_1[X_k(z)],$$

where  $S_1$  is the reflection about the  $x_1$ -axis. This shows that  $X_k$  sends the line  $C_1$  defined by:

$$C_1(t) = 2t\omega_3, \quad t \in ]0, 1[$$

to  $D_1$  which is the  $x_1$ -axis in the horizontal plane  $P_1 = \{x_3 = 0\}$ .

In the same way if  $J$  is the map on  $T^2$  defined by:

$$\forall z \in T^2, \quad J(z) = 2\omega_1 - \bar{z},$$

we can show that:

$$\forall z \in T^2 - \{0, \omega_2\}, \quad X_k(J(z)) = S_2[X_k(z)],$$

where  $S_2$  is the reflection about  $D_2$ , a straight line parallel to  $D_1$  in a horizontal plane  $P_2$ ,  $P_2$  distinct from  $P_1$ . Then  $X_k$  sends the line  $C_2$  of  $T^2$  on  $D_2$  in  $\mathbf{R}^3$  where

$$C_2(t) = \omega_1 + 2t\omega_3, \quad t \in [0, 1], \quad t \neq \frac{1}{2}.$$

Let


$$A = \{z \in T^2 / 0 \leq \operatorname{Re}(z) \leq \omega_1, z \neq 0, z \neq \omega_2\}.$$

As  $\Phi_3 = 1$  and hence:

$$(X_k)_3(z) = \operatorname{Re}(z), \quad \forall z \in T^2 - \{0, \omega_2\},$$

we see that  $X_k$  sends  $A \cap T^2$  to the slab of  $\mathbf{R}^3$  bounded by  $P_1$  and  $P_2$ .

Of course  $X_k(A)$  is a minimal immersed annulus in  $\mathbf{R}^3$  between  $P_1$  and  $P_2$  and bounded by  $D_1 \cup D_2$ .



Finally denoting  $(X_k)_i$  by  $X_i$ ,  $i = 1, 2, 3$  for simplicity, we have:

$$\forall z \in T^2 - \{0, \omega_2\}, \quad (X_1 - iX_2)(z) = \int_{z_0}^z \eta_k - \int_{z_0}^z \overline{\eta_k g_k^2},$$

therefore, for  $z$  near 0 we have

$$\begin{aligned} (X_1 - iX_2)(z) &\simeq \int_{z_0}^z \lambda_k \cdot \frac{dz}{z^{2(2k+1)}} \\ &\simeq -\frac{\overline{\lambda_k}}{4k+1} \left( \frac{1}{z^{4k+1}} \right) \end{aligned}$$

and so the projection of the end near 0 on a horizontal plane is a  $(4k + 1)$  to 1 map.

Near the other end  $\omega_2$  we also have:

$$\begin{aligned} (X_1 - iX_2)(z) &\simeq \int_{z_0}^z \left( \frac{2}{p''(\omega_2)} \right)^{2k+1} \frac{1}{\lambda_k} \cdot \frac{dz}{(z - \omega_2)^{2(2k+1)}} \\ &\simeq \frac{\lambda_k}{3k+1} \cdot \frac{1}{(z - \omega_2)^{4k+1}}, \end{aligned}$$

and again the projection of the end  $\omega_2$  on a horizontal plane is a  $(4k + 1)$  to 1 map.

So it remains to show that conditions (1) hold. As we have  $\Phi_3 = 1$  we deduce:

$$\begin{aligned} &\Leftrightarrow \operatorname{Re} \int_\gamma \Phi_1 = 0 \quad \text{and} \quad \operatorname{Re} \int_\gamma \Phi_2 = 0 \\ &\Leftrightarrow \int_\gamma \eta_k = \int_\gamma \overline{\eta_k g_k^2}. \end{aligned} \tag{1}$$

As  $\eta_k = dz/g_k(z)$  we have:

$$\begin{aligned} \Leftrightarrow \int_{\gamma} \frac{dz}{g_k(z)} &= \int_{\gamma} \overline{g_k(z)} dz \\ \Leftrightarrow \int_{\gamma} \frac{dz}{(P(z) - P(\omega_2))^{2k+1}} &= \lambda_k \bar{\lambda}_k \int_{\gamma} \overline{(P(z) - P(\omega_2))^{2k+1}} dz, \end{aligned} \tag{1}$$

but:

$$\frac{1}{P(z) - P(\omega_2)} = \frac{2}{P''(\omega_2)} (P(z - \omega_2) - P(\omega_2)).$$

So:

$$\begin{aligned} \Leftrightarrow \lambda_k \bar{\lambda}_k \int_{\gamma} \overline{(P(z) - P(\omega_2))^{2k+1}} dz &= \left(\frac{2}{P''(\omega_2)}\right)^{2k+1} \int_{\gamma} (P(z - \omega_2) - P(\omega_2))^{2k+1} dz \\ \Leftrightarrow \lambda_k \bar{\lambda}_k \sum_{q=0}^{2k+1} C_{2k+1}^q (-P(\omega_2))^{2k+1-q} \int_{\gamma} \overline{P(z)^q} dz & \\ = \left(\frac{2}{P''(\omega_2)}\right)^{2k+1} \sum_{q=0}^{2k+1} C_{2k+1}^q (-P(\omega_2))^{2k+1-q} \cdot \int_{\gamma} P(z - \omega_2)^q dz. & \end{aligned} \tag{1}$$

The following lemma shows the last equality is true.

LEMMA 1. *For every positive integer q we have:*

$$\int_{\gamma} P(z)^q dz = - \int_{\gamma} P^q(z - \omega_2) dz.$$

Assuming Lemma 1 we have:

$$\Leftrightarrow \lambda_k \bar{\lambda}_k = - \left(\frac{2}{P''(\omega_2)}\right)^{2k+1}$$

which is true as  $\lambda_k = i\sqrt{(-2/P''(\omega_2))^{2k+1}}$ . So it just remains to prove Lemma 1.

*Proof of Lemma 1.* As  $T^2$  is a rectangular torus we have the following Laurent series for the Weierstrass  $P$ -function, see [2], [8] or the Appendix.

$$P(z) = \frac{1}{z^2} + \sum_{n=1}^{\infty} a_n z^n$$

with  $a_n$  real numbers.

We deduce that for any positive integer  $q$  the Laurent series of  $P^q(z)$  has the following type:

$$P^q(z) = \sum_{n=1}^q b_{-n} z^{-2n} + \sum_{n=0}^{+\infty} b_n z^n,$$

with  $b_j$  real numbers,  $j = 1, \dots, n$ .

Also if  $P^{(k)}(z)$  is the  $k$ th-derivative of  $P$  we have:

$$P^{(k)}(z) = (-1)^k \frac{(k+1)!}{z^{k+2}} + F_k(z),$$

where  $F_k$  is a holomorphic map near 0. We deduce that:

$$P^q(z) - \sum_{n=1}^q \frac{b_{-n}}{(2n-1)!} P^{(2n-2)}(z)$$

is an elliptic function on  $T^2$  without poles, so this function is constant and taking  $z = \omega_1$ , we see that this constant  $c$  must be real, so:

$$P^q(z) = \sum_{n=1}^q \frac{b_{-n}}{(2n-1)!} P^{(2n-2)}(z) + c$$

with  $b_{-n}$  and  $c$  real numbers. Then:

$$\begin{aligned} \int_{\gamma} P^q(z) dz &= \sum_{n=1}^q \frac{b_{-n}}{(2n-1)!} \int_{\omega_1/2}^{\omega_1/2 + 2\omega_3} P^{(2n-2)}(z) dz + c \int_{\omega_1/2}^{\omega_1/2 + 2\omega_3} dz \\ &= b_{-1} \int_{\omega_1/2}^{\omega_1/2 + 2\omega_3} P(z) dz + 2\omega_3 c \end{aligned}$$

since for  $k \geq 2$ :

$$\int_{\omega_1/2}^{\omega_1/2 + 2\omega_3} P^{(k)}(z) dz = P^{(k-1)}\left(\frac{\omega_1}{2} + 2\omega_3\right) - P^{(k-1)}\left(\frac{\omega_1}{2}\right) = 0.$$

Then, as  $\zeta'(z) = -P(z)$  we have:

$$\begin{aligned} \int_{\gamma} P^q(z) dz &= b_{-1} \left[ -\zeta\left(\frac{\omega_1}{2} + 2\omega_3\right) + \zeta\left(\frac{\omega_1}{2}\right) \right] + 2\omega_3 c \\ &= 2\eta_3 b_{-1} + 2\omega_3 c. \end{aligned}$$

In the same way, as:

$$P^q(z - \omega_2) = \sum_{n=1}^q \frac{b_{-n}}{(2n-1)!} P^{(2n-2)}(z - \omega_2) + c,$$

we also have:

$$\int_{\gamma} P^q(z - \omega_2) dz = -2\eta_3 b_{-1} + 2\omega_3 c.$$

This concludes the proof because  $c$  and  $b_{-1}$  are real numbers and  $\omega_3$  and  $\eta_3$  are purely imaginary numbers.

#### *Remarks*

(1) Following the arguments of Lemmas 1 and 2 in §2, it is easy to show that the surfaces  $S_k$  are the only minimal immersed surfaces between  $P_1$  and  $P_2$  bounded by  $D_1 \cup D_2$  with finite total curvature.

(2) We do not know if there exists surfaces like  $S_k$  which are bounded by two horizontal lines  $D_1, D_2$  and make a non zero angle  $\theta$ . Of course for  $k = 0$ , Theorem 1 shows that such a surface does not exist.

#### **§4. Appendix**

Let  $T^2$  be a torus  $\mathbf{C}/\Gamma$  where  $\Gamma$  is the lattice of  $\mathbf{C}$  given by:

$$\Gamma = \{p \cdot 2\omega_1 + q \cdot 2\omega_3, p, q \in \mathbf{Z}, \omega_1 \in \mathbf{R}_*^+, \operatorname{Im}(\omega_3) \in \mathbf{R}_*^+\}.$$

The Weierstrass  $P$ -function is a special function defined on  $T^2$  which is a meromorphic function on  $\mathbf{C}$  such that:

$P$  has a pole of order two at each point of  $\Gamma$  and:

$$\forall z \in \mathbf{C} - \Gamma, \quad P(z + 2\omega_i) = P(z), \quad i = 1, 3.$$

$$\text{If } \omega_2 = \omega_1 + \omega_3, \quad P(\omega_1) + P(\omega_2) + P(\omega_3) = 0.$$

$P$  has the following Laurent series:

$$P(z) = \frac{1}{z^2} + \sum_{\omega \in \Gamma - \{0\}} \left\{ \frac{1}{(z - \omega)^2} - \frac{1}{\omega^2} \right\}.$$

$$P \text{ is an even function: } \forall z \in \mathbf{C} - \Gamma, \quad P(-z) = P(z).$$

There are two other functions related to  $P$ , namely the  $\zeta$  and  $\sigma$ -functions: The  $\zeta$ -function satisfies:

$$\forall z \in \mathbf{C} - \Gamma, \quad \zeta'(z) = -P(z).$$

If  $\eta_i = \zeta(\omega_i)$ ,  $i = 1, 2, 3$ , then  $\eta_2 = \eta_1 + \eta_3$ , so  $\zeta$  is uniquely defined on  $\mathbf{C} - \Gamma$ . We have:

$$\forall z \in \mathbf{C} - \Gamma, \quad \zeta(z + 2\omega_i) = \zeta(z) + 2\eta_i, \quad i = 1, 2, 3,$$

and then  $\zeta$  is not defined on  $T^2$  because  $z$  and  $(z + 2\omega_i)$  represent the same point on  $T^2$ .

$$\zeta \text{ is an odd function: } \forall z \in \mathbf{C} - \Gamma, \quad \zeta(-z) = -\zeta(z).$$

The following Legendre relation holds.

$$\omega_2 \eta_1 - \omega_1 \eta_2 = i \frac{\pi}{2}.$$

The  $\sigma$ -function satisfies:

$$\forall z \in \mathbf{C} - \Gamma, \quad \frac{\sigma'}{\sigma}(z) = \zeta(z),$$

and then  $\sigma$  is an holomorphic function on  $\mathbf{C}$ , furthermore:

$$\lim_{z \rightarrow 0} \frac{\sigma(z)}{z} = 1,$$

so that  $\sigma$  is uniquely determined on  $\mathbf{C}$ .

We also have:

$$\forall z \in \mathbf{C}, \quad \sigma(z + 2\omega_i) = -e^{2\eta_i(z + \omega_i)} \sigma(z). \quad (\#)$$

$\sigma$  is an odd function:  $\forall z \in \mathbf{C}, \sigma(-z) = -\sigma(z)$ .

Furthermore if  $T^2$  is a rectangular torus, that is if  $\omega_3 \in i\mathbf{R}_*^+$ , we have:

$$\forall z \in \mathbf{C} - \Gamma, \quad P(\bar{z}) = \overline{P(z)}; \quad \zeta(\bar{z}) = \overline{\zeta(z)}.$$

$$\forall z \in \mathbf{C}, \quad \sigma(\bar{z}) = \overline{\sigma(z)}.$$

For more details about those functions see Gerretsen–Sansone [2] or Molk–Tannery [8].

#### REFERENCES

- [1] G. DARBOUX. *Théorie des surfaces*. Vol I, 2nd édition, Paris, 1914, pp. 461–644, §252.
- [2] J. GERRETSEN, G. SANSONE. *Lectures on the theory of functions of a complex variable*. P. Noordhoff–Groningen, 1960.
- [3] D. HOFFMAN, H. KARCHER, H. ROSENBERG. *A characterization of Riemann examples*. Preprint.
- [4] D. HOFFMAN, W. H. MEEKS. *The global theory of properly embedded minimal surfaces, Theorem 10.1*. Preprint, University of Massachusetts at Amherst.
- [5] H. B. LAWSON. *Lectures on minimal submanifolds*. Vol I. Publish or Perish.
- [6] W. H. MEEKS, H. ROSENBERG. *The geometry of periodic minimal surfaces*. Preprint.
- [7] W. H. MEEKS, B. WHITE. *Minimal surfaces bounded by convex curves in parallel planes*. Preprint.
- [8] J. MOLK, J. TANNERY. *Éléments de la théorie des fonctions elliptiques*. Bronx, N.Y., Chelsea, 1972.
- [9] B. RIEMANN. *Oeuvres complètes*. Tome XIII des mémoires de la société royale de Goettingen (1867), p. 305.
- [10] M. SCHIFFMAN. *On surfaces of stationary area bounded by two circles, or convex curves, in parallel planes*. *Annals of Math* 63 (1956), 77–90.
- [11] E. TOUBIANA. *Géométrie des surfaces minimales de  $\mathbf{R}^3$* . Thèse de doctorat, Université Paris VII, 1986.

203 rue du Faubourg  
Saint-Antoine  
75011 Paris

Received April 15, 1991