

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 67 (1992)

Artikel: On the semicontinuity of curvatures.
Autor: Lutwak, Erwin
DOI: <https://doi.org/10.5169/seals-51116>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 03.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the semicontinuity of curvatures

ERWIN LUTWAK*

Dedicated to Prof. Dr. Kurt Leichtweiß on the occasion of his Sixty-fifth Birthday

In this article, a convex hypersurface is the boundary of a convex body (compact, convex subset with non-empty interior) of Euclidean n -space, \mathbf{R}^n . Let \mathcal{C}^n denote the space of convex hypersurfaces, endowed with the topology induced by the Hausdorff metric (see, for example, Busemann [1] or Leichtweiß [4]). Let \mathcal{C}_2^n denote the subset of \mathcal{C}^n consisting of the regular C^2 -hypersurfaces with everywhere positive principal curvatures. For $Q \in \mathcal{C}_2^n$, and $x \in Q$, let $K(Q, x)$ denote the Gauss curvature of Q at the point x , and let

$$K^+(Q) = \max_{x \in Q} K(Q, x) \quad \text{and} \quad K^-(Q) = \min_{x \in Q} K(Q, x).$$

A special case of the main result of this article is that,

$$K^+ : \mathcal{C}_2^n \rightarrow (0, \infty) \quad \text{is lower semicontinuous,}$$

while

$$K^- : \mathcal{C}_2^n \rightarrow (0, \infty) \quad \text{is upper semicontinuous.}$$

Let $r_1(x), \dots, r_{n-1}(x)$ denote the principal radii of curvature at the point $x \in Q$. The normalized j -th elementary symmetric function of the principal radii of curvature at x will be denoted by $s_j(x)$; i.e.,

$$\binom{n-1}{j} s_j(x) = \sum_{1 \leq i_1 < \dots < i_j \leq n-1} r_{i_1}(x) \cdots r_{i_j}(x).$$

* Research supported, in part, by NSF Grants DMS–8902550 and DMS–9123571.

For $Q \in \mathcal{C}_2^n$, define

$$C_j^+(Q) = \max_{x \in Q} s_j(x) \quad \text{and} \quad C_j^-(Q) = \min_{x \in Q} s_j(x).$$

Thus, $1/C_{n-1}^+(Q) = K^-(Q)$, and $1/C_{n-1}^-(Q) = K^+(Q)$. It will be shown that the functionals C_j^+ and C_j^- may be extended to all of \mathcal{C}^n in such a manner that

$$C_j^+ : \mathcal{C}^n \rightarrow [0, \infty] \quad \text{is lower semicontinuous,}$$

and

$$C_j^- : \mathcal{C}^n \rightarrow [0, \infty] \quad \text{is upper semicontinuous.}$$

More general results will be established along the way. One of these results will establish a conjecture of Wm. J. Firey (see [2, p. 257]) regarding the upper semicontinuity of a certain functional for convex hypersurfaces.

Some preliminary notation and results will be helpful. Let $C^+(S^{n-1})$ denote the set of positive continuous functions on the unit sphere S^{n-1} . For $f \in C^+(S^{n-1})$, and $\alpha \in \mathbf{R}$, such that $\alpha \neq 0$, define

$$|f|_\alpha = \left\{ \frac{1}{\kappa_n} \int_{S^{n-1}} f(u)^\alpha dS(u) \right\}^{1/\alpha}$$

where the integration is with respect to spherical Lebesgue measure, S , on S^{n-1} , and $\kappa_n = S(S^{n-1})$ denotes the surface area of S^{n-1} . From the Hölder inequality it follows that,

$$|f|_{\alpha_1} \leq |f|_{\alpha_2} \quad \text{whenever } \alpha_1 \leq \alpha_2. \tag{1}$$

For $f \in C^+(S^{n-1})$, and μ a Borel measure on S^{n-1} , let

$$\langle f, \mu \rangle = \frac{1}{\kappa_n} \int_{S^{n-1}} f(u) d\mu(u).$$

For $Q \in \mathcal{C}_2^n$ and $u \in S^{n-1}$, write $(Q, u) \in Q$ for the point of Q at which the outer unit normal is u . Let $r_1(Q, u), \dots, r_{n-1}(Q, u)$ denote the principal radii of curvature at (Q, u) , and let $s_j(Q, u)$ denote the normalized j -th elementary symmetric function of the principal radii of curvature at (Q, u) . It will be convenient to view $s_j(Q, \cdot)$ as a function on S^{n-1} .

Suppose $Q \in \mathcal{C}_2^n$. For $j = 1, \dots, n-1$, and $\alpha \in \mathbf{R}$, such that $\alpha \neq 0$, define

$$\psi_j^\alpha(Q) = |s_j(Q, \cdot)|_\alpha.$$

When $\alpha = 1$ the superscript will be suppressed. The $\psi_j(Q)$ are just the classical integrals of mean curvature of Q . For $\alpha = -\infty, 0$, or ∞ , define

$$\psi_j^\alpha(Q) = \lim_{r \rightarrow \alpha} \psi_j^r(Q).$$

Thus,

$$\psi_j^\infty(Q) = \max_{u \in S^{n-1}} s_j(Q, u) = C_j^+(Q),$$

while

$$\psi_j^{-\infty}(Q) = \min_{u \in S^{n-1}} s_j(Q, u) = C_j^-(Q).$$

For arbitrary convex hypersurfaces there are well-known extensions of the (indefinite integrals of the) elementary symmetric functions of the principal radii of curvature. Specifically, for each $Q \in \mathcal{C}_2^n$, and each j , there exists a Borel measure $S_j(Q, \cdot)$ on S^{n-1} , such that if $Q \in \mathcal{C}_2^n$, then $S_j(Q, \cdot)$ is absolutely continuous with respect to spherical Lebesgue measure, and

$$\frac{dS_j(Q, \cdot)}{dS} = s_j(Q, \cdot), \quad (2)$$

where the derivative is a Radon–Nikodym derivative.

As will be seen, the functional ψ_j^α , on \mathcal{C}_2^n , can be extended to a functional Ψ_j^α , defined on all of \mathcal{C}^n . For $\alpha = 1$, define

$$\Psi_j(Q) = \Psi_j^1(Q) = \frac{1}{\kappa_n} \int_{S^{n-1}} dS_j(Q, u).$$

Thus, $\kappa_n \Psi_{n-1}(Q)$ is the surface area of Q , and the $\Psi_j(Q)$ are essentially the classical Quermassintegrals Q (see, for example, Leichtweiß [4]).

For $\alpha < 1$, and $\alpha \neq 0$, define $\Psi_j^\alpha(Q)$ by

$$\Psi_j^\alpha(Q) = \inf \{ \langle g, S_j(Q, \cdot) \rangle / |g|_{\alpha/(\alpha-1)} : g \in C^+(S^{n-1}) \}, \quad (3a)$$

and for $\alpha > 1$, define $\Psi_j^\alpha(Q)$ by

$$\Psi_j^\alpha(Q) = \sup \{ \langle g, S_j(Q, \cdot) \rangle / |g|_{\alpha/(\alpha-1)} : g \in C^+(S^{n-1}) \}. \tag{3b}$$

For $Q \in \mathcal{C}^n$, and all j ,

$$\Psi_j^{\alpha_1}(Q) \leq \Psi_j^{\alpha_2}(Q), \quad \text{whenever } \alpha_1 \leq \alpha_2. \tag{4}$$

To see this, note that if $\alpha_1 \leq 1 \leq \alpha_2$, then (4) follows from taking g to be a constant function in (3a) and (3b). If $\alpha_1 < \alpha_2 < 1$, or $1 < \alpha_1 < \alpha_2$, then (4) follows directly from (1) and definitions (3a) and (3b).

For $\alpha = -\infty$, or ∞ , define Ψ_j^α on \mathcal{C}^n by:

$$\Psi_j^\alpha(Q) = \lim_{r \rightarrow \alpha} \Psi_j^r(Q),$$

and define

$$\Psi_j^0(Q) = \lim_{r \rightarrow 0^+} \Psi_j^r(Q).$$

Thus (4) shows that

$$\Psi_j^\infty(Q) = \sup_{1 < r < \infty} \Psi_j^r(Q), \tag{5a}$$

$$\Psi_j^{-\infty}(Q) = \inf_{-\infty < r < 0} \Psi_j^r(Q), \tag{5b}$$

and

$$\Psi_j^0(Q) = \inf_{0 < r < 1} \Psi_j^r(Q). \tag{5c}$$

PROPOSITION. For $j = 1, \dots, n - 1$, and $-\infty \leq \alpha \leq \infty$,

$$\Psi_j^\alpha(Q) = \psi_j^\alpha(Q), \quad \text{whenever } Q \in \mathcal{C}_2^n. \tag{6}$$

To see this first note that from (2) it follows that for $Q \in \mathcal{C}_2^n$ and $g \in C^+(S^{n-1})$,

$$\langle g, S_j(Q, \cdot) \rangle = \frac{1}{\kappa_n} \int_{S^{n-1}} g(u) s_j(Q, u) dS(u). \tag{7}$$

The case $\alpha = 1$ of (6) follows by taking $g = 1$ in (7). Suppose that $-\infty < \alpha < 1$, and $\alpha \neq 0$. From (7), and the Hölder inequality [3, p. 140], it follows that

$$\psi_j^\alpha(Q) = |s_j(Q, \cdot)|_\alpha \leq \langle g, S_j(Q, \cdot) \rangle / |g|_{\alpha/(\alpha-1)},$$

with equality if and only if the function $g/s_j(Q, \cdot)^{\alpha-1}$ is a constant function on S^{n-1} . This proves (6) when $-\infty < \alpha < 1$, and $\alpha \neq 0$. The case $1 < \alpha < \infty$ of (6) is established in exactly the same way. The cases of (6) where $\alpha = -\infty, 0$, or ∞ now follow since in these cases both $\Psi_j^\alpha(Q)$ and $\psi_j^\alpha(Q)$ were defined as limits of $\Psi_j^r(Q) = \psi_j^r(Q)$, for real $r \neq 0$.

THEOREM. *For all $j = 1, \dots, n-1$, the functional*

$$\Psi_j^\alpha : \mathcal{C}^n \rightarrow [0, \infty], \quad \text{is upper semicontinuous when } -\infty \leq \alpha \leq 1,$$

while the functional

$$\Psi_j^\alpha : \mathcal{C}^n \rightarrow [0, \infty], \quad \text{is lower semicontinuous when } 1 \leq \alpha \leq \infty.$$

Proof. From the weak continuity of the measures S_j (see for example, Schneider [8]), it follows that for fixed $g \in C^+(S^{n-1})$, the function $\Gamma_g : \mathcal{C}^n \rightarrow (0, \infty)$, defined for $Q \in \mathcal{C}^n$ by

$$\Gamma_g(Q) = \langle g, S_j(Q, \cdot) \rangle / |g|_{\alpha/(\alpha-1)},$$

is continuous. For $1 < \alpha < \infty$, the functional Ψ_j^α is lower semicontinuous, on \mathcal{C}^n , since it is just the supremum of the continuous (on \mathcal{C}^n) functionals Γ_g . The case $\alpha = \infty$ now follows since by (5a), the functional Ψ_j^∞ is just a supremum (over all $r > 1$) of the lower semicontinuous functionals Ψ_j^r . The cases where $-\infty \leq \alpha \leq 1$ follow in exactly the same manner.

For $\alpha = n/(n+1)$, the upper semicontinuity of $\psi_{n-1}^\alpha : \mathcal{C}_2^n \rightarrow (0, \infty)$ was established in [7]. The case of the Theorem where $n = 3$, $\alpha = -1$, and $j = n-1$, was conjectured by Wm. J. Firey at the 1974 Oberwolfach meeting on convex bodies (see [2, p. 257]). The case where $\alpha = n/(n+1)$ and $j = n-1$ is due to Leichtweiß [5] (see also Leichtweiß [6]).

REFERENCES

1. H. BUSEMANN, *Convex Surfaces*, Interscience, New York, 1958.
2. P. M. GRUBER and R. SCHNEIDER, *Problems in geometric convexity*, Contributions to Geometry (J. Tölke and J. M. Wills, eds.), Birkhäuser, Basel, 1979, pp. 255–278.

3. G. HARDY, J. LITTLEWOOD, and G. PÓLYA, *Inequalities*, Cambridge Univ. Press, Cambridge, 1934.
4. K. LEICHTWEIß, *Konvexe Mengen*, Springer, Berlin, 1980.
5. K. LEICHTWEIß, *Bemerkungen zur Definition einer erweiterten Affinoberfläche von E. Lutwak*, *Manuscripta Math.* 65 (1989), 181–197.
6. K. LEICHTWEIß, *On the history of the affine surface area for convex bodies*, *Results in Math.* 20 (1991), 650–656.
7. E. LUTWAK, *Extended affine surface area*, *Adv. in Math.* 85 (1991), 39–68.
8. R. SCHNEIDER, *Boundary structure and curvature of convex bodies*, *Contributions to Geometry* (J. Tölke and J. M. Wills, eds.), Birkhäuser Verlag, Basel, 1979, pp. 13–59.

Department of Mathematics
Polytechnic University
Brooklyn, NY 11201

Received November 27, 1991