Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 68 (1993)

Erratum: Erratum: Complexity and rank double cones and tensor products decompositions.

Autor: Panyushev, D.I.

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Comment. Math. Helvetici **68** (1993) 673-674 0010-2571/93/040673-02$1.50 + 0.20/0
© 1993 Birkhäuser Verlag, Basel

Erratum: D.I. Panyushev, *Complexity and rank of double cones and tensor product decompositions,* Comment. Math. Helvetici *68* (1993), 455–468.

In preparing the above-mentioned paper for printing, the diagrams on pages 461 and 463 were erroneously omitted. The correct versions read as follows:

Page 461

(1.8) *Remarks.* 1. It may happen that $\Gamma(Z) \neq \mathbf{Z}\Gamma(Z) \cap \mathscr{X}(T)_+$, i.e. the canonical embedding $S_{\lambda\mu} \subset G$ does not determine $\Gamma(Z)$ completely.

2. The subgroups $S_{\lambda\mu}$ and $S_{\lambda\mu^*}$ are isomorphic and conjugated in G by corollary 1, nevertheless, they may have *different* canonical embeddings. For example, let $G = \mathrm{E}_6$ and $\mu = \lambda = \varphi_1$, $\mu^* = \varphi_5$ are the fundamental weights with numeration as in [10]. Then $S_{\lambda\mu} \cong S_{\lambda\mu^*} \cong \mathrm{A}_3$, but their canonical embeddings are described by the following pictures:

$S_{\lambda\mu}$:  $S_{\lambda\mu^*}$: 

(The black vertices indicate the simple roots of $S_{\lambda\mu}$ and $S_{\lambda\mu^*}$.)

(1.9) The previous results show that in order to compute the complexity and the rank semigroup of a double cone one has to find the canonical embedding of $S_{\lambda\mu}$ and $\tilde{S}_{\lambda\mu}$. The next result explains how this can be done.

Page 463

(2.2) EXAMPLE. $G = \mathrm{E}_8$, $Z = Z(1, 1)$.

Here $(L_1', m_1) = (\mathrm{E}_7, 2\varphi_1 + 2)$. According to [1] we have Lie $S_{11} = \mathrm{D}_4$. Therefore $r = 4$, $2c + r = 12$ and $c = 4$. In particular, we find out that $\dim k[Z]^U = 8$.

Next, $(L_1, m_1) = (\mathrm{E}_7 \times k^*, \varphi_1 \otimes \varepsilon + \varphi_1 \otimes \varepsilon^{-1} + \varepsilon^2 + \varepsilon^{-2})$. Therefore Lie $\tilde{S}_{11} =$ Lie S_{11} and $\tilde{c} = c - 2 = 2$. According to the corollary 3(ii) the canonical embedding $S_{11} \subset E_8$ implies the embedding of the Dynkin diagrams. Here it can be done in a unique way. Hence, the canonical embedding is described by the following picture:

That is, $\alpha_4, \alpha_5, \alpha_6, \alpha_8$ is the system of simple roots of S_{11} and according to Corollary 3 $\Gamma(Z(1, 1))$ is contained in $M = \langle \alpha_4, \alpha_5, \alpha_6, \alpha_8 \rangle^{\perp} = \langle \tilde{\varphi}_1, \tilde{\varphi}_2, \tilde{\varphi}_3, \tilde{\varphi}_7 \rangle$. This means, that for any n, m the tensor product decomposition $\tilde{\varphi}_1^n \otimes \tilde{\varphi}_1^m$ contains highest weights only from M.

ul. akad. Anokhina
d.30, kor. 1, kv. 7
117602 Moscow, Russia