

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 70 (1995)

Artikel: Local constant of Ind...1.
Autor: Saito, Takeshi
DOI: <https://doi.org/10.5169/seals-53010>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Local constant of $Ind_K^L 1$

TAKESHI SAITO

Abstract. Henniart has computed the local constant $\varepsilon(Ind_K^L 1)$ for an extension L over K of local fields of odd degree in [H]. In this paper, we show that his formula is a consequence of results of Serre [S4] and of Deligne [D2]. Further we compute the local constant for an extension of even degree, assuming the residual characteristic is not equal to 2.

1. Local constant of $Ind_K^L 1$

Let K be a local field. Namely K is a complete discrete valuation field with finite residue field F of order q . We assume that the characteristic of F is not equal to 2. For a separable extension L of degree n over K , let $V = V_{L/K} = Ind_K^L 1$ be the induced representation of the absolute Galois group $G_K = \text{Gal}(K^{sep}/K)$ from the unit representation 1 of G_L . Let $d = d_{L/K} \in K^\times/K^{\times 2}$ be the discriminant of L over K and $\delta = \delta_{L/K} = \det V_{L/K}$ be the character $G_K \rightarrow \mathbb{Z}/2$ corresponding to $d_{L/K} \in K^\times/K^{\times 2} = H^1(K, \mathbb{Z}/2)$. We consider the local constant ([D1])

$$\varepsilon(V^0) = \begin{cases} \varepsilon(V, \psi, \mu) \cdot \varepsilon(\delta, \psi, \mu)^{-n} & n \text{ odd} \\ \varepsilon(V, \psi, \mu) \cdot \varepsilon(1, \psi, \mu)^{-(n-1)} \cdot \varepsilon(\delta, \psi, \mu)^{-1} & n \text{ even} \end{cases}$$

for the virtual representation

$$V^0 = \begin{cases} V - n\delta & n \text{ odd} \\ V - ((n-1)1 + \delta) & n \text{ even.} \end{cases}$$

Since $\dim V^0 = 0$ and $\det V^0 = 1$, the local constant $\varepsilon(V^0)$ is independent of an additive character ψ or a Haar measure μ and we drop them in the notation. Since V is an orthogonal representation, the Artin conductor

$$\alpha(V^0) = \begin{cases} a(V) - na(\delta) & n \text{ odd} \\ a(V) - ((n-1)a(1) + a(\delta)) & n \text{ even} \end{cases}$$

is an even integer by [S3] Théorème 1. We put

$$\bar{\epsilon}(V^0) = \epsilon(V^0)q^{-(a(V^0)/2)}.$$

It is known to be ± 1 . As usual $(\cdot_F) : F^\times/F^{\times 2} \rightarrow \{\pm 1\}$ denotes the Legendre symbol and $(\cdot, \cdot)_K : K^\times/K^{\times 2} \times K^\times/K^{\times 2} \rightarrow \{\pm 1\}$ denotes the Hilbert symbol.

THEOREM. *Let L be a separable extension of degree n of a local field K with residual characteristic $\neq 2$.*

I. ([H] Proposition 2) *If n is odd, for the virtual representation $V_{L/K}^0 = V_{L/K} - n \delta_{L/K}$, we have*

$$\bar{\epsilon}(V_{L/K}^0) = (d_{L/K}, 2)_K.$$

II. *Assume n is even. Let $V_{L/K}^0$ be the virtual representation $V_{L/K} - ((n-1)1 + \delta_{L/K})$. Let $D_{L/K}$ be the different of L over K and put $D = \text{ord}_L D_{L/K}$.*

1. *If D is even, we have*

$$\bar{\epsilon}(V_{L/K}^0) = 1.$$

2. *Assume D is odd. Let e be the ramification index, $f = n/e$ be the residual degree and π_K be an arbitrary prime element of K .*

2A. *If e is odd, then f is even and we have*

$$\bar{\epsilon}(V_{L/K}^0) = \left(\frac{-1}{F} \right)^{\binom{n}{2}} \times (d_{L/K}, \pi_K)_K.$$

2B. *Assume e is even. Let K_1 be the maximum unramified extension of K in L and E be the residue field of L . Let π_L be a prime element of L and g be the minimal polynomial of π_L over K_1 . The class $\alpha = g'(\pi_L)/\pi_L^D \in E^\times/E^{\times 2}$ is independent of π_L and we put $d' = N_{E/F}(\alpha) \cdot d_{E/F} \in F^\times/F^{\times 2}$. Then*

$$\bar{\epsilon}(V_{L/K}^0) = \left(\frac{(-1)^{\binom{f}{2}} d'}{F} \right) \times \begin{cases} \left(\frac{-1}{F} \right)^{e/2-1} \cdot (d_{L/K}, 2)_K & f \text{ odd} \\ (d_{L/K}, \pi_K)_K & f \text{ even.} \end{cases}$$

Before starting the proof we briefly recall [D2] and [S4]. For a continuous orthogonal representation V of the absolute Galois group G_K of a field K , its Stiefel–Whitney class $w_i(V) \in H^i(K, \mathbb{Z}/2)$ is defined [D2] (1.3). The first one

$w_1(V) \in H^1(K, \mathbb{Z}/2)$ is the determinant character regarded as an element in $\text{Hom}(G_K, \mathbb{Z}/2) = H^1(K, \mathbb{Z}/2)$. The total Stiefel–Whitney class $w(V) = 1 + w_1(V) + w_2(V) + \dots \in 1 + H^1(K, \mathbb{Z}/2) + H^2(K, \mathbb{Z}/2) + \dots$ satisfies the multiplicativity $w(V) = w(V_1) \cdot w(V_2)$ for the orthogonal direct sum $V = V_1 \oplus V_2$. Hence the Stiefel–Whitney class is defined for a virtual orthogonal representation $V = V_1 - V_2$ by $w(V) = w(V_1)w(V_2)^{-1}$. When K is a local field of characteristic $\neq 2$, the second Stiefel–Whitney class is related to the local constant as follows. We identify $H^2(K, \mathbb{Z}/2)$ with $\{\pm\}$ by the isomorphism inv_K .

THEOREM D. ([D2] Théorème (1.5), [S3] Théorème 1) *Let K be a local field of characteristic $\neq 2$ with residue field of order q and let V^0 be a virtual orthogonal representation of the absolute Galois group G_K . Assume that $\dim V^0 = 0$ and $\det V^0 = 1$. Then the Artin conductor $a(V^0)$ is an even integer and the local constant $\bar{\epsilon}(V^0) = \epsilon(V^0) \cdot q^{-a(V^0)/2}$ is ± 1 and*

$$\bar{\epsilon}(V^0) = w_2(V^0).$$

For a field K of characteristic $\neq 2$, we call a quadratic K -module a K -vector space of finite dimension with a non-degenerate quadratic form. For a quadratic K -module W , its Stiefel–Whitney class $w_i(W) \in H^i(K, \mathbb{Z}/2)$ is defined ([S3] 1.2). The first one $w_1(W) \in H^1(K, \mathbb{Z}/2)$ is the discriminant regarded as an element in $K^\times/K^{\times 2} = H^1(K, \mathbb{Z}/2)$. The total Stiefel–Whitney class $w(W) = 1 + w_1(W) + w_2(W) + \dots \in 1 + H^1(K, \mathbb{Z}/2) + H^2(K, \mathbb{Z}/2) + \dots$ satisfies the multiplicativity $w(W) = w(W_1) \cdot w(W_2)$ for the orthogonal direct sum $W = W_1 \oplus W_2$. Hence the Stiefel–Whitney class is defined for a virtual orthogonal representation $W = W_1 - W_2$ by $w(W) = w(W_1)w(W_2)^{-1}$. For elements $a, b \in K^\times$, let $\{a\}$ denote the class of a in $H^1(K, \mathbb{Z}/2) = K^\times/K^{\times 2}$ and let $\{a, b\} = \{a\} \cup \{b\} \in H^2(K, \mathbb{Z}/2)$ denote the cup-product. For a quadratic K -module (W, Q) with an orthogonal basis $(e_i)_i$, its total Stiefel–Whitney class is $w(W) = \prod_i (1 + \{Q(e_i)\})$ by definition. The Stiefel–Whitney class of the representation $V_{L/K}$ is related to the Stiefel–Whitney class of a quadratic form as follows.

THEOREM S. ([S4] Théorème 1) *Let L be a finite separable extension of a field K of characteristic $\neq 2$. Let $V = Ind_{G_K}^{G_L} 1$ be the induced orthogonal representation and W be the quadratic K -module $(L, \text{Tr}_{L/K}(x^2))$. Then by putting $d = d_{L/K} = w_1(W) \in H^1(K, \mathbb{Z}/2)$, we have*

$$w_2(V) = w_2(W) + \{d, 2\}$$

in $H^2(K, \mathbb{Z}/2)$.

In the rest of this section, we deduce Theorem for odd n from Theorems D and S. The proof for even n is more complicated and will be given at the end of the next section.

We prepare some terminology for quadratic K -modules. For $c \in K^\times$, let (c) denote the quadratic K -module (K, cx^2) . The dimension of a maximal totally isotropic subspace of a quadratic K -module W is called the index of W . If $\dim W = 2$ index W , the quadratic module W is called hyperbolic.

Proof of Theorem for odd n . We assume n is odd. By Theorem D, we have $\bar{e}(V^0) = w_2(V^0)$ in $H^2(K, \mathbb{Z}/2) = \{\pm 1\}$. Since $(a, b)_K = \text{inv}_K\{a, b\}$, it is sufficient to show that $w_2(V^0) = \{d, 2\} \in H^2(K, \mathbb{Z}/2)$. Let W^0 be the virtual quadratic K -module $W - n(d)$. We show Theorem S implies $w_2(V^0) = w_2(W^0) + \{d, 2\}$. In fact $w(V^0) = w(V)w(n(d))^{-1} = (w(W) + \{d, 2\})w(n(d))^{-1} = w(W^0) + \{d, 2\}$. Note $H^i(K, \mathbb{Z}/2) = 0$ for $i > 2$. Therefore it is sufficient to prove that $w(W) = w(n(d))$. By $\{d, d\} = \{d, -1\}, \{-1, -1\} = 0$ and $n \equiv 1$, we have

$$\begin{aligned} w(n(d)) &= (1 + \{d\})^n = 1 + \{d\} + \frac{n-1}{2} \{d, -1\} \\ &= (1 + \{-1\})^{(n-1)/2} (1 + \{(-1)^{(n-1)/2} d\}). \end{aligned}$$

Namely we have $w(n(d)) = w(W')$ where $W' = (\text{hyperbolic of dimension } n-1) \oplus ((-1)^{(n-1)/2} d)$ is the orthogonal direct sum. Hence it is sufficient to show that $W \simeq W'$. Since the discriminants are equal, it is sufficient to show that the index of W is also $(n-1)/2$. Let W_a be the quadratic K -module $(L, \text{Tr}_{L/K}(ax^2))$ for $a \in L^\times$. The isomorphism class of W_a depends only on the class of a in $L^\times/L^{\times 2}$. If $a \in K^\times$, it is isomorphic to $(a) \otimes W$. Since $K^\times/K^{\times 2} \rightarrow L^\times/L^{\times 2}$ is an injection of finite groups of the same order, it is an isomorphism. Hence for any $a \in L^\times$, there is some $b \in K^\times$ such that $W_a \simeq (b) \otimes W$ and the index of W_a is independent of $a \in L^\times$. Therefore it is sufficient to show that the index of W_a is $(n-1)/2$ for single a . It follows from

LEMMA 1. *Let L be a separable extension of a field K of degree n , t be a primitive element of L and g be the minimal polynomial of t . Put $a = g'(t)^{-1}$. Then the index of the quadratic K -module $W_a = (L, \text{Tr}_{L/K}(ax^2))$ is equal to the integral part of half of n*

$$\text{index } W_a = \left[\frac{n}{2} \right].$$

Proof. We know $\text{Tr}_{L/K}(t^i/g'(t)) = 0$ for $0 \leq i \leq n-2$ and $= 1$ for $i = n-1$ [S1] Chap. III Lemme 2. Hence the subspace spaned by $(t^i)_{0 \leq i \leq [(n-2)/2]}$ is isotropic and is of dimension $[n/2]$. Since $\dim \geq 2$ index, Lemma is proved.

Thus the proof of Theorem for odd n is completed.

Remark. By [S2] Chap. IV Théorème 7, $w(W) = w(n(d))$ implies $W \simeq n(d)$.

2. $\text{Tr}_{L/K}(ax^2)$

In this section, let K be a complete discrete valuation field with residue field F . We do not assume that F is finite but keep the assumption that characteristic of F is not equal to 2. First we consider a totally ramified extension L of K and compute the quadratic K -module $W_a = (L, \text{Tr}_{L/K}(ax^2))$ for $a \in L^\times$.

PROPOSITION 1. *Let L be a separable totally ramified extension of degree e of K and $a \in L^\times$. Let W_a be the quadratic K -module $(L, \text{Tr}_{L/K}(ax^2))$ and $d_a = N_{L/K}(a) \cdot d_{L/K} \in K^\times / K^{\times 2}$ be its discriminant.*

A. *Assume $e = 2m + 1$ is odd. Then there is an orthogonal decomposition*

$$W_a \simeq ((-1)^m d_a) \oplus (\text{hyperbolic}).$$

B. *Assume $e = 2m$ is even and let D be the valuation of the different $D_{L/K}$.*

B1. *If $\text{ord}_L a \equiv D \pmod{2}$, then W_a is hyperbolic.*

B2. *Assume $\text{ord}_L a \equiv D + 1 \pmod{2}$. Let π be a prime element of L and g be the minimal polynomial of π over K . Then the class $\alpha_a = a \cdot g'(\pi)\pi^{-1} \in L^\times / L^{\times 2}$ is independent of π and is in $F^\times / F^{\times 2}$. Let α_a also denote a unit of K whose class in $K^\times / K^{\times 2}$ is α_a . Then there is an orthogonal decomposition*

$$W_a \simeq (\alpha_a) \oplus ((-1)^{m-1} d_a / \alpha_a) \oplus (\text{hyperbolic}).$$

Proof.

A. Assume e is odd. Then $K^\times / K^{\times 2} \rightarrow L^\times / L^{\times 2}$ is an isomorphism. Therefore by the same argument as in the proof of Theorem for odd n above, we see that the index is m . By comparing the discriminant, we obtain the result.

B. Assume e is even. Let π be a prime element of L and g be its minimal polynomial. First we prove the assertion for $a_0 = g'(\pi)^{-1}$ and $a_1 = \pi \cdot g'(\pi)^{-1}$. Note that $D = -\text{ord}_L a_0$ and the class of a_1 in $L^\times / L^{\times 2}$ is independent of choice of π . In fact, it is the image of the refined different $\mathcal{D}(L/K) \in L^\times / 1 + m_L [K]$ Section 2 by the equality 1.16 p. 322 loc. cit.. By Lemma 1, the quadratic module W_{a_0} is

hyperbolic. We show $W_{a_1} \simeq (1) \oplus ((-1)^{m-1} d_{a_1}) \oplus (\text{hyperbolic})$. By the formula in the proof of Lemma 1, the subspace spanned by π^{m-1} is isomorphic to (1) and the subspace spanned by $(\pi^i)_{0 \leq i \leq m-2}$ is totally isotropic and perpendicular to π^{m-1} . Hence we have $W_{a_1} \simeq (1) \oplus (\text{hyperbolic}) \oplus (\text{dimension 1})$. By comparing the discriminant, it is proved.

We consider general $a \in L^\times$. The image of $K^\times/K^{\times 2} \rightarrow L^\times/L^{\times 2}$ is $F^\times/F^{\times 2}$ and is of index 2. Hence the condition $\text{ord}_L a \equiv D$ (resp. $D+1$) is equivalent to that the class of a/a_0 (resp. a/a_1) in $L^\times/L^{\times 2}$ is in the image of $K^\times/K^{\times 2}$. It further implies $W_a \simeq (b) \otimes W_{a_0}$ (resp. $W_a \simeq (b) \otimes W_{a_1}$) for some $b \in K^\times$. Therefore $\text{ord}_L a \equiv D$ implies W_a is hyperbolic. Assume $\text{ord}_L a \equiv D+1$. Since $a = \alpha_a \cdot a_1$ in $L^\times/L^{\times 2}$ and $\alpha_a \in K^\times$, we have $W_a \simeq (\alpha_a) \otimes W_{a_1}$. Hence $W_a \simeq (\alpha_a) \oplus (\text{hyperbolic}) \oplus (\text{dimension 1})$ and comparing the discriminant, we obtain the assertion.

For a general extension, we compute the image of $w_2(\text{Tr}_{L/K}(ax^2)) \in H^2(K, \mathbb{Z}/2)$ by the boundary map $\partial: H^2(K, \mathbb{Z}/2) \rightarrow H^1(F, \mathbb{Z}/2)$. The spectral sequence $H^i(F, H^j(K^{nr}, \mathbb{Z}/2)) \Rightarrow H^{i+j}(K, \mathbb{Z}/2)$ induces an exact sequence

$$0 \rightarrow H^i(F, \mathbb{Z}/2) \rightarrow H^i(K, \mathbb{Z}/2) \xrightarrow{\partial} H^{i-1}(F, \mathbb{Z}/2) \rightarrow 0$$

for an integer i . The pairing with $\{\pi\}$ for a prime element π gives a section of ∂ . For $i=2$, we have a commutative diagram

$$\begin{array}{ccc} K^\times \times K^{\times 2} & \xrightarrow{\text{tame symbol}} & F^\times \\ \{, \} \downarrow & & \downarrow \{\} \\ H^2(K, \mathbb{Z}/2) & \xrightarrow{\partial} & H^1(F, \mathbb{Z}/2). \end{array}$$

For an element $c \in K^\times$ of even valuation, the class $\{c\} \in K^\times/K^{\times 2} = H^1(K, \mathbb{Z}/2)$ is in the subgroup $F^\times/F^{\times 2} = H^1(F, \mathbb{Z}/2)$.

PROPOSITION 2. *Let L be a separable extension of K of degree n . We assume the extension of the residue field E over F is separable of degree f and the characteristic of F is > 2 . Let $a \in L^\times$ and W_a be the quadratic K -module $(L, \text{Tr}_{L/K}(ax^2))$.*

1. *If $\text{ord}_L a \equiv D \pmod{2}$, the boundary of the total Stiefel–Whitney class $\partial w(W_a)$ is 0.*
2. *Assume $\text{ord}_L a \equiv D+1 \pmod{2}$. Let $d_a = N_{L/K}(a) \cdot d_{L/K} \in K^\times/K^{\times 2}$ be the discriminant of W_a .*

2A. *If the ramification index e is odd, we have*

$$\partial w_2(W_a) = \binom{n}{2} \{-1\} + \begin{cases} 0 & f \text{ odd} \\ \{d_a\} & f \text{ even.} \end{cases}$$

2B. Assume e is even. Let K_1 be the maximum unramified subextension of K in L , $\alpha_a \in E^\times/E^{\times 2}$ be as in Proposition 1 for the totally ramified extension L over K_1 and $d'_a = N_{E/F}(\alpha_a) \cdot d_{E/F} \in F^\times/F^{\times 2}$. Then we have

$$\partial w_2(W_a) = \binom{f}{2} \{-1\} + \{d'_a\} + \begin{cases} \left(\frac{e}{2} - 1\right) \{-1\} & f \text{ odd} \\ \{d_a\} & f \text{ even.} \end{cases}$$

Proof. To show $\partial w(W_a)$ is 0, it is enough to find a non-degenerate \mathcal{O}_K -lattice. In fact, then for an orthogonal basis over \mathcal{O}_K , the value of the quadratic form at each element of the basis is a unit. Let K_1 be the maximum unramified subextension of K in L . For a quadratic K_1 -module (W, Q) , let $\text{Tr}_{K_1/K} W$ denote the quadratic K -module $(W, \text{Tr}_{K_1/K} \circ Q)$. Since \mathcal{O}_{K_1} is a non-degenerate \mathcal{O}_K -lattice of $\text{Tr}_{K_1/K}((u))$ for a unit u of K_1 , a non-degenerate \mathcal{O}_{K_1} -lattice of quadratic K_1 -module W is a non-degenerate \mathcal{O}_K -lattice of $\text{Tr}_{L/K} W$.

1. Assume $\text{ord}_L a \equiv D$. By the argument above, it is enough to find a non-degenerate \mathcal{O}_K -lattice of W_a by assuming $K = K_1$. Namely we may assume L is totally ramified. If e is even, W_a is hyperbolic by Proposition 1 B1, and has a non-degenerate \mathcal{O}_K -lattice. Assume e is odd. By Proposition 1 A, we have $W_a \simeq ((-1)^m d_a) \oplus (\text{hyperbolic})$. Since the valuation $\text{ord}_K d_a = \text{ord}_K (N_{L/K}(a) \cdot d_{L/K}) = \text{ord}_L(a) + D$ is even, the quadratic K -module W_a has a non-degenerate \mathcal{O}_K -lattice. Therefore we have $\partial w(W_a) = 0$.

2. We prove the case $\text{ord}_L a \equiv D + 1 \pmod{2}$, by using the following Lemma proved later.

LEMMA 2. Let $W = W_1 \oplus W_2$ be the orthogonal direct sum of quadratic K -modules W_1 and W_2 . Assume W_1 and $(\pi) \otimes W_2$ have non-degenerate \mathcal{O}_K -lattices for a prime element π . Let d and d_i be the discriminants of W and W_i and r_2 be the dimension of W_2 . Then we have

$$\partial w_2(W) = \{d_1\} + \binom{r_2}{2} \{-1\} + \begin{cases} 0 & r_2 \text{ odd} \\ \{d\} & r_2 \text{ even.} \end{cases}$$

We complete the proof using Lemma. Assume first that e is odd. By Proposition 1A, we have $W_a \simeq \text{Tr}_{K_1/K}(\text{hyperbolic}) \oplus \text{Tr}_{K_1/K}((-1)^{(e-1)/2} d_a^1)$ for $d_a^1 = N_{L/K_1}(a) \cdot d_{L/K_1}$. Here $W_1 = \text{Tr}_{K_1/K}(\text{hyperbolic})$ is also hyperbolic and has a non-degenerate \mathcal{O}_K -lattice. Since the valuation $\text{ord}_{K_1} d_a^1 = \text{ord}_L a + D$, is odd, the quadratic K_1 -module $(\pi) \otimes ((-1)^{(e-1)/2} d_a^1)$ has a non-degenerate \mathcal{O}_{K_1} -lattice. Hence the quadratic K_1 -module $(\pi) \otimes W_2$ has a non-degenerate \mathcal{O}_K -lattice for $W_2 = \text{Tr}_{K_1/K}((-1)^{(e-1)/2} d_a^1)$. Therefore by applying Lemma 2 and using $d_1 = (-1)^{(e-1)/2} f$, $r_2 = f$ and $d = d_a$, we have

$$\partial w_2(W_a) = \left(\frac{(e-1)f}{2} + \binom{f}{2} \right) \{-1\} + \begin{cases} 0 & f \text{ odd} \\ \{d_a\} & f \text{ even.} \end{cases}$$

Since e is odd, we have

$$\frac{(e-1)f}{2} + \binom{f}{2} \equiv \frac{(e-1)f}{2} + e \binom{f}{2} = \frac{f(n-1)}{2} \equiv \binom{n}{2}$$

mod 2 and Proposition 2 is proved in this case.

Finally assume e is even. By Proposition 1 B2, we have an orthogonal decomposition $W_a = W_1 \oplus W_2$ where $W_1 = \text{Tr}_{K_1/K}((\alpha_a) \oplus \text{hyperbolic})$ and $W_2 = \text{Tr}_{K_1/K}((-1)^{m-1} d_a^1 / \alpha_a)$. Similarly as above, it is easily checked to satisfy the assumption of Lemma 2. The discriminant $d_1 \in F^\times / F^{\times 2} \subset K^\times / K^{\times 2}$ is $(-1)^{((e/2)-1)/2} \times \text{disc}(\text{Tr}_{K_1/K}(\alpha_a))$ and $\text{disc}(\text{Tr}_{K_1/K}(\alpha_a)) = \text{disc}(\text{Tr}_{E/F}(\alpha_a)) = d'_a$. Hence Lemma 2 gives us

$$\partial w_2(W_a) = \left(\left(\frac{e}{2} - 1 \right) f + \binom{f}{2} \right) \{-1\} + \{d'_a\} + \begin{cases} 0 & f \text{ odd} \\ \{d_a\} & f \text{ even.} \end{cases}$$

Thus Proposition 2 is proved.

Proof of Lemma 2. By the assumption that W_1 has a non-degenerate lattice, $w(W_1) \in H^*(K, \mathbb{Z}/2)$ is in $H^*(F, \mathbb{Z}/2) \subset H^*(K, \mathbb{Z}/2)$. We have $\partial w(W) = w(W_1) \cdot \partial w(W_2) \in H^*(F, \mathbb{Z}/2)$. We have $w(W_1) \equiv 1 + \{d_1\} \pmod{\text{degree} > 1}$. Let $W'_2 = (\pi^{-1}) \otimes W_2$. It has a non-degenerate \mathcal{O}_K -lattice and $w(W'_2) \in H^*(F, \mathbb{Z}/2)$. We have

$$w(W_2) = w((\pi) \otimes W'_2) = \sum_j w_j(W'_2) (1 + \{\pi\})^{r_2 - j} = \sum_{i,j} \binom{r_2 - j}{i} w_j(W'_2) \{\pi\}^i.$$

Since $\{\pi, \pi\} = \{\pi, -1\}$ and $\{-1, -1\} = 0$, we have $\{\pi\}^3 = 0$ and

$$\partial w_k(W_2) = (r_2 - k + 1) w_{k-1}(W'_2) + \binom{r_2 - k + 2}{2} \{w_{k-2}(W'_2), -1\}.$$

Hence we obtain

$$\begin{aligned} \partial w(W) &\equiv (1 + \{d_1\}) \left(r_2 + (r_2 - 1) w_1(W'_2) + \binom{r_2}{2} \{-1\} \right) \pmod{\text{degree} > 1} \\ &= r_2 + \{d_1\} + (r_2 - 1)(\{d_1\} + w_1(W'_2)) + \binom{r_2}{2} \{-1\}. \end{aligned}$$

By

$$(r_2 - 1)(\{d_1\} + w_1(W'_2)) = \begin{cases} 0 & r_2 \text{ odd} \\ \{d_1\} + \{d_2\} = \{d\} & r_2 \text{ even,} \end{cases}$$

Lemma 2 is proved.

Proof of Theorem for even n . Assume n is even. By Theorem D, we have $\bar{e}(V^0) = w_2(V^0)$. Further by $w(V^0) = w(V)w(\delta)^{-1}(w(\delta) + w_2(V))w(\delta)^{-1} = 1 + w_2(V)$, we have $w_2(V^0) = w_2(V)$. Hence by Theorem S, we have $\bar{e}(V^0) = w_2(V) = w_2(W) + (d, 2)_K$. Since $\text{inv}_K = (F) \circ \partial: H^2(K, \mathbb{Z}/2) \simeq \{\pm 1\}$, it is enough to compute the boundary $\partial w_2(W) \in H^1(F, \mathbb{Z}/2)$. We check that Theorem is now a special case of Proposition 2 where F is finite and $a = 1$. In fact if f or D is even, the valuation of d is even and $(d, 2)_K = 1$ and $(\{d\}/F) = (d, \pi_K)_K$ for a prime element π_K of K . Thus the proof of Theorem is completed.

REFERENCES

- [D1] DELIGNE, P., *Les constantes des équations fonctionnelles des fonctions L*, in Modular functions of one variable II, Lect. Notes in Math 349 (1972), 501–597, Springer-Verlag, Berlin–Heidelberg–New York.
- [D2] DELIGNE, P., *Les constantes locales de l'équation fonctionnelle de la fonction L d'Artin d'une représentation orthogonale*, Inventiones Math. 35 (1976), 299–316.
- [H] HENNIART, G., *Galois ϵ -factors modulo roots of unity*, Inventiones Math. 78 (1984), 117–126.
- [K] KATO, K., *Swan conductors with differential values*, Adv. Studies in Pure Math. 12 (1987), Kinokuniya, Tokyo, 315–342.
- [S1] SERRE, J.-P., *Corps Locaux*, Hermann, Paris, 1968.
- [S2] SERRE, J.-P., *Cours d'arithmétique*, Presses Universitaire de France, Paris, 1970.
- [S2] SERRE, J.-P., *Conducteurs d'Artin des caractères réels*, Inventiones Math. 14 (1971), 173–183.
- [S4] SERRE, J.-P., *L'invariant de Witt de la forme $\text{Tr}_{L/K}(x^2)$* , Comm. Math. Helv. 59 (1984), 651–675.

Department of Mathematical Sciences
 University of Tokyo
 Tokyo, 113
 Japan

Received September 21, 1993; November 15, 1994