Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	71 (1996)
Artikel:	On the Malcev completion of Kähler groups.
Autor:	Amorós, Jaume
DOI:	https://doi.org/10.5169/seals-53841

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On the Malcev completion of Kähler groups*

JAUME AMORÓS

Introduction

The study of compact Kähler manifolds made by Hodge and others shows that a Kähler structure imposes very strong conditions on the homotopy type of a compact complex manifold X. In particular, unlike in the case of compact differentiable or closed complex manifolds, not every finitely presented group G is the fundamental group of a compact Kähler manifold. Such groups are called Kähler groups.

This note has been inspired by the recent work of F. Johnson and E. Rees ([JR]) and M. Gromov ([G]), showing that free products, and in particular free groups, are not Kähler. It has been our purpose to extend this result and find other restrictions on the presentations of Kähler groups. This is done by translating properties of cup products in $H^*(X)$ into properties of the group bracket in $\pi_1 X$, an idea that came out of [JR], and also by examining the Albanese map $X \to Alb(X)$ after [C]. We describe an algorithm derived from [St] to compute $\Gamma_1/\Gamma_2 G$, $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$ from a given presentation of a group G, and use it to give three conditions for the groups to be Kähler: The Lie algebra $\mathscr{L}_2 G$, equivalent to the holonomy algebra, cannot be free (3.3); one- or two-relator Kähler groups either have a torsion abelianized or have a Malcev completion isomorphic to that of a compact Riemann surface group (4.7); nonfibered Kähler groups must satisfy certain lower bounds for the number of their defining relations, equivalently upper bounds for the rank of $\Gamma_2/\Gamma_3 G$ (5.6, 5.7).

In §1 we recall the real Malcev completion $G \otimes \mathbb{R}$ of a group G, its equivalent Lie algebra $\mathscr{L}G$, and a 2-step nilpotent Lie algebra $\mathscr{L}_2G \cong (\Gamma_1/\Gamma_2G \otimes \mathbb{R}) \oplus$ $(\Gamma_2/\Gamma_3G \otimes \mathbb{R})$, which is determined by $(\Gamma_1/\Gamma_3G)/_{\text{Torsion}}$ and is equivalent to the cup products $\bigwedge^2 H^1(X) \to H^2(X)$. This algebra is actually equivalent to the holonomy algebra of G (cf. [Ch], [Ko]), and is more convenient for our computations. By [M2], [DGMS], when G is a Kähler group the algebra \mathscr{L}_2G determines the Malcev

^{*}Supported by the Generalitat de Catalunya, Partially supported by DGCYT grant PB93-0790.

completion $G \otimes \mathbb{R}$. In §2 we briefly recall Sullivan's 1-minimal model of X, its duality with $\mathscr{L}\pi_1 X$, and how the algebra $\mathscr{L}_2\pi_1 X$ and the product map \cup : $\bigwedge^2 H^1(X) \to H^2(X)$ are equivalent.

In §3 we use these results to show that if $\mathscr{L}_2 G \cong \mathscr{L}_2 F_n$, where F_n is a finite rank free group then G is not Kähler. This is a strong quantitative restriction, since the generic group presented with few relations verifies $\mathscr{L}_2 \cong \mathscr{L}_2 F_n$ for some n (see Remark 1.15).

The groups with the simplest presentation after free groups are 1- and 2-relator groups. In Theorem 4.7 we determine the Malcev completions of 1- and 2-relator Kähler groups, which to a great extent characterize the groups themselves. The mean to do this is to bound from above dim $\Gamma_2/\Gamma_3\pi_1 X \otimes \mathbb{R}$ for any Kähler group G by a function of the dimension of the image Y of X by its Albanese map α : $X \to Alb(X)$. A desingularization \tilde{Y} of $\alpha(x)$ has been shown by F. Campana ([C]) to verify $\pi_1 X \otimes \mathbb{R} \cong \pi_1 \tilde{Y} \otimes \mathbb{R}$. It turns out of our work that as dim Y increases linearly, dim $\bigwedge^2 H^1(X) - \dim \Gamma_2/\Gamma_3\pi_1(X) \otimes \mathbb{R}$ grows quadratically (Prop. 4.6).

Finally, in §5, we have established a distinction between fibered and nonfibered Kähler groups, and used the mentioned techniques to give upper bounds for dim $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$ for nonfibered groups, which imply lower bounds for their number of defining relations (Prop. 5.6, Cor. 5.7). Specifically, in Cor. 5.7 we give the following lower bound for the difference between the number of generators *n* and of relations *s* of any presentation of a nonfibered Kähler group

$$s-n \ge 4q-7$$

where $q = \frac{1}{2}b_1(G)$. This improves a bound given by Green and Lazarsfeld ([GL]).

To proof our results and make them effective, we give in §1 an algorithm for computing $\Gamma_1/\Gamma_2 G \otimes \mathbb{R}$, $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$ and $\mathscr{L}_2 G$ from a given presentation of G. This algorithm, which is easy to implement by means of the Magnus algebra of free groups, is derived from a spectral sequence given in [St], and was communicated to the author by M. Hartl. We use it in Cor. 1.14 and Rmk. 1.15 to show many cases in which the hypothesis of Thm. 3.3 are fulfilled.

To illustrate our results, we give throughout the paper examples of groups that cannot be Kähler, most of them previously unknown to the author.

Acknowledgements

The author wishes to thank V. Navarro Aznar, for originally proposing the problem and many interesting discussions on it. Manfred Hartl for his valuable help in §1, F. Campana and the members of the Cigrons Seminar of the Universitat de Barcelona are also to be thanked.

§1. Nilpotent Lie algebras related to a group

We will recall here the concept of Malcev completion of a group and some related nilpotent Lie algebras $\mathcal{L}_n G$. We also give an algorithm to compute $\mathcal{L}_2 G$ derived from [St].

Let G be a finitely presented group. We can functorially assign to it a tower of nilpotent groups

$$\cdots \to \Gamma_1/\Gamma_3 G \to \Gamma_1/\Gamma_2 G \to 1$$

where $\Gamma_1 G = G$, $\Gamma_n G = [\Gamma n - 1G, G]$ and $\Gamma_1 / \Gamma_n G = G / \Gamma_n G$.

A group G is said to be uniquely divisible when for any pair $g \in G$, $n \in \mathbb{Z}$, g has a unique nth root in G. The category $n \cdot \mathbb{Q} \cdot \mathscr{G}r$ of uniquely divisible nilpotent groups is included in the category $n \cdot \mathscr{G}r$ of nilpotent groups, and the inclusion functor has a left adjoint, the Malcev completion functor $\otimes \mathbb{Q}: n \cdot \mathscr{G}r \to n \cdot \mathbb{Q} \cdot \mathscr{G}r$. The functor $\otimes \mathbb{Q}$ is the ordinary tensor product on abelian groups. For two alternative ways of defining the Malcev completion, see [HMR] Part I, or App. A of [Q].

The Baker-Campbell-Hausdorff formula gives a categorical equivalence between finitely generated groups in n-Q-Gr and finite-dimensional nilpotent Q-Lie algebras. In the latter category there are $\otimes \mathbb{R}$, $\otimes \mathbb{C}$ functors, and crossing back and forth in this manner we may define a $\otimes \mathbb{R}$ functor over n-Gr. Thus we naturally associate to G a tower

$$\cdots \to \Gamma_1/\Gamma_3 G \otimes \mathbb{R} \to \Gamma_1/\Gamma_2 G \otimes \mathbb{R} \to 1$$

of uniquely divisible nilpotent Lie groups, and its corresponding tower

$$\cdots \to \mathscr{L}_2 G \to \mathscr{L}_1 G \to 0 \tag{1.1}$$

of nilpotent \mathbb{R} -Lie algebras.

Denote the lower central series of a Lie algebra \mathscr{L} as $\mathscr{L}^{(1)} = \mathscr{L}$, $\mathscr{L}^{(n)} = [\mathscr{L}^{(n-1)}, \mathscr{L}]$. There is another tower of nilpotent \mathbb{R} -Lie algebras naturally associated to a group $G: Gr_n G \otimes \mathbb{R} = \bigoplus_{i=1}^n \Gamma_i / \Gamma_{i+1} G \otimes \mathbb{R}$, with bracket induced by the group bracket. We sum up the properties of the tower of Lie algebras (1.1) that we will apply (see [L], [MKS] §5, [S] LA §4):

PROPOSITION 1.2.

(i) The Lie algebras $\mathcal{L}_n G$ have nilpotency class nil $\mathcal{L}_n G = n$.

(ii) The tower maps $\mathscr{L}_{n+1}G \to \mathscr{L}_nG$ induce isomorphisms $\mathscr{L}_{n+1}G/\mathscr{L}_{n+1}G^{(n+1)} \cong \mathscr{L}_nG$.

(iii) There are isomorphisms of \mathbb{R} -vector spaces $\mathscr{L}_n^{(n)}G \cong \Gamma_n/\Gamma_{n+1}G \otimes \mathbb{R}$.

(iv) The graduation of $\mathcal{L}_n G$ by its lower central series produces a natural tower of isomorphisms of graded Lie algebras $Gr \mathcal{L}_n G \cong Gr_n G \otimes \mathbb{R}$.

We will call the *Malcev algebra* and denote $\mathscr{L}G$ the pronilpotent algebra $\lim_{n \to \infty} \mathscr{L}_n G$, which is equivalent to the Malcev completion $G \otimes \mathbb{R}$ by the Baker-Campbell-Hausdorff formula.

When the group G is the fundamental group of a topological space X, the abelian algebra \mathscr{L}_1G is just $H_1(X; \mathbb{R})$. We will consider in this note the following simplest algebra, $\mathscr{L}_2G \cong \Gamma_1/\Gamma_2G \otimes \mathbb{R} \oplus \Gamma_2/\Gamma_3G \otimes \mathbb{R}$. The algebra \mathscr{L}_2G is the quotient of the Malcev algebra $\mathscr{L}G$ by its third commutator ideal $\mathscr{L}G^{(3)}$, and is also the quotient of the holonomy algebra of $G \mathfrak{g}_G$ (cf. [Ch], [Ko]) by its third commutator ideal.

The groups G we will study will be given by finite presentations $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$. This means that G is defined by

$$1 \to N \to F \to G \to 1 \tag{1.3}$$

where F is the free group generated by the generator set $\{x_1, \ldots, x_n\}$, and N is the normal subgroup of F spanned by the relation set $\{r_1, \ldots, r_s\} \subset F$.

We describe the above constructions in a case which is fundamental for our purposes:

Example 1.4. Free groups.

Let $G = F_n = F_{\{x_1, \ldots, x_n\}}$. Its Malcev completion and Lie algebras $\mathscr{L}_n F_n$ may be computed by means of its group algebra (cf. [MKS], [Q], [S]). The conclusion is that, denoting by $\mathscr{L}(S)$ the free \mathbb{R} -Lie algebra spanned by a set S, there are isomorphisms

$$\mathscr{L}_m F_n \cong \mathscr{L}(\{X_1, \ldots, X_n\}) / \mathscr{L}(\{X_1, \ldots, X_n\})^{(m+1)}$$

In particular, $\Gamma_1/\Gamma_2 F_n \otimes \mathbb{R} \cong \mathbb{R} x_1 \oplus \cdots \oplus \mathbb{R} x_n$, $\Gamma_2/\Gamma_3 F_n \otimes \mathbb{R} \cong \mathbb{R}(x_1, x_2) \oplus \cdots \oplus \mathbb{R}(x_{n-1}, x_n)$ and the brackets in $\mathcal{L}_2 F_n$ are the group ones in $\Gamma_1/\Gamma_2 F_n$ and zero all others.

The Lie algebra \mathscr{L}_2G for a finitely presented G may be obtained from its presentation and \mathscr{L}_2F . We will use an algorithm for computing them derived from [St], where a spectral sequence that computes all J_G^m/J_G^{m+1} is described, and communicated to the author by Manfred Hartl.

Consider a group presentation $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$, which induces the exact sequence given in (1.3). Let $\mathbb{R}F,\mathbb{R}G$ be the \mathbb{R} -group algebras of F,G, and denote by J_F, J_G their respective augmentation ideals. The sequence (1.3) induces an exact sequence of \mathbb{R} -algebras

$$0 \to K \to \mathbb{R}F \to \mathbb{R}G \to 0 \tag{1.5}$$

where K is the two-sided ideal generated by the R-vector space $D = \langle r_1 - 1, \dots, r_s - 1 \rangle \subset J_F$. This sequence restricts to exact sequences

$$0 \to K \to J_F^m + K \to J_G^m \to 0 \tag{1.6}$$

for all $m \ge 1$. We will compute J_G/J_G^2 , J_G^2/J_G^3 from those sequences:

PROPOSITION 1.7. Consider the linear map $f: \bigoplus_{i=1}^{s} \mathbb{R}r_i \rightarrow J_F$ determined by $r_i \mapsto r_i - 1$.

(i) Let $d_0: \bigoplus \mathbb{R}r_i \to J_F/J_F^2$ be the projection of f. Then coker $d_0 \cong J_G/J_G^2$.

(ii) The map f induces a linear map

$$d_1 : ker \ d_0 \to J_F^2 / (J_F^3 + J_F \cdot D + D \cdot J_F)$$
$$\sum \lambda_i r_i \mapsto \sum \lambda_i (r_i - 1)$$

and coker $d_1 \cong J_G^2 / J_G^3$.

Proof. (i) The exact sequences of (1.6) induce an isomorphism $J_G/J_G^2 \cong J_F/J_F^2 + K$. As K is the two-sided ideal spanned by D and $\mathbb{R}F \cong \mathbb{R}F \cong \mathbb{R} \oplus J_F$, actually $J_F^2 + K = J_F^2 + D$, and thus $J_G/J_G^2 \cong J_F/J_F^2 + D$. By its construction, Im $d_0 = D$, and this proves (i).

(ii) Again by (1.6) we have

$$J_G^2/J_G^3 \cong (J_F^2/(J_F^2 \cap K))/(J_F^3/(J_F^3 \cap K)) \cong J_F^2/(J_F^3 + J_F^2 \cap K)$$

The last denominator is $J_F^3 + J_F^2 \cap K = J_F^3 + J_F \cdot D + D \cdot J_F + D \cap J_F^2$. Obviously $f(\ker d_0) \subset J_F^2$ and thus d_1 is well defined. Moreover, its image is precisely $D \cap J_F^2$, and (ii) follows from this.

We now relate the computed modules J_G/J_G^2 , J_G^2/J_G^3 with the sought ones Γ_1/Γ_2 , $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$ applying a theory by D. Quillen ([Q2]):

THEOREM. Let G be a group, k a field of characterisitc zero, kG the group algebra and j: $\bigoplus \Gamma_n / \Gamma_{n+1} G \otimes k \rightarrow \bigoplus J_G^n / J_G^{n+1}$ given by $g \mapsto g - 1$ over the homogeneous components.

Then j induces an isomorphism of algebras $U(\oplus \Gamma_n/\Gamma_{n+1}G \otimes \mathbb{R}) \cong \oplus J^n_G/J^{n+1}_G$.

We may define the wedge product, or alternating product, of the associative algebra $\mathbb{R}G$ as

$$x \wedge y := xy - yx$$

The wedge product of two linear subspaces A, $B \subset \mathbb{R}G$ is the linear subspace

$$A \wedge B = \left\{ w = \sum \lambda_i a_i \wedge b_i \in \mathbb{R}G \mid a_i \in A, \, b_i \in B, \, \lambda_i \in \mathbb{R} \right\}$$

Quillen's theorem implies that the Lie algebra $\oplus \Gamma_n / \Gamma_{n+1} G \otimes \mathbb{R}$ is contained in the J_G -adic graduate of the group algebra, $\oplus J_G^n / J_G^{n+1}$. This inclusion sends the brackets of the Lie algebra to wedge products in $\oplus J_G^n / J_G^{n+1}$. In the cases n = 1, 2 this means:

COROLLARY 1.8.

(i) $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} \cong J_G/J_G^2$.

(ii) Consider the inclusion $J_G \wedge J_G \hookrightarrow J_G^2$. Then

$$\Gamma_2/\Gamma_3 G \otimes \mathbb{R} \cong (J_G \wedge J_G + J_G^3)/J_G^3 \subset J_G^2/J_G^3$$

Corollary 1.8 allows us to adapt the algorithm of Prop. 1.7 to compute Γ_1/Γ_2 , $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$:

LEMMA 1.9. The image of the restriction $f: \ker d_0 \to J_F^2$ lies in $J_F \wedge J_F + J_F^3 \subset J_F^2$.

Proof. Denote F_s the free group generated by $\{y_1, \ldots, y_s\}$, and the map $r: F_s \to F$ sending y_i to r_i . The map $d_0: \bigoplus \mathbb{R}r_i \to J_F/J_F^2 \cong \Gamma_1/\Gamma_2 F \otimes \mathbb{R}$ is the map induced by r, $\Gamma_1/\Gamma_2(r) \otimes \mathbb{R}$: $\Gamma_1/\Gamma_2 F_s \otimes \mathbb{R} \to \Gamma_1/\Gamma_2 F \otimes \mathbb{R}$. Furthermore ker $(\Gamma_1/\Gamma_2(r) \otimes \mathbb{R}) \cong \ker(\Gamma_1/\Gamma_2(r)) \otimes \mathbb{R}$, as $\Gamma_1/\Gamma_2 F_s$ is a free abelian group. Thus ker d_0 admits a basis $\bar{w}_1, \ldots, \bar{w}_k$, with the w_i words in F_s mapping to $\Gamma_2 F$ by r.

Now, the map $\Gamma_2 F \to J_F^2$ sends a bracket (a, b) to (a - 1)(b - 1) - (b - 1)(a - 1) + terms in J_F^3 , and a product $\Pi(a_i, b_i)$ to $\Sigma(a_i - 1)(b_i - 1) - (b_i - 1)(a_i - 1) +$ terms in J_F^3 . Therefore, all the $w_i = \Pi(a_{j_i}, b_{j_i})$ map to $J_F \wedge J_F + J_F^3$.

Lemma 1.9 allows us to define a map d_1 : ker $d_0 \rightarrow \bigwedge^2(\Gamma_1/\Gamma_2 G \otimes \mathbb{R})$ by composing

$$\ker d_0 \to (J_F \wedge J_F + J_F^3) / J_F^3 \cong \bigwedge^2 (\Gamma_1 / \Gamma_2 F \otimes \mathbb{R}) \to \bigwedge^2 (\Gamma_1 / \Gamma_2 G \otimes \mathbb{R})$$

PROPOSITION 1.10. coker $d_1 \cong \Gamma_2 / \Gamma_3 G \otimes \mathbb{R}$.

Proof. As we have previously explained, $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} \cong J_G/J_G^2 \cong J_F/(J_F^2 + K) \cong J_F/(J_F^2 + D)$. Thus $\bigwedge^2(\Gamma_1/\Gamma_2 G \otimes \mathbb{R}) \cong (J_F \wedge J_F + (J_F^3 + J_F \cdot D + D \cdot J_F))/2$

 $(J_F^3 + J_F \cdot D + D \cdot J_F)$. Also $f(\ker d_0) = D \cap J_F^2 \subset J_F \wedge J_F + J_F^3$ by Lemma 1.9, so

 $\begin{aligned} \operatorname{coker} d_1 &\cong (J_F \wedge J_F + J_F^3 + J_F \cdot D + D \cdot J_F + D \cap J_F^2) / (J_F^3 + J_F \cdot D + D \cdot J_F \\ &+ D \cap J_F^2) \\ &\cong (J_F \wedge J_F + J_F^3 + K \cap J_F^2) / (J_F^3 + K \cap J_F^2) \\ &\cong (J_G \wedge J_G + J_G^3) / J_G^3 \cong \Gamma_2 \Gamma_3 G \otimes \mathbb{R}, \end{aligned}$

the last isomorphism being given by Cor. 1.8.

COROLLARY 1.11. $\dim \Gamma_2/\Gamma_3 G \otimes \mathbb{R} = \begin{pmatrix} \dim \Gamma_1/\Gamma_2 G \otimes \mathbb{R} \\ 2 \end{pmatrix} - \dim \ker d_0 + \dim \ker d_1.$

 \Box

We are now able to determine the structure of the 2-step nilpotent Lie algebra \mathscr{L}_2G of a finitely presented group $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$:

PROPOSITION 1.12. Let $\bigwedge^{\leq 2}(\Gamma_1/\Gamma_2 G \otimes \mathbb{R})$ be the free exterior algebra generated by $\Gamma_1/\Gamma_2 G \otimes \mathbb{R}$ modulo the ideal $\bigwedge^{\geq 3}(\Gamma_1/\Gamma_2 G \otimes \mathbb{R})$ generated by wedges of length 3 or more. There is an isomorphism

$$\mathscr{L}_2 G \cong \left(\bigwedge^{\leq 2} \left(\Gamma_1 / \Gamma_2 G \otimes \mathbb{R} \right) \right) / (\ker d_0 / \ker d_1)$$

Proof. There is an obvious map of exterior algebras, which is a linear isomorphism in every degree by the above results.

Thus $\mathscr{L}_2 G$ is the quotient of a free 2-step nilpotent \mathbb{R} -Lie algebra $\wedge {}^{\leq 2}(H_1(G; \mathbb{R}))$ by a subspace of 2-brackets ker d_0 /ker d_1 , which corresponds to the relations of the holonomy algebra. We have stated in Ex. 1.4 the case of free groups. Let us examine this structure in some other simple cases:

COROLLARY 1.13. Let $G = \langle x_1, \ldots, x_n; r \rangle$ be a group admitting a presentation with a single relation. Then:

- (i) If $r \notin \Gamma_2 F$, there is an isomorphism $\mathcal{L}_2 G \cong \mathcal{L}_2 F_{n-1}$ with F_{n-1} a free group of rank n-1.
- (ii) If $r \in \Gamma_2 F \setminus \Gamma_3 F$, there is an isomorphism $\mathscr{L}_2 G \cong \mathscr{L}_2 F / d_1(r)$.
- (iii) If $r \in \Gamma_3 F$, there is an isomorphism $\mathscr{L}_2 G \cong \mathscr{L}_2 F$.

Proof. All cases are found by applying Prop. 1.12.

(i) In this case $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} \cong \Gamma_1/\Gamma_2 F_{n-1} \otimes \mathbb{R}$, and as $r \notin \Gamma_2 F$, ker $d_0 = \{0\}$.

- (ii) In this case the map F→G induces an isomorphism Γ₁/Γ₂F⊗ ℝ ≅ Γ₁/Γ₂G ⊗ ℝ, ker d₀ = ℝr, and as r ∉ Γ₃F, the coincidence of the lower central series and augmentation ideal power filtrations in free groups ([MKS] 5.12, [S]) shows that r − 1 ∉ J_F³, hence d₁(r) ≠ 0.
- (iii) In this case, ker $d_0 = \mathbb{R}r$ and again by the above coincidence of filtrations, $d_1(r) = 0$.

COROLLARY 1.14. Let $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$ be a finitely presented group such that its defining relations may be divided in two sets: $\langle r_1, \ldots, r_k \rangle$ such that $\bar{r}_1, \ldots, \bar{r}_k$ are linearly independent in $\Gamma_1/\Gamma_2 F \otimes \mathbb{R}$ and $\{r_{k+1}, \ldots, r_s\}$ which belong to $\Gamma_3 F$. Then there is an isomorphism $\mathscr{L}_2 G \cong \mathscr{L}_2 F_{n-k}$, where F_{n-k} is a free group of rank n - k.

Proof. In this case $\Gamma_1/\Gamma_2 G \otimes \mathbb{R}$ has rank n-k, ker $d_0 = \mathbb{R}r_{k+1} \oplus \cdots \oplus \mathbb{R}r_n$ because those r_j are commutators and the other relations form a basis of Im f, and ker $d_1 = \ker d_0$ because $r_{k+1}, \ldots, r_n \in \Gamma_3 F$.

Remark 1.15. We will be interested in this note in which groups G have a free 2-step nilpotent Lie algebra \mathscr{L}_2G , which by Prop. 1.12 is equivalent to ker $d_0 = \ker d_1$.

Generic presentations with less relations than generators produce a free $\mathcal{L}_2 G$. The reason is that given a group presentation $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$ with a number of relations $s \leq n$, ker $d_0 = 0$ and therefore $\mathcal{L}_2 G$ is free, unless the classes $\bar{r}_1, \ldots, \bar{r}_s \in \Gamma_1/\Gamma_2 F_n \otimes \mathbb{R}$ are linearly dependent. But the sets of linearly dependent $\bar{r}_1, \ldots, \bar{r}_s$ form a codimension n - s + 1 closed subset of $(\Gamma_1/\Gamma_2 F_n \otimes \mathbb{R})^s$.

The hypotheses of Corollary 1.14 may be weakened by requiring only that $\{r_1, \ldots, r_k\}$ map on a basis of Im d_0 , and the remaining relations $\{r_{k+1}, \ldots, r_s\}$ belong to $\Gamma_3 F \cdot N_k$, where N_k is the normal closure in F of $\{r_1, \ldots, r_k\}$.

§2. Sullivan's 1-minimal models, brackets and cup products

We sum up for the reader's convenience some basic facts on Sullivan's 1-minimal models, its equivalence with the Malcev completion of the fundamental group and its relation with cup products.

Let X be now a differentiable manifold, and $\mathcal{E}^*(X)$ its De Rham complex.

The theory of minimal models developed by Sullivan shows that $\mathcal{E}^*(X)$ has a 1-minimal model. This is a certain minimal commutative differential graded algebra (cdga) M(2, 0)(X), defined as the limit of an inductive system of cdga $M(1, 1) \subseteq M(1, 2) \subseteq M(1, 3) \subseteq \cdots$, together with an algebra morphism $\rho: M_X \to \mathcal{E}^*(X)$ such that $H^0(\rho)$, $H^1(\rho)$ are isomorphisms and $H^2(\rho)$ is a monomorphism. For a more complete and elementary account of the theory of minimal models we remit ourselves to [GM], which treats 1-minimal models in Chap. XII. We will only review the construction up to the second step M(1, 2), which will be used to relate $\pi_1(X, *)$ and cup products on $H^1(X)$.

Define $M(1, 1) = \bigwedge (V_1^1)$, the free exterior algebra generated by the \mathbb{R} -vector space $V_1^1 = H^1(X)$. Every element of V_1^1 is defined to have degree one and boundary zero, and the map $\rho: M(1, 1) \to \mathcal{E}^*(X)$ sends every $x \in V_1^1$ to its image in a prefixed \mathbb{R} -vector space section $H^1(X) \to (\operatorname{cocycles})^1$.

The (1,2)-minimal model is defined as an extension of M(1,1): $M(1,2) = \bigwedge (V_1^1 \oplus V_2^1)$, where $V_2^1 = \ker(H^2\rho; H^2M(1,1) \to H^2\mathcal{E}^*(X))$ For any $v \in V_2^1$ we define dv as the element of $\bigwedge^2 V_1^1$ defining its cohomology class, and if $dv = \sum x_i y_i$, $\rho(v)$ is a linearly varying primitive of $\sum \rho(x_i)\rho(y_i)$ in $\mathcal{E}^*(X)$.

Remark 2.1. By its definition, $H^2M(1, 1)$ is the exterior product $\bigwedge^2 H^1(X)$, and as ρ is a cdga morphism, there is an isomorphism $V_2^1 = (\ker H^2M(1, 1) \rightarrow H^2\mathcal{E}^*(X)) \cong \ker(\bigcup: \bigwedge^2 H^1(X) \rightarrow H^2(X)).$

The following steps M(1, n) are constructed in a likewise manner, defining V_n^1 as ker $H^2M(1, n-1) \rightarrow H^2\mathcal{E}^*(X)$, and d, ρ on it as on V_2^1 . The inductive limit is denoted M(2, 0) and is the 1-minimal model of X.

Remark 2.2. Among the properties of the 1-minimal model let us remark:

- It is well defined up to isomorphism.
- It is functorial up to homotopy, i.e., any cdga morphism $\mathcal{E}^* Y \to \mathcal{E}^* X$ may be lifted to a morphism $M(2, 0)(Y) \to M(2, 0)(X)$, and all its liftings are homotopic in the cdga category.
- As a consequence of its uniqueness, if a map $f: X \to Y$ induces isomorphisms H^0f , H^1f and a monomorphism H^2f , then it induces an isomorphism of 1-minimal models $M(2, 0)(Y) \cong M(2, 0)(X)$.
- The 1-minimal model of X may be computed replacing $\mathcal{E}^*(X)$ by any other cdga quasi-isomorphic to it. Thus if X is a complex manifold, we may compute it from its holomorphic De Rham complex, logarithmic complexes, Dolbeault complexes, etc..

We recall now the dualizing process between Lie algebras and free commutative differential graded algebras generated by elements of degree one.

Let L be a finite-dimensional \mathbb{R} -Lie algebra. Its bracket is a bilinear alternating map

$$[\cdot, \cdot]: \bigwedge^2 L \to L \tag{2.3}$$

Dualizing on both sides, $[\cdot, \cdot]$ has an adjoint map

$$d: L^{\vee} \to \bigwedge^2 L^{\vee} \tag{2.4}$$

The map d may be extended as a graded derivation to the free graded algebra $\bigwedge L^{\vee}$, defining the degree of elements in V^{\vee} to be one. The Jacobi identity satisfied by $[\cdot, \cdot]$ dualizes then as $d^2 = 0$.

Reciprocally, if $M = \bigwedge W$ is a free cdga and deg W = 1, the differential restricts to a map d: $W = M^1 \rightarrow M^2 = \bigwedge^2 W$, which dualizes to a map $[\cdot, \cdot]$: $\bigwedge^2 W^{\vee} \rightarrow W^{\vee}$, and the fact $d^2 = 0$ in M translates as the Jacobi identity in W^{\vee} .

DEFINITION 2.5. A Lie algebra L and a free cdga generated by elements of degree one are *dual* when each one yields the other by the above processes.

The following result is due to D. Sullivan. The reader will find a complete proof of it in [BG].

THEOREM 2.6. (Sullivan) Let X be an arc-connected topological space with a finitely presented fundamental group $\pi_1(X, *)$. The inductive system $M(1, 1) \hookrightarrow M(1, 2) \hookrightarrow \cdots$ formed by the (1, n)-minimal models of X and the projective system $\cdots \to \mathscr{L}_2 \pi_1 X \to \mathscr{L}_1 \pi_1 X$ described in (1.1) are dual.

This theorem has important consequences for our purposes. The most obvious is about the duality as vector spaces:

COROLLARY 2.7. $V_n^1 \cong (\Gamma_n / \Gamma_{n+1}(\pi_1 X) \otimes \mathbb{R})^{\vee}$

The duality Lie algebra-cdga also has consequences:

LEMMA 2.8. The diagram

 $\begin{array}{cccc} \wedge^2 V_1^1 & \to & H^2 M(1,2) \\ \wedge^2 \rho \downarrow & & \downarrow H^2 \rho \\ \wedge^2 H^1(X) & \stackrel{\cup}{\to} & H^2(X) \end{array}$

is commutative, and the first column is an isomorphism.

Remark 2.9. Lemma 2.8 implies that the cup product between 1-classes is determined by the brackets in $\mathscr{L}_2 \pi_1 X$. This is due to the fact that it factors through M(1, 2), which is dual to $\mathscr{L}_2 \pi_1 X$.

Another particular consequence of Theorem 2.6 we will use is:

COROLLARY 2.10. dim ker(\cdots : $\bigwedge^2 H^1(X) \to H^2(X)$) = dim $\mathscr{L}_2 \pi_1 X^{(2)}$ = dim $\Gamma_2 / \Gamma_3 \pi_1(X, *) \otimes \mathbb{R}$.

Proof. Both spaces are isomorphic to V_2^1 .

§3. The \mathcal{L}_2 algebras of Kähler groups

In this chapter we prove that groups with a free \mathcal{L}_2 algebra cannot be Kähler (Thm. 3.3), which by Remark 1.15 is the generic situation in groups with few defining relations.

We recall now the definition of the objects of our interest:

DEFINITION 3.1. Let G be a group. It is a Kähler group when $G \cong \pi_1(X, *)$, where X is a compact Kähler manifold.

Some restrictions on Kähler groups are well known: They are finitely presented, their rank is even. Sullivan's theory of minimal models, completed by Morgan, ..., yields subtle conditions Kähler groups must satisfy, due to the formality of compact Kähler manifolds ([DGMS]), and to the mixed Hodge structure its 1-minimal model supports, which is an extension of the Hodge structure on the cohomology ring ([M1], [M2]). The formality condition it imposes is in our case:

PROPOSITION 3.2. ([M1], 9.4) Let G be a Kähler group. The Lie algebra $Gr \mathscr{L}G$ is the quotient of a free Lie algebra by an ideal generated by sums of length two brackets.

This is equivalent to the Malcev and holonomy algebras of G being isomorphic (cf. [Ko]). For instance, the group $G = \langle x, y; ((x, y), y) \rangle$ cannot be Kähler because $\mathscr{L}G$ is the quotient of a free algebra by the ideal generated by a length three bracket $[[\bar{x}, \bar{y}], \bar{y}]$.

In this note we will look in another direction through the results of the preceding section. Johnson and Rees rule out in [JR] free groups as Kähler groups because they have finite index subgroups of odd rank. In this paragraph we establish this by studying cup products, and in this way the result holds for groups

with a free 2-step nilpotent Lie algebra \mathscr{L}_2 . This is a consequence of the Lefschetz decomposition and the nondegenerate alternating pairing Q that the real cohomology $H^*(X)$ of a compact Kähler manifold X supports. Our standard reference for these properties will be [W], Chap. V.

THEOREM 3.3. Let G be a finitely presented group such that $\mathcal{L}_2 G \cong \mathcal{L}_2 F_n$ for some n. Then G is not Kähler.

Proof. Suppose $\pi_1(X, *) \cong G$. As $\mathscr{L}_2 G \cong \mathscr{L}_2 F_n$, and using Example 1.4, $\dim \Gamma_1/\Gamma_2 G \otimes \mathbb{R} = \dim \Gamma_1/\Gamma_2 F_n \otimes \mathbb{R} = n$, and $\dim \Gamma_2/\Gamma_3 G \otimes \mathbb{R} = \dim \mathscr{L}_2 G^{(2)} = \dim \mathscr{L}_2 F_n = n(n-1)/2$. Thus by Cor. 2.10 ker $\cup = \bigwedge^2 H^1(X)$, all cup products are zero. But cohomology classes of rank one are primitive, and the pairing Q is non-degenerate (see [W], 5.6). This leads to a contradiction, hence G cannot be Kähler.

To use Theorem 3.3, one needs to know when $\mathscr{L}_2 G \cong \mathscr{L}_2 F_n$. This can be established from a presentation of G as we have shown in section §1 (see Remark 1.15). We will now remark the case of one-relator groups. The consequence for them of [M1], 9.4 is that if G admits a presentation $G = \langle x_1, \ldots, x_n; r \rangle$, then its single defining relation r does not lie in $\Gamma_3 F\{x_1, \ldots, x_n\}$. We complete this with

COROLLARY 3.4. Let $G = \langle x_1, \ldots, x_n; r \rangle$ be a Kähler group presented with a single relation. Then $r \in \Gamma_2 F\{x_1, \ldots, x_n\} \setminus \Gamma_3 F\{x_1, \ldots, x_n\}$.

Proof. By Corollary 1.13, $r \in \Gamma_2 F \setminus \Gamma_3 F$ is the only case in which $\mathscr{L}_2 G$ is not free.

Examples 3.5.

- (i) ([JR]) Free groups cannot be Kähler.
- (ii) The group G = ⟨x, y; ((x, y), y)⟩ cannot be Kähler because ((x, y), y) ∈ Γ₃F and therefore L₂G ≅ L₂F₂. This group was known to be non-Kähler by [M1], [M2].
- (iii) The group $G = \langle x, y, z, t; x^3y^{-4}z^2y, y^2z^2 \rangle$ is not Kähler because the two defining relations are linearly independent in $\Gamma_1/\Gamma_2 F \otimes \mathbb{R} \cong \mathbb{R}^4$, and thus by Cor. 1.14 $\mathscr{L}_2 G \cong \mathscr{L}_2 F_2$.
- (iv) The group $G = \langle x_1, \ldots, x_5; x_1 x_2^2 x_1, x_2 x_3^2 x_2, x_5 x_4^2 x_5 \rangle$ has also a linearly independent relation set, and thus $\mathcal{L}_2 G \cong \mathcal{L}_2 F_2$, and G cannot be Kähler either.
- (v) Compact Riemann surfaces provide examples showing that one-relator groups with a defining relation $r \in \Gamma_2 F/\Gamma_3 F$ are possible.

§4. One- and two-relator Kähler groups

In this chapter we give a lower bound for the number of defining relations that a presentation of a Kähler group $\pi_1 X$ must have, determined by the dimension of the Albanese image of X (Prop. 4.6). We apply this to fully determine the Malcev completions of one- and two-relator Kähler groups in Thm. 4.7. Our starting point 4.3 is due to F. Campana ([C]).

The Albanese variety and Albanese map are another feature of compact complex manifolds which is very useful to study its fundamental group. Let us briefly recall it.

PROPOSITION 4.1. Let X be a compact Kähler manifold, and let $q = \dim H^0(\Omega^1_X)$. There is a complex torus Alb(X) of dimension q (the Albanese torus) and a holomorphic map $\alpha_X : X \to Alb(X)$ (the Albanese map) such that α_X induces an isomorphism $H^1(Alb(X); \mathbb{Z}) \to H^1(X; \mathbb{Z})$. The pair $(Alb(X), \alpha_X)$ is determined up to isomorphism by this property. Moreover $\alpha_X(X)$ is a generating set for Alb(X) as an abelian group.

Let us fix our notation: Let X be compact Kähler, $\alpha_X \colon X \to Alb(X)$ its Albanese map, and denote $Y = \alpha_X(X)$ its image, which may be singular. We consider a desingularization $\varepsilon \colon \tilde{Y} \to Y$, and a desingularization \tilde{X} of the pullback of α_X :

It is clear that the manifold \tilde{X} is also compact Kähler, and that the map ε_X induces an isomorphism ε_{X*} : $\pi_1 \tilde{X} \to \pi_1 X$.

We will call the map $\tilde{\alpha}_X \colon \tilde{X} \to \tilde{Y}$ a smoothing of the Albanese map of X. The properties of the original Albanese map α_X relate X, \tilde{X} and \tilde{Y} for our purposes:

PROPOSITION 4.3. ([C]) Let X be compact Kähler, and $\tilde{\alpha}_X \colon \tilde{X} \to \tilde{Y}$ be a smoothing of the Albanese map of X. Then $\tilde{\alpha}_X$ induces an isomorphism $\mathscr{L}(\pi_1 \tilde{X}) \xrightarrow{\sim} \mathscr{L}(\pi_1 \tilde{Y})$.

Proof. (Cf. [C]) As ε_X induces an isomorphism of fundamental groups, hence ε_X^* : $H^1(X) \to H^1(\tilde{X})$ is also an isomorphism. This implies that Alb(X) is the Albanese torus of \tilde{X} and $\alpha_X \circ \varepsilon_X = \varepsilon \circ \tilde{\alpha}_X$ its Albanese map. As a consequence $\tilde{\alpha}_X^* \colon H^1(\tilde{Y}) \to H^1(\tilde{X})$ is onto. As α_X itself is also onto, $\tilde{\alpha}_X^*$ is also injective for H^* . Therefore $\tilde{\alpha}_X$ induces an isomorphism $H^1(\tilde{Y}) \cong H^1(\tilde{X})$ and an injection

 $H^2(\tilde{Y}) \hookrightarrow H^2(\tilde{X})$. Thus by universality of the 1-minimal model, $\tilde{\alpha}_X$ induces an isomorphism $M(2,0)(\tilde{Y}) \cong M(2,0)(\tilde{X})$; our statement is its dual.

Thus the study of Malcev completions of Kähler groups may be reduced to the study of smoothings of its Albanese images. This is specially convenient in the following case:

COROLLARY 4.4. ([C]) Let X be compact Kähler, with $q = \frac{1}{2}b_1(X) = 1$. Then the Malcev algebra of X is $\mathcal{L}\pi_1(X) \otimes \mathbb{R} \cong (\mathbb{R}^2, +)$.

Proof. The Albanese torus E = Alb(X) has dimension 1 in this case. Therefore α_X is onto, has smooth image and so equals $\tilde{\alpha}_X$.

The elliptic curve E has abelian fundamental group $\pi_1(E) \cong \mathbb{Z}^2$, thus our statement is a direct consequence of Prop. 4.3.

But rather than follow this line, we will derive from it consequences on $H^*(X)$, which will determine Malcev completions of Kähler groups with one or two defining relations.

LEMMA 4.5. Let X be compact Kähler, Y the Albanese image of X and $m = \dim_{\mathbb{C}} \tilde{Y}$. Then the graded algebra $H^*(X; \mathbb{C})$ contains a free graded exterior algebra $\bigwedge(V)$, where V is a complex vector space of dimension m and degree 1 spanned by holomorphic forms.

Proof. (cf. [Be] V. 18) Let $y \in Alb(X)$ be a regular point of the Albanese image $Y = \alpha_X(X)$. As dim Y = m, there are local coordinates u_1, \ldots, u_n of Alb(X) in a neighbourhood U of y such that $Y \cap U$ is defined as $u_{m+1} = 0, \ldots, u_n = 0$. The holomorphic forms du_1, \ldots, du_m are defined on U and, as Alb(X) is parallelizable, the forms in $\bigwedge (du_1, \ldots, du_m)$ extend to global holomorphic forms on Alb(X). Its direct image $\bigwedge (\alpha_X^* du_1, \ldots, \alpha_X^* du_m)$ defines a subalgebra of holomorphic cohomology classes in $H^*(X)$ which is free over y, hence is free.

The above Lemma together with the correspondence of 2.10 may be used to bound from below the number of defining relations for Kähler groups, and to study those admitting a one- or two-relation presentation.

PROPOSITION 4.6. Let G be a Kähler group, X a compact Kähler manifold such that $\pi_1 X \cong G$ and Y its Albanese image. Then:

(i) If dim Y = 1, there is an isomorphism $\mathscr{L}G \cong \mathscr{L}\pi_1 C_g$ with C_g a compact Riemann surface of genus g, induced by a group map $G \to \pi_1 C_g$.

 (ii) If dim Y = m > 1, dim ker(d₀: ℝr₁⊕····⊕ ℝr_s → Γ₁/Γ₂F⊗ ℝ) ≥ 2(^m₂) + 1. In particular, any presentation G = ⟨x₁,...,x_n; r₁,...,r_s⟩ must have defining relations r₁,...,r_k such that they form a basis of Im f and at least another 2(^m₂) + 1 defining relations.

Proof.

- (i) is just Prop. 4.3. with \tilde{Y} as C_g .
- (ii) By Lemma 4.5, the algebra $H^*(X; \mathbb{C})$ contains a free algebra $\bigwedge(V)$ generated by *m* linearly independent holomorphic 1-forms. By the Hodge structure of $H^*(X)$ it contains an isomorphic algebra $\bigwedge(\overline{V})$ spanned by *m* independent antiholomorphic 1-forms. Both algebras being free, one obtains the lower bound dim $\operatorname{Im} \cup : \bigwedge^2 H^1(X) \to H^2(X) \ge 2\binom{m}{2}$ considering either holomorphic or antiholomorphic products alone. Finally, due to the properties of the *Q* pairing in $H^1(X)$ ([*W*] 5.6), the product of a holomorphic 1-form with its conjugate cannot be zero, so dim $(\operatorname{Im} \cup) \cap H^{1,1}(X) \ge 1$. By the correspondence of 2.10 this produces the sought bound.

Remark. If X is compact Kähler and satisfies (i) in the above Prop. 4.6, it is not hard to check that the map $\pi_1 X = G \rightarrow \pi_1 C_g$ is onto. If the genus g is ≥ 2 , such groups G are examples of what we will call fibered Kähler groups, to be defined in 5.1.

We are now able to complete our study of compact Kähler groups with one or two defining relations begun in Cor. 3.4.

THEOREM 4.7. Let G be a Kähler group admitting a presentation with only one or two defining relations. Then either $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} = 0$ or $\mathcal{L}G \cong \mathcal{L}\pi_1 C_g$ with C_g a compact Riemann surface.

Proof. If $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} \neq 0$, then $\mathscr{L}_2 G \neq 0$, and by Prop. 4.6 any presentation of G must have at least $2\binom{\dim Y}{2} + 1$ defining relations, with Y = Alb(X). Thus the only possible case is dim Y = 1, and Prop. 4.6 (i) completes the proof.

Remark.

- The 1-relator groups G with $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} = 0$ are exactly the $G \cong \mathbb{Z}/n\mathbb{Z}$.
- The 2-relator groups G with $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} = 0$ are those with a presentation $\langle x_1, x_2; r_1, r_2 \rangle$ with \bar{r}_1, \bar{r}_2 linearly independent in $\Gamma_1/\Gamma_2 F\{x_1, x_2\}$. This is immediately derived from the exact sequence (1.5).

Examples 4.8. Denote C_g a compact Riemann surface of genus g.

(i) The group G defined in Ex. 3.5 (iii) can also be seen not to be Kähler by Thm. 4.7, as $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} \cong \mathbb{R}^2$ but $\mathscr{L}_2 G \not\cong \mathscr{L}_2 \pi_1(C_1)$.

(ii) The group G = ⟨x₁, x₂, x₃, x₄; (x₁x₂, x₃²), (x₁x₃x₁, x₄³)⟩ has a Malcev algebra which fulfills the quadratic presentation condition imposed by Morgan (Prop. 3.2). Yet G cannot be Kähler because Γ₁/Γ₂G ⊗ ℝ ≅ ℝ⁴ but dim Γ₂/Γ₃G ⊗ ℝ = 4 ≠ 5 = dim Γ₂/Γ₃π₁(C₂), contradicting Thm. 4.7.

§5. Nonfibered Kähler groups

Here we establish a dicothomy between fibered and nonfibered Kähler groups, arising from a result by A. Beauville and Y. T. Siu on the existence of irregular pencils on compact Kähler manifolds. We skip the fibered case, and we give in Prop. 5.6 an upper bound for dim $\Gamma_2/\Gamma_3 G \otimes \mathbb{R}$ in the case of nonfibered groups. This translates as a lower bound for the number of relations that their presentations must have.

Let $G = \pi_1(X, *)$ be a fundamental group. By Corollary 2.10 dim $\Gamma_2/\Gamma_3 G \otimes \mathbb{R} = \dim \bigwedge^2 H^1(X) - \dim \operatorname{Im}(\cup: \bigwedge^2 H^1(X) \to H^2(X))$. We have seen in §3 that if X is compact Kähler, $\operatorname{Im} \cup$ must be nonzero. Now we will establish a lower bound on its dimension in the case of nonfibered manifolds, by recalling a result of Casteln-uovo-De Franchis and its extension to arbitrary dimension.

DEFINITION 5.1. Let G be a Kähler group.

- (i) We call G a fibered Kähler group when $G = \pi_1(X, *)$ with X compact Kähler admitting a nonconstant holomorphic map $f: X \to C_g$, with C_g a compact Riemann surface of genus $g \ge 2$.
- (ii) We call G a nonfibered Kähler group when $G = \pi_1(X, *)$ with X compact Kähler not admitting any nonconstant holomorphic map to a compact Riemann surface of genus $g \ge 2$.

A. Beauville and Y. T. Siu independently proved that the above definitions make sense:

PROPOSITION 5.2. ([Be2], [Siu]) Let X be a compact Kähler manifold, write $G = \pi_1(X, *)$, and let $g \ge 2$ be an integer. Then X admits a nonconstant holomorphic map to a compact Riemann surface of a genus $h \ge g$ if and only if there is an epimorphic group morphism $G \rightarrow \pi_1(C_g, *)$ the fundamental group of a compact Riemann surface of genus g.

Prop. 5.2 means that a Kähler group G is either fibered or nonfibered, and that the former are characterised by admitting a $\pi_1(C_g)$ as a quotient.

Remark. If we have an onto map $G \to H \to 1$, it induces onto maps $\Gamma_n/\Gamma_{n+1}G \otimes \mathbb{R} \to \Gamma_n/\Gamma_{n+1}H \otimes \mathbb{R} \to 0$ for all *n*. This together with the fact that the

lower central series quotients of the $\pi_1 C_g$ have all nonzero rank shows that nilpotent or rationally nilpotent Kähler groups must be nonfibered.

We now study the cup products of 1-forms in the case of nonfibered compact Kähler manifolds. We begin with an extension of a classical result (see [Ca]):

PROPOSITION 5.3. (Castelnuovo-De Franchis) Let X be a compact Kähler manifold. If there exist ω_1, ω_2 linearly independent holomorphic 1-forms such that $\omega_1 \wedge \omega_2 = 0$ then there is a holomorphic map $f: X \to C$ with C a curve of genus $g(C) \ge 2$, such that ω_1, ω_2 belong to Im f^* .

Remark. The form equality $\omega_1 \wedge \omega_2 = 0$ is equivalent to $\omega_1 \wedge \omega_2$ being exact. This is a result of Hodge theory, showing that a nonzero holomorphic form over a compact Kähler manifold cannot be exact.

The Castelnuovo-De Franchis theorem together with the conic structure of the set of products in $H^{2,0}(X)$ yield the following corollary (see [BPV] IV, Prop. 4.2):

COROLLARY 5.4. If X is a nonfibered compact Kähler manifold, then dim $Im(\cup: \bigwedge^2 H^{1,0}(X) \to H^{2,0}(x)) \ge 2 \dim H^{1,0}(X) - 3.$

Cor. 5.4 gives a bound for the products of holomorphic 1-forms, and by conjugation, of antiholomorphic 1-forms. The dimension of products of holomorphic-antiholomorphic 1-forms has been bounded for compact complex surfaces in [BPV], IV, Prop. 4.3. We slightly alter their proof to extend it to compact Kähler manifolds of arbitrary dimension:

PROPOSITION 5.5. Let X be a nonfibered compact Kähler manifold. Then dim $Im(\cup: H^{1,0}(X) \otimes H^{0,1}(X) \to H^{1,1}(X)) \ge 2 \dim H^{1,0}(X) - 1.$

Proof. Denote $n = \dim X \ge 2$, $V = \operatorname{Im} \cup : H^{1,0}(X) \otimes H^{0,1}(X) \to H^{1,1}(X)$ and fix ω a fundamental Kähler form on X. We begin by showing that the pairing $\cup : H^{1,0}(X) \otimes H^{0,1}(X) \to V$ becomes injective when we fix a nonzero $\xi \in H^{1,0}(X)$ or $\bar{\eta} \in H^{0,1}(X)$.

Suppose there are holomorphic 1-forms ξ , η such that $\xi \wedge \bar{\eta} = d\alpha$. Then obviously $\xi \wedge \eta \wedge \bar{\xi} \wedge \bar{\eta} = d\alpha'$, and

$$\int_X \xi \wedge \eta \wedge \overline{\xi} \wedge \overline{\eta} \wedge \omega^{n-2} = 0$$

By the properties of the pairing Q of compact Kähler manifolds (see [W] 5.6), this implies that $\xi \wedge \eta = 0$, thus by the Castelnuovo-De Franchis theorem ξ and η are

linearly dependent. Take $\xi = a\eta$, with $a \in \mathbb{C}^*$. Then $0 = \xi \land \overline{\eta} = a\eta \land \overline{\eta}$. Again by the properties of the pairing Q, this means that $\xi, \eta = 0$.

Thus a map may be defined

$$\mathbb{P}(H^{1,0}(X)) \times \mathbb{P}(H^{0,1}(X)) \to \mathbb{P}(V)$$

with injective restrictions fixing a point in either factor of the source. We apply now the following result from [RV]:

PROPOSITION. Let $\varphi: \mathbb{P}^m(\mathbb{C}) \times \mathbb{P}^k(\mathbb{C}) \to \mathbb{P}^l(\mathbb{C})$ be a holomorphic mapping, with l < m + k. Then φ factors through one of the projections $\mathbb{P}^m \times \mathbb{P}^k \to \mathbb{P}^m$, $\mathbb{P}^m \times \mathbb{P}^k \to \mathbb{P}^k$.

In our case, \cup cannot factor through any of the projections because it is fiberwise injective in both cases, so it holds that dim $V \ge 2 \dim H^{1,0}(X) - 1$ as was wanted.

We have now all the required pieces to study $\Gamma_2/\Gamma_3 \otimes \mathbb{R}$ of nonfibered groups. We return to the notations defined in §1.

PROPOSITION 5.6. Let X be a nonfibered compact Kähler manifold with $q = \frac{1}{2} \dim H^1(X) = \frac{1}{2} \dim \Gamma_1/\Gamma_2 \pi_1(X, *) \otimes \mathbb{R}$. Then:

(i) If $q = 0, 1, \dim \Gamma_2 / \Gamma_3 \pi_1(X, *) \otimes \mathbb{R} = 0.$

(ii) If $q \ge 2$, dim $\Gamma_2/\Gamma_3 \pi_1(X, *) \otimes \mathbb{R} \le 2q^2 - 7q + 7$.

Proof. We have seen in Cor. 2.10 that dim $\Gamma_2/\Gamma_3\pi_1(X, *) \otimes \mathbb{R} = \dim \bigwedge^2 H^1(X) - \dim \operatorname{Im}(\cup: \bigwedge^2 H^1(X) \to H^2(X)) = (2q(2q-1)/2) - \dim \operatorname{Im} \cup.$

Thus if q = 0, dim $\Gamma_2/\Gamma_3 \pi_1(X, *) \otimes \mathbb{R} = 0$.

If q = 1, dim Im $\cup \leq 1$, so dim $\Gamma_2/\Gamma_3\pi_1(X, *) \otimes \mathbb{R} \leq 1$. Let a, b be a basis of $\Gamma_1/\Gamma_2\pi_1(X, *) \otimes \mathbb{R}$. The equality dim $\Gamma_2/\Gamma_3\pi_1(X, *) \otimes \mathbb{R} = 1$ would imply that $(a, b) \neq 0$ in $\mathscr{L}_2\pi_1(X, *)$ by Prop. 1.2 (iv). Therefore there would be an isomorphism $\mathscr{L}_2F_2 \xrightarrow{\approx} \mathscr{L}_2\pi_1(X, *)$ sending the generators X_1, X_2 of \mathscr{L}_2F_2 to a, b respectively. By Thm. 3.3 this would mean that $\pi_1(X, *)$ is not Kähler, leading to a contradiction. Hence our statement follows.

For $q \ge 2$, we break $H^{1}(X)$ into its Hodge components. By Cor. 5.4 $\dim(\operatorname{Im}(\bigwedge^{2}H^{1,0}(X) \to H^{2,0}(X))) \ge 2q - 3$. The same holds by conjugation for $\bigwedge^{2}H^{0,1}(X) \to H^{0,2}(X)$. Prop. 5.5 gives the inequality dim $\operatorname{Im}(H^{1,0}(X) \otimes H^{0,1}(X) \to H^{1,1}(X))) \ge 2q - 1$ and our statement follows from the addition of bounds. \Box

Prop. 5.6 roughly means that nonfibered Kähler groups need many defining relations. M. Green and R. Lazarsfeld give a bound ([GL], Thm. 5.4), establishing that given X nonfibered compact Kähler in the sense of Def. 5.1, that is admitting

no pencil of genus $g \ge 2$, and a presentation of its fundamental group $\pi_1(X) = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$, then

 $s \ge n-3$

Prop. 5.6 above allows us to establish a more accurate bound:

COROLLARY 5.7. Let $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$ be a finite group presentation. If $G = \pi_1(X)$, with X nonfibered compact Kähler, and writing $q = \frac{1}{2}b_1(X)$, the total number of relations must satisfy

- (i) If q = 0, $s \ge n$.
- (ii) If q = 1, $s \ge n 1$.
- (iii) If $q \ge 2$, $s \ge n + 4q 7$.

Proof. The group presentation $G = \langle x_1, \ldots, x_n; r_1, \ldots, r_s \rangle$ induces an exact sequence $1 \to N \to F \to G \to 1$ described in (1.3). Let $d_0: \bigoplus \mathbb{R}r_i \to J_F/J_F^2 \cong \Gamma_1/\Gamma_2 F \otimes \mathbb{R}$ be the map defined in Prop. 1.7. We may suppose the relations r_1, \ldots, r_s ordered so that the images of r_1, \ldots, r_k , with $k \leq s$ form a basis of $\operatorname{Im} d_0 \cong N/N \cap \Gamma_2 F \otimes \mathbb{R} \subseteq \Gamma_1/\Gamma_2 F \otimes \mathbb{R}$.

By Prop. 1.7 and Cor. 1.8 (i), dim $\Gamma_1/\Gamma_2 G \otimes \mathbb{R} = \dim \Gamma_1/\Gamma_2 F \otimes \mathbb{R} - \dim N/N$ $\cap \Gamma_2 F \otimes \mathbb{R}$, so there is an equality

$$n = k + 2q$$

Let us remark also that dim ker $d_0 = s - k$.

Thus if q = 0 we have $n = k \le s$ as was wanted. If q = 1, by Cor. 1.11 and Prop. 5.6 (i) we have that

$$0 = \dim \Gamma_2 / \Gamma_3 G \otimes \mathbb{R} = \binom{2}{2} - (s - k) + \dim \ker d_1$$

 $= 1 - s + k + \dim \ker d_1 \ge 1 - s + k$

As n = k + 2 in this case, this yields the sought bound. If n > 2 again by Cor 1.11 and Prop 5.6

If $q \ge 2$, again by Cor. 1.11 and Prop. 5.6,

dim
$$\Gamma_2/\Gamma_3 G \otimes \mathbb{R} = \binom{2q}{2} - (s-k) + \ker d_1 \le 2q^2 - 7q + 7$$

which implies

$$k + 2q + \dim \ker d_1 \le s - 4q + 7$$

and as n = k + 2q,

$$s \ge n + 4q - 7$$

Examples 5.8.

- (i) A group $G = \langle x_1, \ldots, x_{2q}; w_1, \ldots, w_s \rangle$ with $w_1, \ldots, w_s \in \Gamma_2 F$ can be nonfibered Kähler only if $s \ge 6q 7$ for $q \ge 2$, and $s \ge 1$ for q = 1.
- (ii) Chain link groups (see [Ro], 3.G) The group $G_{2q} = \langle x_1, \ldots, x_{2q}; (x_1, x_2), \ldots, (x_{2q-1}, x_{2q}), (x_{2q}, x_1) \rangle$ is the fundamental group of a link of 2q circumferences forming a circular chain, for $q \ge 2$. This group verifies $k = \dim \Gamma_1/\Gamma_2 F \otimes \mathbb{R} \dim \Gamma_1/\Gamma_2 G_{2q} \otimes \mathbb{R} = 0$, and s = 2q < 6q 7, and therefore G_{2q} cannot be nonfibered Kähler. Broadly speaking, if a link is not very intertwined, its group is not going to be nonfibered Kähler. The group G_4 verifies that dim $\Gamma_2/\Gamma_3 G_4 \otimes \mathbb{R} = 2$, and therefore it cannot be fibered Kähler either, as it cannot map onto $\pi_1(C_g, *)$ for any $g \ge 2$. The groups G_{2q} with $q \ge 3$ do admit onto mappings to $\pi_1(C_2, *)$, and the author does not know if they are fibered Kähler.
- (iii) The fundamental group of a compact Riemann surface of genus $g \ge 2$ admits a presentation $\langle a_1, \ldots, a_g, b_1, \ldots, b_g; (a_1, b_1) \cdots (a_g, b_g) \rangle$. In this presentation k = 0, and s = 1 < 6g - 7. Therefore, it can only be the fundamental group of a fibered Kähler manifold. This is a particular case of Prop. 5.2.
- (iv) Let $G = \langle x_1, \ldots, x_5; x_1^2 x_2^{-2} x_4^2, (x_1, x_2), (x_2, x_3), (x_3, x_4), (x_4, x_5) \rangle$. In this case n = 5, k = 1, q = 2 as $\operatorname{Im} d_0 = \langle 2\bar{x}_1 2\bar{x}_2 + 2\bar{x}_4 \rangle$, and s = 5 < n + 4q 7 = 6. Therefore G cannot be nonfibered Kähler. The group G cannot either map onto $\pi_1(C_g)$, with C_g a smooth projective curve of genus $g \ge 2$ because dim $\Gamma_2/\Gamma_3 G \otimes \mathbb{R} = 2$, dim $\Gamma_2/\Gamma_3 \pi_1(C_g) \otimes \mathbb{R} = (2g(2g-1)/2) 1 \ge 5$, so we reach the conclusion that G cannot be Kähler.

REFERENCES

- [Be] A. BEAUVILLE, Surfaces algébriques complexes (3^{me} Ed.), Astérisque 54, S.M.F. 1978.
- [Be2] A. BEAUVILLE, Appendix to [Ca].
- [BG] A. BOUSFIELD, V. GUGENHEIM, On PL De Rham theory and rational homotopy type, Mem. Amer. Math. Soc. 179, A.M.S. 1976.
- [BPV] W. BARTH, C. PETERS, A. VAN DE VEN, Compact complex surfaces, Ergeb. Math. Grenzgeb. (3) 4, Springer-Verlag 1984.
- [C] F. CAMPANA, Remarques sur les groupes de Kähler nilpotents, C. R. Acad. Sci. Paris 317 (1993) 777-780; to appear in Ann. Sci. École Norm. Sup.
- [Ca] F. CATANESE, Moduli and classification of irregular Kaehler manifolds (and algebraic varieties) with Albanese general type fibrations, Invent. Math. 104 (1991) 263-289.
- [Ch] K. T. CHEN, Iterated integrals of differential forms and loop space cohomolgy, Ann. of Math. 97 (1973) 217-246.
- [DGMS] P. DELIGNE, P. GRIFFITHS, J. MORGAN, D. SULLIVAN, Real Homotopy Theory of Kähler Manifolds, Invent. Math. 29 (1975) 245-274.

JAUME AMORÓS

- [GL] M. GREEN, R. LAZARSFELD, Higher obstructions to deforming cohomology groups of line bundles, J. Amer. Math. Soc. 4 (1991) 87-103.
- [G] M. GROMOV, Sur le groupe fondamental d'une variété Kählérienne, C. R. Acad. Sci. Paris 308 (1989) 67-70.
- [GM] P. GRIFFITHS, J. MORGAN, Rational Homotopy Theory and Differential Forms, Progr. Math. 16, Birkhäuser 1981.
- [HMR] P. HILTON, G. MISLIN, J. ROITBERG, Localization of Nilpotent Groups and Spaces, Math. Studies 15, North Holland 1975.
- [JR] F. JOHNSON, E. REES, On the fundamental group of a complex algebraic manifold, Bull. London Math. Soc. 19 (1987) 463-466.
- [JR2] F. JOHNSON, E. REES, The fundamental groups of algebraic varieties, in S. Jackowsky et altri (Eds.) Algebraic Topology Poznan 1989, Lecture Notes in Math. 1474, Springer-Verlag 1991, 75-82.
- [Ko] T. KOHNO, Série de Poincaré-Koszul associée aux groupes de tresses pures, Invent. Math. 82 (1985) 57-75.
- [L] M. LAZARD, Sur les groupes nilpotents et les anneaux de Lie, Ann. scient. Éc. Norm. Sup. 71 (1954) 101–190.
- [M1] J. MORGAN, The algebraic topology of smooth algebraic varieties, Inst. Hautes Études Sci. Publ. Math. 48 (1978) 137-204.
- [M2] J. MORGAN, Hodge theory for the algebraic topology of smooth algebraic varieties, Proc. Sympos. Pure Math. 32 AMS 1978, 119-128.
- [Q] D. QUILLEN, Rational Homotopy Theory, Ann. of Math. (2) 90 (1969) 205-295.
- [Q2] D. QUILLEN, On the Associated Graded Ring of a Group Ring, J. Algebra 10 (1968) 411-418.
- [RV] R. REMMERT, A. VAN DE VEN, Zur Funktionentheorie homogener komplexer Mannigfaltigkeiten, Topology 2 (1963) 137-157.
- [Ro] D. ROLFSEN, Knots and Links, Math. Lecture Ser. 7, Publish or Perish 1990.
- [S] J.-P. SERRE, Lie algebras and Lie groups, W. A. Benjamin 1965.
- [Siu] Y. T. SIU, Strong Rigidity for Kähler Manifolds and the Construction of Bounded Holomorphic Functions, in R. Howe (Ed.), *Discrete Groups in Geometry and Analysis*, Progr. Math. 67 Birkhäuser 1987, 124–151.
- [St] J. R. STALLINGS, Quotients of the powers of the augmentation ideals in a group ring, in L. P.. Neuwirth, Knots, Groups and 3-Manifolds, Ann. of Math. Stud. 84, Princeton University Press 1975, 101-118.
- [WO] R. O. WELLS, Comparison of De Rham and Dolbeault cohomology for proper surjective mappings, Pacific J. Math. 53 (1974) 281-300.
- [W] R. O. WELLS, Differential Analysis on Complex Manifolds, Graduate Texts in Math. 65, Springer-Verlag 1980.

Universitat de Barcelona

Dep. d'Algebra i Geometria-Fac. de Matemàtiques Gran Via 585 E-08007 Barcelona Spain amoros@cerber.mat.ub.es Current address: Universitat Politècnica de Catalunya Departament de Matemàtica Aplicada I, ETSEIB Diagonal 647 E-08028 Barcelona (Spain) amoros@ma1.upc.es

Received August 11, 1994; June 23, 1995