

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 71 (1996)

**Artikel:** Compactifying coverings of 3-manifolds.  
**Autor:** Mihalik, Michael L.  
**DOI:** <https://doi.org/10.5169/seals-53848>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 20.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Compactifying coverings of 3-manifolds

MICHAEL L. MIHALIK

**Abstract.** If a finitely presented group  $G$  is negatively curved, automatic or asynchronously automatic then  $G$  has an asynchronously bounded “almost prefix closed” combing. Results in [Br<sub>1</sub>] and [E] imply that the fundamental group of any closed 3-manifold satisfying Thurston’s geometrization conjecture has an asynchronously bounded, almost prefix closed combing.

**MAIN THEOREM.** If  $M$  is a compact  $P^2$ -irreducible 3-manifold,  $\pi_1(M)$  has an asynchronously bounded, almost prefix closed combing, and  $H$ , a subgroup of  $\pi_1(M)$ , is quasiconvex with respect to this combing, then the cover of  $M$  corresponding to  $H$  is a missing boundary manifold.

### §1. Introduction

A 3-manifold  $M$  is a *missing boundary manifold* if there is a compact manifold  $\bar{M}$ , and a subset  $K$  of the boundary of  $\bar{M}$  such that  $\bar{M} - K$  is homeomorphic to  $M$ . If  $M$  is a compact  $P^2$ -irreducible 3-manifold and  $\hat{M}$  is the cover of  $M$  corresponding to a finitely generated subgroup of  $\pi_1(M)$ , then J. Simon has conjectured that  $\hat{M}$  is a missing boundary manifold (see [Si]).

In §2, we define a geometric group theory notion “1-tame”, for pairs of groups, motivated by the combing ideas for groups introduced in [MT].

The connection between compactifying 3-manifolds and 1-tame pairs is described by the following result, proved in §4.

If  $G$  is the fundamental group of a compact  $P^2$ -irreducible 3-manifold  $M$ , and  $H$  is a finitely generated subgroup of  $G$  then, the pair  $(H, G)$  is 1-tame if and only if the cover of  $M$  corresponding to  $H$  is a “missing boundary” manifold.

In [Si] Simon verified his conjecture for  $M$  any compact  $P_2$ -irreducible 3-manifold and  $H$  the fundamental group of a boundary component of  $M$ . In [J] W. Jaco generalized this to the case where  $H$  is a finitely generated peripheral subgroup of  $\pi_1(M)$ . W. Thurston showed in [T] that if  $M$  admits a geometrically finite, complete hyperbolic structure of infinite volume, then  $M$  is a missing boundary manifold. F. Bonahon [Bo] showed that any hyperbolic-3-manifold  $M$  with finitely generated fundamental group is a missing boundary manifold, provided that  $\pi_1(M)$  is not a

free product. In [HRS] J. Hass, H. Rubinstein and P. Scott show that if  $M$  is a closed  $P^2$ -irreducible 3-manifold such that  $\pi_1(M)$  contains a subgroup  $A$  isomorphic to  $\mathbb{Z} \times \mathbb{Z}$ , then the cover of  $M$  with fundamental group  $A$  is a missing boundary manifold.

Our results on the Simon conjecture are obtained through an examination of combings of groups. If a finitely presented group  $G$  is word hyperbolic, automatic or asynchronously automatic then  $G$  has an asynchronously bounded “almost prefix closed” combing. Results in [Br<sub>1</sub>] and [E] imply that the fundamental group of any closed 3-manifold satisfying Thurston’s geometrization conjecture has an asynchronously bounded, almost prefix closed combing.

Our main result in this paper is

**MAIN THEOREM.** *If  $M$  is a compact  $P^2$ -irreducible 3-manifold,  $\pi_1(M)$  has an asynchronously bounded, almost prefix closed combing, and  $H$ , a subgroup of  $\pi_1(M)$ , is quasiconvex with respect to this combing, then the cover of  $M$  corresponding to  $H$  is a missing boundary manifold.*

M. Bridson has observed that if  $G$  is the 3-dimensional Heisenberg group then any finitely generated subgroup of  $G$  is quasiconvex with respect to some asynchronously bounded almost prefix closed combing. If  $G$  is a CAT(0) group acting properly discontinuously, cocompactly and by isometries on the geodesic metric space  $X$  and  $H$  is an abelian subgroup of  $G$  then by the “Flat Torus Theorem”,  $H$  acts on an  $n$ -flat in  $X$  and so  $H$  is quasiconvex with respect to geodesics in  $X$ . In particular, if  $G$  is a knot or link group (then  $G$  is CAT(0) see [BH] and [L]) and  $H$  a peripheral subgroup then  $H$  is quasiconvex in a bounded almost prefix closed combing of  $G$ . More generally, if  $M$  is a compact  $P_2$ -irreducible 3-manifold satisfying Thurston’s geometrization conjecture and  $H$  the fundamental group of a boundary component of  $M$ , then  $H$  is quasiconvex in some bounded almost prefix closed combing of  $\pi_1(M)$  (see [Br<sub>2</sub>]). Combining this with our main theorem gives Simon’s result [Si].

In [R], L. Reeves shows that cubed 3-manifolds have biautomatic fundamental groups. These manifolds have canonical immersed surfaces and the corresponding surface groups are quasiconvex (rational) in the fundamental group of the cubed manifolds with respect to the biautomatic structure. Hence (by the above theorem) the covers of cubed manifolds corresponding to these surface groups are missing boundary manifolds.

In [GS] S. Gersten and H. Short study the quasiconvex subgroups of biautomatic groups.

In the case that  $G$  is the fundamental group of a closed irreducible 3-manifold  $M$ , the pair  $(G, H)$  is 1-tame and  $H$  is the trivial group, results in [BT], [MT] and

[Tu] imply that the universal cover of  $M$  is homeomorphic to  $\mathbf{R}^3$ . It is an easy matter to see that if  $G$  is “almost convex” then the pair  $(G, 1)$  is 1-tame. This gives a different proof of the main result of V. Poénaru’s paper [P]. The relation between A. Casson’s  $C_2$  condition, the  $QSF$  condition developed by S. Brick, S. Gersten and J. Stallings and tame pairs  $(G, 1)$  is discussed in [MT].

## §2. Definitions, and statements of results

While our main interest in this paper is in compactifying covers of compact 3-manifolds we will work with more general spaces, namely polyhedra. If  $T$  is a triangulation of the pair of polyhedra  $(X, C)$ , then the largest subcomplex of the first barcentric subdivision of  $T$ , whose realization is contained in  $X - C$ , is a strong deformation retract of  $X - C$ . This can be used to show,

LEMMA 1. *If  $(X, C)$  is a pair of polyhedra with  $C$  compact, then the following are equivalent.*

- (1)  $\pi_1(X - C)$  is finitely generated (I.e. each component of  $X - C$  has finitely generated fundamental group).
- (2)  $\pi_1(Cl(X - C))$  is finitely generated.
- (3) For each component  $K$  of  $X - C$ ,  $\pi_1(Cl(K))$  is finitely generated.
- (4) For any vertex  $v \in X - C$ , there is a finite subcomplex  $B$  of  $X$  containing  $v$  such that any loop in  $X - C$ , based at  $v$ , is homotopic rel  $\{0, 1\}$  to a loop in  $B$  by a homotopy in  $X - C$ .

By (4), if  $C_1 \subset C_2$  are finite subcomplexes of  $X$  and  $\pi_1(X - C_2)$  is finitely generated then  $\pi_1(X - C_1)$  is finitely generated. So to show that  $\pi_1(X - C)$  is finitely generated for all finite subcomplexes  $C$  of  $X$ , it suffices to show  $\pi_1(X - C)$  is finitely generated for sufficiently large  $C$ .

DEFINITION. *Let  $X$  be a compact polyhedra,  $H$  a finitely generated subgroup of  $\pi_1(X)$  and  $*$  a vertex of  $\tilde{X}$  ( $\equiv$  the universal cover of  $X$ ). The pair  $(X, H)$  is 1-tame if for each integer  $N$  there is an integer  $M$  such that for any edge path  $\alpha$  in  $Cl(\tilde{X} - St^N(H*))$  with  $\alpha(0), \alpha(1) \in St^N(H*)$ ,  $\alpha$  is homotopic rel  $\{0, 1\}$  to an edge path  $\beta$  in  $St^M(H*)$ , by a homotopy in  $Cl(\tilde{X} - St^N(H*))$ .*

This definition is easily seen to be independent of base point  $*$ , and triangulation of  $\tilde{X}$ .



**THEOREM 1.** *If  $X_1$  and  $X_2$  are compact polyhedra,  $H_1$  is a finitely generated subgroup of  $\pi_1(X_1)$  and  $f(\pi_1(X_1), H_1) \rightarrow (\pi_1(X_2), H_2)$  is an isomorphism of pairs then  $(X_1, H_1)$  is 1-tame if and only if  $(X_2, H_2)$  is 1-tame.*

**DEFINITION.** *If  $G$  is a finitely presented group and  $H$  a finitely generated subgroup of  $G$ , then the pair  $(G, H)$  is 1-tame if for some (any) finite polyhedra  $X$ , with  $\pi_1(X) = G$ , we have  $(X, H)$  is 1-tame.*

**DEFINITION.** *A 3-manifold  $M$  is a missing boundary manifold if there is a compact manifold  $\bar{M}$ , and a subset  $K$  of the boundary of  $\bar{M}$  such that  $\bar{M} - K$  is homeomorphic to  $M$ .*

In [Tu], T. Tucker shows that if  $M$  is a noncompact  $P^2$ -irreducible 3-manifold, and for each finite subcomplex  $C$  of (some triangulation of)  $M$ ,  $\pi_1(M - C)$  is finitely generated then  $M$  is a missing boundary manifold.

**THEOREM 2.** *If  $X$  is a finite polyhedra with  $\pi_1(X) = G$  and  $H$  is a finitely generated subgroup of  $G$ , then  $(G, H)$  is 1-tame iff for each finite subcomplex  $C$  of  $H \setminus \tilde{X}$ ,  $\pi_1((H \setminus \tilde{X}) - C)$  is finitely generated.*

**COROLLARY 3.** *If  $M$  is a compact  $P^2$ -irreducible 3-manifold and  $H$  is a finitely generated subgroup of  $\pi_1(M)$  then  $H \setminus \tilde{X}$  is a missing boundary manifold iff  $(\pi_1(M), H)$  is 1-tame.*

In the next section we define combings of groups.

If a finitely presented group  $G$  is negatively curved, automatic or asynchronously automatic then  $G$  has an asynchronously bounded “almost prefix closed” combing. Results in [Br<sub>1</sub>] and [E] imply that the fundamental group of any closed 3-manifold satisfying Thurston’s geometrization conjecture has an asynchronously bounded, almost prefix closed combing.

Our main result is an easy corollary of the following more general result.

**THEOREM 4.** *If a group  $G$  has an asynchronously bounded, almost prefix closed combing, and  $H$ , a subgroup of  $G$ , is quasiconvex with respect to this combing, then the pair  $(G, H)$  is 1-tame.*

### §3. Algebraic preliminaries

Our approach to combings is a geometric one. If  $A$  is a finite set of generators for a group  $G$ , then  $\Gamma(G, A)$ , the Cayley graph of  $G$  with respect to  $A$ , is a

1-complex with 0-skeleton equal to  $G$  and a directed edge from vertex  $v$  to vertex  $w$  if  $va = w$  for some  $a \in A$ . Making each edge of  $\Gamma$  isometric to the unit interval gives a metric  $d$ , on  $\Gamma$ . Let  $*$  be the identity vertex of  $\Gamma$ . A *combing* of  $G$  (with respect to  $A$ ) is a set of edge paths  $\{\rho_v: [0, T_v] \rightarrow \Gamma\}$  such that  $v \in G$ ,  $\rho_v(0) = *$ ,  $\rho_v(T_v) = v$ . As a convenience  $\rho_v$  is extended to  $[0, \infty)$  by setting  $\rho_v(x) = v$  for all  $x > T_v$ . A combing is *asynchronously bounded* if there is an integer  $K$  such that for each pair of adjacent vertices  $u, v$  in  $\Gamma$ , there are non-decreasing surjections  $\alpha, \beta: [0, \infty) \rightarrow [0, \infty)$  each of which, when restricted to an interval between adjacent integers, is either constant or an isometry and so that  $d(\rho_u(\alpha(x)), \rho_v(\beta(x))) \leq K$  for all  $x \in [0, \infty)$ . We call  $(\alpha, \beta)$  an *asynchronous adjustment* for  $(\rho_u, \rho_v)$ . The constant  $K$  is called the *asynchronous fellow traveler constant*.

A combing  $\{\rho_v\}$  of  $G$  (with respect to  $A$ ) defines a map  $r$  of  $G$  into  $A^*$ , the free monoid. The group  $G$  is *asynchronously automatic* if  $\{\rho_v\}$  is asynchronously bounded and  $r(G)$  is the language accepted by some finite state automaton.

A combing  $\{\rho_v\}$  is *almost prefix closed* if there is an integer  $K$  such that for each  $v \in G$  and integer  $n \in [0, T_v]$  there is a  $w \in G$  such that  $\rho_w(T_w) \in St^K(\rho_v(n))$  and the prefix  $\rho_v([0, n])$  is a subset of  $St^K(im(\rho_w))$ .

Say  $G$  is asynchronously automatic. Then  $r(G) \subset A^*$  is the accept language of a finite state automaton. Let  $K$  be an integer so that if  $S$  is a state of this automaton that can be joined to an accept state, then it can be joined by a path of length  $\leq K$ . Then any prefix of a combing path given by  $r$  can be extended by  $K$  or fewer edges to an actual combing path. Hence any asynchronously automatic group has an asynchronously bounded, almost prefix closed combing. By Theorem 8.28 of [E] the fundamental group of a Haken 3-manifold need not be asynchronously automatic.

If  $G$  is a finitely generated group with combing  $\{\rho_v\}$ , then the subgroup  $H$  of  $G$  is *quasi-convex* with respect to  $\{\rho_v\}$  if there is an integer  $L$  such that for each  $h \in H$ ,  $im(\rho_h) \subset St^L(H) \subset \Gamma(G, A)$ .

#### §4. Proofs of Theorems 1 and 2

The following lemmas are straightforward.

**LEMMA 4.1.** *If  $X$  is a finite simplicial complex and  $X^{(2)}$  its 2-skeleton, then  $(X, H)$  is 1-tame iff  $(X^{(2)}, H)$  is 1-tame.*

Hence we need only consider the case  $X_1$  and  $X_2$  are 2-dimensional simplicial complexes.

LEMMA 4.2. *If  $(X_1, *_1)$  and  $(X_2, *_2)$  are finite complexes such that  $(\pi_1(X_1, *_1), H_1)$  and  $(\pi_1(X_2, *_2), H_2)$  are isomorphic pairs of groups, then there are maps  $f_1: (X_1, *_1) \rightarrow (X_2, *_2)$  and  $f_2: (X_2, *_2) \rightarrow (X_1, *_1)$  such that:*

- (i)  $(f_1 \circ f_2)_\#$  and  $(f_2 \circ f_1)_\#$ , the induced maps on fundamental groups, are the identity.
- (ii)  $(f_1)_\#(H_1) = H_2$  (and so  $(f_2)_\#(H_2) = H_1$ ).
- (iii) For each edge  $e$  of  $X_1(X_2)$ ,  $f_1(e)(f_2(e))$  is an edge path.

LEMMA 4.3. *Say  $f_i, X_i, H_i$  and  $*_i$  are as in Lemma 4.2. Let  $\tilde{f}_1$  be the lift of  $f_1$  to  $(\tilde{X}_1, \tilde{*}_1)$  that takes  $\tilde{*}_1$  to  $\tilde{*}_2$  and let  $\tilde{f}_2$  be the lift of  $f_2$  to  $(\tilde{X}_2, \tilde{*}_2)$  that takes  $\tilde{*}_2$  to  $\tilde{*}_1$ . Then there is an integer  $K_1$  such that if  $x \in \tilde{X}_2$ , then  $\tilde{f}_1 \circ \tilde{f}_2(x) \in \text{St}^{K_1}(x)$  and, if  $x \in \tilde{X}_1$  then  $\tilde{f}_2 \circ \tilde{f}_1(x) \in \text{St}^{K_1}(x)$ .*

### Proof of Theorem 1

Assume the notation of Lemma 4.3, and that  $(X_2, H_2)$  is 1-tame. The group  $\pi_1(X_i, *_i)$  acts on the left of  $\tilde{X}_i (i \in \{1, 2\})$ . By elementary covering space theory, if  $g_i \in \pi_1(X_i, *_i)$ , then  $\tilde{f}_1(g_1 \tilde{*}_1) = ((f_1)_\#(g_1)) \tilde{*}_2$ , and  $\tilde{f}_2(g_2 \tilde{*}_2) = ((f_2)_\#(g_2)) \tilde{*}_1$ . In particular,  $\tilde{f}_2 \circ \tilde{f}_1(g_1 \tilde{*}_1) = g_1 \tilde{*}_1$ ,  $\tilde{f}_1 \circ \tilde{f}_2(g_2 \tilde{*}_2) = g_2 \tilde{*}_2$ ,  $\tilde{f}_1(H_1 \tilde{*}_1) = H_2 \tilde{*}_2$  and  $\tilde{f}_2(H_2 \tilde{*}_2) = H_1 \tilde{*}_1$ . Note that if  $K_2$  is an integer larger than the length of the edge path  $\tilde{f}_i(e)$  for all edges  $e$  in  $X_i$ , then for any edge  $e$  in  $\tilde{X}_1(\tilde{X}_2)$ ,  $\tilde{f}_2 \tilde{f}_1(e)(\tilde{f}_1 \tilde{f}_2(e))$  has length  $\leq K_2^2$ .

Now choose an integer  $K_3$  such that any edge loop in  $\tilde{X}_1$  or  $\tilde{X}_2$  of length  $\leq 2K_1 + K_2^2 + 1$  is homotopically trivial in  $\text{St}^{K_3}(v)$  for any vertex  $v$  of the loop.

Let  $A$  be an integer and  $\gamma$  an edge path in  $\text{Cl}(\tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1))$  such that  $\gamma(0), \gamma(1)$  are elements of  $\text{St}^A(H_1 \tilde{*}_1)$ . Then  $\gamma$  decomposes into edge subpaths each of which is either a path in  $\text{St}^{A+K_3}(H_1 \tilde{*}_1)$  or an edge path  $\alpha$  such that  $\alpha(0), \alpha(1)$  are elements of  $\text{St}^{A+K_3}(H_1 \tilde{*}_1)$  and  $\text{im}(\alpha) \subset \text{Cl}(\tilde{X}_1 - \text{St}^{A+K_3}(H_1 \tilde{*}_1))$ . To see that  $(X_1, H_1)$  is 1-tame, it suffices to find an integer  $J$  so that any such  $\alpha$  is homotopic rel $\{0, 1\}$  to an edge path in  $\text{St}^J(H_1 \tilde{*}_1)$  by a homotopy in  $\text{Cl}(\tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1))$ .

Let  $\alpha$  be as above. Let  $v_0$  be the initial point of  $\alpha$  and  $v_1$  the end point of  $\alpha$ . Choose  $\delta_0(\delta_1)$ , an edge path of length  $\leq K_1$  from  $v_0$  to  $\tilde{f}_2 \circ \tilde{f}_1(v_0)$  ( $v_1$  to  $\tilde{f}_2 \circ \tilde{f}_1(v_1)$ ). By the definition of  $K_3$ ,  $\alpha$  is homotopic rel $\{0, 1\}$  to  $\langle \delta_0, \tilde{f}_2 \circ \tilde{f}_1(\alpha), \delta_1^{-1} \rangle$  by a homotopy in  $\text{St}^{K_3}(\text{im}(\alpha)) \subset \text{Cl}(\tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1))$ . We have reduced our problem to showing:

There is an integer  $J$  such that for any  $\alpha$ , as above,  $\tilde{f}_2 \circ \tilde{f}_1(\alpha)$  is homotopic rel $\{0, 1\}$  to an edge path in  $\text{St}^J(H_1 \tilde{*}_1)$  by a homotopy in  $\text{Cl}(\tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1))$ .

If  $N$  is a positive integer choose  $L_1(N)$  such that  $\tilde{f}_1(\text{St}^N(\tilde{*}_1)) \subset \text{St}^{L_1(N)}(\tilde{*}_2)$  and  $L_2(N)$  such that  $\tilde{f}_2(\text{St}^N(\tilde{*}_2)) \subset \text{St}^{L_2(N)}(\tilde{*}_1)$ . Then  $\tilde{f}_1(\text{St}^N(H_1 \tilde{*}_1)) \subset \text{St}^{L_1(N)}(H_2 \tilde{*}_2)$  and

$\tilde{f}_2(\text{St}^N(H_2 \tilde{*}_2)) \subset \text{St}^{L_2(N)}(H_1 \tilde{*}_1)$ . Furthermore, we may assume that  $L_i$  is an increasing function.

*Note* that if  $x \in \tilde{X}_2 - \text{St}^{K_1+L_1(N)}(H_2 \tilde{*}_2)$ , then  $\tilde{f}_2(x) \in \tilde{X}_1 - \text{St}^N(H_1 \tilde{*}_1)$ . (If  $\tilde{f}_2(x) \in \text{St}^N(H_1 \tilde{*}_1)$ , then  $\tilde{f}_1 \tilde{f}_2(x) \in \text{St}^{L_1(N)}(H_2 \tilde{*}_2)$ , but  $x \in \text{St}^{K_1}(\tilde{f}_1 \circ \tilde{f}_2(x))$  implies that  $x \in \text{St}^{K_1+L_1(N)}(H_2 \tilde{*}_2)$ .) Similarly if  $x \in \tilde{X}_1 - \text{St}^{K_1+L_2(N)}(H_1 \tilde{*}_1)$ , then  $\tilde{f}_1(x) \in \tilde{X}_2 - \text{St}^N(H_2 \tilde{*}_2)$ .

As  $\tilde{f}_1(\text{St}^{A+K_3}(H_1 \tilde{*}_1)) \subset \text{St}^{L_1(A+K_3)}(H_2 \tilde{*}_2)$ , we have  $\tilde{f}_1(\alpha(0))$  and  $\tilde{f}_1(\alpha(1))$  in  $\text{St}^{L_1(A+K_3)}(H_2 \tilde{*}_2)$ . Decompose  $\tilde{f}_1(\alpha)$  as  $\langle \beta_1, \alpha_1, \beta_2, \alpha_2, \dots, \beta_n, \alpha_n, \beta_{n+1} \rangle$  where  $\alpha_i$  and  $\beta_i$  are sub-edge paths of  $\tilde{f}_1(\alpha)$ , such that  $\text{im}(\beta_i) \subset \text{St}^{K_1+L_1(A+K_3)}(H_2 \tilde{*}_2)$  and  $\text{im}(\alpha_i) \subset \text{Cl}(\tilde{X}_2 - \text{St}^{K_1+L_1(A+K_3)}(H_2 \tilde{*}_2))$  with  $\alpha_i(0)$  and  $\alpha_i(1)$  in  $\text{St}^{K_1+L_1(A+K_3)}(H_2 \tilde{*}_2)$ , for all  $i$ . Since  $(X_2, H_2)$  is 1-tame, there is an integer  $M(K_1 + L_1(A + K_3))$  and for each  $i$ , a homotopy  $F_i$  of  $\alpha_i$ ,  $\text{rel}\{0, 1\}$ , to an edge path  $\gamma_i$ , in  $\text{St}^M(H_2 \tilde{*}_2)$  such that  $\text{im}(F_i) \subset \text{Cl}(\tilde{X}_2 - \text{St}^{K_1+L_1(A+K_3)}(H_2 \tilde{*}_2)) \subset \tilde{X}_2 - \text{St}^{K+1+L_1(A)}(H_2 \tilde{*}_2)$ . By the above note,  $\text{im}(\tilde{f}_2 \circ F_i) \subset \tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1)$ .

Let  $M_1 = \max\{M(K_1 + L_1(A + K_3)), K_1 + L_1(A + K_3)\}$ . Then  $\tilde{f}_2 \circ \tilde{f}_1(\alpha)$  is homotopic  $\text{rel}\{0, 1\}$  to an edge path in  $\tilde{f}_2(\text{St}^{M_1}(H_2 \tilde{*}_2)) \subset \text{St}^{L_2(M_1)}(H_1 \tilde{*}_1)$  by a homotopy in  $\tilde{X}_1 - \text{St}^A(H_1 \tilde{*}_1)$ , and we can select  $J = L_2(M_1)$ .

Similarly if  $(X_1, H_1)$  is 1-tame, we have  $(X_2, H_2)$  is 1-tame.  $\square$

*Proof of Theorem 2.* Say  $(G, H)$  is 1-tame. Let  $*$  be a vertex of  $X$  and  $\tilde{*}$  a vertex of  $\tilde{X}$  over  $*$ . Let  $q: \tilde{X} \rightarrow H \setminus \tilde{X}$  be the quotient map and let  $\hat{*} = q(\tilde{*})$ . To help distinguish which space a star occurs in, we use  $\text{St}_H^N$  for the  $N$ -th star in  $H \setminus \tilde{X}$ . If  $N$  is a positive integer and  $v$  is a vertex of  $\text{Bd}(\text{St}_H^N(\hat{*}))$  then we show there is an integer  $M(N)$  such that any loop at  $v$  in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  is homotopic  $\text{rel}\{0, 1\}$  to a loop in  $\text{St}_H^M(\hat{*})$ , by a homotopy in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ . As  $\pi_1(\text{Cl}(\text{St}_H^M(\hat{*}) - \text{St}_H^N(\hat{*})))$  is finitely generated, we will have  $\pi_1(\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*})))$  is finitely generated. Let  $\alpha$  be an edge loop at  $v$  with image in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ . Choose  $\tilde{v} \in \tilde{X}$  over  $v$  and let  $\tilde{\alpha}$  be the lift of  $\alpha$  to  $\tilde{v}$ . Observe that for any integer  $K$ ,  $q^{-1}(\text{St}_H^K(\hat{*})) = \text{St}^K(H \tilde{*})$ . Hence  $\tilde{\alpha}$  has image in  $\text{Cl}(\tilde{X} - \text{St}^N(H \tilde{*}))$  and  $\tilde{\alpha}(0), \tilde{\alpha}(1)$  are elements of  $\text{St}^N(H \tilde{*})$ .

Since  $(G, H)$  is 1-tame, there is an integer  $M$  such that  $\alpha$  is homotopic  $\text{rel}\{0, 1\}$  to an edge path in  $\text{St}^M(H \tilde{*})$  by a homotopy in  $\text{Cl}(\tilde{X} - \text{St}^N(H \tilde{*}))$ . Composing this homotopy with  $q$  gives a homotopy of  $\alpha$   $\text{rel}\{0, 1\}$ , to an edge loop in  $\text{St}_H^M(\hat{*})$  by a homotopy in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ .

Conversely, assume for each integer  $N$ , each component of  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  has finitely generated fundamental group. For each component of  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  (there are only finitely many) choose a vertex base point in the boundary of this component. If  $C$  is such a component and  $v$  is the base point of  $C$ , then as  $\pi_1(C, v)$  is finitely generated, we may choose  $M_{C,v}$  an integer such that  $\text{St}_H^M(\hat{*})$  contains a finite set of edge loops which represent generators of  $\pi_1(C, v)$ .

Now any loop in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  based at  $v$  is homotopic  $\text{rel}\{0, 1\}$  to a loop in  $\text{St}_H^M(\hat{*})$  by a homotopy in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ . Choose  $M_1$  such that if  $v$  and  $w$  are vertices of  $\text{Bd}(\text{St}_H^N(\hat{*}))$  which are in the same component of  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  then there is an edge path between  $v$  and  $w$ , in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  of length  $\leq M_1$ . Let  $M$  be an integer larger than  $M_1$  and each of the (finitely many)  $M_{C,v}$  above. Now by a change of base point argument any loop in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  based at a vertex in  $\text{Bd}(\text{St}_H^N(\hat{*}))$  is homotopic  $\text{rel}\{0, 1\}$  to a loop in  $\text{St}_H^M(\hat{*})$  by a homotopy in  $\text{Cl}(\tilde{X} - \text{St}_H^N(\hat{*}))$ .

Let  $\alpha$  be an edge path in  $\text{Cl}(\tilde{X} - \text{St}^N(H\tilde{*}))$  such that  $\alpha(0), \alpha(1)$  are vertices of  $\text{St}^N(H\tilde{*})$ . Then  $q \circ \alpha$  is an edge path in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$  with vertices in  $\text{Bd} \text{St}_H^N(\hat{*})$ . Let  $\beta$  be an edge path of length  $\leq M_1$  from  $q\alpha(1)$  to  $q\alpha(0)$  in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ . Now  $\langle q\alpha, \beta \rangle$  is homotopic  $\text{rel}\{0, 1\}$  to an edge loop  $\gamma$  at  $q\alpha(0)$  where  $\gamma$  is in  $\text{St}_H^M(\hat{*})$  and the image of this homotopy is in  $\text{Cl}(H \setminus \tilde{X} - \text{St}_H^N(\hat{*}))$ . Let  $F$  be such a homotopy.

We have  $q \circ \alpha$  is homotopic  $\text{rel}\{0, 1\}$  to  $\langle \beta, \gamma^{-1} \rangle$  by a homotopy with the same image as  $F$ . Call this homotopy  $W$ . Lifting  $W$  gives a homotopy of  $\alpha$   $\text{rel}\{0, 1\}$  to an edge path (the lift of  $\langle \gamma, \beta^{-1} \rangle$  to  $\alpha(0)$ ) in  $\text{St}^M(H\tilde{*})$ . Furthermore this homotopy has image in  $\text{Cl}(\tilde{X} - \text{St}^N(\tilde{*}))$ , so  $(G, H)$  is 1-tame.  $\square$

## §5. Proof of the main theorem

Let  $P$  be a presentation for  $G$ . Let  $X$  be the standard 2-complex corresponding to  $P$ . Let  $\tilde{X}$  be the universal cover of  $X$ . The 1-skeleton of  $\tilde{X}$  is the Cayley graph of  $P$ , and so the vertices of  $\tilde{X}$  are the elements of  $G$ . Let  $*$  represent the identity vertex. Let  $q: \tilde{X} \rightarrow H \setminus \tilde{X}$  be the quotient map, and  $* \in H \setminus \tilde{X} = q(*)$ . We have an asynchronously bounded, almost prefix closed combing for  $G$ . For each vertex  $v$  of  $\tilde{X}$ , let  $\rho_v: [0, \infty) \rightarrow \tilde{X}$  be the combing path for  $v$ . Let  $L$  be the quasiconvex constant for  $H$  with respect to  $\{\rho_v\}$ , and  $\delta$  be the asynchronous fellow traveler constant. Let  $I$  be the constant that arises from  $\{\rho_v\}$  being almost prefix closed. I.e. if  $\alpha: [0, n] \rightarrow \tilde{X}$  is a prefix of a combing path, then there exists  $v \in G$  such that  $d(v, \alpha(n)) \leq I$  and  $\text{im}(\alpha) \subset \text{St}^I(\text{im}(\rho_v))$ . Let  $Q$  be an integer such that any edge loop  $\alpha$  in  $\tilde{X}$  of length at most  $2\delta + 2$  is homotopically trivial in  $\text{St}^Q(v)$  for all vertices  $v \in \alpha$ .

**LEMMA 5.1.** *If  $M$  is an integer,  $\alpha: [0, n] \rightarrow \tilde{X}$  is an initial segment of a combing path and  $\alpha(n) \in \text{St}^M(H)$ , then  $\text{im}(\alpha) \subset \text{St}^{I+L+(I+M)\delta}(H)$ .*

*Proof.* Choose  $v$  such that  $d(v, \alpha(n)) \leq I$  and  $\text{im}(\alpha) \subset \text{St}^I(\text{im}(\rho_v))$ . Let  $\langle e_1, \dots, e_k \rangle$  be an edge path in  $\tilde{X}$  such that  $e_1(0) \in H$ ,  $e_k(1) = v$  and  $k \leq I + M$ . We have  $\text{im}(\rho_{e_1(0)}) \subset \text{St}^L(H)$  (since  $H$  is quasiconvex). Also,  $\text{im}(\rho_{e_1(1)}) \subseteq$

$\text{St}^\delta(\text{im}(\rho_{e_1(0)}))$  and  $\text{im}(\rho_{e_i(1)}) \subset \text{St}^\delta(\text{im}(\rho_{e_{i-1}(1)}))$  for  $i \in \{2, 3, \dots, k\}$  (by the definition of  $\delta$ ). Hence  $\text{im}(\rho_v) \subseteq \text{St}^{L+(I+M)\delta}(H)$  and so  $\text{im}(\alpha) \subset \text{St}^{I+L+(I+M)\delta}(H)$ .  $\square$

**LEMMA 5.2.** *If  $\alpha$  and  $\beta$  are terminal segments of combing paths,  $\rho_v$  and  $\rho_w$  respectively, such that there is an edge  $e$  from  $v$  to  $w$ , the initial points of  $\alpha$  and  $\beta$  correspond under the asynchronous matching, and  $\gamma$  is an edge path of length  $\leq \delta$  from the initial point of  $\alpha$  to the initial point of  $\beta$  then  $\langle \alpha, e, \beta^{-1}, \gamma^{-1} \rangle$  is homotopically trivial in  $\text{St}^Q(\text{im}(\alpha))$ .*

*Proof.* Choose  $0 = x_1 < x_2 < \dots < x_k = 1$ . Define  $F: I \times I \rightarrow \tilde{X}$  as follows:

(1)  $F|_{[0,1] \times \{0\}}$  and  $F|_{[0,1] \times \{1\}}$  are respectively, a reparametrization of  $\alpha$  composed with an asynchronous adjustment and a reparametrization of  $\beta$  composed with an asynchronous adjustment, so that  $F|_{[x_i, x_{i+1}] \times \{0\}}$  is an edge of  $\alpha$  or constant,  $F|_{[x_i, x_{i+1}] \times \{1\}}$  is an edge of  $\beta$  or constant, and  $d(F(x_i, 0), F(x_i, 1)) \leq \delta$  for all  $i$ . (This is possible since  $\alpha$  and  $\beta$  are asynchronous  $\delta$ -fellow travelers.)

(2)  $F|_{\{x_i\} \times [0,1]}$  is an edge path of length  $\leq \delta$ .

(3)  $F|_{\{0\} \times [0,1]} = \gamma$  and  $F|_{\{1\} \times [0,1]} = e$ .

Now consider the edge loop  $\langle F|_{[x_i, x_{i+1}] \times \{0\}}, F|_{\{x_{i+1}\} \times I}, (F|_{[x_i, x_{i+1}] \times \{1\}})^{-1}, (F|_{\{x_i\} \times I})^{-1} \rangle$ . The length of this loop is  $\leq 2\delta + 2$ . Hence it is homotopically trivial in  $\text{St}^Q(F(x_i, 0)) \subset \text{St}^Q(\text{im}(\alpha))$ . Define  $F$  on  $[x_i, x_{i+1}] \times I$  to realize this homotopy. Then  $\text{im}(F) \subset \text{St}^Q(\text{im}(\alpha))$ , and the result follows.  $\square$

Let  $\alpha$  be an edge path in  $\text{Cl}(\tilde{X} - \text{St}^N(H))$  such that  $\alpha(0), \alpha(1) \subset \text{St}^N(H)$ . Choose  $M > Q + N$ . It suffices to show that  $\alpha$  is homotopic rel  $\{0, 1\}$  to a path in  $\text{St}^{I+(M+I)\delta+L+\delta}(H)$  by a homotopy in  $\tilde{X} - \text{St}^N(H)$ . The path  $\alpha$  decomposes into sub-edge paths,  $\alpha_1, \beta_1, \alpha_2, \beta_2, \dots, \alpha_{n-1}, \beta_{n-1}, \alpha_n$  where  $\text{im}(\alpha_i) \subset \text{St}^M(H)$  and  $\text{im}(\beta_i) \subset \text{Cl}(\tilde{X} - \text{St}^M(H))$ . It suffices to show that  $\beta(\equiv \beta_i)$  is homotopic rel  $\{0, 1\}$  to a path with image in  $\text{St}^{I+(M+I)\delta+L+\delta}(H)$  by a homotopy in  $\tilde{X} - \text{St}^N(H)$ .

Let  $e_1, e_2, \dots, e_n$  be the edges of  $\beta$ ,  $\rho_i$  be the combing path corresponding to  $e_i(1)$ ,  $\rho_0$  the combing path corresponding to  $e_1(0)$  and  $T_i \equiv T_{e_i(1)}$ .

If  $t_i$  is the last point of  $[0, T_i]$  such that  $\rho_i(t_i) \in \text{St}^M(H)$ , then let  $a_i$  be a point of  $[0, T_i]$  such that  $a_i$  corresponds (under the asynchronous tracking) to  $\rho_{i-1}(t_{i-1})$ ,  $\gamma_i$  be an edge path of length  $\leq \delta$  from  $\rho_{i-1}(t_{i-1})$  to  $\rho_i(a_i)$  and  $\xi_i$  be the subpath of  $\rho_i$  and  $a_i$  to  $t_i$ .

It suffices to show that for each  $i$ .

(1)  $\text{Im} \langle \gamma_i, \xi_i \rangle \subset \text{St}^{I+(M+I)\delta+L+\delta}(H)$  and

(2)  $\langle \gamma_i, \xi_i \rangle$  is homotopic rel  $\{0, 1\}$ , by a homotopy in  $\tilde{X} - \text{St}^N(H)$ , to  $\langle \rho_{i-1}|_{[t_{i-1}, T_{i-1}]}, e_i, (\rho_i|_{[t_i, T_i]})^{-1} \rangle$ .

For then, patching together these homotopies gives a homotopy rel  $\{0, 1\}$  of  $\beta \equiv \langle e_1, e_2, \dots, e_n \rangle$  to  $\langle \gamma_1, \xi_1, \gamma_2, \dots, \gamma_n, \xi_n \rangle$ , in  $\tilde{X} - \text{St}^N(H)$ .



If  $a_i \leq t_i$ , then  $\xi_i = \rho_i|_{[a_i, t_i]}$ . In this case, by Lemma 5.1,  $\rho_i|_{[0, t_i]}$  has image in  $\text{St}^{I+(M+I)\delta+L}(H)$ , and  $\text{im}(\gamma_i) \subset \text{St}^\delta(\rho_{i-1}(t_{i-1})) \subset \text{St}^{\delta+M}(H)$ , so  $\text{im}\langle \xi_i, \gamma_i \rangle \subset \text{St}^{I+(M+I)\delta+L+\delta}(H)$ . By applying Lemma 5.2 with  $\rho_{i-1}|_{[t_{i-1}, T_{i-1}]}$  in place of  $\alpha$ ,  $e_i$  in place of  $e$ ,  $\gamma_i$  in place of  $\gamma$  and  $\rho_i|_{[a_i, T_i]}$  in place of  $\beta$  we have  $\langle \gamma_i, \xi_i \rangle$  is homotopic rel  $\{0, 1\}$ , by a homotopy in  $\text{St}^Q(\text{im}(\rho_{i-1}|_{[t_{i-1}, T_{i-1}]})$ , to  $\langle \rho_{i-1}|_{[t_{i-1}, T_{i-1}]}, e_i, (\rho_i|_{[t_i, T_i]})^{-1} \rangle$ . As  $\text{im}(\rho_{i-1}|_{[t_{i-1}, T_{i-1}]} \subset \text{Cl}(\tilde{X} - \text{St}^M(H))$  and  $M > N + Q$ , the image of this homotopy does not intersect  $\text{St}^N(H)$ .

If  $t_i \leq a_i$ , then again  $\text{im}(\gamma_i) \subset \text{St}^\delta(\rho_{i-1}(t_{i-1})) \subset \text{St}^{I+(M+I)\delta+L+\delta}(H)$ . Observe that  $\xi_i = (\rho_i|_{[t_i, a_i]})^{-1}$  and  $\text{im}(\xi_i) \subseteq \text{St}^\delta(\text{im}(\rho_{i-1}|_{[0, t_{i-1}]})$ . Hence by Lemma 5.1  $\text{im}(\xi_i) \subseteq \text{St}^{I+(M+I)\delta+L+\delta}(H)$ . By Lemma 5.2,  $\gamma_i$  is homotopic rel  $\{0, 1\}$  to  $\langle \rho_{i-1}|_{[t_{i-1}, T_{i-1}]}, e_i, (\rho_i|_{[a_i, T_i]})^{-1} \rangle$  by a homotopy in  $\text{St}^Q(\text{im}(\rho_{i-1}|_{[t_{i-1}, T_{i-1}]}) \subset \tilde{X} - \text{St}^N(H)$ . As  $\text{im}(\xi_i) \subset \text{Cl}(\tilde{X} - \text{St}^M(H)) \subset \tilde{X} - \text{St}^N(H)$ ,  $\langle \gamma_i, \xi_i \rangle$  is homotopic rel  $\{0, 1\}$  to  $\langle \rho_{i-1}|_{[t_{i-1}, T_{i-1}]}, e_i, (\rho_i|_{[t_i, T_i]})^{-1} \rangle$  by a homotopy in  $\tilde{X} - \text{St}^N(H)$ .  $\square$

## Acknowledgement

This paper was written while the author was at Center de Recerca Matemàtica, Institut d'Estudis Catalans, in Spain. We are grateful for the hospitality received at the Center. A special thanks is also expressed to G. A. Swarup who encouraged the author to consider the problems solved in this paper. Finally we wish to thank the referee for several helpful suggestions.

## REFERENCES

- [Bo] F. BONAHO, *Bouts des variétés hyperboliques de dimension 3*, Annals of Math. 124 (1986), 71–158.
- [Br<sub>1</sub>] M. BRIDSON, *Combings of semidirect products and 3-manifold groups*, Geom. and Funct. Anal. 3 No. 3 (1993), 263–278.
- [Br<sub>2</sub>] M. BRIDSON, Personal communication.
- [BH] M. BRIDSON and A. HAEFLIGER, *Metric spaces of non-positive curvature*, preprint.
- [BT] M. BRIN and T. THICKSTUN, *3-manifolds which are end 1-movable*, Memoirs of the AMS 81 No. 411 (1989).
- [E] D. B. A. EPSTEIN, J. W. CANNON, D. F. HOLT, S. V. F. LEVY, M. S. PATTERSON and W. P. THURSTON, *Word processing in groups*, Jones and Barlett, 1992.
- [GS] S. GERSTEN and H. SHORT, *Rational subgroups of biautomatic groups*, Annals of Math. 134 (1991), 125–158.
- [HRS] J. HASS, H. RUBINSTEIN and P. SCOTT, *Compactifying coverings of closed 3-manifolds*, J. Diff. Geom. 30 (1989), 817–832.
- [J] W. JACO, *Lectures on three-manifold topology*, American Math Soc. CBMS Lecture Note Series, 1980.
- [L] B. LEEB, *3-Manifolds with(out) metrics of non-positive curvature*, Invent. Math. 122 No. 2 (1995), 277–290.
- [MT] M. L. MIHALIK and S. T. TSCHANTZ, *Tame combings and the quasi-simply-filtered condition for groups*, Trans. Amer. Math. Soc., to appear.

- [P] V. POÉNARU, *Almost convex groups, Lipschitz combing, and  $\pi_1^\infty$  for universal covering spaces of closed 3-manifolds*, J. Diff. Geom. 35 (1992), 103–130.
- [R] L. REEVES, *Rational subgroups of cubed 3-manifold groups*, Michigan Math. J. 42 No. 1 (1995), 109–126.
- [Si] J. SIMON, *Compactification of covering spaces of compact 3-manifolds*, Mich. Math. J. 23 (1976), 245–256.
- [T] W. THURSTON, *Geometry and topology of 3-manifolds*, Lecture Notes, Princeton University, Princeton, NJ, 1978–9.
- [Tu] T. W. TUCKER, *Non-compact 3-manifolds and the missing boundary problem*, Topology (13) (1974), 267–273.

*Michael L. Mihalik*  
*Department of Mathematics*  
*Vanderbilt University*  
*Nashville TN 37240*  
*U.S.A.*  
*E-mail address:* mihalikm@ctrvax.vanderbilt.edu

Received January 23, 1994; February 12, 1996