**Zeitschrift:** Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

**Band:** 71 (1996)

**Artikel:** Local structure of the moduli space of vector bundles over curves.

Autor: Laszlo, Yves

**DOI:** https://doi.org/10.5169/seals-53849

#### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 11.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Local structure of the moduli space of vector bundles over curves

YVES LASZLO\*

#### 0. Introduction

Let X be a smooth, projective and connected curve (over an algebraically closed field of characteristic zero) of genus  $g(X) \ge 2$ . Let x be a (closed) point of X and  $SU_X(r, d)$  the moduli space of semi-stable vector bundles on X of rank  $r \ge 2$  and determinant  $\mathcal{O}(dx)$ . As usual, the geometric points of  $SU_X(r, d)$  correspond to polystable bundles, namely direct sums  $E = \bigoplus E_i$  where  $E_i$  is stable of slope  $\mu(E_i) = d/r$  (and  $\bigotimes_i \det(E_i) = \mathcal{O}(dx)$ ).

DEFINITION. The number of stable summands in the preceding sum is called the length of the polystable bundle E.

The singular locus of  $SU_X(r, d)$  consists exactly of the non stable points (except if r = g(X) = 2 and d even). In this case,  $SU_X(r, d) = \mathbb{P}^3$  [N-R1]). In particular, except in the exceptional case above,  $SU_X(r, d)$  is smooth if and only if r and d are relatively prime. General facts about the action of reductive groups ensure that  $SU_X(r, d)$  is Cohen-Macaulay [E-H], normal and that the singularities are rational [B]. The principal aim of this paper is to give additional information about the singularities, essentially the description of the completion of the local ring at a singular point of  $SU_X(r, d)$  and to compute the multiplicity and the tangent cones at those singular points E which are not too bad, i.e. l(E) = 2 (or equivalently  $Aut(E) = G_m \times G_m$ ). Further, we give a complete description in the rank 2 case (corollaries III.2 and III.3 and theorem IV.4).

As recalled with some details (following a request of the referee), Narasimhan and Ramanan [N-R2] have a long time ago discovered the link between the rank 2 vector bundles on a canonical curve X of genus 3 and the geometry of the Kummer variety  $\kappa(X)$  of JX. The crucial point is that  $SU_X(2)$  is in this case canonically isomorphic to the so called *Coble quartic*. We get the local form of this

<sup>\*</sup> The author was partially supported by the European Science Project "Geometry of Algebraic Varieties", Contract no. SCI-0398-C(A).

quartic and prove that the  $\kappa(X)$  is schematically defined by 8 cubics, the partials derivatives of the Coble quartic (theorem IV.6), although the corresponding homogeneous ideal is not generated by these cubics.

One could also give partial information at least if  $\operatorname{Aut}(E)$  is a torus, or by using results of [P], if  $\operatorname{Aut}(E) = \operatorname{Gl}_r(k)$  (the latter case meaning that E is a twist of the trivial bundle). But it seems to be difficult (and somewhat messy) to calculate for instance the multiplicity. In the last part of the paper, we compute the multiplicity of a generalized theta divisor of  $\operatorname{SU}_X(2, \emptyset)$  at a point  $[L \oplus L^{\vee}]$ , where  $L^2 \neq \emptyset$ . In fact, this computation could be done with only minor changes for a point E of any rank with  $\det(E) = \emptyset$  and  $\operatorname{Aut}(E) = G_m \times G_m$ .

Let us also mention that similar results could be obtained exactly in the same way for certain surfaces. But, all the future applications that we have in mind as well as the applications that we have in our hand are for curves. Therefore, we have restricted ourselves to the case of curves.

#### Notations and conventions

All the schemes are of finite type over an algebraically closed field k of characteristic  $p \neq 2$ . Except in the first section, it will be assumed of characteristic zero. By point we mean closed point. The scheme P(V) is the projective space P(V) of lines of V and V will also denote the pointed affine space P(V) = P(V) = P(V) (notice that P(V) = P(V)) is the coordinate ring of P(V).

The notations X (resp. x) will denote a smooth projective connected curve over k of genus  $g \ge 2$  (resp. a closed point x of X). If  $\mathscr L$  is a line bundle on a scheme, the linear system  $PH^0(\mathscr L)$  is denoted by  $|\mathscr L|$  and the dual projective space by  $|\mathscr L|^{\vee}$ . Fix also two integers  $r \ge 2$  and d. Finally, E will always denote a rank r polystable bundle on X of determinant dx and  $SU_X(r,d)$  is the moduli space of rank r semi-stable vector bundles of determinant  $\mathscr O(dx)$  on X.

# I. Around the Coble quartic

In this section, we first recall for the non-expert reader some facts about the geometry of the Kummer variety (essentially due to Mumford, Kempf and Khaled). We explain then the link between the Coble quartic and the Kummer of the jacobian JX of a canonical genus 3 curve X on one hand, and, in the other hand the identification of the Coble quartic and  $SU_X(2,0)$ . Finally, we state our result about the Coble quartic.

Generalities on linear system on abelian varieties

Let A be an abelian variety and L an ample line bundle on it. It is well-known that L<sup>n</sup> is very ample if  $n \ge 3$ . A more subtle result due to Kempf (see [Kh1]) says that the products

$$H^0(A, L^n) \otimes H^0(A, L^m) \rightarrow H^0(A, L^{n+m})$$

are surjective if  $n \ge 2$  and  $m \ge 3$ . In particular, the complete linear system  $A \to |L^n|^\vee$  is projectively normal. Let  $I_n$  be the corresponding homogeneous ideal

$$I_n = \operatorname{Ker} \left( \bigoplus_{k \geq 0}^{\infty} \operatorname{Sym}^k H^0(A, L^n) \to \bigoplus_{k \geq 0}^{\infty} H^0(A, L^{nk}) \right).$$

Kempf and Khaled (cf. [Kh1]) have also shown that  $I_n$  is generated by quadrics and cubics if  $n \ge 3$  and by quadrics if  $n \ge 4$ .

The situation is very different for n = 2. In this case, the linear system  $|L^2|$  is in general only base point free.

### From now on, A is a principally polarized irreducible abelian variety

Let  $\Theta$  be a symmetric divisor representing the polarization. Then the line bundle  $\mathcal{L} = \mathcal{O}(2\Theta)$  does not depend on the particular choice of such a  $\Theta$ . Then, the canonical morphism

$$\kappa_{\mathbf{A}} \colon \mathbf{A} \to |\mathcal{L}|^{\vee}$$

induces an embedding of  $A/\pm 1$  in  $|\mathcal{L}|^{\vee}$ . The image  $\kappa(A)$  of  $\kappa_A$  is the Kummer variety of A. The line bundle  $\mathcal{L}$  is canonically  $\{\pm 1\}$ -linearized (as the pull-back of  $\kappa_A^*\mathcal{O}(1)$  and one has

$$\mathrm{H}^0(\kappa(\mathrm{A}),\,\mathcal{O}(n))=\mathrm{H}^0(\mathrm{A},\,\mathcal{L}^n)^+$$

where  $H^0(A, \mathcal{L}^n)^+$  is the invariant part of  $H^0(A, \mathcal{L}^n)$  under -1.

Remark. If  $k = \mathbb{C}$ , the link with the classical geometry of theta functions can be explained as follows. The Chern class  $c_1(\mathcal{L}) \in H^2(A, \mathbb{Z})$  give a symplectic non degenerate Z-bilinear form on  $H_1(A, \mathbb{Z})$  which depends only on the polarization  $[\mathcal{L}]$ . The abelian variety A is the quotient of  $T_0(A)$  by the lattice  $\Gamma = h_1(A, \mathbb{Z})$ .

Choose a symplectic basis  $(\gamma_1, \ldots, \gamma_{2g})$  of  $\Gamma$ . Then  $(\gamma_1, \ldots, \gamma_g)$  is a basis of  $T_0(A)$  which therefore will be identified to  $C^g$ . The matrix  $\tau$  of coordinates of  $(\gamma_{g+1}, \ldots, \gamma_{2g})$  in  $C^g$  is called the period matrix and one has by construction  $A = C^g/(Z^g \oplus \tau Z^g)$ . With this description of A, the space  $H^0(A, \mathcal{L}^n)$  becomes the space of order 2n theta functions (with respect of  $\tau$ ). The sections of  $H^0(\kappa(A), \mathcal{O}(n))$  are the even theta functions of order 2n.

Notice the formula dim  $H^0(\kappa(A), \mathcal{O}(n)) = 2^{g-1}(n^g+1)$  (see [C], page 56 in the case  $k = \mathbb{C}$  and formula (+) below in general).

The theorem of Mumford-Koizumi [Ko] says that the Kummer variety is projectively normal if and only if A has no vanishing theta-nulls. Geometrically, this condition means that among the  $2^{2g}$  symmetric theta divisors  $\Theta$  representing the principal polarization, there is no  $\Theta$  vanishing at the origin with even multiplicity. In the case where A is the jacobian of a smooth curve X, this condition simply means that there is no effective even theta-characteristic on X.

### Suppose now that A has no one vanishing theta-null

The number h(n, g) of independent degree n hypersurfaces containing  $\kappa(A)$  is

$$h(n,g) = {2^g + n - 1 \choose n} - 2^{g-1}(n^g + 1).$$

In particular  $\kappa(A)$  is never contained in a quadric. If g=2, one has h(3,g)=0 and h(4,g)=1: the Kummer surface is a quartic surface with  $2^{2g}=16$  nodes. Suppose now  $g \ge 3$ ; then h(3,g)>0 and  $\kappa(A)$  is always contained in cubics. Let  $I_A$  (resp.  $\mathscr{I}_A$ ) be the homogeneous ideal (resp. the sheaf of ideal) of  $\kappa(A)$ ; by definition, one has

$$I_{A} = \operatorname{Ker} \left( \bigoplus_{k \geq 0}^{\infty} \operatorname{Sym}^{k} \operatorname{H}^{0}(A, \mathcal{L}) \twoheadrightarrow \bigoplus_{k \geq 0}^{\infty} \operatorname{H}^{0}(A, \mathcal{L}^{k})^{+} \right)$$

and the dimension of the degree  $\delta$  component  $I_A[d] = \dim H^0(\kappa_A, \mathcal{O}(\delta))$  of  $I_A$  is  $h(g, \delta)$ . One has the

THEOREM 1 ([Kh 2]). With the notations above,  $I_A$  is generated by quartics in degree  $\geq 4$ .

In particular, the ideal  $\mathcal{I}_A$  is itself generated by quartics.

Remark. When g = 3 or A is generic, this theorem was known from Wirtinger [Wi].

If g=3: the Kummer variety is embedded in  $\mathbf{P}^7=|\mathcal{L}|^\vee$ . In this case, dim  $I_A[3]=8$  and dim  $I_A[4]=70>64=\dim I_A[3]$ . dim  $H^0(\mathbf{P}^7,\mathcal{O}(1))$  which shows that  $I_A[4]$  is not generated by cubics.

- (2) One can ask the following natural questions:
- (2.1) Is the Kummer  $\kappa(A)$  variety set-theoretically defined by the cubics of  $I_A[3]$ ?
- (2.2) Is the Kummer  $\kappa(A)$  variety schematically defined by the cubics of  $I_A[3]$ ? In other words, is the canonical map  $I_A[3] \otimes \mathcal{O}(-3) \to \mathscr{I}_A$  surjective?

The dimension 3 case: the Coble quartic

Let us first recall general facts from [M] about the Mumford group  $G(\mathcal{L})$ , the group of pairs  $(\alpha, a)$  where  $\alpha$  is an isomorphism of  $\tau_a^*(\mathcal{L})$  on  $\mathcal{L}$  ( $\tau_a$  is the translation by a). The morphism  $(\alpha, a) \mapsto a$  surjects onto  $A_2$  and one has a canonical exact sequence

$$1 \to k^* \to G(\mathcal{L}) \to A_2 \to 0$$

with  $k^*$  central. Let V be a  $G(\mathcal{L})$ -module. We say that V is of level n if  $0 < \dim(V) < \infty$  and  $k^*$  acts on V by  $t \mapsto t^n$ . If V is irreducible, there exist an unique integer n such that V is of level n. The basic fact about the representation theory of  $G(\mathcal{L})$  is the following classical observation:

LEMMA 3. Let w be an odd number. There exists only one (up to isomorphism) irreducible  $G(\mathcal{L})$ -module  $V_w$  of level w. Moreover, if V is any  $G(\mathcal{L})$ -module of level w, then V is a sum of copies of V(w).

Let us explicit the representation V(w). Let G(2) be the group which is set theoretically the product  $k^* \times \mathbf{F}_2^g \times \widehat{\mathbf{F}}_2^g$  with the group law

$$(\lambda, \varepsilon, \hat{\varepsilon}) \cdot (\mu, \eta, \hat{\eta}) = ((-1)^{\langle \hat{\eta}, \varepsilon \rangle} \lambda \mu, \varepsilon + \eta, \hat{\varepsilon} + \hat{\eta}).$$

Because the characteristic is  $\neq 2$ , the group  $G(\mathcal{L})$  is (non canonically) isomorphic to G(2): an isomorphism is called a level 2 theta structure on A.

Remark. If  $k = \mathbb{C}$ , the polarization can be seen as a symplectic non degenerate form on  $H_1(A, \mathbb{Z})$  (for instance, it is the Chern class of a symmetric divisor  $\Theta$  as above). The choice of a symplectic basis  $\mathscr{B}$  of  $H_1(A, \mathbb{Z})$  defines the period matrix  $\tau$  of A and identifies A and  $\mathbb{C}^g/(\mathbb{Z}^g + \tau \mathbb{Z}^g)$ . Moreover  $\mathscr{B}$  defines canonically a theta structure on  $\mathscr{L}$ .

### From now on, A is endowed with a level 2 structure

Then V(w) can be viewed as  $Map(\mathbf{F}_2^g, k)$  with the action given by the formula

$$(\lambda, \varepsilon, \hat{\varepsilon}) \cdot f(v) = (-1)^{\langle \hat{\varepsilon}, v + \varepsilon \rangle} \lambda^{w} f(v + \varepsilon), \qquad v \in \mathbf{F}_{2}^{g}, \qquad f \in \mathbf{Map}(\mathbf{F}_{2}^{g}, k).$$

Of course,  $H^0(A, \mathcal{L})$  is a level 1 module of dimension  $2^g = \dim \operatorname{Map}(\mathbf{F}_2^g, k)$ . Therefore, there exists a  $G(\mathcal{L})$  isomorphism  $V(1) \tilde{\to} H^0(A, \mathcal{L})$  well defined up to a non zero scalar. The action on  $\pm 1$  is given by the *inverse formula* (p. 331 of [M]): this is just the multiplication by -1 on  $\mathbf{F}_2^g$ . More generally, using a level 2n theta structure,  $H^0(A, \mathcal{L}^n)^+$  can be identified with the **even** functions on  $(\mathbf{Z}/2n\mathbf{Z})^g$  and we get the formula

$$\dim H^0(A, \mathcal{L}^n)^+ = 2^{g-1}(n^g + 1). \tag{+}$$

For  $v \in \mathbb{F}_2^g$ , let  $\mathbb{Z}_v \in \mathbb{V}(w)$  be the characteristic function of  $\{v\}$ . We have then

$$(\lambda, \varepsilon, \hat{\varepsilon}) \cdot Z_v = (-1)^{\langle \hat{\varepsilon}, \varepsilon + v \rangle} \lambda^w Z_{v + \varepsilon}.$$

Remark. If  $k = \mathbb{C}$  and  $A = \mathbb{C}^g/(\mathbb{Z}^g \oplus \tau \mathbb{Z}^g)$  as above,  $H^0(A, \mathcal{L})$  is canonically identified with the space of level 2 theta function with respect of  $\tau$  and  $\mathbb{Z}_v$  corresponds to the theta function with characteristic (v, 0)

$$\theta \begin{bmatrix} v \\ 0 \end{bmatrix} (2z, 2\tau).$$

We now focus our attention to the 3-dimensional case. Because A is assumed to be irreducible with no vanishing theta-nulls, A is the jacobian JX of canonical curve X of genus 3.

Then  $I_{JX}[3]$  is a level 3  $G(\mathcal{L})$ -module of dimension 8 and therefore isomorphic to V(3). Let  $C_v \in I_{JX}[3]$  be the cubic corresponding to  $Z_v$ . The cubic  $C_0$  is invariant under the subgroup  $I = \{1\} \times \{0\} \times \widehat{\mathbf{F}}_2^g$  of  $G(2) = G(\mathcal{L})$ . Notice that a monomial  $Z_u Z_v Z_w$  is invariant under I if and only if w = u + v. We think to  $\mathbf{F}_2^3$  as  $\mathbf{P}^2(\mathbf{F}_2) \cup 0$ . It follows that  $C_0$  is of the form

$$\sum_{\alpha_v \in \mathbf{F}_2^3} \mathbf{Z}_0 \mathbf{Z}_v^2 + \sum_{d \in \mathbf{P}^2 \mathbf{F}_2^{\vee}} \alpha_d \mathbf{Z}^d$$

where  $Z^d = \prod_{0 \neq v \in d} Z_v$ . The group I acts on  $C_v$  by  $(1, 0, \hat{\epsilon}) \mapsto (-1)^{\langle \hat{\epsilon}, v \rangle}$  and therefore  $Z_v \cdot C_v$  is also invariant under I and can be written using the coefficients  $\alpha_v$  and  $\alpha_d$  of  $C_0$ . Let C be the quartic

$$\sum_{v \in \mathbf{F}_2^g} \mathbf{Z}_v \cdot \mathbf{C}_v.$$

By direct computation, one observes  $\partial C/\partial Z_0 = 4C_0$ . By construction, C is invariant under  $\{1\} \times \mathbf{F}_2^g \times \{0\}$  which gives  $\partial C/\partial Z_p = 4C_p$ 

for all  $c \in \mathbb{F}_2^g$ . In particular, there exists a quartic containing  $\kappa(JX)$  in its singular locus. Notice that C being invariant under I as  $\mathbb{Z}_v \cdot \mathbb{C}_v$  is, the quartic div(C) of  $|\mathcal{L}|^{\vee}$  is invariant under the whole  $G(\mathcal{L})$ . Let us prove the

LEMMA 4. The Coble quartic is the unique  $G(\mathcal{L})$ -invariant quartic in  $|\mathcal{L}|^{\vee}$  containing  $\kappa(JX)$  in his singular locus.

Remark. This is probably the meaning of the assertion (6) page 106 of [C]. Proof of the lemma: let F be an equation of a  $G(\mathcal{L})$ -invariant quartic. There exists a character  $\chi$  of  $G(\mathcal{L})$  such that (with the notations above)

$$F((\lambda, \varepsilon, \hat{\varepsilon}) \cdot Z_v) = \chi((\lambda, \varepsilon, \hat{\varepsilon})) \cdot F(Z_v). \tag{1}$$

Let  $p \in \mathbb{F}_2^g$  such that  $\chi(1, 0, \hat{\epsilon}) = (-1)^{t_p \cdot \hat{\epsilon}}$ . Differentiating (1) with respect of  $\mathbb{Z}_v$  gives that there exists a scalar  $\lambda_v$  such that

$$\partial \mathbf{F}/\partial \mathbf{Z}_{v} = \lambda_{v} \, \partial \mathbf{C}/\partial \mathbf{Z}_{v+p}. \tag{2}$$

Because  $p \neq 2$ , the endomorphism given by  $Z_v \rightarrow Z_{v+p}$  is semi-simple and the system (2) becomes

$$d\mathbf{F} = d\mathbf{C} \circ \mathbf{M} \tag{3}$$

where M is a semi-simple matrix. After eventually replacing F and C by projectively equivalent hypersurfaces, one can assume that M is *diagonal*. In this case, (3) gives that F and some multiple of C differs by a  $p^{th}$ -power of some form G. By degree consideration (using  $p \neq 2$ ), one obtains G = 0.

To go further, one has to construct geometrically the Coble quartic: this has been done in [N-R2].

Rank 2-vector bundles and the Coble quartic

Let X be a canonical curve of genus 3 (recall that  $p = \text{char}(k) \neq 2$ ). Let A be the jacobian JX. Let  $\Theta$  be the (canonical) theta divisor on  $\text{Pic}^{g-1}(X)$ . If  $[E] \in SU_X(2, 0)$ , the locus

$$\Theta_{E} = \left\{ L \in \operatorname{Pic}^{g-1}(X) \mid H^{0}(X, E \otimes L) \neq 0 \right\}$$

is a divisor in PH<sup>0</sup>(Pic<sup>g-1</sup>(X),  $\mathcal{O}(2\Theta)$ ) and the G( $\mathcal{O}(2\Theta)$ )-equivariant morphism  $\tilde{\varphi}_X$ 

$$\tilde{\varphi}_{\mathbf{X}} : \begin{cases} \mathrm{SU}_{\mathbf{X}}(2,0) & \to & \mathbf{P}\mathrm{H}^0(\mathrm{Pic}^{g-1}(\mathbf{X}),\,\mathcal{O}(2\Theta)) \\ \mathrm{E} & \mapsto & \Theta_{\,\mathrm{E}} \end{cases}$$

is an embedding onto a quartic C(X) by [N&R]. The Riemann's bilinear relations furnish a canonical identification  $|\mathcal{L}|^{\vee} = PH^0(Pic^{g-1}(X), \mathcal{O}(2\Theta))$  and the corresponding  $G(\mathcal{L})$ -equivariant morphism  $\varphi_X: SU_X(2,0) \hookrightarrow |\mathcal{L}|^{\vee}$  restricts on  $Sing(SU_X(2,0)) = A/\pm 1$  to the Kummer morphism  $\kappa_{JX}$ .

Exercise. Suppose that k is of characteristic 0. Using the Verlinde formula, prove directly that  $\varphi_X$  is an embedding. Notice that this formula was unknown when Narasimhan and Ramanan stated and proved their theorem

Notice that  $\varphi_X[Sing(SU_X(2,0))] = \kappa(JX)$ . In particular C(X) is a quartic singular exactly along  $\kappa(JX)$ . The  $G(\mathcal{L})$ -invariance and the lemma above shows then that C(X) is the Coble quartic. In fact, at least if k is of characteristic zero, a stronger statement due to Beauville than the lemma 4 is true.

PROPOSITION 5 (Beauville). Assume that  $p = \operatorname{char}(k)$  is zero. The Coble quartic is the unique quartic in  $|\mathcal{L}|^{\vee}$  containing  $\kappa(JX)$  in its singular locus.

Sketch of proof. let F be a quartic singular along  $\kappa(JX)$ . Notice that C is not a cone. The condition to be a cone being algebraic, the generic member of the pencil  $(C + tF)_{t \in k}$  is not a cone. In particular, one can assume that F is not a cone. The partial derivatives of C are then a basis of the cubics containing the Kummer. This fact together with the fact that F is not a cone proves that F and C have the same jacobian ideal. A result of [Do] ensures that F and C are related by an invertible linear transformation T. Because C is singular exactly along  $\kappa(JX)$ , the linear morphism T leaves  $\kappa(X)$  invariant. An easy adaptation of the Torelli theorem proves that T is induced by an element of Aut(X) which leaves obviously  $\operatorname{div}(C) = \varphi_X(SU_X(2,0))$  invariant.

The beautiful corollary of the equality  $Sing(C(X)) = \kappa(JX)$  is the

THEOREM 6 [N-R2]. The Kummer variety of the jacobian of a genus 3 canonical curve is set theoretically defined by cubics.

This gives a positive answer to the question (2.1).

More surprisingly (recall that  $I_{JX}[3]$  does not generate  $I_{JX}[4]$ ), the main result of the present paper is

THEOREM IV.6. Assume that p = char(k) is zero. The scheme defined by the eight partial derivatives of the Coble quartic is reduced.

The proposition and the fact that these eight cubics defines set-theoretically the Kummer variety of  $\kappa(JX)$  proves the

COROLLARY. Assume that  $p = \operatorname{char}(k)$  is zero. Let X be a canonical genus 3 curve over. The sheaf of ideals  $\mathcal{I}_{JX}$  of the Kummer variety  $\kappa(JX)$  is generated by cubics.

These facts will be proved in section IV and give a positive answer to the question (2.2).

### II. Local structure of $SU_X(r, d)$ and classical invariant theory

From now, k is of characteristic zero. If  $(X_i)_{1 \le N}$  (resp.  $(n_i)_{1 \le N}$ ) are indeterminates (resp. non negative integers), let me denote by  $\underline{X}$ ,  $\underline{n}$  and  $X^n$  the N-tuple  $\underline{X} = (X_i)$ , the multi-index  $\underline{n} = (n_i)$  and the product  $X^n = \prod_{i=1}^N x_i^{n_i}$  respectively. For

V a finite dimensional vector space with dual  $V^{\vee}$ , the ring k[V] (resp. k[[V]]) is the polynomial ring Sym  $V^{\vee}$  (resp. its completion at the origin).

It is well known (see [S] for instance) that the key ingredient to analyse the local structure of  $SU_x(r, d)$  is the étale slice theorem of Luna. Let us recall this analysis.

Take *n* big enough such that E(nx) is globally generated and has no  $H^1$  for every semi-stable rank *r* vector bundle E of degree *d* (every *n* such that rn + d > r(2g - 1) has this property). Let  $\chi = \chi(E(nx))$  be the corresponding Euler-characteristic. In Grothendieck's scheme *Quot* which parametrizes quotients

$$\mathcal{O}(-nx)^{\oplus \chi} \rightarrow E$$

let  $\mathcal{Q}$  be the open set whose closed points correspond to such quotients with the following properties:

- (i) E is semi-stable of rank r and degree d.
- (ii)  $H^1(X, E(nx)) = 0$  and the natural map  $H^0(X, \mathcal{O}^{\oplus x}) \to H^0(X, E(nx))$  is onto.

Let  $\mathscr{E}$  be the universal quotient bundle on  $\mathscr{Q}$ . The scheme  $\mathscr{Q}$  is smooth and the semi-simple group  $G = PGL_{\chi}$  acts on it. The moduli space  $SU_{\chi}(r, d)$  is the GIT quotient of  $\mathscr{Q}/G$ .

For  $(E_i)$  be a set of a vector bundles over X, the kernel of the trace map

$$\operatorname{Ker}(\bigoplus \operatorname{Ext}^{1}(\operatorname{E}_{i},\operatorname{E}_{i}) \xrightarrow{\bigoplus \operatorname{Tr}_{i}} \operatorname{H}^{1}(\operatorname{X},\mathscr{O}))$$

will be denoted by  $(\bigoplus Ext^1(E_i, E_i))_0$ .

Let  $q = [\mathcal{O}(-nx)^{\oplus \chi} \rightarrow E]$  a point of  $\mathcal{Q}$  and gr the corresponding graded object. By the very definition of semi-stability, there exists a G-stable open affine neighborhood  $\Omega$  and the fibre of  $\Omega \rightarrow \Omega/G$  at [E] contains a unique closed orbit G(q). This orbit is either characterized as being of minimal dimension, or as having an isotropy group  $G_q = \operatorname{Aut}(E)$  of maximal dimension: one checks that this exactly means that E is *polystable*. The corresponding orbit is closed. The closedness of the orbit and the smoothness of  $\Omega$  corresponding to E allows us to use the Luna étale slice theorem (more precisely the remarque page 97 of [Lu]) which gives precisely the:

THEOREM 1 (Luna). Let E be a polystable of rank r and determinant  $\mathcal{O}(dx)$ . There exists an étale neighborhood of  $(\operatorname{Ext}_0^1(E,E)/\operatorname{Aut}(E),0)$  (resp.  $(\operatorname{SU}_X(r,d),E)$ ) which are isomorphic.

Remarks 2. (a) The group Aut(E) acts by functoriality on both arguments of  $Ext_0^1(E, E)$ . Notice that the scalars acts trivially and therefore

$$\operatorname{Ext}_0^1(E, E)/\operatorname{Aut}(E) = \operatorname{Ext}_0^1(E, E)/\operatorname{G}_E$$

where

$$G_E = Ker\{Aut(E) \xrightarrow{\det} G_m\}.$$

- (b) We'll say for short that  $SU_X(r, d)$  is étale locally isomorphic to  $Ext_0^1(E, E)/G_E$  at E.
  - (c) By definition, one has

$$\operatorname{Ext}_0^1(E, E)/G_E = \operatorname{Spec}(A_E)$$

where  $A_E$  is the ring of polynomial maps on  $Ext_0^1(E, E)$  invariant under  $G_E$ .

### Unless otherwise stated, all bundles will be polystable

COROLLARY. The local ring  $\widehat{\mathcal{O}}_{\mathrm{SU}_{\mathbf{X}}(r,d),[E]}$  depends only on the numerical invariants of  $\mathbf{X}$  and  $\mathbf{E}$ .

Suppose once for all that E is non stable (l(E) > 1). One has of course the inequalities

$$1 \le \dim G_{\mathcal{E}} \le r^2 - 1 \tag{1}$$

with equality on the left hand side (resp. right hand side) of (1) if  $G_E = G_m$  (Case 1) (resp.  $G_E = Sl_r$  (Case 2)). Let's examine these 2 cases.

# III. Case 1: $G_E = G_m$

In this case, E is a direct sum  $E = E_1 \oplus E_2$  where  $E_i$  is stable of slope d/r, rank  $r_i \neq 0$  and  $E_1 \approx E_2$ . Each element  $(\alpha_1, \alpha_2) \in G_E(k)$  acts by multiplication by  $\alpha_j \cdot \alpha_i^{-1}$  on each factor  $Ext^1(E_i, E_i)$  of

$$\operatorname{Ext}^1(\operatorname{E},\operatorname{E}) = \bigoplus \operatorname{Ext}^1(\operatorname{E}_i,\operatorname{E}_j).$$

Let

$$d_{i,j} = \dim \operatorname{Ext}^{1}(\mathbf{E}_{i}, \mathbf{E}_{j}) = \begin{cases} r_{i}r_{j}(g-1) & \text{if } i \neq j \\ r_{i}r_{j}(g-1) + 1 & \text{if } i = j \end{cases}$$

and  $X_{i,j}^k$ ,  $k = 1, ..., d_{i,j}$  a basis of  $\operatorname{Ext}_0^1(E_i, E_j)^{\vee}$ . The ring

$$A_E \subset k[Ext^1(E, E)] = k[X_{i,j}^k, 1 \le i, j \le n, 1 \le k \le d_{i,j}]$$

is the ring generated by  $(Ext^1(E_1, E_1) \oplus Ext^1(E_2, E_2))_0^{\vee}$  and the products

$$\langle X_{1,2}^k \cdot X_{2,1}^l, 1 \le k, l \le d_{i,j} \rangle$$
.

Let  $\mathcal{S}$  be the affine cone of the Segre variety

$$P(Ext^{1}(E_{1}, E_{2})) \times P(Ext^{1}(E_{2}, E_{1})) \subset P(Ext^{1}(E_{1}, E_{2}) \otimes_{k} Ext^{1}(E_{2}, E_{1})).$$

PROPOSITION 1. There is an isomorphism

$$\operatorname{Spec}(A_{E}) \stackrel{\sim}{\to} (\operatorname{Ext}^{1}(E_{1}, E_{1}) \oplus \operatorname{Ext}^{1}(E_{2}, E_{2}))_{0} \times \mathscr{S}.$$

Note that  $Spec(A_E)$  is a cone.

COROLLARY 2. With the previous notations,  $SU_X(r, d)$  is étale locally isomorphic to  $(Ext^1(E_1, E_1) \oplus Ext^1(E_2, E_2))_0 \times \mathscr{S}$  at  $E = E_1 \oplus E_2$ .

Using furthermore that the multiplicity at the origin of the affine cone of a projective variety is just its degree, one gets the

COROLLARY 3. The Zariski tangent space is

$$(\operatorname{Ext}^{1}(E_{1}, E_{1}) \oplus \operatorname{Ext}^{1}(E_{2}, E_{2}))_{0} \oplus (\operatorname{Ext}^{1}(E_{1}, E_{2}) \otimes_{k} \operatorname{Ext}^{1}(E_{2}, E_{1})).$$

Moreover the multiplicity of  $SU_X(r, d)$  at [E] is

$$\text{mult}_{[E]}(SU_{X}(r, d)) = \begin{pmatrix} 2 \cdot d^{1,2} - 2 \\ d_{1,2} - 1 \end{pmatrix}$$

(recall that  $d_{1,2} = r_1 \cdot r_2(g-1)$ ).

Remark 4. Note that  $\binom{2 \cdot d_{1,2} - 2}{d_{1,2} - 1} = 1$  if and only if g = 2 and  $r_1 = r_2 = 1$ . Using that the singular locus of Sing  $SU_X(r, d)$  is closed in  $SU_X(r, d)$ , one obtains easily in this way another proof of the fact that Sing  $SU_X(r, d)$  is the non stable locus, except if g = r = 2 and d = 0.

# IV. Case 2: $G_E = Sl_r$

In this case,  $E = L^{\oplus r}$  where  $L^{\otimes r} = \emptyset$ . Using a translation by  $L^{-1}$  which induces an automorphism of  $SU_X(r, 0)$ , one may assume  $L = \emptyset$ . The ring  $A_E$  is the ring of polynomial maps on  $M_0(r)^g$  invariant under  $Sl_r$ . This group  $Sl_r$  acts diagonally by conjugation on each factor  $M_0(r)$ , which is the space of traceless matrices of size r. For general r, Procesi [P] and Rasmyslev [Ras] have obtained the following description of the first 2 syzigies of  $A_E$ :

- Generators: for every sequence  $\underline{i} = (i_1, \dots, i_{N(\underline{i})})$  of integers of  $[1, \dots, g]$ , let  $t_{\underline{i}}$  be the invariant polynomial map

$$t_i : \begin{cases} M_0(r)^g & \to & k \\ (X_1, \dots, X_g) & \mapsto & Tr(X_{i_1}, \dots, X_{i_N}) \end{cases}$$

Then the  $t_i$  with  $N(\underline{i}) \le 2^g - 1$  form a system of generators.

- Relations between all the  $t_i$ 's: Let  $P_X$  be the characteristic polynomial of the general matrix X. The homogeneous polynomial  $X \mapsto Tr(X \cdot P_X(X))$  gives by polarization (namely by taking the total differential of order g+1) a multilinear map  $F(H_1, \ldots, H_{g+1})$  where the  $H_i$ 's runs in the set of all possible monomials in the  $X_i$ .

Although this description is quite explicit, it looks difficult to obtain a complete **finite** set of relations between the (finite) set of generators constructed above.

As far as I know, the only case where such a finite description is available is for r = 2(\*).

<sup>(\*)</sup> According to some experts of invariant theory, it is more or less hopeless to obtain such a finite description of  $A_E$  in the general case.

#### Assume moreover that r=2

In this case,  $A_E$  can be described by using classical results of the geometric invariant theory of  $SO_3(k)$ . Following [LeB], I.4, let me briefly explain this description.

For  $X \in M_0(2)$ , let  $u(X) = (u_1(X), u_2(X), u_3(X)) \in k^3$  be defined by the equality

$$\mathbf{X} = \begin{pmatrix} u_1(\mathbf{X}) & u_2(\mathbf{X}) - \sqrt{-1}u_3(\mathbf{X}) \\ u_2 + \sqrt{-1}u_3(\mathbf{X}) & -u_1(\mathbf{X}) \end{pmatrix}.$$

By theorem 4.1 of [LeB] the isomorphism

$$\begin{cases} M_0(2) & \to & k^3 \\ X & \mapsto & u(X) \end{cases}$$

induces an identification of  $A_E$  with the polynomial maps of  $(k^3)^{\oplus g}$  invariant under the canonical diagonal action of  $SO_3(k)$ .

Let  $T_{i,j}$  be the invariant function corresponding to  $(u_1, \ldots, u_n) \mapsto (u_i \cdot u_j)$  (scalar product), namely  $T_{i,j}(X_1, \ldots, X_g) = \frac{1}{2} Tr(X_i X_j)$ .

Let  $T_{i,j,k}$  be the invariant function corresponding to  $(u_1, \ldots, u_n) \mapsto u_i \wedge u_j \wedge u_k$  (the wedge product lives in  $\wedge^3 k^3 \stackrel{\text{can}}{=} k$ ), namely  $T_{i,j,k}(X_1, \ldots, X_g) = Tr(X_i X_j X_k)$ . With some abuse of notation, one can now use the results of H. Weyl [W], theorem (2.9 A) and (2.17 B) and it's sequel on page 77 which says the following:

- Generators for the invariants maps under  $O_3(k)$ : the set

$$\langle T_{i,j} \rangle$$

- Relations between the generators: the 4-minors of the  $g \times g$ -symmetric matrix

$$\left\{
\begin{array}{ccc}
\vdots \\
\cdots & \mathbf{T}_{i,j} & \cdots \\
\vdots \\
\vdots & \vdots
\end{array}
\right\}$$

and the relations  $T_{i,j} = T_{j,i}$ . One recognizes the coordinate ring of the (affine) cone C of symmetric matrices of rank  $\leq 3$ . This scheme is well understood: it

is integral and normal [easy], Cohen-Macaulay [H-R], it multiplicity at the origin (or the degree of the projectivization PC) is known [H-T]...

- Generators for the  $SO_3(k)$ -invariants maps are: the  $T_{i,j}$ 's and the  $T_{i,j,k}$ 's.
- Relations: the previous 4-minors and:

$$T_{i_1,i_2,i_3} \cdot T_{j_1,j_2,j_3} = \det([T_{i_n,j_m}]_{1 \le n,m \le 3})$$
(1)

$$T_{i_0,i_4}T_{i_1,i_2,i_3} - T_{i_1,i_4}T_{i_0,i_2,i_3} + T_{i_2,i_4}T_{i_0,i_1,i_3} - T_{i_3,i_4}T_{i_0,i_1,i_2}$$
 (2)

and the relations given by the symmetry of  $T_{i,j,k}$  and the skew symmetry of  $T_{i,j,k}$  in the indices.

Let  $\bar{C}$  be the tangent cone of  $Spec(A_E)$  at the origin. It is the subscheme of

$$C \times \operatorname{Spec} k[T_{i,j,k}]$$

whose ideal is generated by

$$T_{i_1,i_2,i_3} \cdot T_{i_1,i_2,i_3}$$
 (3)

and

$$T_{i_0,i_4}T_{i_1,i_2,i_3} - T_{i_1,i_4}T_{i_0,i_2,i_3} + T_{i_2,i_4}T_{i_0,i_1,i_3} - T_{i_3,i_4}T_{i_0,i_1,i_2}$$
(4)

and relations given by the skew symmetry of  $T_{i,j,k}$  in the indices. The ideal described above is the ideal of initial forms of the ideals given by (1) and (2).

Let k(C) be the function field of C and K its algebraic closure. Note that, according to (3), the ideal  $I_{C/\bar{C}}$  of C in  $\bar{C}$  is nilpotent. This implies by [F] example 4.3.4, the formula

$$\operatorname{mult}_0 \bar{C} = \operatorname{length} \mathcal{O}_{C,\bar{C}} \cdot \operatorname{mult}_0(C). \tag{5}$$

The next formula is clear

$$\operatorname{length} \mathcal{O}_{C,\bar{C}} = 1 + \dim_{k(C)} I_{C/\bar{C}} \otimes_{\ell_{\bar{C}}} k(C) = 1 + \dim_{K} I_{C/\bar{C}} \otimes_{\ell_{\bar{C}}} K.$$
 (6)

One therefore has to compute the dimension over K of the sub-vector space  $V_T$  of the dual space of  $W = \bigoplus K \cdot T_{i,j,k}$  of equations given by (4) and the skew symmetry condition for the  $T_{i,j,k}$ . This vector space is isomorphic to  $I_{C/\bar{C}} \otimes_{\ell \in \bar{C}} K$ .

LEMMA 2. The dimension of  $V_T$  depends only on the conjugation class of T. Proof. The symmetric matrix

$$T = [T_{i,j}] \in M_g(K)$$

acts on the dual vector space V of  $K^g$ . Let  $(e_i)_{1 \le i \le g}$  be the canonical basis and  $(e_i^{\vee})_{1 \le i \le g}$  its dual basis. The map

$$\begin{cases} \mathbf{W} & \to & \wedge^{3}\mathbf{V} \\ \mathbf{T}_{i,j,k} & \mapsto & e_{i} \wedge e_{j} \wedge e_{k} \end{cases}$$

identifies W and  $\wedge$  <sup>3</sup>V $^{\vee}$ . With this identification, the relations (4) become

$$-T(e_{i_4}^{\vee}) \perp e_{i_0} \vee e_{i_1} \vee e_{i_2} \vee e_{i_3}$$

and dim V<sub>T</sub> is the corank of

$$\begin{cases} \mathbf{V}^{\vee} \otimes \wedge^{4} \mathbf{V} & \to & \wedge^{3} \mathbf{V} \\ x^{\vee} \otimes y & \mapsto & \mathbf{T}(x^{\vee}) \, \bot y \end{cases}$$

This map depends only on the conjugation class of T.

One can therefore assume that T is diagonal of eigenvalues  $\lambda_i$  with  $\lambda_i = 0$  for  $3 < i \le g$  and  $\lambda_i \ne 0$  if  $i \le 3$ .

Let us prove this simple lemma

LEMMA 3. The dimension of  $V_T$  is [g(g-1)(g-2)/6] - [(g-3)(g-4)(g-5)/6] if  $g \ge 3$  and 0 if  $g \le 2$ .

*Proof.* If  $g \le 2$ , the vector space  $\wedge^3 V$  is zero and so is  $V_T$ . Suppose  $g \ge 3$ . Consider an equation

$$T_{i_0,i_4}T_{i_1,i_2,i_3} - T_{i_1,i_4}T_{i_0,i_2,i_3} + T_{i_2,i_4}T_{i_0,i_1,i_3} - T_{i_3,i_4}T_{i_0,i_1,i_2}$$
(4)

defining  $V_T$ . If  $i_4 \notin \{1, 2, 3\}$  or  $i_4 \notin \{i_0, i_1, i_2, i_3\}$  then the equation (4) is trivial. Let me suppose that  $i_4 \in \{1, 2, 3\}$  and for instance that  $i_4 = i_0$ . If  $i_4 \in \{i_1, i_2, i_3\}$ , the equation is just a consequence of the skew symmetry of  $T_{i_1, i_2, i_3}$ . To get a

new relation, one has therefore further to suppose further that  $i_4 \notin \{i_1, i_2, i_3\}$  and the equation becomes

$$\lambda_{i_0} \cdot T_{i_1,i_2,i_3} = 0.$$

Of course, the other cases are obtained by symmetry. One has proved the following: the equations (4) are non trivial if and only if

$$\{i_0, i_1, i_2, i_3\} = \{i, j, k\} \cup \{i_4\}$$
 and  $i_4 \notin \{i, j, k\}$ .

In this case the relation (4) becomes

$$T_{i,i,k}=0$$
,

or equivalently

$$T_{i,i,k} = 0$$
 if  $\{i, j, k\} \cap \{1, 2, 3\} \neq \emptyset$ .

In particular this corank is

$$\dim \wedge^{3}V - A_{g-3}^{3} = \frac{g(g-1)(g-2)}{6} - \frac{(g-3)(g-4)(g-5)}{6}.$$

The degree  $d_g^r$  of the locus (\*) of  $g \times g$ -symmetric matrices of corank  $\geq r$  is computed in [H-T], proposition 12.b:

$$d_g^r = \deg \mathbf{PC} = \prod_{\alpha=0}^{r-1} \frac{\binom{g+\alpha}{r-\alpha}}{\binom{2\alpha+1}{\alpha}}.$$

Using the formulas (5) and (6) and the lemma 3, one obtains the

<sup>(\*)</sup> In [H-T], this locus is endowed with the reduced scheme structure. But it is known in full generality that the natural scheme structure given by the vanishing of the (g-r+1)-minors is Cohen-Macaulay [J] and generically reduced [easy] and therefore reduced. In our case (r=g-3), this reduceness is obvious, because C is a ring of invariants.

**THEOREM 4.** The multiplicity of  $[\mathcal{O} \oplus \mathcal{O}]$  in  $SU_x(2,0)$  is

$$\left(1+\frac{g(g-1)(g-2)}{6}-\frac{(g-3)(g-4)(g-5)}{6}\right)\cdot d_g^{g-3}$$

if  $g \ge 3$  and 1 if g = 2.

Remarks 5.

1. The preceding discussion gives a precise description of the tangent cone  $\bar{C}$  of  $SU_X(2, 0)$ . In particular, the Zariski tangent space at the trivial bundle is

$$T_{[\emptyset \oplus \emptyset]}SU_{\mathbf{X}}(2,0) = Sym^2 V \oplus \wedge^3 V.$$

2. One recovers the smoothness of  $SU_X(2, 0)$  if g = 2.

In the case of a non hyperelliptic genus 3 curve, the generalized  $\Theta$  divisor embeds  $SU_X(2,0)$  as the Coble quartic  $SU_X(2,0)$  in  $PH^0(J^2, 2 \cdot \Theta_{J^2})$  (see [N-R2] and [D-O] pages 184–185).

THEOREM 6. The scheme defined by the eight partial derivatives of the Coble quartic C(X) is reduced.

This theorem is the consequence of the

THEOREM 6 bis. (i) The (étale) local equation of the Coble quartic at the trivial bundle is

$$T^2 = \det([T_{i,i}]_{1 \le i,i \le 3})$$

in the affine space  $A^7$  with coordinates T,  $T_{i,j}$  with  $T_{i,j} = T_{j,i}$ .

- (ii) The (étale) local equation of the Coble quartic at  $E = E_1 \oplus E_2$  with  $E_1 \neq E_2$  is a rank 4 quadric in  $A^7$ .
- (iii) The ideal generated by the 8 cubic equations which are the partials derivatives of the Coble quartic is prime.

*Proof.* The first 2 points are clear from proposition II.2 and (1), (2). Let me prove (iii). Let  $\tilde{K}$  the scheme defined by the partial derivatives of an equation of C(X). The Kummer variety K(X) is the reduced scheme of  $\tilde{K}$ . It is therefore enough to prove that the completion of  $\tilde{K}$  at each non stable point E of  $K(X) \subset \tilde{K}$  is *reduced*. Because of the invariance of the Coble quartic under the

Theta group of  $2 \cdot \Theta_{J^2}$ , one has 2 cases to examine: either E is trivial, or  $E = E_1 \oplus E_2$  with  $E_1 \neq E_2$ . In the first case, by (i), the equations in  $k[[T, T_{i,j}]]$  of the completion of  $\tilde{K}$  at  $[\mathcal{O} \oplus \mathcal{O}]$  are T and the  $2 \times 2$ -minors of  $[T_{i,j}]_{1 \leq i,j \leq 3}$ . It is precisely the (completion at the origin) of the cone over the Veronese surface in  $P^5$  (with homogeneous coordinates  $T_{i,j}$ ) and K is therefore reduced. The second case is even simpler,  $\tilde{K}$  being (the completion of) a 3-plane in  $A^7$  (the tangent of K(X)).

### V. The case $SU_X(3,0)$ for of a genus 2 curve

Suppose in this section that X has genus 2 and let  $\mathcal{M} = SU_X(3,0)$  be the moduli space of rank 3 semi-stable vector bundles on X with trivial determinant. Consider a non stable point E of  $\mathcal{M}$ .

The case  $G_E = G_m$  has been treated in section II: in this case, the completion of  $\mathcal{M}$  at E is the completion at the origin of a rank 4 quadric in  $A^9$ .

Suppose now that  $G_E = G_m \times G_m$  which means

$$E = E_1 \oplus E_2 \oplus E_3$$
 with  $E_i \neq E_j$  for  $i \neq j$  and  $deg(E_i) = 0$ .

Let  $X_{i,j}$  be a basis of  $\operatorname{Ext}^1(E_i, E_j)^{\vee}$  for  $i \neq j$ . Let  $\mathscr{A}_E = k[X_{i,j}]^{G_E}$  be the ring of invariants of  $k[X_{i,j}]$  under  $G_E$  with the action defined by the following rule

$$\underline{\alpha} \cdot \mathbf{X}^{\underline{n}} = \prod \left( \frac{\alpha_i}{\alpha_j} \right)^{n_{i,j}} \mathbf{X}^{\underline{n}}$$

for  $(\alpha_1, \sigma_2, \alpha_3) \in G_E(k) = (k^*)^3/k^*$ . The following equality is easy

$$A_E = k[(\bigoplus Ext^1(E_i, E_i))_0] \otimes \mathscr{A}_E.$$

A polynomial  $\sum p_n X^n$  is in  $\mathcal{A}_E$  if and only if

$$\sum_{j} n_{j,i} = \sum_{j} n_{i,j} \qquad \text{for } i = 1, 2, 3$$
 (1)

if  $p_n \neq 0$ . Therefore,  $\mathscr{A}_{\mathrm{E}}$  is generated by the monomials

 $X^n$  such that n satisfies (1).

LEMMA 1. The ring  $\mathcal{A}_E$  is generated by

$$X_{3,2}X_{2,1}X_{1,3}$$
,  $X_{1,2}X_{2,3}X_{3,1}$  and  $X_{i,j}X_{j,i}$ ,  $i < j$ .

*Proof.* put  $\delta_{i,j} = n_{i,j} - n_{j,i}$  and let  $\underline{n}$  be a multi-index satisfying (1). The relations (1) become

$$\delta_{1,2} + \delta_{1,3} = 0$$
,  $\delta_{1,2} = \delta_{2,3}$ ,  $\delta_{1,3} + \delta_{2,3} = 0$ .

If  $\delta^+ = \delta_{1,2} \ge 0$ , we write  $\underline{n} = (n_{2,1} + \delta^+, n_{2,1}, n_{1,3}, n_{1,3} + \delta^+, n_{3,2} + \delta^+, n_{3,2})$  and use the monomial  $X_{1,2}X_{2,3}X_{3,1}$  corresponding to  $\underline{n}_0 = (1, 0, 0, 1, 1, 0)$  to write  $\underline{n} = \underline{m} + \delta^+ \cdot \underline{n}_0$ . This allows us to write

$$X^n = (X_{1,2}X_{2,3}X_{3,1})^{\delta+} \prod_{i < j} (X_{i,j} \cdot X_{j,i})^{m_{i,j}}$$

with  $m_{i,j} \ge 0$ . In the same way, when  $\delta^- = -\delta_{1,2} > 0$ , one has an equality

$$X^n = (X_{3,2}X_{2,1}X_{1,3})^{\delta-} \prod_{i < j} (X_{i,j} \cdot X_{j,i})^{m_{i,j}}$$

with  $m_{i,j} \geq 0$ .

For  $i \in I = \{1, 2, 3\}$ , put

$$x_i = X_{j,k},$$
  $x_{i+3} = X_{k,j},$   $\zeta_i = x_i x_{i+3}$ 

with  $I = \{i, j, k\}$  and j < k. Let  $\zeta_4 = X_{3,2}X_{2,1}X_{1,3}$  and  $\zeta_5 = X_{1,2}X_{2,3}X_{3,1}$ . There is an equality

$$\zeta_4\zeta_5=\zeta_1\zeta_2\zeta_3.$$

PROPOSITION 2. The natural morphism

$$f : \begin{cases} k[\mathbf{X}_i] & \to & \mathcal{A}_{\mathbf{E}} \\ \mathbf{X}_i & \mapsto & \zeta_i \end{cases}$$

gives an isomorphism

$$k[X_i]/(X_4X_5-X_1X_2X_3) \stackrel{\sim}{\to} \mathscr{A}_E$$
.

*Proof.* Let p be the (prime) ideal generated by  $X_4X_5 - X_1X_2X_3$ . Let

$$\mathbf{P} = \sum \alpha_n \mathbf{X}^{\underline{n}}$$

be an element of Ker(f). Then, one finds by simple expansion

$$f(\mathbf{P}) = \sum \alpha_n \prod x^{\phi(n)} = \sum_{m} x^m \sum_{\phi(n) = m} \alpha_n$$

with

$$\phi(\underline{n}) = (n_1 + n_5, n_2 + n_4, n_3 + n_5, n_1 + n_4, n_2 + n_5, n_3 + n_5)$$

which implies

$$\sum_{\phi(n)=m} \alpha_n = 0. \tag{2}$$

(Here  $X^n = \prod_i X_i^{n_i}$  and  $x^m = \prod_i x_i^{m_i}$   $\underline{n} = (n_i)_{1 \le i \le 5}$  and  $\underline{m} = (m_i)_{1 \le i \le 6}$  are multi-indices).

The kernel of  $\phi$  is generated by (1, 1, 1, -1, -1): if  $\phi(\underline{m}') = \phi(\underline{m})$ , there exists  $\alpha \in \mathbb{Z}_+$  such that

$$\pm \alpha(1, 1, 1, 0, 0) + m' = \pm \alpha(0, 0, 0, 1, 1) + m.$$

In particular, one has the congruence

$$(X_4 \cdot X_5)^{\alpha} \cdot X^{\underline{m}'} \equiv (X_4 \cdot X_5)^{\alpha} \cdot X^{\underline{m}} \bmod \mathfrak{p}. \tag{3}$$

According to (2) and (3), we get the existence of a positive integer a such that

$$(\mathbf{X}_4 \cdot \mathbf{X}_5)^a \cdot \mathbf{P} \equiv 0 \bmod \mathfrak{p}.$$

Since the ideal  $\mathfrak{p}$  is prime and  $(X_4 \cdot X_5) \notin \mathfrak{p}$  and therefore  $P \in \mathfrak{p}$ .

COROLLARY 3. Let E a point of  $\mathcal{M}$  satisfying  $G_E = G_m \times G_m$ . Then  $\mathcal{M}$  is étale locally isomorphic at E to

$$(\bigoplus \text{Ext}^{1}(\mathbf{E}_{i}, \mathbf{E}_{i}))_{0} \times \text{Spec}(k[\mathbf{X}_{i}]/(\mathbf{X}_{4}\mathbf{X}_{5} - \mathbf{X}_{1}\mathbf{X}_{2}\mathbf{X}_{3})).$$

Its tangent cone is a rank 2 quadric in the Zariski tangent space  $T_{[E]}\mathcal{M} = A^9$ .

In particular, there exists a family E of semi-stable bundles of trivial determinant over a germ of curve such that:

- (i) The group  $G_{E_n}$  of the generic bundle  $E_{\eta}$  is  $G_m \otimes_k k(\eta)$ .
- (ii) The group of  $G_{\mathbf{E}_s}$  the special bundle  $\mathbf{E}_s$  is  $G_m \times G_m$ .
- (iii) The multiplicity of  $\mathcal{M}$  at  $[\mathbf{E}_n]$  and  $[\mathbf{E}_s]$  are the same.

This shows that 2 points of  $\mathcal{M}$  can have the same multiplicity without having the same group of automorphisms.

When the E has 3 summands for which at least 2 are isomorphic, or equivalent if  $G_E$  is not a torus, the calculations are very intricate (but seem to be possible). In fact, one can in spite of this obtain the following

PROPOSITION 4. The tangent cone at each non stable point E such that  $G_E$  is not a torus is a quadric in  $A^9$  of rank  $\leq 2$ .

*Proof.* let E be a semi-stable vector bundle of rank 3 and determinant  $\mathcal{O}$  and  $\mathcal{O}_{J^1}$  the canonical theta divisor on Pic<sup>1</sup>(X). Using the corollary 1.7.4 of [Ray], the determinantal locus  $\mathcal{O}_E$  in Pic<sup>1</sup>(X)

$$\Theta_E = \{L \in Pic^1(X) \text{ such that } H^0(X, E \otimes L) \neq 0\}$$

is a divisor in  $PH^0(J^1, \mathcal{O}(3\Theta_{J^1}))$ . The Picard group of  $\mathcal{M}$  is cyclic with ample generator  $\mathcal{O}(\Theta)$  [D-N]. By [B-N-R], the inverse image of  $\mathcal{O}(1)$  by the morphism

$$\pi : \begin{cases} \mathscr{M} & \to & \mathbf{P}\mathbf{H}^0(\mathbf{J}^1, \mathcal{O}(3\Theta_{\mathbf{J}^1})) \\ [E] & \mapsto & \Theta_{\mathbf{E}} \end{cases}$$

is  $\mathcal{O}(\Theta)$  and  $\pi^*$  is a (canonical) isomorphism

$$|\Theta|^{\vee} \stackrel{\sim}{\to} PH^0(J^1, \mathcal{O}(3\Theta_{J^1})).$$

Using this isomorphism,  $\pi$  becomes the morphism given by the complete linear system  $|\Theta|$  (in particular,  $|\Theta|$  has no base point in this case!). There are various ways to prove this simple lemma, but the following one can be generalized for the higher rank case.

# LEMMA 5. The morphism $\pi$ is finite of degree 2 over $\mathbf{P}^8$ .

Proof of the lemma. Using the isomorphism  $\pi^*\mathcal{O}(1) \xrightarrow{\sim} \mathcal{O}(\Theta)$ , we get that  $\pi$  is finite of degree  $c_1(\Theta)^8$  onto  $\mathbf{P}^8$ . One therefore has to compute the degree of  $\pi$ . Although there exists a general beautiful formula due to Witten to evaluate the volume

$$\frac{c_1(\Theta)^{\dim \mathrm{SU}_{\mathbf{X}}(r0)}}{(\dim \mathrm{SU}_{\mathbf{X}}(r,0))!},$$

we give a simplest (and elementary) method to get this volume for  $\mathcal{M}$ . One has to prove that the leading term of the Hilbert polynomial  $n \mapsto P(n) = \chi(X, \Theta^n)$  is 2/8!. The canonical divisor of  $\mathcal{M}$  is  $\Theta^{-6}$  [D-N]. Serre duality implies therefore the symmetry

$$P(n) = P(-6-n).$$
 (4)

The Grauert-Reimenschneider vanishing theorem (recall that  $\mathcal{M}$  has rational singularities) gives the equality

$$P(n) = \dim H^0(\mathcal{M}, \Theta^n)$$
 for  $n \ge -5$ .

One therefore obtains the values

$$P(n) = 0$$
 for  $n = -5, ..., -1$ ,  $P(0) = 1$  and  $P(1) = 9$ . (5)

By (4) and (5), one obtains

$$P(X) = \lambda(X+5)(X+2)(X+3)^{2}(X+2)(X+1)(X-\alpha)(X+6+\alpha).$$

The equalities P(0) = 1 and P(1) = 9 imply

$$\alpha = -3 \pm \sqrt{-47}$$
 and  $\lambda = \frac{2}{8!}$ .

One has proved that the morphism  $\pi$  is finite of degree 2 onto  $|\Theta|^{\vee} = \mathbb{P}^8$ . Since  $\mathscr{M}$  is Cohen-Macaulay and  $\mathbb{P}^8$  smooth, this double covering is flat ([EGA IV]15.4.2) and is given locally by an equation

$$t^2 = f(\underline{x})$$

where  $\underline{x}$  are local coordinates on  $\mathbf{P}^8$ . This implies that the multiplicity of each point of  $\mathcal{M}$  is  $\leq 2$ .

Take a point  $E \in \mathcal{M}$  such that  $G_E$  is not a torus: it is a non smooth point of  $\mathcal{M}$ , therefore the tangent cone is a quartic (the initial term of f is not linear). But, such a point is a specialization of a point  $E_{\eta}$  such that  $G_{E_{\eta}} = G_m \times G_m$ : by the (obvious) semi-continuity of the rank of the quartic cone of E, the inequality rank  $\leq 2$  follows from the corollary 3.

Remark 6 (Dolgacev). The morphism  $\pi$  is ramified along a sextic S(X) in  $PH^0(J^1, \mathcal{O}(3\Theta_{J^1}))$ . Consider the embedding

$$\mathbf{J}^1 \hookrightarrow |3\boldsymbol{\Theta}_{\mathbf{J}^1}|^{\vee} = \mathbf{P}^8.$$

The variety  $J^1$  is contained in 9 quadrics. Using the action of the Mumford group as in I, it can be shown that there exists a unique cubic  $\mathscr{C}(X)$  of  $|3\Theta_{J^1}|^{\vee}$  which is both invariant under the Mumford group and singular along  $J^1$ .

QUESTION (Dolgacev). Are the sextic S(X) and the cubic C(X) dual to each other?

# VI. Multiplicity of the theta divisor (rank 2 case)

Recall that there exists a (Cartier) divisor  $\Theta$  on  $SU(2, \omega_X)$  which is characterized by the following universal property [D-N]: let S be a k-scheme and E a family of semi-stable vector bundles over  $X_S = X \times_k S$  of determinant  $\omega_X$ . Let  $\pi: S \to SU_X(2, \omega_X)$  be the classifying map corresponding to E. Then, one has the equality

$$\pi^*(\Theta) = \operatorname{div}(\det Rp_*E)^{\vee}.$$

By construction, the geometric points of  $\Theta$  are the classes  $[E] \in SU(2, \omega_X)$  such that  $H^0(X, E) \neq 0$ .

Remark. The vanishing of  $H^0(X, E)$  depends on the S-equivalence class [E] of E. Let E be a semi-stable vector bundle of determinant  $\omega_X$ . Recall ([La], theorem III.3) that for E stable

$$\operatorname{mult}_{[E]} \Theta = \operatorname{dim} H^0(X, E)$$

and that the tangent cone is defined in  $\operatorname{Ext}_0^1(E, E)$  by the ideal of the determinant of linear forms defined by the cup-product

$$H^0(X, E) \otimes Ext_0^1(E, E) \rightarrow H^1(X, E)$$
.

These facts can be generalized formally (using the universal property of  $\Theta$ ) as follows. Let  $[E] \in Su(2, \omega_X)$  a non stable point of  $\Theta$  of graded object  $E = E_1 \oplus E_2$  and let

$$h = \frac{1}{2} \dim H^0(X, E) = \dim H^0(X, E_1) = \dim H^0(X, E_2)$$

(note that  $E_1 \otimes E_2 = \omega_X$  which implies by Serre duality and Riemann-Roch the equality

$$\dim H^0(X, E_2) = \dim H^1(X, E_2) = \dim H^0(X, E_1). \tag{1}$$

With the notations of I, let  $V = \operatorname{Spec} k[[\operatorname{Ext}_0^1(E, E)]]$  be a (formal) étale slice of  $\mathcal{Q}$  at E and

$$\pi: V \to V/Aut(E) \xrightarrow{\text{étale}} SU(2, \omega_X)$$

the canonical morphism. Then, the induced map

$$\pi^*(\Theta)/\operatorname{Aut}(E) \to \Theta$$

is étale. The tangent cone of  $\pi^*(\Theta)$  is given by the determinant

$$d_{\rm E} \in \operatorname{Sym}^{2h} \operatorname{Ext}_0^1({\rm E},{\rm E})^{\vee}$$

defined (up to a non zero scalar) by the cup-product

$$H^0(X, E) \otimes Ext_0^1(E, E) \rightarrow H^1(X, E)$$

In particular, a point  $e \in \operatorname{Ext}_0^1(E, E)$  is in the tangent cone of  $\pi^*(\Theta)$  at [E] if and only if the cup-product

$$\cup e: H^0(X, E) \rightarrow H^1(X, E)$$

is not onto.

**PROPOSITION** 1. Assume that  $E_1 \neq E_2$ . Then, the multiplicity of  $\Theta$  at [E] is

$$\text{mult}_{[E]} \Theta = \frac{1}{2} \text{dim } H^0(X, E) \cdot \text{mult}_{[E]} SU_X(2, 0).$$

*Proof.* With the notation of the second section, the completion of  $SU(2, \omega_X)$  at [E] is the completion at the origin of

$$(\operatorname{Ext}^{1}(\operatorname{E}_{1},\operatorname{E}_{1}) \oplus \operatorname{Ext}^{1}(\operatorname{E}_{2},\operatorname{E}_{2}))_{0} \times \mathscr{S}$$

where  $\mathcal{S}$  is the cone over the Segre variety

$$P(Ext^{1}(E_{1}, E_{2})) \times P(Ext^{1}(E_{2}, E_{1})) \subset P(Ext^{1}(E_{1}, E_{2})) \otimes_{k} Ext^{1}(E_{2}, E_{1})).$$

Fix coordinates

$$\underline{\mathbf{X}} = (\mathbf{X}_{i,i}^k)$$
 on  $\mathrm{Ext}^1(\mathbf{E}_i, \mathbf{E}_i)$  for  $i \neq j$  and  $(\underline{\mathbf{Y}}) = \mathbf{Y}_i^k$  on  $\mathrm{Ext}^1(\mathbf{E}_i, \mathbf{E}_i)$ .

The equation F of  $\pi^*(\Theta)$  is of the form

$$\mathbf{F} = d_{\mathbf{E}} + \mathbf{G}_{2h+1}$$

where  $G_{2h+1}$  vanishes at the origin with order  $\geq 2h+1$ . The polynomials  $d_E$  and  $G_{2h+1}$  are  $G_E$  invariant and therefore (see section 2) can be written in terms of  $\underline{Y}$  and  $z^{k,l} = X_{1,2}^k \cdot X_{2,1}^l$ . Let me decompose  $d_E$  as

$$d_{\mathbf{E}} = \sum_{i=0}^{2h} \mathbf{Q}_i(\underline{\mathbf{X}}) \mathbf{P}_{2h-i}(\underline{\mathbf{Y}})$$

where the degree of  $Q_i$  (resp.  $P_{2h-i}$ ) is i (resp. 2h-i) and  $P_0=1$ . Using the degrees, one finds the following properties:

- If i is odd, then  $Q_i(\underline{X})P_{2h-i}(\underline{Y}) = 0$ .
- If  $P_{2h-2i} \neq 0$ , then  $Q_{2i}$  is invariant and therefore

$$Q_{2i}(\underline{X}) = R_i(\underline{z})$$

with  $R_i$  is a polynomial in  $\underline{z}$  of degree i which is defined up to the ideal of the Segre cone  $\mathscr{S}$ .

It follows that the equation of  $\pi^*(\Theta)$  can therefore be written as

$$R_h(\underline{z}) + S(\underline{z}, \underline{Y}) \tag{2}$$

where S vanishes at the origin with order  $\geq h+1$  at the origin.

LEMMA 2. The polynomial  $Q_{2h}(\underline{X})$  is non zero.

Proof of the lemma. According to the previous discussion, one just has to prove the existence of  $e \in \operatorname{Ext}^1(E_1, E_2) \oplus \operatorname{Ext}^1(E_2, E_1) \subset \operatorname{Ext}^1_0(E, E)$  such that the cup product  $\cup e \colon H^0(X, E) \to H^1(X, E)$  is onto. By symmetry, one only has to prove the existence of  $e_1 \in \operatorname{Ext}^1(E_1, E_2)$  such that the cup product  $\cup e_1 \colon H^0(X, E_1) \to H^1(X, E_2)$  is onto. This is classical (see [La], lemma II.8): let  $\Gamma$  be the variety

$$\Gamma = \{(k \cdot s, k \cdot e) \in \mathbf{PH}^0(\mathbf{X}, \mathbf{E}_1) \times \mathbf{P} \, \mathrm{Ext}^1(\mathbf{E}_1, \mathbf{E}_2) \, \mathrm{such that } s \cup e = 0\}$$

and p (resp. q) the first (resp. second) projection. Let  $0 \neq s \in H^0(X, E_1)$  and D = div(s) its zero divisor. The canonical surjection  $\cup s : \mathcal{H}om(E_1, E_2) \rightarrow E_2(-D)$  gives a surjection

$$\cup s: \operatorname{Ext}^{1}(\operatorname{E}_{1}, \operatorname{E}_{2}) \twoheadrightarrow \operatorname{H}^{1}(\operatorname{X}, \operatorname{E}_{2}). \tag{3}$$

By (3) the dimension of  $p^{-1}(k \cdot s)$  is

$$\dim p^{-1}(k \cdot s) = \dim \mathbf{P} \operatorname{Ext}^{1}(\mathbf{E}_{1}, \mathbf{E}_{2}) - \mathbf{P} \dim \mathbf{H}^{0}(\mathbf{X}, \mathbf{E}_{1}) - 1$$

Therefore dim 
$$\Gamma = \dim \mathbf{P} \operatorname{Ext}^1(\mathbf{E}_1, \mathbf{E}_2) - 1$$
 and  $q(\Gamma) \neq \Gamma$ .

The polynomial  $R_h$  can be thought of as an element of

$$H^{0}(\mathbf{P} \operatorname{Ext}^{1}(E_{1}, E_{2}) \times \mathbf{P} \operatorname{Ext}^{1}(E_{2}, E_{1}), \mathcal{O}(h, h))$$

which by the lemma 2 is non zero. According to (2), the completion of the tangent cone is therefore the hypersurface of  $(Ext^1(E_1, E_1) \oplus Ext^1(E_2, E_2))_0 \times \mathcal{S}$  given by  $R_h$ . The proposition follows.

### Acknowledgements

We would like to thank A. Beauville for useful discussions and M. Van Der Put for helping me to correct the english of a preliminary version. We thank also the referee for valuable comments.

#### REFERENCES

- [B-N-R] A. BEAUVILLE, M. S. NARASIMHAN and M. S. RAMANAN, Spectral curves and the generalized theta divisor, J. reine angew. Math. 398 (1989) 169-179.
- [B] J. F. BOUTOT, Singularites rationelles et quotients par les groupes reductifs, Inv. Math. 88 (1987) 65-68.
- [C] A. B. Coble, Algebraic geometry and Theta functions, Amer. Math. Soc. (1929).
- [D-N] J. M. DREZET and M. S. NARASIMHAN, Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques, Invent. Math. 97 (1989) 53-94.
- [D-O] I. DOLGACEV and D. ORTLAND, Point sets in projective spaces and theta functions, Astérisque, 165 (1988).
- [EGA IV] A. GROTHENDIECK, Elements de géométrie algébrique, avec la collaboration de J. Dieudonne, Publ. Math. I.H.E.S. 24 (1965).
- [F] W. FULTON, *Intersection theory*, Ergeb. der Math. und ihrer Grenzgebiete, 3. Folge Band 2, Springer-Verlag, Berlin Heidelberg New York Tokyo (1984).
- [H-R] M. HOCHSTER, J. ROBERTS, Rings of invariants of reductive groups acting on regular local rings are Cohen-Macaulay, Adv. in Math. 13 (1974) 115-175.
- [H-T] J. HARRIS and L. W. Tu, On symmetric and skew symmetric determinantal varieties, Topology 23 (1984) 71-84.
- [Ha] R. HARTSHORNE, Algebraic Geometry, GTM 52, Springer Verlag, Berlin-Heidelberg-New York (1977).
- [J] T. JOZEFIAK, *Ideals generated by minors of a symmetric matrix*, Comment. Math. Helvetici 53 (1978) 595-607.
- [Kh1] A. KHALED, Projective normality and equations of Kummer varieties, J. reine angew. Math. 465 (1995), 197-217.
- [Kh2] A. KHALED, Equations des variétés de Kummer, Math. Annal. 295 (1993) 685-701.
- [Ko] S. KOIZUMI, Theta relations and projective normality of Abelian varieties, Amer. Journ. of Math. 98 (1976) 865-889.
- [La] Y. LASZLO, Un théorème de Riemann pour les diviseurs Thêta généralisés sur les espaces de modules de fibrés stables sur une courbe, Duke Math. Journ. 64 (1991) 333-347.
- [LeB] L. LE BRUYN, Trace rings of generic 2 by 2 matrices, Mem. A.M.S. 66 (1987).
- [Lu] D. Luna, Slices étales, Mém. Soc. Math. France 33 (1973) 81-105.
- [M] D. MUMFORD, On equations defining Abelian varieties, Invent. Math. 1 (1966) 287-354.
- [N-R1] M. S. NARASIMHAN and M. S. RAMANAN, Moduli of vector bundles on a compact Riemann surface, Ann. of Math. 89 (1969) 14-51.
- [N-R2] M. S. NARASIMHAN and M. S. RAMANAN, 2θ-linear system on abelian varieties, in Vector bundles and algebraic varieties, Oxford University Press, (1987) 415-427.
- [P] C. PROCESI, The invariant theory of n by n matrices, Adv. in Math. 19, 306-381.
- [Ras] J. RASMYSLEV, Trace identities of full matrix algebras over a field of characteristic zero, Izv. Akad. Nauk. U.S.S.R. 8 (1974) 727-760.
- [Ray] M. RAYNAUD, Sections des fibrs vectoriels sur une courbe, Bull. Soc. Math. France 110 (1982) 103-125.
- [S] C. S. SESHADRI, Fibrés vectoriels sur les courbes algébriques (rédigé par J. M. Drezet), Astérisque 96 (1982).

[W] H. WEYL, The classical groups, Princeton Univ. Press, Princeton, N.J. (1946).

[W] H. WIRTINGER, Untersuchung über thetafunctionen, Teubner, Leipzig (1935).

Yves Laszlo Ecole Normale Supérieure, D.M.I. 45, rue d'Ulm 75005 Paris France e-mail: laszlo@dmi.ens.fr

Received May 30, 1994