

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 72 (1997)

Artikel: On Macdonald's formula for the volume of a compact Lie group
Autor: Hashimoto, Yoshitake
DOI: <https://doi.org/10.5169/seals-54611>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On Macdonald's formula for the volume of a compact Lie group

Yoshitake Hashimoto

Abstract. We give a simple proof of Macdonald's formula for the volume of a compact Lie group.

Mathematics Subject Classification (1991). 22E15.

Keywords. Compact Lie group, Todd class.

It is well-known that a compact Lie group G of rank r is rational homotopy equivalent to a product of spheres

$$S = S^{2m_1+1} \times \dots \times S^{2m_r+1}$$

where (m_1, \dots, m_r) is called the exponent of G . In [3] I. G. Macdonald calculated the volume of G with respect to the Haar measure μ induced from a Lebesgue measure λ on the Lie algebra \mathfrak{g} of G . The result is as follows:

$$\mu(G) = \lambda(\mathfrak{g}/\mathfrak{g}_{\mathbb{Z}})\sigma(S)$$

where $\mathfrak{g}_{\mathbb{Z}}$ is the Chevalley lattice and σ is the product of the superficial measure of the unit spheres S^{2m_i+1} in \mathbb{R}^{2m_i+2} , that is

$$\sigma(S^{2m+1}) = \frac{2\pi^{m+1}}{m!}.$$

In this note we shall give a simpler proof of this formula.

Let T be a maximal subgroup of G , \mathfrak{t} its Lie algebra and $\mathfrak{t}_{\mathbb{Z}}$ a lattice in \mathfrak{t} such that the kernel of the exponential map from \mathfrak{t} to T is $2\pi\mathfrak{t}_{\mathbb{Z}}$. We fix a positive definite inner product on \mathfrak{g} which is invariant under the adjoint action of G such that the induced volume form coincides with λ . This inner product induces Lebesgue measures on \mathfrak{t} and $\mathfrak{g}/\mathfrak{t}$ (also denoted by λ), which determine a Haar measure on T and a G -invariant measure on the flag manifold G/T (also denoted by μ). Then clearly

$$\mu(T) = (2\pi)^r \lambda(\mathfrak{t}/\mathfrak{t}_{\mathbb{Z}}), \quad \mu(G) = \mu(T)\mu(G/T).$$

The main problem is to compute the volume of the flag manifold. We compute at a point of the flag manifold the ratio of the volume form μ and another G -invariant top form, the top term of the Todd class, whose integral value is known to be 1 [1]. The adjoint action of G on \mathfrak{g} induces trivialization of the direct sum of the tangent bundle of G/T and a trivial \mathfrak{t} bundle. Hence the rational Pontrjagin classes of G/T all vanish and the Todd class equals $\exp \frac{1}{2}c_1(G/T)$. The top term is $\frac{1}{n!}(\frac{1}{2}c_1)^n$ where n is the complex dimension of G/T . Let $X_\alpha \in \mathfrak{g} \otimes \mathbb{C}$ be an eigenvector for a root α such that $\frac{1}{2}[X_\alpha, X_{-\alpha}] = \alpha^\vee$, the coroot for α , then

$$\frac{1}{2}c_1(X_\alpha, X_\beta) = \begin{cases} \frac{\langle \rho | \alpha^\vee \rangle}{\pi}, & \beta = -\alpha, \\ 0, & \beta \neq -\alpha. \end{cases}$$

where

$$\rho = \frac{1}{2} \sum_{\alpha > 0} \alpha,$$

on the other hand

$$\langle X_\alpha, X_{-\alpha} \rangle = |\alpha^\vee|^2.$$

Evaluating the volume form and $\frac{1}{n!}(\frac{1}{2}c_1)^n$ for $2n$ vectors $(X_\alpha, X_{-\alpha}; \alpha > 0)$, we obtain

$$\prod_{\alpha > 0} |\alpha^\vee|^2, \quad \prod_{\alpha > 0} \frac{\langle \rho | \alpha^\vee \rangle}{\pi}$$

respectively. Since α^\vee is the sum of $\langle \rho | \alpha^\vee \rangle$ simple coroots [2], it holds that

$$\prod_{\alpha > 0} \langle \rho | \alpha^\vee \rangle = \prod_{i=1}^r (m_i!).$$

Hence

$$\mu(G/T) = \frac{\sigma(S)}{(2\pi)^r} \prod_{\alpha > 0} |\alpha^\vee|^2.$$

The final result follows from

$$\lambda(\mathfrak{g}/\mathfrak{g}_{\mathbb{Z}}) = \lambda(\mathfrak{t}/\mathfrak{t}_{\mathbb{Z}}) \prod_{\alpha > 0} |\alpha^\vee|^2.$$

The author would like to thank K. Abe, A. Hattori and K. Ono for useful comments and encouragement.

References

- [1] A. Borel and F. Hirzebruch, Characteristic classes and homogeneous spaces II, *Amer. J. Math.* **81** (1959), 315–382.
- [2] N. Bourbaki, *Groupes et algèbres de Lie, chapitres 4, 5 et 6*, Hermann, Paris 1968.
- [3] I. G. Macdonald, The volume of a compact Lie group, *Invent. math.* **56** (1980), 93–95.

Yoshitake Hashimoto
Department of Mathematics
Osaka City University
Osaka, Japan

(Received: May 10, 1997)