

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 72 (1997)

**Artikel:** A theory of cobordism for non-spherical links  
**Autor:** Blablœil, Vincent / Michel, Françoise  
**DOI:** <https://doi.org/10.5169/seals-54576>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 05.02.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## A theory of cobordism for non-spherical links

Vincent Blanlœil and Françoise Michel

**Abstract.** We define an equivalence relation, called algebraic cobordism, on the set of bilinear forms over the integers. When  $n \geq 3$ , we prove that two  $2n-1$  dimensional, simple fibered links are cobordant if and only if they have algebraically cobordant Seifert forms. As an algebraic link is a simple fibered link, our criterion for cobordism allows us to study isolated singularities of complex hypersurfaces up to cobordism.

**Mathematics Subject Classification (1991).** 57R, 57R80, 57R90, 57M25, 57Q45, 32S, 32S55, 14B05.

**Keywords.** Knots and links, knot-cobordism, algebraic links, singularities.

### 0. Introduction

In this work we present a cobordism theory for links which is motivated by the study of the topology of isolated singularities of complex hypersurfaces. Let us be more precise:

(0.1) Let  $f : (\mathbb{C}^{n+1}, 0) \rightarrow (\mathbb{C}, 0)$ , be a holomorphic germ with an isolated singular point at the origin. We denote by  $D_\delta^{2k}$  the compact ball of radius  $\delta$  centred at 0 in  $\mathbb{C}^k$ , and by  $S_\delta^{2k-1}$  its boundary. The orientation-preserving homeomorphism class of the pair  $(D_\varepsilon^{2n+2}, f^{-1}(0) \cap D_\varepsilon^{2n+2})$  does not depend on the choice of a sufficiently small  $\varepsilon$ , by definition it is the *topological type* of  $f$ . The orientation preserving diffeomorphism class of the pair  $(S_\varepsilon^{2n+1}, K(f))$ , where  $K(f) = (f^{-1}(0)) \cap S_\varepsilon^{2n+1}$  is the link of  $f$ . The Milnor's conic structure theorem (see [M3, 68]) shows that the link  $K(f)$  determines the topological type of  $f$ . Moreover, J. Milnor has also proved that:

1.  $f/|f| : S_\varepsilon^{2n+1} \setminus K(f) \rightarrow S^1$  is a differentiable fibration which is trivial on  $U \setminus K(f)$ , when  $U$  is a sufficiently "small" open tubular neighbourhood of  $K(f)$ .
2. The manifold  $K(f)$  is  $(n-2)$ -connected.
3. The adherence  $F$  of a fiber of  $f/|f|$  is a compact, oriented,  $(n-1)$ -connected

smooth submanifold of  $S_\varepsilon^{2n+1}$  having  $K(f)$  as boundary. By definition  $F$  is the *Milnor fiber* of  $K(f)$ .

(0.2) More generally, we will say that a *link* is a  $(n-2)$ -connected, oriented, smooth, closed,  $(2n-1)$  dimensional submanifold of  $S^{2n+1}$ . A *knot* is a spherical link (i.e. a link abstractly homeomorphic to  $S^{2n-1}$ ). It is well-known that, for any link  $K$ , there exists a smooth, compact, oriented  $2n$ -submanifold  $F$  of  $S^{2n+1}$ , having  $K$  as boundary ; such a manifold  $F$  is called a *Seifert surface* for  $K$ .

(0.3) Following M. Kervaire [K1, 65], we say that two links  $K_0$  and  $K_1$ , abstractly diffeomorphic to the same manifold  $\mathcal{K}$ , are *cobordant* if there exists an embedding  $\Phi, \Phi : \mathcal{K} \times [0, 1] \rightarrow S^{2n+1} \times [0, 1]$ , such that:

$$\Phi(\mathcal{K} \times \{0\}) = K_0 \text{ and } \Phi(\mathcal{K} \times \{1\}) = -K_1,$$

where  $-K_1$  is the link  $K_1$  with the orientation reversed.

(0.4) Let  $F$  be a  $2n$  dimensional oriented smooth manifold of  $S^{2n+1}$ , and let  $G$  be the quotient of  $H_n(F, \mathbb{Z})$  by its  $\mathbb{Z}$ -torsion.

The *Seifert form* associated to  $F$  is the bilinear form  $A : G \times G \rightarrow \mathbb{Z}$  defined as follows (see also [K2, 70] p.88 or [L2, 70], p.185): let  $(x, y)$  be in  $G \times G$ , then  $A(x, y)$  is the linking number in  $S^{2n+1}$  of  $x$  and  $i_+(y)$ , where  $i_+(y)$  is the cycle  $y$  "pushed" in  $(S^{2n+1} \setminus F)$  by the positively oriented vector field normal to  $F$  in  $S^{2n+1}$ .

By definition a *Seifert form for a link*  $K$  is the Seifert form associated to a Seifert surface for  $K$ .

When  $n \geq 2$ , J. Levine ([L1, 69]) and M. Kervaire ([K2, 70]) gave a complete characterization of cobordism classes of knots in terms of Witt-equivalence classes of Seifert forms.

(0.5) A *simple link* is a link which has a  $(n-1)$ -connected Seifert surface. A link  $K$  is a simple fibered link if there exists a differentiable fibration  $\varphi : S^{2n+1} \setminus K \rightarrow S^1$ ,  $\varphi$  being trivial on  $U \setminus K$ , where  $U$  is a "small" open tubular neighbourhood of  $K$ , and having  $(n-1)$ -connected fibers, the adherence of which are Seifert surfaces for  $K$ . In this paper we define in §1 (see (1.2)) an equivalence relation on integral bilinear forms which is much more sophisticated than "Witt-equivalence" and the theorems 2 and 3, stated in §1, imply:

**Theorem A.** *If  $n \geq 3$ , two simple fibered links are cobordant if and only if they have algebraically cobordant Seifert forms.*

(0.6) By definition an *algebraic link* is a link  $K(f)$  associated, as described above, to a holomorphic germ  $f$  with an isolated singularity. Furthermore, Milnor's theory of singular complex hypersurfaces implies that algebraic links are simple fibered links. So theorem 2' and 3 stated in §1 imply:

**Theorem B.** *If  $n \geq 3$ , two algebraic links are cobordant if and only if the Seifert forms associated to their Milnor's fibers are algebraically cobordant.*

In [Lê, 72], D.T. Lê showed that two cobordant algebraic links of plane curves (i.e. when  $n = 1$ ) are isotopic. In [DB-M, 93], P. du Bois and F. Michel found (using the classical cobordism theory for knots of M. Kervaire and J. Levine), for all  $n \geq 3$ , examples of non isotopic but cobordant algebraic knots. But in general algebraic links are not spherical links. So theorem B gives a cobordism theory for algebraic links.

Furthermore, having algebraically cobordant Seifert forms is also a necessary condition of cobordism for simple fibered links when  $n$  is 1 or 2. So we obtain in §5, without any restriction of dimension, a "Fox-Milnor" relation (see [F-M, 66]) for the Alexander polynomials of cobordant simple fibered links which implies:

(0.7) **Corollary.** *Let  $K_0$  and  $K_1$  be two algebraic links having respectively  $\Delta_0$  and  $\Delta_1$  as characteristic polynomials of monodromy. If  $K_0$  and  $K_1$  are cobordant then the product  $\Delta_0 \cdot \Delta_1$  is a square in  $\mathbb{Z}[X]$ .*

(0.8) **Comments.** In [V1, 77] and [V2, 78] R. Vogt gave, when  $n \geq 3$ , a sufficient, but not necessary, condition of cobordism for simple links having torsion free homology groups. As shown in [DB-M, 93] the sufficient condition of cobordism for algebraic links given in [Sz, 89] by S. Szczepanski, cannot be true. So the problem of finding a criterion for cobordism of simple fibered links was largely open. Our definition of algebraic cobordism for Seifert forms solves the problem.

(0.9) In this paper we use the following **notations**: If  $X$  is a differentiable manifold we denote by  $\partial X$  its boundary, by  $\overset{\circ}{X}$  its interior and by  $H_k(X)$  the  $k^{th}$ -homology group of  $X$  with coefficients in  $\mathbb{Z}$ . If  $a$  is a  $k$ -cycle of  $X$  we denote by  $[a]$  its homology class in  $H_k(X)$ . If  $G$  is an abelian group let  $\text{rk}(G)$  be the rank of  $G$ , and  $\text{Tors}(G)$  be the torsion subgroup of  $G$ .

## 1. Definitions and statement of results

Let  $\mathcal{A}$  be the set of bilinear forms defined on free  $\mathbb{Z}$ -modules  $G$  of finite rank.

Let  $\varepsilon$  be  $+1$  or  $-1$ .

(1.1) If  $A$  is in  $\mathcal{A}$ , let us denote by  $A^T$  the transpose of  $A$ , by  $S$  the  $\varepsilon$ -symmetric form  $A + \varepsilon A^T$  associated to  $A$ , by  $S^* : G \rightarrow G^*$  the adjoint of  $S$  ( $G^*$  being the dual  $\text{Hom}_{\mathbb{Z}}(G; \mathbb{Z})$  of  $G$ ), by  $\overline{S} : \overline{G} \times \overline{G} \rightarrow \mathbb{Z}$  the  $\varepsilon$ -symmetric non degenerated form induced by  $S$  on  $\overline{G} = G / \text{Ker } S^*$ . A submodule  $M$  of  $G$  is pure if  $G/M$  is torsion free. If  $M$  is any submodule of  $G$  let us denote by  $M^\wedge$  the smallest pure submodule of  $G$  which contains  $M$ . In fact  $M^\wedge$  is equal to  $(M \otimes \mathbb{Q}) \cap G$ . For a submodule  $M$  of  $G$  we denote by  $\overline{M}$  the image of  $M$  in  $\overline{G}$ .

**Definition.** *Let  $A : G \times G \rightarrow \mathbb{Z}$  be a bilinear form in  $\mathcal{A}$ . The form  $A$  is Witt associated to 0 if the rank  $m$  of  $G$  is even and if there exists a pure submodule  $M$  of rank  $\frac{m}{2}$  in  $G$  such that  $A$  vanishes on  $M$  ; such a module  $M$  is called a*

metabolizer for  $A$ .

(1.2) **Definition.** Let  $A_i : G_i \times G_i \rightarrow \mathbb{Z}$ ,  $i=0,1$ , be two bilinear forms in  $\mathcal{A}$ . Let  $G$  be  $G_0 \oplus G_1$  and  $A$  be  $(A_0 \oplus -A_1)$ . The form  $A_0$  is algebraically cobordant to  $A_1$  if there exists a metabolizer  $M$  for  $A$  such that  $\overline{M}$  is pure in  $\overline{G}$ , an isomorphism  $\varphi$  from  $\text{Ker } S_0^*$  to  $\text{Ker } S_1^*$  and an isomorphism  $\theta$  from  $\text{Tors}(\text{Coker } S_0^*)$  to  $\text{Tors}(\text{Coker } S_1^*)$  which satisfy the two following conditions:

- c.1:  $M \cap \text{Ker } S^* = \{(x, \varphi(x)); x \in \text{Ker } S_0^*\}$ ,
- c.2:  $d(S^*(M)^\wedge) = \{(x, \theta(x)); x \in \text{Tors}(\text{Coker } S_0^*)\}$ , where  $d$  is the quotient map from  $G^*$  to  $\text{Coker } S^*$ .

In §2 (see (2.3)) we prove:

**Theorem 1.** Algebraic cobordism is an equivalence relation on the set  $\mathcal{A}$ .

(1.3) From now on,  $A_0$  and  $A_1$  will always be two Seifert forms associated to some  $(n-1)$ -connected Seifert surfaces  $F_0$  and  $F_1$ , of two simple links  $K_0$  and  $K_1$ . Let us justify the definition of algebraic cobordism. As a generalization of the Kervaire-Levine theory of knot cobordism we obtain in §3 (see (3.10)):

**Proposition.** If  $K_0$  and  $K_1$  are cobordant simple links, then  $A = A_0 \oplus -A_1$  has a metabolizer.

**Remark.** Let  $\varepsilon$  be  $(-1)^n$ , then for  $i=0,1$ ,  $S_i = A_i + \varepsilon A_i^T$  is the intersection form on  $H_n(F_i)$ ,  $\text{Ker } S_i^*$  is the image of  $H_n(K_i)$  in  $H_n(F_i)$  and  $\text{Coker } S_i^*$  is isomorphic to  $\tilde{H}_{n-1}(K_i)$ . So for spherical links, both  $\text{Ker } S_i^*$  and  $\text{Coker } S_i^*$  are zero, and conditions c.1 and c.2 in definition (1.2) vanish. Then, for spherical links, two Witt associated Seifert forms are algebraically cobordant, and we recover the Kervaire-Levine criterion for cobordism.

In the non-spherical case, the topology of the cobordism implies that the restriction of  $A_0$  on  $\text{Ker } S_0^*$  is isomorphic (on  $\mathbb{Z}$ ) to the restriction of  $A_1$  on  $\text{Ker } S_1^*$  (it is easy to check it directly, and it is also implied by the more general proposition (3.10)). This necessary condition for cobordism is not implied by the fact that  $A_0 \oplus -A_1$  is Witt associated to 0, but by condition c.1 in definition (1.2). The topology of the cobordism also implies that the linking forms on  $\text{Tors}(H_{n-1}(K_i))$  are isomorphic. This necessary condition for cobordism is contained in point c.2 of definition (1.2).

(1.4) The major result of this work is theorem 2 proved in §3 (see (3.10) and (3.13)):

**Theorem 2.** Let  $K_0$  and  $K_1$  be two cobordant simple links. If  $K_0$  and  $K_1$  have  $(n-1)$ -connected Seifert surfaces  $F_0$  and  $F_1$  with unimodular Seifert forms  $A_0$

and  $A_1$ , then  $A_0$  is algebraically cobordant to  $A_1$ .

**Remark.** Let  $i$  be 0 or 1. Let us suppose that  $K_i$  is a simple fibered link and let  $F_i$  be a  $(n-1)$ -connected fiber of a fibration  $\varphi_i : S^{2n+1} \setminus K_i \rightarrow S^1$ ; then, the Seifert form  $A_i$  associated to  $F_i$  is unimodular. Conversely, if  $n \geq 3$  and if  $A_i$  is unimodular then  $K_i$  is a simple fibered link (see [K-W, 77] chap. V, §3, p.118).

So, theorem 2 implies:

**Theorem 2'.** *Let  $K_0$  and  $K_1$  be two simple fibered links having  $F_0$  and  $F_1$  as  $(n-1)$ -connected fibers of differentiable fibrations  $\varphi_0$  and  $\varphi_1$ . If  $K_0$  is cobordant to  $K_1$ , then the Seifert forms  $A_0$  and  $A_1$ , associated respectively to  $F_0$  and  $F_1$ , are algebraically cobordant.*

(1.5) Using classical methods of surgery, we prove in §4 (see (4.4) and (4.5)):

**Theorem 3.** *Let  $n$  be greater or equal to 3 and let  $K_0$  and  $K_1$  be two  $2n-1$ -dimensional simple links. If the Seifert forms  $A_0$  and  $A_1$ , associated to some  $(n-1)$ -connected Seifert surfaces  $F_0$  and  $F_1$  of  $K_0$  and  $K_1$ , are algebraically cobordant then  $K_0$  is cobordant to  $K_1$ .*

(1.6) Proposition (3.10), which does not use (as remarked in (3.12)) any hypothesis on the Seifert forms, gives:

**Theorem 4.** *Let  $K_0$  and  $K_1$  be two cobordant simple links. If  $A_0$  (resp.  $A_1$ ) is a Seifert form associated to any  $(n-1)$ -connected Seifert surface for  $K_0$  (resp.  $K_1$ ), then  $A_0 \oplus -A_1$  has a metaboliser  $M$  such that  $M \cap \text{Ker } S^* = \{(x, \varphi(x)) ; x \in \text{Ker } S_0^*\}$ , where  $\varphi$  is an isomorphism between  $\text{Ker } S_0^*$  and  $\text{Ker } S_1^*$ .*

## 2. Algebraic cobordism

(2.0) Let  $A_0$  and  $A_1$  be two algebraically cobordant forms, let  $A$  be the form  $A_0 \oplus -A_1$  defined on  $G = G_0 \oplus G_1$  and  $S$  be  $A + \varepsilon A^T$ . In this section we prove proposition (2.1) which shows that the algebraic cobordism between  $A_0$  and  $A_1$  allows us to describe  $S$ ; this characterization of  $S$  is fundamental to prove theorem 3 (see §4). Let  $M$ ,  $\varphi$  and  $\theta$  be as in (1.2), let  $m$  be  $\text{rk}(G)$  and  $r$  be  $\text{rk}(\text{Ker } S_0^*)$ . Then definition (1.2) implies that  $s = \text{rk}(S^*(M)) = \frac{1}{2} \text{rk}(S^*(G))$  and  $\text{rk}(M) = r + s = \frac{m}{2}$ .

We use the following notations: if  $E$  is any subset of  $G$  we denote by  $\langle E \rangle$  the submodule of  $G$ , generated by  $E$ . If  $L$  is any submodule of  $G$  then:

$$L^\perp = \{x \in G \text{ s.t. } S(x, l) = 0 \ \forall l \in L\}$$

$$\text{Hom}_{\mathbb{Z}}(G|_L, \mathbb{Z}) = \{f \in G^* \text{ s.t. } f(l) = 0 \ \forall l \in L\}$$

Moreover if  $L_1$  and  $L_2$  are two submodules of  $G$ , orthogonal for  $S$ , we denote by  $L_1 \oplus^\perp L_2$  their (orthogonal) direct sum.

**Lemma.** *We have:  $S^*(G) \cap S^*(M)^\wedge = S^*(M^\perp)$ .*

*Proof.* Let  $r$  be the rank of  $\text{Ker } S_0^*$  and  $s$  be the rank of  $S^*(M)$ . As  $M$  is a metabolizer for  $S$  which fulfills condition c.1 in (1.2) we have:

$\text{rk}(\text{Ker } S^*) = 2 \text{rk}(M \cap \text{Ker } S^*) = 2r$ ,  $\text{rk}(S^*(G)) = 2s$  and  $\text{rk}(M^\perp) = s + 2r$ . Hence  $M^\perp = (M + \text{Ker } S^*)^\wedge$  and  $S^*(M^\perp) \subset S^*(G) \cap S^*(M)^\wedge$ .

Moreover,  $S^*(M)$  is of finite index in  $\text{Hom}_{\mathbb{Z}}(G|_{M^\perp}; \mathbb{Z})$ . As  $\text{Hom}_{\mathbb{Z}}(G|_{M^\perp}; \mathbb{Z})$  is a pure submodule of  $G^*$ , we get  $S^*(M)^\wedge = \text{Hom}_{\mathbb{Z}}(G|_{M^\perp}; \mathbb{Z})$ . So if  $S^*(x) \in S^*(M)^\wedge$ , then  $S^*(x, l) = 0$  for all  $l$  in  $M^\perp$  and  $x$  is in  $M^\perp$ .  $\square$

Since  $S^*(M)$  is of finite index in  $S^*(M)^\wedge$ , one can write  $(S^*(M)^\wedge)/S^*(M) \cong \bigoplus_{i=1}^s \mathbb{Z}/a_i \mathbb{Z}$  where  $a_i \in \mathbb{N} \setminus \{0\}$  and  $a_i$  divides  $a_{i+1}$  (we do not exclude that there exists an integer  $l$  such that  $a_i = 1$  for  $i = 1, \dots, l$ ).

**Proposition.** *The submodule  $\overline{M}$  is pure in  $\overline{G}$  if and only if  $S^*(M^\perp) = S^*(M)$ .*

*Proof.* We suppose that  $\overline{M}$  is pure in  $\overline{G}$ . As  $M \cap \text{Ker } S^* = \Delta(\varphi)$  has rank  $r$ , the rank of  $M + \text{Ker } S^*$  is  $s + 2r$ . So  $M + \text{Ker } S^*$  is of finite index in  $M^\perp$ . Let  $x$  be in  $M^\perp$ ; there exists a positive integer  $k$  such that  $kx = y + m$ , where  $y$  is in  $\text{Ker } S^*$ ,  $m$  is in  $M$ ; so  $\overline{m} = k\overline{x}$ . Since  $\overline{M}$  is pure in  $\overline{G}$  then  $\overline{x}$  is in  $\overline{M}$ , so there exists  $y'$  in  $\text{Ker } S^*$  such that  $x + y'$  is in  $M$ . Finally  $S^*(x) = S^*(x + y') \in S^*(M)$ , and  $S^*(M^\perp) \subset S^*(M)$ . But  $M \subset M^\perp$  so  $S^*(M^\perp) = S^*(M)$ .

We suppose that  $S^*(M) = S^*(M^\perp)$ . First we prove that  $\overline{M^\perp}$  is pure in  $\overline{G}$ . Let  $z$  be in  $M^\perp$  with  $\overline{z} = k\overline{x}$  where  $x$  is in  $G$  and  $k$  is a positive integer. So there exists  $y$  in  $\text{Ker } S^*$  such that  $kx = z + y$ . For all  $m$  in  $M$  we have  $S(kx, m) = S(z + y, m) = 0$ , so  $S(x, m) = 0$  and  $x$  is in  $M^\perp$ . Now we prove that  $S^*(M^\perp) = S^*(M)$  implies  $\overline{M} = \overline{M^\perp}$ . Let  $z$  be in  $M^\perp$ . If  $S^*(z) = f$  there exists  $m$  in  $M$  such that  $S^*(m) = f$ . So  $z - m = y$  is in  $\text{Ker } S^*$ , and  $\overline{z} = \overline{m}$  is in  $\overline{M}$ . Finally, since  $\overline{M^\perp}$  is pure in  $\overline{G}$  and  $M^\perp \subset \overline{M}$  we get  $\overline{M^\perp} = \overline{M}$  is pure in  $\overline{G}$ .  $\square$

By definition (1.2)  $\overline{M}$  is pure in  $\overline{G}$ , so lemma (2.0) and proposition (2.0), and, conditions c.1 and c.2 in definition (1.2) imply that  $\text{Coker } S^*$  is isomorphic to

$$\mathbb{Z}^{2r} \oplus \left( \bigoplus_{i=1}^s \mathbb{Z}/a_i \mathbb{Z} \right)^2.$$

(2.1) **Proposition.** *There exists a basis  $\mathcal{B} = \{m_i, m_i^*; i=1, \dots, s+r\}$  of  $G$  such that:*

1.  $\{m_i; i=1, \dots, s+r\}$  is a basis of  $M$ ,
2.  $\{m_i, m_i^*; i=s+1, \dots, s+r\}$  is a basis of  $\text{Ker } S^*$  and  $\{m_i^*; i=s+1, \dots, s+r\}$  is a basis of  $\text{Ker } S_0^*$ ,
3. the submodules  $\langle m_i, m_i^* \rangle, i=1, \dots, s+r$ ; are orthogonal for  $S$ , i.e.:  $G = \bigoplus_{1 \leq i \leq s+r} \langle m_i, m_i^* \rangle^\perp$ ,
3. when  $i=1, \dots, s$ ,  $S(m_i, m_i^*) = a_i$ .

**Definition.** *Such a basis is called a good basis of  $G$  associated to  $M$ .*

The form  $S = A + \varepsilon A^T$  is always an even form. Moreover, when the  $a_i$  are odd we get the following corollary:

**Corollary.** *When the  $a_i$  are odd, the isomorphic class of  $S$  is given by  $m = \text{rk}(G)$  and the isomorphic class of  $\text{Coker } S^*$ .*

*Proof of proposition (2.1).* In (2.0) we have seen that  $S^*(M)^\wedge = \text{Hom}_{\mathbb{Z}}(G|_{M^\perp}; \mathbb{Z})$ . Let  $M_0$  be any direct summand complement of  $(M \cap \text{Ker } S^*)$  in  $M$ . There exists a basis  $\{m_i; i=1, \dots, s\}$  of  $M_0$  and a basis  $\{h_i; i=1, \dots, s\}$  of  $\text{Hom}_{\mathbb{Z}}(G|_{M^\perp}; \mathbb{Z})$  such that  $S^*(m_i) = a_i h_i$  where  $a_i \in \mathbb{N} \setminus \{0\}$  and  $a_i$  divides  $a_{i+1}$ . Let  $m_1^*$  be any element in  $G$  such that  $G = \text{Ker } h_1 \oplus \langle m_1^* \rangle$  and  $h_1(m_1^*) = S(m_1, m_1^*) \cdot a_1^{-1} = 1$ .

**Claim.** For all  $x$  in  $G$ ,  $a_1$  divides  $S(x, m_1^*)$ .

If  $a_1 = 1$  it is obvious. If  $a_1 > 1$ , condition c.2 in (1.2) implies that  $(S^*(G)^\wedge)/S^*(G)$  is isomorphic to  $(S^*(M)^\wedge)/S^*(M)$ .  $(S^*(M)^\wedge)/S^*(M) \cong (\bigoplus_{i=1}^s \mathbb{Z}/a_i \mathbb{Z})^2$  and the rank of  $S^*(G)$  is  $2s$ .

So  $a_1$  divides  $S^*(x)$  for all  $x$  in  $G$ .

Now, we will construct an orthogonal complement  $(M_1 \oplus R_1)$  for  $\langle m_1, m_1^* \rangle$  in  $G$  such that:

- i)  $M = \langle m_1 \rangle \oplus M_1$ ,
- ii)  $\text{Ker } h_1 = M \oplus R_1$ .

Let  $M_1$  be the submodule of  $M$  generated by  $m_i' = m_i - a_1^{-1} S(m_i, m_1^*) \cdot m_1$ ,  $2 \leq i \leq s$ , and  $M \cap \text{Ker } S^*$ . By construction  $M_1$  is orthogonal to  $\langle m_1, m_1^* \rangle$  and  $M = \langle m_1 \rangle \oplus M_1$ .

By construction  $\text{Ker } h_1$  is orthogonal to  $m_1$  and  $M$  is in  $\text{Ker } h_1$ .

If  $\{x_i; i=2, \dots, s+r\}$  is a basis of any direct summand complement of  $M$  in  $\text{Ker } h_1$ , let  $R_1$  be the submodule of  $\text{Ker } h_1$  generated by  $x_i'$  where:  $x_i' = x_i - a_1^{-1} S(x_i, m_1^*) \cdot m_1$ . Then  $\text{Ker } h_1 = \langle m_1 \rangle \oplus M_1 \oplus R_1$  and  $R_1$  is orthogonal to  $m_1^*$ .

Now we have an orthogonal decomposition of  $G$  in  $\langle m_1, m_1^* \rangle \oplus^\perp (M_1 \oplus R_1)$ . By

induction on  $s$  we obtain an orthogonal decomposition:

$$G = (\oplus^{\perp} \langle m_i, m_i^* \rangle) \oplus^{\perp} (M_s \oplus R_s) \text{ where } \text{Ker } S^* = M_s \oplus R_s.$$

Let  $\{m_{s+1}, \dots, m_{s+r}\}$  be any basis of  $\text{Ker } S^* \cap M$ . Thanks to condition c.1,  $\text{Ker } S^* \cap M = \{(x, \varphi(x)); x \in \text{Ker } S_0^*\}$ . So we can choose any basis  $\{m_{s+1}^*, \dots, m_{s+r}^*\}$  of  $\text{Ker } S_0^*$  to build up a basis of  $G$  which fulfills proposition (2.1).  $\square$

(2.2) Now, we use the notations established in §1 and the following convention: if  $f : R \rightarrow S$  is an isomorphism of  $\mathbb{Z}$ -modules,  $\Delta(f)$  is the submodule  $\{(x, f(x)); x \in R\}$  in  $R \oplus S$ . To prove theorem 1, we need the following proposition which gives an equivalent definition of algebraic cobordism.

**Proposition.** *Let  $A_0$  and  $A_1$  be in  $\mathcal{A}$ . Then  $A_0$  is algebraically cobordant to  $A_1$  if and only if there exists a pure submodule  $H$  of  $G = G_0 \oplus G_1$  on which  $A = A_0 \oplus -A_1$  vanishes, an isomorphism  $\varphi$  from  $\text{Ker } S_0^*$  to  $\text{Ker } S_1^*$  and an isomorphism  $\theta$  from  $\text{Tors}(\text{Coker } S_0^*)$  to  $\text{Tors}(\text{Coker } S_1^*)$  such that:*

- c.11:  $\Delta(\varphi) \subset H$ ,
- c.12: the image  $\overline{H}$  of  $H$  in  $\overline{G} = G / \text{Ker } S^*$  is a metabolizer for  $\overline{S} = \overline{S_0} \oplus -\overline{S_1}$ ,
- c.2:  $d(S^*(H)^\wedge) = \Delta(\theta)$ .

*Proof.* Let  $M, \varphi, \theta$  be as in definition (1.2). Then  $M$  satisfies c.1 and c.2. The existence of  $\varphi$  shows that  $\text{Ker } S_0^*$  and  $\text{Ker } S_1^*$  have the same rank,  $r$ . So the rank of  $\overline{G}$  is  $(m_0 + m_1 - 2r)$ . By c.1  $M \cap \text{Ker } S^* = \Delta(\varphi)$  and  $\text{rk}(M) = \frac{m_0 + m_1}{2}$  because  $M$  is a metabolizer for  $A$ . So  $\text{rk}(\overline{M}) = \frac{m_0 + m_1}{2} - r$  and  $\overline{S}$  vanishes on  $\overline{M}$ . It implies that  $\overline{M}$  is a metabolizer for  $\overline{S}$ .

Conversely let  $H, \varphi$  and  $\theta$  be as in the statement of proposition (2.1). As  $\Delta(\varphi)$  is pure in  $H$  and in  $\text{Ker } S^*$ , there exists a direct sum decomposition  $H \cap \text{Ker } S^* = \Delta(\varphi) \oplus M_0$ . As  $\text{Ker } S^*$  is pure in  $G$ , there exists also a direct sum decomposition  $H = M_1 \oplus (H \cap \text{Ker } S^*)$ . Let  $M$  be  $M_1 \oplus \Delta(\varphi)$ . By construction  $A$  vanishes on  $M$ ,  $M \cap \text{Ker } S^* = \Delta(\varphi)$  and  $S^*(M) = S^*(H)$ . So  $M, \varphi$  and  $\theta$  satisfy c.1 and c.2 of definition (1.2). Furthermore,  $\overline{H} = \overline{M_1} = \overline{M}$  and by c.12 the rank of  $\overline{H}$  is  $\frac{m_0 + m_1}{2} - r$ . But  $M_1$  being isomorphic to  $\overline{M_1}$ , the rank of  $M$  is  $\frac{m_0 + m_1}{2}$  and  $M$  is a metabolizer for  $A$ .  $\square$

(2.3) *Proof of theorem 1.* The only non trivial property to check is the transitivity of the relation "algebraic cobordism".

(2.4) **Lemma.** *Let  $B_i : G_i \times G_i \rightarrow \mathbb{Z}$  be in  $\mathcal{A}$ ,  $i = 0, 1, 2$ . Let  $m_i$  be the rank of  $G_i$ . If there exists a metabolizer  $H_{01}$  (resp.  $H_{12}$ ) for  $B_0 \oplus -B_1$  (resp.  $B_1 \oplus -B_2$ ) and if the  $B_i$  are non-degenerate, the form  $B_0 \oplus -B_2$  vanishes on  $H_{02} = \pi(L)$  and  $\text{rk } H_{02} = \frac{1}{2} \text{rk}(G_0 \oplus G_2)$ , where:  $G = G_0 \oplus G_1 \oplus G_1 \oplus G_2$ ,  $H = H_{01} \oplus H_{12}$ ,*

$\Delta = \{(y, y) \in G_1 \oplus G_1 ; y \in G_1\}$ ,  $L = H \cap (G_0 \oplus \Delta \oplus G_2)$  and  $\pi$  is the projection of  $G$  on  $G_0 \oplus G_2$ .

*Proof.* As  $B_0 \oplus -B_2$  vanishes on  $H_{02}$  by construction, it is sufficient to prove that the rank of  $H_{02}$  is  $\frac{m_0+m_1}{2}$ . The definition of  $H_{02}$  gives the following exact sequence:

$$0 \rightarrow L \cap \Delta \xrightarrow{i} L \xrightarrow{\pi} H_{02} \rightarrow 0.$$

So we get:

$$(*) \quad \text{rk}(L) = \text{rk}(L \cap \Delta) + \text{rk}(H_{02}).$$

If  $v$  is in  $H$ , there exists unique  $x$  in  $G_0$ ,  $y_1$  and  $y_2$  in  $G_1$  and  $z$  in  $G_2$  such that  $v = (x, y_1, y_2, z)$ . Let  $\rho : H \rightarrow G_1 \oplus G_1$  be defined by  $\rho(v) = (y_1 - y_2, 0)$ . Let us denote by  $L_1$  the image  $\rho(H)$ . By construction  $L$  is the kernel of  $\rho$  and we get the exact sequence:  $0 \rightarrow L \xrightarrow{i} H \xrightarrow{\rho} L_1 \rightarrow 0$ . Both this sequence and  $(*)$  show:

$$(**) \quad \frac{m_0 + m_2 + 2m_1}{2} - \text{rk}(L_1) = \text{rk}(L \cap \Delta) + \text{rk}(H_{02}).$$

**Claim.** By  $(B_1 \oplus -B_1)$ ,  $\Delta \cap L$  is orthogonal to  $L_1 \oplus \Delta$ .

Indeed,  $\Delta$  is self-orthogonal ; if  $(y, y)$  is in  $\Delta \cap L$ , then  $(0, y)$  is in  $H_{01}$  and  $(y, 0)$  is in  $H_{12}$ . On the other hand, an element of  $L_1$  is of the form  $(y_1, -y_2)$  where there exists  $(x, y_1)$  in  $H_{01}$  and  $(y_2, z)$  in  $H_{12}$ . So  $B_1(y, y_1) = B_1(y_1, y) = 0$  and  $-B_1(y, y_2) = -B_1(y_2, y) = 0$ .

The rank of  $L_1 \oplus \Delta$  is  $m_1 + \text{rk}(L_1)$ . The claim implies that the rank of the restriction of  $B_1 \oplus -B_1$  to  $(\Delta \cap L) \times (G_1 \oplus G_1)$  is smaller or equal to  $m_1 - \text{rk}(L_1)$ . But  $B_1 \oplus -B_1$  is non-degenerate by hypothesis, so:  $\text{rk}(\Delta \cap L) \leq m_1 - \text{rk}(L_1)$ . By  $(**)$  it implies:  $\frac{m_0+m_2}{2} \leq \text{rk}(H_{02})$ .

As  $B_0$  and  $B_2$  are non-degenerate by hypothesis and as  $B_0 \oplus -B_2$  vanishes on  $H_{02}$ ,  $\text{rk}(H_{02}) \leq \frac{m_0+m_2}{2}$ . It ends the proof of the lemma.  $\square$

Let us go back to the proof of theorem 1. Let  $A_i$  be algebraically cobordant to  $A_{i+1}$ ,  $i = 0, 1$ . Let  $M_{i,i+1}$  be a metabolizer for  $A_i \oplus -A_{i+1}$  with the isomorphisms  $\varphi_i$  and  $\theta_i$  fulfilling conditions c.1 and c.2 in definition (1.2).

Let us take the following notations:  $G = G_0 \oplus G_1 \oplus G_1 \oplus G_2$ ,  $S_{02} = S_0 \oplus -S_2$ ,  $G_{02} = G_0 \oplus G_2$ ,  $S = S_0 \oplus -S_1 \oplus S_1 \oplus -S_2$ ,  $\Delta = \{(x, x) ; x \in G_1\} \subset G_1 \oplus G_1$ ,  $d$  be the quotient map from  $G$  to  $\text{Coker } S^*$  and  $d_{02}$  the quotient map from  $G_{02}^*$  to  $\text{Coker } S_{02}^*$ . Let  $\pi$  (resp.  $\tilde{\pi}$ ) be the obvious projection from  $G$  (resp.  $\text{Coker } S^*$ ) to  $G_0 \oplus G_2$  (resp.  $\text{Coker } S_{02}^*$ ). Since  $\overline{M}_{i,i+1}$  is pure in  $\overline{G}_i \oplus \overline{G}_{i+1}$  we have the following decompositions  $M_{i,i+1}^\perp = \Delta(\varphi_i) \oplus \text{Ker } S_i^* \oplus R_{i,i+1}$  with  $M_{i,i+1} = \Delta(\varphi_i) \oplus R_{i,i+1}$ , and  $\overline{R}_{i,i+1}$  is pure in  $\overline{G}_i \oplus \overline{G}_{i+1}$ . Let  $Q_{i,i+1}$  be any direct summand complement of  $M_{i,i+1}^\perp$  in  $G_i \oplus G_{i+1}$ . If  $T_{i,i+1} = R_{i,i+1} \oplus Q_{i,i+1}$ , then we have the following decomposition  $G = \text{Ker } S_{01}^* \oplus \text{Ker } S_{12}^* \oplus T_{01} \oplus T_{12}$ . Let us denote by  $T_0$  (resp.  $T_1$ ,  $T'_1$ ,  $T_2$ ) the projection of  $T_{01}$  (resp.  $T_{01}$ ,  $T_{12}$ ,  $T_{12}$ ) to  $G_0$  (resp.  $G_1$ ,  $G_1$ ,  $G_2$ ). We

modify  $R_{12}$  and  $Q_{12}$  by adding to them some elements of  $\Delta(\varphi_1)$  in order to have  $T_1 = T'_1$ . Moreover, we have the following equalities:  $G_i = \text{Ker } S_i^* \oplus T_i$   $i = 0, 1, 2$ .

Let  $T_{02}$  be  $T_{02} = \pi(T_{01} \oplus T_{12}) = T_0 \oplus T_2$ . Let  $R_{02}$  be the smallest pure submodule of  $T_{02}$  which contains the projection of  $(R_{01} \oplus R_{12}) \cap (G_0 \oplus \Delta \oplus G_2)$  on  $T_{02}$ :  $R_{02} = (\pi((R_{01} \oplus R_{12}) \cap (G_0 \oplus \Delta \oplus G_2)))^\wedge$ ; and let  $A$  be  $A_0 \oplus -A_2$ ,  $\varphi$  be  $\varphi_1 \circ \varphi_0$  and  $\theta$  be  $-(\theta_1 \circ \theta_0)$ .

By proposition (2.2), to prove that  $A_0$  is algebraically cobordant to  $A_2$  it is sufficient to prove that  $H = \Delta(\varphi) \oplus R_{02}$  is a metabolizer for  $A_0 \oplus -A_2$ , and,  $H$  fulfill conditions c.11, c.12 and c.2 of (2.2). First we remark that  $H$  fulfills c.11 by definition.

(2.5) **Lemma.** *We have the equality  $d_{02}(S_{02}^*(H))^\wedge = \Delta(-\theta_1 \circ \theta_0)$ .*

(2.6) **Lemma.** *The submodule  $H$  is a metabolizer for  $A$ , and  $\overline{H}$  is a metabolizer for  $\overline{S}_0 \oplus -\overline{S}_2$ .*

*Proof of lemma (2.5).* By construction:  $d(S^*(G))^\wedge = \text{Tors}(\text{Coker } S^*)$  and  $d_{02}(S_{02}^*(H))^\wedge = \tilde{\pi}(d(S^*(L))^\wedge)$ . But c.2 implies:

$d(S^*(L))^\wedge = (\Delta(\theta_0) \oplus \Delta(\theta_1)) \cap d(S^*(G_0 \oplus \Delta \oplus G_2))^\wedge$ , so:

$$d(S^*(L))^\wedge = \{(x, \theta_0(x), y, \theta_1(y)); x \in \text{Tors}(\text{Coker } S_0^*), y = -\theta_0(x)\}.$$

Finally:  $d_{02}(S_{02}^*(H))^\wedge = \{(x, -\theta_1 \circ \theta_0(x)); x \in \text{Tors}(\text{Coker } S_0^*)\} = \Delta(-\theta_1 \circ \theta_0)$ .  $\square$

*Proof of lemma (2.6).* The restriction  $S_{i,i+1}|_{T_{i,i+1}}$  on  $T_{i,i+1}$ , of the  $\varepsilon$ -symetric bilinear form  $S_{i,i+1}$ , is non-degenerate; and the submodule  $R_{i,i+1}$  is a metabolizer for  $S_{i,i+1}|_{T_{i,i+1}}$ ,  $i = 0, 1$ . By construction  $T_0$  (resp.  $T_1, T_2$ ) is the projection of  $T_{01}$  (resp.  $T_{01}, T_{12}$ ) onto  $G_0$  (resp.  $G_1, G_2$ ). So we have  $S_{i,i+1}|_{T_{i,i+1}} = S_i|_{T_i} \oplus -S_{i+1}|_{T_{i+1}}$ . We use lemma (2.4) replacing  $B_i$  by  $S_i|_{T_i}$ , so  $S_{02}|_{T_{02}}$  vanishes on  $R_{02}$  and  $\text{rk } R_{02} = \frac{1}{2}\text{rk } T_{02}$ . Since the pure submodule  $H$  of  $G_{02} = \text{Ker } S_{02}^* \oplus T_{02}$  is defined by the equality  $H = \Delta(\varphi) \oplus R_{02}$  then  $\text{rk } H = \frac{1}{2}\text{rk } G_{02}$ . Moreover for all  $h_1, h_2$  in  $H$  there exist two integers  $a_1$  and  $a_2$  such that for  $i = 1, 2$  we have:  $a_i h_i = \pi(m_i)$  and  $m_i = (x_i, \varphi_0(x_i), \varphi_0(x_i), \varphi(x_i)) + (m_{0,i}, m_{1,i}, m_{1,i}, m_{2,i})$  is in  $M_{01} \oplus M_{12}$ . So  $A(h_1, h_2) = \frac{1}{a_1 a_2} (A_{01} \oplus -A_{12})(m_1, m_2) = 0$ , so  $A$  vanishes on the pure submodule  $H$  of  $G_{02}$ . Finally  $H$  is a metabolizer for  $A$ . By construction  $S_{02}|_{T_{02}}$  is isomorphic to  $\overline{S}_{02}$ , so as  $R_{02}$  is pure in  $T_{02}$  then  $\overline{R}_{02}$  is a metabolizer for  $\overline{S}_{02}$ .  $\square$

The above properties of  $H$ , and, lemmas (2.5) and (2.6) imply conditions c.12 and c.2 of proposition (2.2), and  $A_0$  is algebraically cobordant to  $A_2$ . This ends the proof of theorem 1.  $\square$

### 3. The necessary condition to have a cobordism

Let  $K_0$  and  $K_1$  be two cobordant links. Let us denote by  $\mathcal{S}$  the product  $S^{2n+1} \times [0, 1]$  and by  $\Sigma$  its oriented boundary. The definition of cobordism gives a submanifold  $C = \Phi(\mathcal{K} \times [0, 1])$  of  $\mathcal{S}$  such that  $\Sigma \cap C = K_0 \coprod (-K_1)$ . Let  $N$  be  $F_0 \cup C \cup (-F_1)$  where  $F_i$  is a Seifert surface for  $K_i$ . By construction  $N$  is a closed, compact, oriented,  $2n$ -submanifold of  $\mathcal{S}$ .

(3.1) **Lemma.** *There exists a smooth oriented, compact, submanifold  $W$  of  $\mathcal{S}$  such that  $N$  is the boundary of  $W$ .*

*Proof.* This lemma is a consequence of classical obstruction theory. If  $n \geq 3$  a proof is written in [L2, 70], p. 183. As the existence of  $W$  is fundamental to obtain theorem 2, we write a proof which works in any dimension.


Let  $C_j$  for  $j = 1, \dots, k$  be the  $k$  connected components of  $C$ . As  $C$  has a trivial normal bundle in  $\mathcal{S}$ , it is possible to choose disjoint, closed, tubular neighbourhoods  $U_j$  of  $C_j$  and a diffeomorphism  $\Psi : C \times D^2 \rightarrow U = \coprod_{1 \leq j \leq k} U_j$ .

Now we have meridians  $m_j$  on  $\partial U_j$  defined by:  $m_j = \Psi(P_j \times S^1)$  where  $P_j$  is some point of  $C_j$  and  $m_j$  is oriented such that the linking number of  $m_j$  and  $C_j$  (in  $\mathcal{S}$ ) is  $+1$ . Let  $X$  be  $\mathcal{S} \setminus \overset{\circ}{U}$ ,  $v$  be the diffeomorphism induced by the inclusion of  $\partial X$  in  $U$ ,  $e$  be the excision isomorphism and  $\partial_X^i$  (resp.  $\partial_X^i$ ) be the connectant homomorphism for the pair  $(\mathcal{S}, U)$  (resp.  $(X, \partial X)$ ). Then we have the following commutative diagram:

$$\begin{array}{ccccccc}
 \xrightarrow{\partial_X^0} & H^1(X, \partial X) & \xrightarrow{\rho} & H^1(X) & \xrightarrow{\sigma} & H^1(\partial X) & \xrightarrow{\partial_X^1} & H^2(X, \partial X) & \rightarrow \\
 \cong \uparrow e & & & \uparrow & & v \uparrow & & \cong \uparrow e & \\
 \xrightarrow{\partial} & H^1(\mathcal{S}, U) & \rightarrow & 0 = H^1(\mathcal{S}) & \rightarrow & H^1(U) & \xrightarrow{\cong \partial^1} & H^2(\mathcal{S}, U) & \rightarrow 0
 \end{array}$$

The commutativity of all the squares of the above diagram implies that the homomorphism  $\rho$  is zero so  $\sigma$  is injective and  $\partial_X^i$  is surjective for  $0 \leq i \leq 2n-1$ . We have the following direct sum decomposition:  $H^1(\partial X) = \sigma(H^1(X)) \oplus v(H^1(U))$ . Any element of  $\sigma(H^1(X))$  is represented by a differentiable map from  $\partial X$  to  $S^1$ , which is, up to homotopy, characterized by its degree on each meridian  $m_j$ , and which has a unique extension to  $X$ . Let  $g : X \rightarrow S^1$  be the unique, up to homotopy, differentiable map which has degree  $+1$  on each meridian. Thanks to the Thom-Pontriagin construction there exists a differentiable map  $f : \Sigma \setminus (K_0 \coprod -K_1) \rightarrow S^1$  which has  $\overset{\circ}{F_0} \coprod (-\overset{\circ}{F_1})$  as regular fiber and  $f$  has degree  $+1$  on the meridians of the connected components of  $K_0 \coprod (-K_1)$ . So  $f$  and  $g$  have homotopic restrictions on  $X \cap \Sigma$  and we can choose  $g$  such that its restriction on  $X \cap \Sigma$  coincides with  $f$ .

Then  $g$  has a regular fiber  $\overline{W}$  such that  $\overline{W} \cap \Sigma = (F_0 \coprod -F_1) \cap X$ . The union of  $\overline{W}$  with a small collar in  $U$  is the manifold  $W$  such that  $N = \partial W$ .  $\square$



(3.2) Let us take  $A_0$  (resp.  $A_1$ ) the Seifert form associated to a  $(n-1)$ -connected Seifert surface  $F_0$  (resp.  $F_1$ ) for  $K_0$  (resp.  $K_1$ ). Let  $\tau : K_0 \rightarrow K_1$  be the diffeomorphism defined by:  $\tau(P) = \Phi(\Phi^{-1}(P) \times \{1\})$  where  $P$  is any point of  $K_0$ . The diffeomorphism  $\tau$  induces isomorphisms  $\theta_j : H_j(K_0) \rightarrow H_j(K_1)$  such that for any  $j$ -cycle  $x$  of  $K_0$ ,  $(x, \theta_j(x))$  is a boundary in  $C = \Phi(\mathcal{K} \times [0, 1])$ . Let  $\chi_i : H_n(K_i) \rightarrow H_n(F_i)$  and  $\lambda_i : H_n(F_i) \rightarrow H_n(N)$ ,  $i = 0, 1$ , be the homomorphisms induced by the inclusions  $K_i \subset F_i \subset N$ . The Mayer-Vietoris exact sequence associated to the decomposition of  $N$  in the union of  $F_0 \cup C$  and  $C \cup (-F_1)$  gives:

$$\rightarrow H_n(K_0) \xrightarrow{\chi} H_n(F_0) \oplus H_n(F_1) \xrightarrow{\lambda} H_n(N) \xrightarrow{\delta} H_{n-1}(K_0) \rightarrow$$

where  $\chi = (\chi_0, \chi_1 \circ \theta_n)$  and  $\lambda = (\lambda_0, \lambda_1)$

(3.3) **Remark.** Let  $m_i$  be  $\text{rk}(H_n(F_i))$ ,  $m$  be  $\text{rk}(H_n(N))$  and  $r$  be  $\text{rk}(\chi(H_n(K_0)))$ . By Poincaré duality  $m = m_0 + m_1$ ,  $r = \text{rk}(\delta(H_n(N)))$  and  $r = \text{rk}(\text{Ker } S_i^*)$  where  $S_i^*$  is the adjoint of the intersection form  $S_i$  on  $H_n(F_i)$ .

(3.4) Construction of the isomorphisms  $\varphi : \text{Ker } S_0^* \rightarrow \text{Ker } S_1^*$  and  $\theta : \text{Tors}(\text{Coker } S_0^*) \rightarrow \text{Tors}(\text{Coker } S_1^*)$ .

Let  $S_{i*} : H_n(F_i) \rightarrow H_n(F_i, K_i)$  and  $\partial : H_n(F_i, K_i) \rightarrow H_{n-1}(K_i)$  be the homomorphisms given by the long exact sequence for the pair  $(F_i, K_i)$ . Let  $U : H^n(F_i) \rightarrow \text{Hom}_{\mathbb{Z}}(H_n(F_i); \mathbb{Z})$  be the universal coefficient isomorphism ( $F_i$  is  $(n-1)$ -connected) and let  $P : H_n(F_i, K_i) \rightarrow H^n(F_i)$  be the Poincaré duality isomorphism. We have the following commutative diagram:

$$\begin{array}{ccccccc}
 0 & \rightarrow & \chi_i(H_n(K_i)) & \rightarrow & H_n(F_i) & \xrightarrow{S_{i*}} & H_n(F_i, K_i) & \xrightarrow{\partial} & \partial(H_n(F_i, K_i)) & \rightarrow & 0 \\
 & & \parallel & & \parallel & & \cong \downarrow U \circ P & & \downarrow \Delta_i & & \\
 0 & \rightarrow & \text{Ker } S_i^* & \rightarrow & H_n(F_i) & \xrightarrow{S_i^*} & \text{Hom}_{\mathbb{Z}}(H_n(F_i); \mathbb{Z}) & \xrightarrow{d} & \text{Coker } S_i^* & \rightarrow & 0
 \end{array}$$

By definition  $\Delta_i : \partial(H_n(F_i, K_i)) \rightarrow \text{Coker } S_i^*$  is the quotient of the isomorphism  $U \circ P$ , so  $\Delta_i$  is an isomorphism.

Let us consider again the isomorphism  $\theta_j : H_j(K_0) \rightarrow H_j(K_1)$ , which is defined in (3.2) thanks to the existence of the cobordism. Since  $F_i$  is  $(n-1)$ -connected then  $\partial(H_n(F_i, K_i)) = \tilde{H}_{n-1}(K_i)$  and  $\theta_n(\text{Ker } \chi_0) = \text{Ker } \chi_1$ , so  $\theta_{n-1} \circ \partial(H_n(F_0, K_0)) = \partial(H_n(F_1, K_1))$ .

Let  $\theta$  be the restriction of the isomorphism  $\Delta_1 \circ \theta_{n-1} \circ \Delta_0^{-1}$  on the  $\mathbb{Z}$ -torsion of  $\text{Coker } S_0^*$ .

Let  $\varphi$  be the restriction of  $\theta_n$  on  $\chi_0(H_n(K_0))$ . As  $\chi_i(H_n(K_i)) = \text{Ker } S_i^*$ , so  $\varphi$  is defined on  $\text{Ker } S_0^*$ .

We denote by  $\Delta(\varphi)$  the submodule  $\{(x, \varphi(x)); x \in \text{Ker } S_0^*\}$  of  $H_n(F_0) \oplus H_n(F_1)$ .

(3.5) **Remark.** By construction  $\varphi$  fulfills:  $\varphi \circ \chi_0 = \chi_1 \circ \theta_n$  and  $\Delta(\varphi) = \chi(H_n(K_0))$  where  $\chi = (\chi_0, \chi_1 \circ \theta_n)$  as in (3.2).

(3.6) To prove theorem 2, we will construct a metabolizer  $M$  (in  $H_n(F_0 \coprod -F_1)$ ) for  $A = A_0 \oplus -A_1$ . This metabolizer  $M$  will fulfill conditions c.1 and c.2 in definition (1.2) of the algebraic cobordism, for the isomorphisms  $\varphi$  and  $\theta$  defined in (3.4). To do that, we have to choose an oriented submanifold  $W$  of  $\mathcal{S}$  with  $\partial(W) = N$  (thanks to (3.1) such a  $W$  exists). Let  $j : H_n(N) \rightarrow H_n(W)$  be the homomorphism induced by the inclusion of  $N$  in  $W$ .

(3.7) **Lemma.** *The form  $A = A_0 \oplus -A_1$  vanishes on  $\lambda^{-1}(\text{Ker } j^\wedge)$ .*

*Proof.* It is sufficient to prove that  $A$  vanishes on  $\lambda^{-1}(\text{Ker } j)$ . Let  $a = [x]$  and  $b = [y]$  be two homology classes in  $\lambda^{-1}(\text{Ker } j)$ . As  $\lambda$  is induced by the inclusion of  $F_0 \coprod -F_1$  in  $N$  (see (3.2)), there exists two  $(n+1)$ -chains  $\alpha$  and  $\beta$  in  $W$  such that  $\partial\alpha = x$  and  $\partial\beta = y$ . Let  $i_+$  be the positively oriented normal vector field to  $W$  in  $\mathcal{S}$ . The intersection of  $\alpha$  and  $i_+(\beta)$  is zero. Hence the linking number in  $\Sigma$  of  $x$  and  $i_+(y)$  is zero. But this linking number is, by definition, equal to  $A(a, b)$ , so  $A(a, b) = 0$  and the lemma is proved.  $\square$

(3.8) **Lemma.** *Let  $m$  be the rank of  $H_n(N)$ . The rank of  $\text{Ker } j$  is  $\frac{m}{2}$ .*

*Proof.* The long exact sequence for the pair  $(W, N)$  gives the exactness of:

$$0 \rightarrow H_{2n+1}(W) \rightarrow H_{2n+1}(W, N) \rightarrow H_{2n}(N) \rightarrow \dots \rightarrow H_{n+1}(W, N) \rightarrow \text{Ker } j \rightarrow 0$$

The alternating sum of the ranks in this exact sequence together with the Poincaré duality give:

$$\text{rk}(\text{Ker } j) = \frac{\text{rk}(H_n(N))}{2} = \frac{m}{2}.$$

$\square$

(3.9) **Lemma.** *There exists a direct summand decomposition of  $\lambda^{-1}(\text{Ker } j^\wedge)$  in*

$\Delta(\varphi) \oplus R_0 \oplus R$  where  $\Delta(\varphi) = \{(x, \varphi(x)); x \in \text{Ker } S_0^*\}$ ,  $R_0 = \lambda^{-1}(\text{Ker } j^\wedge) \cap \text{Ker } S_0^*$ , and  $R$  is any direct summand complement of  $\lambda^{-1}(\text{Ker } j^\wedge) \cap \text{Ker } S^*$  in  $\lambda^{-1}(\text{Ker } j^\wedge)$ .

*Proof.* As the considered submodules of  $\lambda^{-1}(\text{Ker } j^\wedge)$  are pure, the lemma comes from the following equalities:

$$\chi(H_n(K_0)) = \text{Ker } \lambda \subset \lambda^{-1}(\text{Ker } j^\wedge) \text{ (see (3.2))},$$

$$\Delta(\varphi) = \chi(H_n(K_0)) \text{ (see (3.5))},$$

$$\text{Ker } S^* = \chi(H_n(K_0)) \oplus \text{Ker } S_0^*. \quad \square$$

(3.10) **Proposition.** *The submodule  $M = \Delta(\varphi) \oplus R$  of  $\lambda^{-1}(\text{Ker } j^\wedge)$  is a metabolizer for  $A = A_0 \oplus -A_1$ , which fulfills:  $M \cap \text{Ker } S^* = \Delta(\varphi)$ .*

*Proof.* By lemma (3.9),  $M \cap \text{Ker } S^* = \Delta(\varphi)$ . By (3.6),  $A$  vanishes on  $M$ . So we only have to show that  $M$  is of rank  $\frac{m}{2}$ . As remarked in (3.3),  $r = \text{rk}(\delta(H_n(N)))$ , so  $\text{rk}(\delta(\text{Ker } j^\wedge)) \leq r$ . Let us consider the following exact sequence induced by (3.2):  $0 \rightarrow \Delta(\varphi) \xrightarrow{\lambda} \lambda^{-1}(\text{Ker } j^\wedge) \xrightarrow{\lambda} \text{Ker } j^\wedge \xrightarrow{\delta} \delta(\text{Ker } j^\wedge) \rightarrow 0$ . This exact sequence together with the equalities:  $\text{rk}(\text{Ker } j^\wedge) = \frac{m}{2}$  (see (3.8)),  $\text{rk}(\Delta(\varphi)) = r$ ; give  $\text{rk}(\lambda^{-1}(\text{Ker } j^\wedge)) = r + \frac{m}{2} - \text{rk}(\delta(\text{Ker } j^\wedge))$ . So  $\text{rk}(\lambda^{-1}(\text{Ker } j^\wedge)) \geq \frac{m}{2}$ .

We can remark that if  $A$  is non degenerated (as supposed in theorem 2) then we have  $\text{rk}(\lambda^{-1}(\text{Ker } j^\wedge)) \leq \frac{1}{2}\text{rk}(H_n(F_0) \oplus H_n(F_1)) = \frac{m}{2}$ , because  $A$  vanishes on  $\lambda^{-1}(\text{Ker } j^\wedge)$  (see (3.6)). So, if  $A$  is non degenerated,  $\text{rk}(\lambda^{-1}(\text{Ker } j^\wedge)) = \frac{m}{2}$ ,  $\text{rk}(\delta(\text{Ker } j^\wedge)) = r$ ,  $\text{rk}(R_0) = 0$  and  $M = \lambda^{-1}(\text{Ker } j^\wedge)$  is a metabolizer for  $A$ .

Come back to the general case. Let  $r_0$  be the rank of  $R_0$ . By construction:  $\text{rk}(M) = \text{rk}(\lambda^{-1}(\text{Ker } j^\wedge)) - r_0 = r + \frac{m}{2} - \text{rk}(\delta(\text{Ker } j^\wedge)) - r_0$ .

(3.11) **Lemma.** *The rank  $l$  of  $\delta(H_n(N)) / \delta(\text{Ker } j^\wedge)$  is greater or equal to  $r_0$ .*

*Proof.* Let  $\{e_j\}$ ,  $j = 1, \dots, r_0$  be a basis of  $R_0$ . Let  $\{e_j^*\}$  be in  $H_n(N) \otimes_{\mathbb{Z}} \mathbb{Q}$  such that  $S_N(\lambda(e_j), e_j^*) = \delta_{ij}$  where  $S_N$  is the intersection form on  $H_n(N) \otimes_{\mathbb{Z}} \mathbb{Q}$ . The  $e_j^*$  exists because  $S_N$  is unimodular. Let  $R^*$  be the submodule of  $H_n(N) \otimes_{\mathbb{Z}} \mathbb{Q}$  generated by  $\{e_j^*\}$ . Since  $R_0 \cap \text{Ker } \lambda = \{0\}$ , then  $\text{rk}(\lambda(R_0)) = r_0$ . As  $S$  vanishes on  $R_0$ , then  $S_N$  vanishes on  $\lambda(R_0)$ . It implies that  $\text{rk}(R^*) = \text{rk}(R_0) = r_0$ , and  $\text{Ker } j \cap R^* = \{0\}$ . Since  $R_0 \subset \text{Ker } S_0^*$ , we have  $S(x, y) = 0$  for all  $x$  in  $R_0$  and all  $y$  in  $H_n(F_0 \coprod -F_1)$ . So  $R^* \cap \lambda(H_n(F_0 \coprod -F_1)) = \{0\}$  and  $\text{rk}(\delta(H_n(N)) / \delta(\text{Ker } j^\wedge)) = l \geq \text{rk}(\delta(R^*)) = \text{rk}(R^*) = r_0$ .  $\square$

In order to end the proof of (3.10), we only have to show that  $\text{rk}(R) = \frac{m}{2} - r$ . But  $\text{rk}(\delta(\text{Ker } j^\wedge)) = r - l$ ; so we already have shown that  $\text{rk}(R) = \text{rk}(M) - r = \frac{m}{2} - (r - l) - r_0$ .

By lemma (3.11), we have  $l - r_0 \geq 0$ , so  $\text{rk}(R) \geq \frac{m}{2} - r$ . But  $R \cap \text{Ker } S^* = \{0\}$  by construction, and the form  $\overline{S}$  induced by  $S$  on  $H_n(F_0 \coprod -F_1) / \text{Ker } S^*$  is non-

degenerate of rank  $m - 2r$ . So  $\text{rk}(R) \leq \frac{m}{2} - r$  because  $\overline{S}$  vanishes on  $\overline{R} = R/(R \cap \text{Ker } S^*)$ .  $\square$

(3.12) **Remark.** We have found a metabolizer  $M = \Delta(\varphi) \oplus R$  for  $A$  which fulfills condition c.1 of the algebraic cobordism without any condition on  $A$ . We already have got theorem 4 (see (1.6)). To prove condition c.2 and  $\overline{M}$  is pure in  $\overline{G}$ , we will have to choose  $(n-1)$ -connected Seifert surfaces  $F_i$  for  $K_i$  on which the Seifert forms  $A_i$  are unimodular. So the following proposition (3.13) together with proposition (3.10) imply theorem 2 stated in (1.4).

Let  $\theta_{n-1}$  be the isomorphism between  $H_{n-1}(K_0)$  and  $H_{n-1}(K_1)$  defined in (3.2), and let  $\theta$  the isomorphism between  $\text{Tors}(\text{Coker } S_0^*)$  and  $\text{Tors}(\text{Coker } S_1^*)$  defined in (3.4). Using the notation of (2.2), let  $\Delta(\theta_{n-1})$  (resp.  $\Delta(\theta)$ ) be the group  $\{(x, \theta_{n-1}(x)) ; x \in \text{Tors}(H_{n-1}(K_0))\}$  (resp.  $\{(x, \theta(x)) ; x \in \text{Tors}(\text{Coker } S_0^*)\}$ ).

(3.13) **Proposition.** *If  $A_0$  and  $A_1$  are unimodular the metabolizer  $M = \Delta(\varphi) \oplus R$  of  $A = A_0 \oplus -A_1$ , fulfills  $d(S^*(M)^\wedge) = \Delta(\theta)$  ; and  $\overline{M}$  is pure in  $H_n(F)/\text{Ker } S^*$ .*

*Proof.* Let us denote  $F_0 \coprod -F_1$  by  $F$ ,  $K_0 \coprod -K_1$  by  $K$ , and  $S_0^* \oplus -S_1^*$  by  $S^*$ . We consider for  $F$  the following commutative diagram already constructed for  $F_i$  in (3.4):

$$\begin{array}{ccccccc} 0 & \rightarrow & \text{Ker } S_* & \hookrightarrow & H_n(F) & \xrightarrow{S_*} & H_n(F, K) & \xrightarrow{\partial} & \partial(H_n(F, K)) & \rightarrow & 0 \\ & & \parallel & & \parallel & & \cong \downarrow U \circ P & & \cong \downarrow \Delta_0 \oplus \Delta_1 & & \\ 0 & \rightarrow & \text{Ker } S^* & \hookrightarrow & H_n(F) & \xrightarrow{S^*} & \text{Hom}_{\mathbb{Z}}(H_n(F); \mathbb{Z}) & \xrightarrow{d} & \text{Coker } S^* & \rightarrow & 0 \end{array}$$

(3.14) **Lemma.** *The equality  $d(S^*(M)^\wedge) = \Delta(\theta)$  is equivalent to the equality  $\partial(S_*(M)^\wedge) = \Delta(\theta_{n-1})$ .*

*Proof.* The lemma is a consequence of the two following statements:

The restriction of  $\Delta_0 \oplus \Delta_1$  on  $\Delta(\theta_{n-1})$  is an isomorphism to  $\Delta(\theta)$  because  $\theta \circ \Delta_0 = \Delta_1 \circ \theta_{n-1}$  by construction (see (3.4)).

The restriction of  $\Delta_0 \oplus \Delta_1$  on  $\partial(S_*(M)^\wedge)$  is an isomorphism to  $d(S^*(M)^\wedge)$  because the commutativity of the above diagram gives  $U \circ P(S_*(M)^\wedge) = S^*(M)^\wedge$ .  $\square$

Let  $\kappa : H_n(N) \rightarrow H_n(N, C)$  be the homomorphism which is defined in the long exact sequence for the pair  $(N, C)$  and  $\rho : H_n(N, C) \rightarrow N_n(F, K)$  be the inverse of the excision isomorphism induced by the inclusion of the pair  $(F, K) \subset (N, C)$ . Let  $\xi = \rho \circ \kappa : H_n(N) \rightarrow H_n(F, K)$  and  $\overline{\theta} = (\text{Id}, \theta_{n-1}) : H_{n-1}(K_0) \rightarrow H_{n-1}(K)$ .

With the notations used in (3.2) we have the following commutative diagram:

$$\begin{array}{ccccccc}
 \rightarrow & H_n(K_0) & \xrightarrow{\chi} & H_n(F) & \xrightarrow{\lambda} & H_n(N) & \xrightarrow{\delta} H_{n-1}(K_0) \rightarrow \\
 (\star) & & & \parallel & (I) & \downarrow \xi & (II) & \downarrow \bar{\theta} \\
 \rightarrow & H_n(K) & \xrightarrow{\chi_0 \oplus \chi_1} & H_n(F) & \xrightarrow{S_*} & H_n(F, K) & \xrightarrow{\partial} H_{n-1}(K) \rightarrow
 \end{array}$$

The square (I) is commutative by functoriality, and (II) is commutative by definition of  $\xi$  and  $\bar{\theta}$ .

(3.15) **Lemma.** *If  $A_0$  and  $A_1$  are unimodular, then we have  $\delta(\text{Ker } j^\wedge) = \tilde{H}_{n-1}(K_0)$ .*

We first show that lemma (3.15) implies proposition (3.13).

We show that  $\bar{M}$  is pure in  $H_n(F)/\text{Ker } S^*$ , which is equivalent to prove that the quotient  $H_n(F)/(\text{Ker } S^* + M)$  is torsion free. Since  $A = A_0 \oplus -A_1$  is non-degenerate  $M = \lambda^{-1}(\text{Ker } j^\wedge)$ . Furthermore by diagram  $(\star)$  we get  $\lambda(\text{Ker } S^*) = \text{Ker } \xi$ . Let  $\text{pr}$  be the projection of  $H_n(N)$  on  $H_n(N)/(\text{Ker } j^\wedge + \text{Ker } \xi)$ , so  $\text{Ker}(\text{pr} \circ \lambda) = M + \text{Ker } S^*$ . The quotient of  $\text{pr} \circ \lambda$  induces an injective map from  $H_n(F)/(\text{Ker } S^* + M)$  into  $H_n(N)/(\text{Ker } j^\wedge + \text{Ker } \xi)$ .

**Claim.** The module  $H_n(N)/(\text{Ker } j^\wedge + \text{Ker } \xi)$  is torsion free.

*Proof of the claim.* There exists  $x_i, i = 1, \dots, r$ , in  $\text{Ker } j^\wedge$  such that  $\tilde{H}_{n-1}(K_0) = \bigoplus_{i=1}^r \langle \delta(x_i) \rangle \oplus \text{Tors}(\tilde{H}_{n-1}(K_0))$ . Let  $(y_i)_{i=1, \dots, r}$  a basis of  $\text{Ker } \xi$  such that  $S_N(x_i, y_j) = \delta_{ij}$ . By induction on  $r$ , we can construct these bases such that  $H_n(N) = T \oplus^\perp T^\perp$  where  $T = \bigoplus_{i=1}^r \langle x_i, y_i \rangle$ . If we denote by  $D$  the module  $D = T^\perp \cap \text{Ker } j^\wedge$  and by  $D^*$  any direct summand complement of  $D$  in  $T^\perp$ , then we get:  $H_n(N)/(\text{Ker } \xi + \text{Ker } j^\wedge) \cong D^*$  which is torsion free.  $\square$

Finally  $H_n(F)/(\text{Ker } S^* + M)$  is torsion free and  $\bar{M}$  is pure in  $H_n(F)/(\text{Ker } S^*)$ .

So if  $n = 1$ , the links  $K_0$  and  $K_1$  have torsion free homology groups ( $\mathcal{K}$  is a one dimensional compact manifold), so  $\text{Tors}(\text{Coker } S^*) = \{0\}$  and we have already proved proposition (3.13).

Now let us take  $n \geq 2$ .

Thanks to lemma (3.14), the equality:  $\Delta(\theta_{n-1}) = \partial(S_*(M)^\wedge)$  gives proposition (3.13). The above diagram  $(\star)$  and lemma (3.15) imply:  $\bar{\theta}(H_{n-1}(K_0)) =$

$\Delta(\theta_{n-1}) \subset \partial(S_*(M)^\wedge)$ . To show that the inclusion:  $\Delta(\theta_{n-1}) \subset \partial(S_*(M)^\wedge)$  is an equality, it is sufficient to take any  $x$  in  $(\partial(S_*(M)^\wedge)) \cap \partial(H_n(F_0, K_0))$ , and to show that such a  $x$  is zero.

Let us denote by  $L$  (resp.  $L_i$ ) the linking form on  $\text{Tors}(H_{n-1}(K))$  (resp.  $\text{Tors}(H_{n-1}(K_i))$ ). By definition (see remark (3.16)) such a form  $L = L_0 \oplus -L_1$  is non degenerated and vanishes on  $\partial(S_*(M)^\wedge)$  because  $S_0 \oplus -S_1$  vanishes on  $M$ . Let  $(y, \theta_{n-1}(y))$  be in  $\Delta(\theta_{n-1})$ . Then  $L(x, (y, \theta_{n-1}(y))) = L_0(x, y) = 0$  for all  $y \in \text{Tors}(H_{n-1}(K_0))$ . The non degeneracy of  $L_0$  implies  $x = 0$ . This ends the proof of proposition (3.13).  $\square$

(3.16) **Remark.** The linking form  $L$  is defined as follows (see [L-L, 75] prop. 2.1): Let  $x, y$  be in  $\text{Tors}(H_{n-1}(K))$  such that  $p$  and  $q$  are the smallest positive integers with  $p.x = q.y = 0$ . Let  $\bar{x}$  and  $\bar{y}$  be in  $H_n(F)$  such that  $\partial(S_*(\bar{x}) \otimes \frac{1}{p}) = x$  and  $\partial(S_*(\bar{y}) \otimes \frac{1}{q}) = y$ . Then:  $L(x, y) \equiv \frac{1}{p, q} S(\bar{x}, \bar{y}) \pmod{\mathbb{Z}}$ .

*Proof of lemma (3.15).* As shown in (3.10), if  $A_0 \oplus -A_1$  is non degenerated,  $M = \lambda^{-1}(\text{Ker } j^\wedge)$  has rank  $\frac{m}{2}$  and is the chosen metabolizer. So  $\lambda$  induces a monomorphism  $\bar{\lambda}$  on  $H_n(F)/M$  to  $H_n(N)/\text{Ker } j^\wedge$  and we get the following exact sequence:

$$0 \rightarrow H_n(F)/M \xrightarrow{\bar{\lambda}} H_n(N)/\text{Ker } j^\wedge \xrightarrow{\bar{\delta}} \tilde{H}_{n-1}(K_0)/\delta(\text{Ker } j^\wedge) \rightarrow 0.$$

As  $\bar{\lambda}$  is injective and  $M$  is pure in  $H_n(F)$  there exists two  $\mathbb{Z}$ -bases  $\{\bar{e}_j; j=1, \dots, \frac{m}{2}\}$  of  $H_n(F)/M$  and  $\{\bar{k}_j; j=1, \dots, \frac{m}{2}\}$  of  $H_n(N)/\text{Ker } j^\wedge$  such that  $\bar{\lambda}(\bar{e}_j) = p_j \cdot \bar{k}_j$  with  $p_j \in \mathbb{Z} \setminus \{0\}$ . Let  $E$  (resp.  $H$ ) be a direct summand complement of  $M$  (resp.  $\text{Ker } j^\wedge$ ) in  $H_n(F)$  (resp.  $H_n(N)$ ). Let also  $\{e_j; j=1, \dots, \frac{m}{2}\}$  (resp.  $\{k_j; j=1, \dots, \frac{m}{2}\}$ ) be a  $\mathbb{Z}$ -basis of  $E$  (resp.  $H$ ) such that  $e_j \equiv \bar{e}_j \pmod{M}$  (resp.  $k_j \equiv \bar{k}_j \pmod{\text{Ker } j^\wedge}$ ). By construction  $\lambda(e_j) - p_j \cdot k_j = x \in \text{Ker } j^\wedge$ . So there exists a  $(n+1)$ -chain  $\gamma$  in  $W$  and a positive integer  $a$  such that:  $\partial\gamma = a\lambda(e_j) - a p_j \cdot k_j$ . Let  $\rho$  be a  $(n+1)$ -chain of  $S^{2n+1} \times [0, 1]$  with  $\partial\rho = k_j$ . So  $a e_j$  is the boundary of  $\gamma + a p_j \cdot \rho$  in  $S^{2n+1} \times [0, 1]$ .

Statement: for all  $m$  in  $M$ ,  $p_j$  divides  $A(e_j, m)$ .

Let  $m$  be in  $M = \lambda^{-1}(\text{Ker } j^\wedge)$  and  $\Delta$  be a  $(n+1)$ -chain in  $S^{2n+1} \times [0, 1]$  such that  $\partial\Delta = i_+(m)$ . By definition  $A(a e_j, m)$  is the intersection in  $S^{2n+1} \times [0, 1]$  of  $\gamma + a p_j \cdot \rho$  and  $\Delta$ . But  $\lambda(a m) \in \text{Ker } j$  so there exists a  $(n+1)$ -chain  $\mu$  in  $W$  such that  $\partial\mu = a m$ . We have  $\partial(i_+(\mu)) = a i_+(m)$ . Since  $\partial(a \Delta) = a i_+(m)$ , we get  $\gamma \cap (a \Delta) = \gamma \cap (i_+(\mu)) = 0$ . But  $a > 0$ , so  $a(\gamma \cap \Delta) = 0$  implies  $\gamma \cap \Delta = 0$ . Finally  $A(a e_j, m) = a p_j \cdot (\rho \cap \Delta)$  and  $p_j$  divides  $A(e_j, m)$ .

If  $A$  is unimodular the statement implies that  $p_j = \pm 1$  for all  $j = 1, \dots, \frac{m}{2}$ . So  $\bar{\lambda}$  is an isomorphism and his cokernel is zero. As asked we have got:  $\delta(\text{Ker } j^\wedge) = \tilde{H}_{n-1}(K_0)$ . This ends the proof of lemma (3.15).  $\square$

(3.17) **Remark.** As above we can also prove that: for all  $m$  in  $M$   $p_j$  divides  $A(m, e_j)$ .

#### 4. The sufficient condition to have a cobordism

(4.1) Let  $K_0$  and  $K_1$  be two  $2n - 1$  dimensional simple links, with  $n \geq 3$ . We suppose that there exists  $(n-1)$ -connected Seifert surfaces  $F_0$  and  $F_1$ , for  $K_0$  and  $K_1$ , such that the associated Seifert forms  $A_0$  and  $A_1$  are algebraically cobordant. We consider  $K_0$  (resp.  $-K_1$ ) as embedded in the sphere  $S^{2n+1} \times \{0\}$  (resp.  $S^{2n+1} \times \{1\}$ ) which are oriented as the boundary of  $S^{2n+1} \times [0, 1]$ .

Let  $x$  be in  $S^{2n+1} \times \{0\}$  such that  $(x \times [0, 1]) \cap (F_0 \coprod -F_1)$  is empty, and let  $U$  be a "small" open ball around  $x$  in  $S^{2n+1} \times \{0\}$ . The boundary  $S$  of the disk  $D = (S^{2n+1} \times [0, 1]) \setminus (U \times [0, 1])$  contains  $F_0 \coprod \overset{\circ}{F_1}$ . Let  $G$  be the closure of the connected sum, in  $S$ , of the interiors  $\overset{\circ}{F_0}$  and  $-\overset{\circ}{F_1}$ . By construction  $A = A_0 \oplus -A_1$  is the Seifert form of  $K_0 \coprod -K_1$ , associated to  $G$ .

(4.2) *Proof of theorem 3.* In order to prove theorem 3 we will do in  $D$ , an embedded surgery on  $G$ , the result of which being a manifold  $\tilde{G}$  diffeomorphic to  $\mathcal{K} \times [0, 1]$ .

By proposition (2.1) we can choose a good basis  $\mathcal{B} = \{(m_i, m_i^*); i=1, \dots, s+r\}$  of  $H_n(G)$ . Thanks to J. Milnor ([M1, 61] lemma 6 p. 50), any cycle of  $G$  can be represented by the image of an embedding of  $S^n$ . Furthermore:

(4.3) **Proposition.** *There exists  $s+r$  disjointed embeddings  $\psi_i : D^{n+1} \times D^n \rightarrow D$  such that for any  $i \in \{1, \dots, s+r\}$  we have*

- 1-  $[\psi_i(S^n \times \{0\})] = m_i$ ,
- 2-  $(\psi_i(D^{n+1} \times D^n)) \cap G = \psi_i(D^{n+1} \times D^n) \cap S = \psi_i(S^n \times D^n)$ .

*Proof.* Let  $\overline{\psi_i} : S^n \rightarrow G$  be an embedding of  $S^n$  which represents  $m_i$ . Let  $i, j$  with  $i \neq j$ , be in  $\{1, \dots, s+r\}$ , then  $m_i$  and  $m_j$  are in the metabolizer  $M$  and we have:  $S(m_i, m_j) = A(m_i, m_j) + (-1)^n A(m_j, m_i) = 0$ . Since  $n \geq 3$ , thanks to Whitney's procedure [Wh, 44] we can choose the  $\psi_i$  such that  $\psi_i(S^n) \cap \overline{\psi_j}(S^n) = \emptyset$ . Since  $n \geq 2$ , the Whitney obstruction to extend  $\overline{\psi_i}$  to disjoint embeddings  $\psi_i$  of  $D^{n+1}$  in the  $(2n+2)$ -disk  $D$ , is the matrix  $A(m_i, m_j)$  which is zero. Furthermore,  $A(m_i, m_i) = 0$  is the classical obstruction to extend  $\psi_i$  to  $\psi_i : D^{n+1} \times D^n \rightarrow D$ . (see [Br, 72] and for details see [Bl, 94] proposition 5.1.2, p.58). We choose this extension  $\psi_i$  such that the restriction to  $S^n \times D^n$  is a tubular neighbourhood of  $\psi_i(S^n)$  in  $G$ .  $\square$

So thanks to proposition (4.3) we obtain a submanifold  $\tilde{G}$  of  $D$  as follows:

$$\tilde{G} = (G \setminus \left( \coprod_{i=1}^{s+r} \psi_i(S^n \times D^n) \right)) \cup \left( \coprod_{i=1}^{s+r} \psi_i(D^{n+1} \times S^{n-1}) \right).$$

(4.4) **Proposition.** *The inclusion  $k_o$  (resp.  $k_1$ ) of  $K_0$  (resp.  $K_1$ ) in  $\tilde{G}$ , induces isomorphisms  $k_{o,j}$  (resp.  $k_{1,j}$ ) from  $H_j(K_0)$  (resp.  $H_j(K_1)$ ) to  $H_j(\tilde{G})$  for all  $j$ .*

(4.5) **Corollary.** *We have  $H_*(\tilde{G}, K_0) = H_*(\tilde{G}, K_1) = 0$ .*

This corollary (4.5) and the h-cobordism theorem imply that  $\tilde{G}$  is diffeomorphic to  $K_0 \times [0, 1]$ . More precisely  $\dim \tilde{G} = 2n \geq 6$  and:

**h-cobordism Theorem** [M2, 65]. *Let  $\mathcal{M}$  be a  $k$ -dimensional differentiable compact manifold with  $\partial\mathcal{M} = \mathcal{M}_0 \coprod \mathcal{M}_1$  such that  $\mathcal{M}$ ,  $\mathcal{M}_0$  and  $\mathcal{M}_1$  are simply connected. If  $H_*(\mathcal{M}, \mathcal{M}_0) = 0$  and  $k \geq 6$  then  $\mathcal{M}$  is diffeomorphic to  $\mathcal{M}_0 \times [0, 1]$ .*

So to end the proof of theorem 3 we only have to prove proposition (4.4).

*Proof of proposition (4.4).* According to proposition (2.1), the intersection form on  $H_n(F)$  splits in an orthogonal sum on the submodules  $\langle m_i, m_i^* \rangle$ ,  $i = 1, \dots, s+r$ . So the proof of (4.4) when  $s+r = 1$  implies the general case.

Let us suppose that  $\text{rk}(M) = 1$  and let  $m$  be a generator of  $M$ , then  $H_n(G) = \langle m, m^* \rangle$ . We denote by  $\psi : D^{n+1} \times D^n \rightarrow D$  an embedding choosen as in proposition (4.3), by  $\eta : S^n \rightarrow G$  an embedding such that  $[\eta(S^n)] = m^*$ , and by  $G_T$  the manifold  $G_T = G \setminus \psi(S^n \times \overset{\circ}{D^n})$ .

(4.6) The Mayer-Vietoris sequence associated to the following decomposition of the manifold:  $G = G_T \cup \psi(S^n \times D^n)$  gives:

$$0 \rightarrow H_n(\psi(S^n \times S^{n-1})) \rightarrow H_n(G_T) \oplus H_n(\psi(S^n \times D^n)) \rightarrow H_n(G)$$

$$\xrightarrow{\delta} H_{n-1}(\psi(S^n \times S^{n-1})) \rightarrow H_{n-1}(G_T) \rightarrow 0.$$

where  $\delta$  is given by the intersection of cycles with  $m$ .

(4.7) The Mayer-Vietoris sequence associated to the following decomposition of the manifold:  $\tilde{G} = G_T \cup \psi(D^{n+1} \times S^{n-1})$  gives:

$$0 \rightarrow H_n(\psi(S^n \times S^{n-1})) \xrightarrow{\alpha} H_n(G_T) \rightarrow H_n(\tilde{G}) \xrightarrow{\gamma} H_{n-1}(\psi(S^n \times S^{n-1}))$$

$$\xrightarrow{\beta} H_{n-1}(\psi(D^{n+1} \times S^{n-1})) \oplus H_{n-1}(G_T) \rightarrow H_{n-1}(\tilde{G}) \rightarrow 0.$$

Remark that the homomorphism  $\beta$  is injective into  $H_{n-1}(\psi(D^{n+1} \times S^{n-1}))$ , hence  $\gamma = 0$  and the sequence (4.7) splits up into:

$$(4.8) \quad 0 \rightarrow H_n(\psi(S^n \times S^{n-1})) \xrightarrow{\alpha} H_n(G_T) \rightarrow H_n(\tilde{G}) \rightarrow 0,$$

$$(4.9) \quad 0 \rightarrow H_{n-1}(\psi(S^n \times S^{n-1})) \xrightarrow{\beta} H_{n-1}(\psi(D^{n+1} \times S^{n-1})) \oplus H_{n-1}(G_T) \rightarrow H_{n-1}(\tilde{G}) \rightarrow 0.$$

Since  $\text{rk}(M) = 1 = s+r$  we have to consider the two following cases:  $s = 0, r = 1$  and  $s = 1, r = 0$ .

★ 1<sup>st</sup> case:  $s = 0$  and  $r = 1$ , then  $\text{Ker } S^* = \langle m, m^* \rangle$ .

In sequence (4.6) we have  $\text{Ker } \delta = \langle m, m^* \rangle$ , then  $H_n(G_T) = \langle [\psi(S^n \times \{1\})], [\eta(S^n)] \rangle$  and  $H_{n-1}(G_T) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$ . In sequence (4.8) we have  $\text{Im } \alpha =$

$\langle [\psi(S^n \times \{1\})] \rangle$ , so  $H_n(\tilde{G}) = \langle [\eta(S^n)] \rangle$ . By construction of the good basis (2.1),  $[\eta(S^n)]$  is a generator of  $\text{Im}(H_n(K_0) \rightarrow H_n(G))$ . So the inclusion of  $K_0$  in  $\tilde{G}$  induces the isomorphism:  $k_{0,n} : H_n(K_0) \xrightarrow{\cong} H_n(\tilde{G})$ .

Since  $H_{n-1}(G_T) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$  in sequence (4.9), we have  $H_{n-1}(\tilde{G}) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$ . Condition c.1 of the algebraic cobordism gives that there exists  $a$  in  $\text{Ker } S_0^*$  such that  $m = (a, \varphi(a))$ . If we denote by  $\gamma_0 : H_n(K_0) \rightarrow H_n(G)$  the homomorphism induced by the inclusion, then we can choose  $b$  in  $H_{n-1}(K_0)$  such that  $H_{n-1}(K_0) = \langle b \rangle$  and  $b$  is the dual of  $\gamma_0^{-1}(a)$  for the intersection form of  $K_0$ . There exists  $B$  in  $H_n(G, K_0)$  such that  $\partial B = b$  and the intersection between  $B$  and  $m$  is +1. The boundary of the  $n$ -chain  $(B - (B \cap \psi(S^n \times \overset{\circ}{D}{}^n)))$  is homologous to the  $(n-1)$ -cycle  $b - (\psi(\{1\} \times S^{n-1}))$ , hence  $b$  and  $[\psi(\{1\} \times S^{n-1})]$  are homologous in  $H_{n-1}(\tilde{G}) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$ . Thus the inclusion of  $K_0$  in  $\tilde{G}$  induces the isomorphism:  $k_{0,n-1} : H_{n-1}(K_0) \xrightarrow{\cong} H_{n-1}(\tilde{G})$ .

\* 2<sup>nd</sup> case:  $s = 1$  and  $r = 0$ , then  $\text{Ker } S^* = \{0\}$  and  $H_n(K_0) = 0$ .

In sequence (4.6) we have  $\text{Ker } \delta = \langle m \rangle$ , then  $H_n(G_T) = \langle [\psi(S^n \times \{1\})] \rangle$  and  $H_{n-1}(G_T) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$ . In sequence (4.8) we have  $\text{Im } \alpha = \langle [\psi(S^n \times \{1\})] \rangle$ . Since  $H_n(G_T) = \langle [\psi(S^n \times \{1\})] \rangle$  we have  $H_n(\tilde{G}) = 0 = H_n(K_0)$ .

– if  $S_*(m)$  is indivisible (i.e.  $H_{n-1}(K_0) = 0$ ), then  $\delta$  in (4.6) is surjective.

Thus  $H_{n-1}(\tilde{G}) = 0 = H_{n-1}(K_0)$ .

– If  $a \neq 1$  is the greatest divisor of  $S_*(m)$  (i.e.  $H_{n-1}(K_0) \cong \mathbb{Z}/a\mathbb{Z}$ ) then condition c.2 of algebraic cobordism together with lemma (3.14) give that there exists  $c$  in  $H_{n-1}(K_0)$  such that  $\partial(\frac{1}{a} S_*(m)) = (c, \theta_{n-1}(c))$ . Let  $b$  in  $H_{n-1}(K_0)$  be the dual of  $c$  for the linking form of  $K_0$ . There exists  $B$  in  $H_n(G, K_0)$  such that  $\partial B = b$  and the intersection between  $B$  and  $m$  is +1. As before the boundary of the  $n$ -chain  $B - (B \cap \psi(S^n \times \overset{\circ}{D}{}^n))$  is the  $n$ -cycle  $b - \psi(\{1\} \times S^{n-1})$ , hence  $b$  and  $[\psi(\{1\} \times S^{n-1})]$  are homologous in  $H_{n-1}(G)$ . Since  $H_{n-1}(G_T) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$  in sequence (4.9) we have  $H_{n-1}(\tilde{G}) = \langle [\psi(\{1\} \times S^{n-1})] \rangle$ . Thus  $b$  and  $[\psi(\{1\} \times S^{n-1})]$  are homologous in  $H_{n-1}(\tilde{G})$  and the inclusion of  $K_0$  in  $\tilde{G}$  induces the isomorphism:  $k_{0,n-1} : H_{n-1}(K_0) \xrightarrow{\cong} H_{n-1}(\tilde{G})$ .

Since  $\tilde{G}$  is obtained by surgery on  $n$ -cycles, this surgery only modifies homology groups of dimensions  $n$  and  $n-1$ . Hence for  $k \neq n, n-1$  we have  $H_k(G) \cong H_k(K_0) \xrightarrow{k_{0,k}} H_k(\tilde{G})$ . By symmetry we also have the same results with  $K_1$ . Finally  $k_{0,j}$  and  $k_{1,j}$  are some isomorphisms for all  $j$ . This ends the proof of proposition (4.4), and the proof of theorem 3.  $\square$

## 5. Appendix – Alexander polynomials of cobordant links.

Let  $K$  be a  $2n-1$  dimensional simple link, and  $\varepsilon = (-1)^n$ . One can associate a polynomial  $\Delta \in \mathbb{Z}[X]$  to any Seifert surface  $F$  for the link  $K$ , defined by:  $\Delta(X) =$

$\det(XA + \varepsilon A^T)$ , where  $A$  is the Seifert form associated to  $F$ . Such a polynomial  $\Delta$  is called a Alexander polynomial for the link  $K$ . Changing the Seifert surface to another multiplies  $\Delta$  by  $\pm X^m$  with  $m$  in  $\mathbb{Z}$ .

For a polynomial  $\gamma$  in  $\mathbb{Z}[X]$  we define the polynomial  $\gamma^*$  by:  $\gamma^*(X) = X^{\deg \gamma} \gamma(X^{-1})$ .

(5.1) **Proposition.** *Let  $K_0$  and  $K_1$  be two cobordant simple  $2n-1$  dimensional links. If  $\Delta_0$  and  $\Delta_1$  are Alexander polynomials for  $K_0$  and  $K_1$ , then there exists  $\gamma$  in  $\mathbb{Z}[X]$  such that:  $\gamma \gamma^* = \pm \Delta_0 \Delta_1$ .*

**Remark.** If  $F$  is the Milnor fiber of an algebraic link  $K$ , then the associated Alexander polynomial is the characteristic polynomial of the monodromy. Hence the above proposition and the monodromy theorem imply corollary (0.7).

*Proof of proposition (5.1).* We denote by  $F_0$  and  $F_1$  two  $(n-1)$ -connected Seifert surfaces for  $K_0$  and  $K_1$ , and by  $A_0$  and  $A_1$  the associated Seifert forms. The links  $K_0$  and  $K_1$  are cobordant so proposition (3.10) implies that the form  $A = A_0 \oplus -A_1$  has a metabolizer  $M$ . Therefore, there exists a basis for  $H_n(F_0) \oplus H_n(F_1)$  such that in this basis the matrix for  $A$  is  $\begin{pmatrix} 0 & B_1 \\ B_2 & B_3 \end{pmatrix}$  where  $B_i$ ,  $i=1,2,3$  are square matrices. We have  $\Delta_0(X) \cdot \Delta_1(X) = \det(XA + \varepsilon A^T)$ , hence  $\Delta_0(X) \cdot \Delta_1(X) = \varepsilon \cdot \det(XB_1 + \varepsilon B_2^T) \cdot \det(XB_2 + \varepsilon B_1^T)$ . Let  $\gamma(X)$  be  $\det(XB_1 + \varepsilon B_2^T)$ , then  $\gamma^*(X) = \det(XB_2 + \varepsilon B_1^T)$ . Finally we get  $\gamma \cdot \gamma^* = \pm \Delta_0 \Delta_1$ .  $\square$

### Acknowledgements

This work has been partly supported by the Fonds National Suisse de la Recherche Scientifique.

We thank D. T. Lê who already drew the attention of the second author on the cobordism of algebraic links in 1980. We also thank M. Kervaire and C. Weber for useful discussions and the University of Geneva for its hospitality.

### References

- [Bl, 94] V. Blanlœil, Cobordisme des entrelacs fibrés simples et forme de Seifert. *Thèse de l'Université de Nantes* (1994).
- [Bl, 95] V. Blanlœil, Cobordisme des entrelacs fibrés simples et forme de Seifert. *Note aux Comptes Rendus de l'Académie des Sciences de Paris* (1995), t. 320, Série I, p. 985–988.
- [Br, 72] W. Browder, Surgery on Simply-connected Manifolds. *Erger. Math.* **65** Springer, 1972.
- [DB-M, 93] P. du Bois et F. Michel, Cobordism of Algebraic Knots via Seifert Forms. *Invent. Math.* **111** (1993), 151–169.
- [D1, 74] A. Durfee, Fibered Knots and Algebraic Singularities. *Topology* **13** (1974), 47–59.
- [D2, 77] A. Durfee, Bilinear and Quadratic Forms on Torsion Modules. *Advances in Mathematics* **25** (1977), 133–164.

- [F-M, 66] R. Fox and J. Milnor, Singularities of 2-spheres in 4-spaces and Cobordism of Knots. *Osaka J. Math.* **3** (1966), 257–267.
- [K1, 65] M. Kervaire, Les noeuds de dimension supérieure. *Bulletin de la Société Mathématique de France* **93** (1965), 225–271.
- [K2, 70] M. Kervaire, Knot Cobordism in Codimension Two. Manifolds Amsterdam 1970, *Lecture Notes* **197**, 83–105.
- [K-W, 77] M. Kervaire, C. Weber, A Survey of Multidimensional knots. Knot Theory, Proceedings. Plans-sur-Bex, Switzerland 1977, *Lecture Notes* **685**, 61–134.
- [L-L, 75] J. Lannes and F. Latour, Forme quadratique d'enlacement et applications. *Société Mathématique de France Astérisque* **26** (1975).
- [L1, 69] J. Levine, Knot Cobordism in Codimension Two. *Comment Math. Helv.* **44** (1969), 229–244.
- [L2, 70] J. Levine, An Algebraic Classification of Some Knots of Codimension Two. *Comment. Math. Helv.* **45** (1970), 185–198.
- [Lê, 72] D. T. Lê, Sur les noeuds algébriques. *Compos. Math.* **25** (1972), 281–321.
- [M1, 61] J. Milnor, A Procedure for Killing Homotopy Groups of Differentiable Manifolds. *Proceeding of Symposia in Pure Math. (A.M.S.)* (1961) t.3, 39–55.
- [M2 65] J. Milnor, Lectures on the h-Cobordism Theorem. Princeton Mathematical Notes, Princeton U. Press, 1965.
- [M3, 68] J. Milnor, Singular Points of Complex Hypersurfaces. *Annals of Math. Studies* **61** (1968).
- [Sa, 74] K. Sakamoto, The Seifert Matrices of Milnor Fiberings defined by Holomorphic Functions. *J. Math. Soc. Japan* **26**(4) (1974), 714–721.
- [V1, 77] R. Vogt, Cobordismus von Knoten. *Lect. Notes in Math.* **685** (1977), 218–226.
- [V2, 78] R. Vogt, Cobordismus von hochzusammenhängenden Knoten. *Inauguraldissertation zur Erlangung des Doktorgrades*. Bonn 1978.
- [W, 70] C.T.C. Wall, *Surgery on Compact Manifolds*. Academic Press, New York 1970.
- [Wh, 44] H. Whitney, The Self-Intersection of a Smooth  $n$ -Manifold in  $2n$ -Spaces. *Annals of Math.* **45** (1944), 220–246.

Vincent Blanlœil  
 Section de Mathématiques  
 Université de Genève  
 2-4, rue du Lièvre  
 Case postale 240  
 1211 Genève 24 Suisse  
 e-mail: blanloeil@sc2a.unige.ch

Françoise Michel  
 Département de Mathématiques  
 Université de Nantes  
 2, rue de la Houssinière  
 44072 Nantes cedex 03 France  
 e-mail: fmichel@math.univ-nantes.fr

(Received: August 24, 1995)