

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 72 (1997)

Artikel: Volume-preserving mean curvature flow of rotationally symmetric surfaces
Autor: Athanassenas, Maria
DOI: <https://doi.org/10.5169/seals-54577>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Volume-preserving mean curvature flow of rotationally symmetric surfaces

Maria Athanassenas

Abstract. A rotationally symmetric n -dimensional surface in \mathbb{R}^{n+1} , of enclosed volume V and with boundary in two parallel planes, is evolving under volume-preserving mean curvature flow. For large volume V , we obtain gradient and curvature estimates, leading to long-time existence of the flow, and convergence to a constant mean curvature surface.

Mathematics Subject Classification (1991). 35K22, 58G11, 53C21.

Keywords. Geometric evolution equations, motion by mean curvature under a volume constraint.

Introduction

Consider n -dimensional hypersurfaces M_t , defined by a one parameter family of smooth immersions $x_t : M^n \rightarrow \mathbb{R}^{n+1}$, with $M_t = x_t(M^n)$. The hypersurfaces M_t are said to move by mean curvature, if $x_t = x(\cdot, t)$ satisfies

$$\frac{d}{dt}x(p, t) = -H(p, t)\nu(p, t), \quad p \in M^n, \quad t > 0. \quad (1)$$

By $\nu(p, t)$ we denote a choice of unit normal of M_t at $x(p, t)$, and by $H(p, t)$ the mean curvature with respect to this normal. The surface area $|M_t|$ of the hypersurfaces is known to decrease under the flow. So the evolution can be used for obtaining minimal surfaces in the limit, if it converges.

Here we are interested in the evolution of compact hypersurfaces M_t enclosing a prescribed volume V . In particular, we consider the evolution equation

$$\frac{d}{dt}x(p, t) = -(H(p, t) - h(t))\nu(p, t), \quad p \in M^n, \quad t > 0. \quad (2)$$

where $h(t)$ is the average of the mean curvature

$$h(t) = \frac{\int_M H dg_t}{\int_M dg_t},$$

and g_t the metric on M_t . As initial surface we choose a compact n -dimensional hypersurface M_0 , with boundary $\partial M_0 \neq \emptyset$. We assume M_0 to be smoothly embedded in the domain

$$G = \{x \in \mathbb{R}^{n+1} : 0 < x_{n+1} < d\}, \quad d > 0,$$

and $\partial M_0 \subset \partial G$. The vector $\nu(p, t)$ is the outer unit normal.

The surface area $|M_t|$ is again decreasing under the flow defined by (2) and in addition the enclosed volume is constant (see Section 1). In this case the hypersurfaces can be expected to converge to a surface of constant mean curvature which solves the isoperimetric problem.

The motion of surfaces by their mean curvature (1) was first studied by Brakke [5], using methods of geometric measure theory. Huisken [22] proves that compact, convex initial surfaces without boundary converge asymptotically to round spheres. Gage and Hamilton [17], Grayson [19] study the problem for curves in the plane. In the noncompact case it is shown by Ecker and Huisken in [13] that entire graphs over \mathbb{R}^n of linear growth 'flatten out'.

The interesting question of the formation of singularities in the nonconvex case is considered by Huisken [24], Grayson [20], Dziuk and Kawohl [9], and more recently by Altschuler, Angenent and Giga [1], Ecker [12].

For the volume-preserving flow (2), Huisken [23] proves long-time existence if the initial hypersurface M_0 is compact, without boundary and uniformly convex; eventually the M_t 's converge to a round sphere enclosing the same volume as M_0 .

The uniform convexity is crucial for the proof; using a maximum principle for parabolic systems developed by Hamilton in ([21], Theorem 9.1), Huisken shows that uniform convexity is preserved for $t > 0$.

The major difficulty in the volume-preserving evolution (2), and its difference to the mean curvature flow (1), is how to control h , which introduces a global character to the problem. Parabolic maximum principles, an important tool in the investigation of evolution equations (see [10]), either fail or become more subtle.

In this paper, except for the volume constraint, we have a free boundary. A convexity assumption is not natural. We replace this by assuming the initial surface to be rotationally symmetric, contained in the region G between two parallel hyperplanes and for a start the surfaces M_t to intersect ∂G orthogonally at the boundary. The motivation is the fact that in solving the isoperimetric problem using methods of the calculus of variations, the minimizers prove to be surfaces of revolution intersecting the obstacle at a right angle [2], [3].

Mean curvature flow (without a volume constraint) for complete rotationally symmetric surfaces has been studied by Simon [27]. He gives gradient and height estimates and discusses pinch off behaviour. His approach is especially interesting for us. Dziuk and Kawohl [9], Grayson [20] and Altschuler, Angenent and Giga [1] consider the question of developing singularities.

The main theorem we prove is

Theorem. *Assume $V, d \in \mathbf{R}$ to be given and $M_0 \subset G$ to be a smooth, rotationally symmetric, initial hypersurface which intersects ∂G orthogonally at the boundary and encloses the volume V . Then the flow defined by (2) will exist for all times $t > 0$ and will converge to the cylinder $C \subset G$ of volume V under the assumption*

$$|M_0| \leq \frac{V}{d}.$$

The paper is organised as follows:

In Section 1 we give some definitions and preliminaries. For ‘large’ volume (see Lemma 1) we prove in Section 2 that the surfaces do not pinch off. Also in this case h is shown to be bounded. Gradient and curvature estimates (Section 4 and 5) lead to long-time existence and convergence to a constant curvature surface (Section 6).

Rotationally symmetric surfaces of constant mean curvature in \mathbb{R}^3 are known as the Delaunay surfaces [8]; they are plane, the sphere, the cylinder, the catenoid, the unduloid and the nodoid. In [2] the author proves that among them only the sphere (hemisphere) or the cylinder can be stable in G , depending on the volume. The present condition on the volume (see Lemma 1) excludes the existence of unduloids in G . The flow can only converge to a cylinder, which affirms the result in [2].

The methods we use here are those introduced by Huisken [22] for the mean curvature flow, and also used for instance in [10], [13], [14], [23].

Mean curvature flow, but not the volume-preserving problem, is also investigated from different points of view. Evans and Spruck [15], [16], Chen, Giga and Goto [6], [7] work with hypersurfaces which are defined as level sets of viscosity solutions of a nonlinear partial differential equation on some domain in \mathbb{R}^{n+1} . Regularity results are given in [15], [16], [18], [25], [26].

The author would like to thank Klaus Ecker for his interest and support, and the Department of Mathematics at the University of Melbourne and the Centre for Mathematics and its Applications at the Australian University for their hospitality.

1. Notations and preliminaries

Let $G = \{x \in \mathbb{R}^{n+1} : 0 < x_{n+1} < d\}$, for given $d > 0$. We denote by Π_i , $i = 1, 2$, the two parallel hyperplanes bounding the domain G .

The n -dimensional manifold M^n is assumed to be smoothly embedded in \mathbb{R}^{n+1} , compact, with boundary ∂M . The one-parameter family of surfaces obtained by the flow is defined by means of the position vector

$$x : M^n \times [0, t_1) \rightarrow \mathbb{R}^{n+1},$$

where x satisfies the evolution equation (2) above.

By M_t we denote the image $M_t = x_t(M^n)$ and M_0 will be a given initial surface. In addition we assume

(i) The hypersurface M_0 is rotationally symmetric about an axis which intersects Π_i orthogonally.

We also use the parametrization

$$\rho_S : [0, d] \rightarrow \mathbb{R}$$

for the generating curve of a surface S of revolution. Actually, the flow preserves rotational symmetry (see Fact 1 below).

(ii) The boundary $x_t(\partial M) = \partial M_t$ is contained in $\partial G = \cup_{i=1,2} \Pi_i$.
 (iii) M_t intersects ∂G orthogonally at the free boundary; i.e. $\dot{\rho}(z) = 0$, for $z = 0$ and $z = d$. (Here $\dot{\rho} = \frac{d\rho}{dz}$.)

By $g = \{g_{ij}\}$ and $A = \{h_{ij}\}$ we denote the metric and the second fundamental form on M_t . For the mean curvature and the norm of the second fundamental form we have

$$H = g^{ij} h_{ij}, \quad |A|^2 = g^{ij} g^{kl} h_{ik} h_{jl}.$$

Let

$$\tilde{C} = g^{ij} g^{kl} g^{mn} h_{ik} h_{lm} h_{nj}.$$

Latin indices are used for n and Greek for $n + 1$ dimensions, if not otherwise specified.

Facts:

1. The flow preserves rotational symmetry. This is clear from the evolution equation, since the mean curvature and the normal are symmetric.
2. The surface area $|M_t|$ is decreasing. To see this we need the evolution equation of the metric

$$\frac{d}{dt} g_{ij} = 2(h - H)h_{ij}$$

(compare ([23], Proposition 1.1), [22])

Therefore

$$\frac{d}{dt} \sqrt{\det g_{ij}} = -H(H - h) \sqrt{\det g_{ij}}$$

and using the mean value property of h

$$\frac{d}{dt} |M_t| = - \int_M H(H - h) dg_t = - \int_M (H - h)^2 dg_t \leq 0.$$

3. The enclosed volume V is preserved. Denote by $E_t \subset G$ the $(n+1)$ -dimensional set, with boundary in G equal to M_t . Then the evolution (2) can be extended to a vector field on the whole of E_t and by the first variation formula and the

divergence theorem we have

$$\begin{aligned} \frac{d}{dt}V_t &= \int_{E_t} \operatorname{div} \frac{dx}{dt} d\mathcal{H}^{n+1} = \int_{\partial E_t} \frac{dx}{dt} \cdot \nu d\sigma \\ &= - \int_M (H - h) dg_t = 0. \end{aligned}$$

By standard parabolic theory the flow exists for some short time $0 < t < t_1$. We write also $[0, T_{\max})$ to indicate the maximal time interval for which the flow exists.

2A. Height estimates

By purely geometric arguments it is possible to show that if the enclosed volume is sufficiently large, the surfaces do not pinch off. The condition on the volume is such that, in \mathbb{R}^3 , there are no parts of unduloids satisfying it and at the same time intersecting the planes perpendicularly.

We will need the following notation: Given an initial surface M_0 , we denote by C the cylinder with same enclosed volume V as M_0 , and height d .

Lemma 1. *If $|M_0| \leq \frac{V}{d}$, then there exists $c_0 > 0$ such that*

$$\rho_{M_t} > c_0, \quad \text{for } 0 \leq z \leq d, \quad t \in [0, t_1].$$

Proof. We recall that $|M_t| \leq |M_0|$ for all $t > 0$. Let us now assume that there is some $t_1 > 0$ such that M_{t_1} pinches off. We project M_{t_1} onto the plane Π_1 , using $\pi : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^n$. Then

$$|M_{t_1}| \geq |\pi(M_{t_1})|.$$

By the volume constraint any M_t has to intersect the cylinder C at least once. Therefore

$$|\pi(M_{t_1})| > |\pi(C)| = \omega_n \rho_C^n = \frac{V}{d}$$

and we obtain a contradiction to M_{t_1} pinching off if only we choose V appropriately. \square

Remarks. (i) Knowing the absolute minimizers in \mathbb{R}^3 (compare [2]), we can easily see that for small enclosed volume V there is no initial surface satisfying the condition of the Lemma. If $V \leq \pi(\frac{2}{3})^4 d^3$ we denote by S the (in this case absolute) minimizing hemisphere. Then

$$|S| \leq |M|$$

for any surface M with the same enclosed volume, but

$$|S| > \frac{V}{d} (= |\pi(C)|).$$

(ii) In \mathbb{R}^3 there are no unduloids intersecting the planes Π_i perpendicularly at the free boundary and satisfying the volume condition of Lemma 1. As is well-known [8], the generating curve of the unduloid is obtained by rolling an ellipse along the z -axis, and tracing the path of either its foci. Let the ellipse be parametrized by $\gamma : [0, 2\pi) \rightarrow \mathbb{R}^3$, and denote by $a, b \in \mathbb{R}$, $0 < b \leq a$ its axes and by $l(\gamma)$ its length. Then the condition for the unduloid to intersect the planes at right angles is

$$k \frac{l(\gamma)}{2} = d, \quad k \in \mathbb{N},$$

$\frac{k}{2}$ the periods, i.e. $2a \leq d$. Consider the cylinder of enclosed volume $V = 4\pi d^3$ (see assumption of Lemma 1, for $n = 3$). Its radius is given by $\rho = 2d$. Any unduloid of the same enclosed volume would have to intersect this cylinder. This is not possible for $2a \leq d$. Thus, assuming $V \geq 4\pi d^3$ (actually, we only need $V \geq \pi d^3$) the flow will never converge to an unduloid in \mathbb{R}^3 .

(iii) Height estimates from above.

Assume there exists an R such that $\rho_{M_t} \geq R$ at some given time t . We would then have

$$|M_0| \geq |M_t| > \omega_n (R - \rho_C)^n$$

by comparing the surface area of M_t to that of the n -dimensional annulus of radii ρ_C and R . Here $\rho_C = (\frac{V}{\omega_n d})^{1/n}$ denotes the radius of the cylinder C of enclosed volume V . Of course M_t has to intersect this cylinder.

We deduce from the above that

$$R \geq \left(\frac{|M_0|}{\omega_n} \right)^{1/n} + \left(\frac{V}{\omega_n d} \right)^{1/n}$$

would contradicts the fact that the evolution decreases surface area.

2B. Estimates on h

Lemma 2. *Assume $M \subset G$ to be a smooth, rotationally symmetric hypersurface, intersecting Π_i orthogonally at the boundary and with radius function $\rho \geq c_0 > 0$. Then the mean value h of the mean curvature satisfies*

$$0 \leq h \leq c_1,$$

where c_1 depends on the dimension and the height estimates.

Proof. M being rotationally symmetric we have for the mean curvature $H = \kappa_1 + (n-1)\kappa_2$, where κ_1 and κ_2 denote the principal curvatures. If we parametrize M by its radius function $\rho \in C^\infty([0, d], [c_0, R])$, then clearly

$$H = -\frac{\ddot{\rho}}{(1+\dot{\rho}^2)^{\frac{3}{2}}} + \frac{n-1}{\rho(1+\dot{\rho}^2)^{\frac{1}{2}}}$$

and

$$h = \frac{1}{|M|} \int_M H dg = \frac{\int_0^d \left(-\frac{\ddot{\rho}}{(1+\dot{\rho}^2)^{\frac{3}{2}}} \rho^{n-1} + (n-1)\rho^{n-2} \right) dz}{\int_0^d \rho^{n-1} \sqrt{1+\dot{\rho}^2} dz}.$$

For the second term we have

$$0 \leq \frac{1}{|M|} \int_M (n-1)\kappa_2 dg \leq c(n, c_0).$$

For the first we remark that

$$\frac{\ddot{\rho}}{(1+\dot{\rho}^2)} = \frac{d}{dz}(\arctan \dot{\rho})$$

and therefore

$$\begin{aligned} \frac{n\omega_n}{|M|} \int_0^d -\frac{\ddot{\rho}}{(1+\dot{\rho}^2)} \rho^{n-1} dz &= \frac{n\omega_n}{|M|} \int_0^d -\frac{d}{dz}(\arctan \dot{\rho}) \rho^{n-1} dz \\ &= \frac{n(n-1)\omega_n}{|M|} \int_0^d (\arctan \dot{\rho}) \dot{\rho} \rho^{n-2} dz. \end{aligned}$$

This is positive, as $(\arctan \dot{\rho}) \dot{\rho} \geq 0$, and bounded

$$\frac{1}{|M|} \int_M \kappa_1 dg \leq \frac{n(n-1)\omega_n \pi}{|M|} \frac{1}{2} \int_0^d \sqrt{1+\dot{\rho}^2} \rho^{n-2} dz \leq \frac{(n-1)\pi}{|M|} \frac{1}{2} \int_M \frac{1}{\rho} dg \leq c'(n, c_0),$$

since $|(\arctan \dot{\rho}) \dot{\rho}| \leq \frac{\pi}{2} |\dot{\rho}| \leq \frac{\pi}{2} \sqrt{1+\dot{\rho}^2}$. \square

3. Evolution equations

Notation. Let $\omega = \frac{\dot{x}}{|x|} \in \mathbb{R}^{n+1}$, $\hat{x} = (x_1, \dots, x_n, 0)$ denote the unit outer normal to the cylinder intersecting M_t at the point $x(p, t)$. Set

$$v = (\omega, \nu)^{-1}.$$

Remark that $v = \sqrt{1 + \rho^2}$, where ρ is the radius of M_t , i.e. the “height” of the generating curve. We call $u = \langle x, \omega \rangle$ the height function of M_t .

Lemma 3. *We have the evolution equations*

- (i) $\left(\frac{d}{dt} - \Delta \right) u = h \langle \nu, \omega \rangle - \frac{n-1}{u}$,
- (ii) $\left(\frac{d}{dt} - \Delta \right) v = -|A|^2 v + \frac{n-1}{u^2} v - 2v^{-1} |\nabla v|^2$,
- (iii) $\left(\frac{d}{dt} - \Delta \right) H = (H - h)|A|^2$,
- (iv) $\left(\frac{d}{dt} - \Delta \right) |A|^2 = -2|\nabla A|^2 + 2|A|^4 - 2h\tilde{C}$,

where $\tilde{C} = g^{ij}g^{kl}g^{mn}h_{ik}h_{lm}h_{nj}$.

Proof. (iii) and (iv) are the same as in [23].

(i) We use the notation D for the gradient in \mathbb{R}^{n+1} and denote by $D\omega$ the $(n+1) \times (n+1)$ matrix of first derivatives of ω . We have

$$(D\omega)_{\alpha\beta} = \begin{cases} \frac{1}{u}(\delta_{\alpha\beta} - \omega_\alpha \omega_\beta) & \text{for } 1 \leq \alpha, \beta \leq n, \\ 0 & \text{for } \alpha = n+1 \text{ or } \beta = n+1. \end{cases}$$

We need the identities

$$\langle \nu, D\omega \cdot \nu \rangle = \frac{1}{u}(\nu_i \nu_i - \langle \omega, \nu \rangle^2) = 0, \quad (1)$$

$$\langle x, D\omega \cdot \nu \rangle = \frac{1}{u}(x_i \nu_i - \langle \omega, x \rangle \langle \omega, \nu \rangle) = 0, \quad (2)$$

$$Du = \omega \quad \text{and} \quad \nabla^M u = \omega - \langle \omega, \nu \rangle \nu. \quad (3)$$

Using (2) we have

$$\frac{d}{dt} u = -(H - h) \langle \omega, \nu \rangle. \quad (4)$$

From (1), (3) and with a suitable choice of basis $\{e_i\}_{1 \leq i \leq n}$ for $T_x M_t$ we obtain

$$\begin{aligned} \operatorname{div}_M \omega &= \operatorname{div} \omega - \langle D\omega \cdot \nu, \nu \rangle = \frac{n-1}{u} \\ \text{and} \quad \operatorname{div}_M (\langle Du, \nu \rangle \nu) &= e_i \langle Du, \nu \rangle \langle e_i \rangle + \langle Du, \nu \rangle \operatorname{div}_M \nu = \langle \omega, \nu \rangle H; \end{aligned} \quad (5)$$

therefore

$$\Delta^M u = \operatorname{div}_M \nabla^M u = \operatorname{div}_M \omega - \operatorname{div}_M (\langle Du, \nu \rangle \nu) = \frac{n-1}{u} - H \langle \omega, \nu \rangle. \quad (6)$$

(4) and (6) imply

$$\left(\frac{d}{dt} - \Delta \right) u = h \langle \nu, \omega \rangle - \frac{n-1}{u}.$$

(ii) As in ([22], Lemma 3.3), we obtain $\frac{d}{dt} \nu = \nabla H$. Therefore, and by (1)

$$\frac{d}{dt} v = -v^2 \langle \nabla H, \omega \rangle. \quad (7)$$

The following computation is as in ([4], Prop. 2.1) adapted to our case. For computing the Laplacian of v we will work with normal coordinates in a neighbourhood of the point $x(p, t)$. Also, we will need the relations

$$\nabla_{e_i} \nu = h_{ij} g^{jk} e_k, \quad \nabla_{e_i} e_j = -h_{ij} \nu + (\nabla_{e_i} e_j)'' , \quad (8)$$

$$[\omega, e_i] = 0, \quad (9)$$

where $''$ denotes the tangential part of a vectorfield and $[,]$ the Lie bracket. Then, using (8) and (9), we have

$$\begin{aligned} \Delta \langle \omega, \nu \rangle &= e_i e_i \langle \omega, \nu \rangle \\ &= e_i \langle \nabla_\omega e_i, \nu \rangle + e_i \langle \omega, g^{jk} h_{ij} e_k \rangle \\ &= \langle \nabla_{e_i} \nabla_\omega e_i, \nu \rangle + h_{ij} g^{jk} \langle \nabla_\omega e_i, e_k \rangle + e_i (h_{ij} g^{jk} \langle \omega, e_k \rangle). \end{aligned}$$

Having normal coordinates at x , \mathbb{R}^{n+1} being flat, and using (8), (9), as well as the Codazzi equations, for exchanging derivatives of the h_{ij} , we obtain

$$\begin{aligned} e_i(h_{ij}g^{jk}\langle\omega, e_k\rangle) &= h_{ij}g^{jk}(\langle\nabla_\omega e_i, e_k\rangle + \langle\omega, \nabla_{e_i}e_k\rangle) + e_i(h_{ij}g^{jk})\langle\omega, e_k\rangle \\ &= h_{ij}\langle\nabla_\omega e_i, e_j\rangle - |A|^2\langle\omega, \nu\rangle + e_j(h_{ij})\langle\omega, e_j\rangle \end{aligned}$$

and

$$\begin{aligned} \langle\nabla_{e_i}\nabla_\omega e_i, \nu\rangle &= \langle\nabla_\omega\nabla_{e_i}e_i, \nu\rangle = \langle\nabla_\omega(-h_{ii}\nu + (\nabla_{e_i}e_i)'''), \nu\rangle \\ &= -\omega(h_{ii}) = -\omega(h_{ij})g^{ij}. \end{aligned}$$

We deduce

$$\begin{aligned} \Delta\langle\omega, \nu\rangle &= -\omega(H) + 2h_{ij}\langle\nabla_\omega e_i, e_j\rangle - |A|^2\langle\omega, \nu\rangle + \langle\omega, \nabla H\rangle \\ &= -\omega(h_{ij})g^{ij} - h_{ij}\omega(g^{ij}) - |A|^2\langle\omega, \nu\rangle + \langle\omega, \nabla H\rangle. \end{aligned}$$

Combining this with (7) we obtain (ii). \square

4. Gradient estimates

General assumption. For the remaining of the paper we always assume the enclosed volume V to be so large that the results of Section 2 hold; i.e. the evolving surfaces do not pinch off and h is bounded.

Remark. Since we have an evolution of rotationally symmetric surfaces, which intersect the hyperplanes Π_i orthogonally, and $v = \sqrt{1 + \rho^2}$ along the generating curve, the following Proposition 4 gives the gradient estimate.

Proposition 4. *Under the assumptions of Section 1 and the large volume of Lemma 1, if we assume $v \leq v_0$ on the initial surface M_0 , then*

$$\max_{t>0} v \leq c_2(n, c_0, R, v_0).$$

Proof. Given the Neumann boundary conditions at $z = 0$ and $z = d$, we can equivalently consider the evolution of periodic surfaces \tilde{M}_t defined along the whole z -axis. We assume that the product u^2v attains a maximum, denoted by K , on \tilde{M}_{t_1} for $t_1 > 0$. The idea is to prove that if K is large enough then $(\frac{d}{dt} - \Delta)u^2v \leq 0$ at this maximum point.

In Lemma 3 we obtained the evolution equations

$$\begin{aligned}\left(\frac{d}{dt} - \Delta\right)v &= -|A|^2v + \frac{n-1}{u^2}v - 2v^{-1}|\nabla v|^2, \\ \left(\frac{d}{dt} - \Delta\right)u &= h\langle\nu, \omega\rangle - \frac{n-1}{u},\end{aligned}$$

and hence

$$\left(\frac{d}{dt} - \Delta\right)u^2 = 2u\left(hv^{-1} - \frac{n-1}{u}\right) - 2|\nabla u|^2.$$

For the following calculation we remark that

$$-2\nabla u^2 \cdot \nabla v = -2v^{-1}\nabla v \cdot \nabla(u^2v) + 2v^{-1}u^2|\nabla v|^2.$$

We have

$$\begin{aligned}\left(\frac{d}{dt} - \Delta\right)u^2v &= u^2\left(\frac{d}{dt} - \Delta\right)v + v\left(\frac{d}{dt} - \Delta\right)u^2 - 2\nabla u^2 \cdot \nabla v \\ &= u^2\left(-|A|^2v + \frac{n-1}{u^2}v\right) + v2u\left(hv^{-1} - \frac{n-1}{u}\right) \\ &\quad - v^2|\nabla u|^2 - 2v^{-1}\nabla v \cdot \nabla(u^2v) \\ &\leq -(n-1)v + 2hu\end{aligned}$$

at a maximum K of the product u^2v ; in particular, $v \geq \frac{K}{R^2}$ at this point, since $c_0 \leq u \leq R$ (Lemma 1). We also have $0 \leq h \leq c_1(n, c_0)$ (Lemma 2).

We deduce that $(\frac{d}{dt} - \Delta)u^2v \leq 0$ at the maximum point whenever

$$K > \frac{2R^3c_1(n, c_0)}{n-1}.$$

Therefore

$$\max_{t>0} u^2v \leq \frac{2R^3c_1(n, c_0)}{n-1},$$

and

$$\max_{t>0} v \leq \max\left(\frac{2R^3c_1(n, c_0)}{(n-1)c_0^2}, v_0\right).$$

□

5. Curvature estimates

In this section we prove that as long as M_t doesn't pinch off and its generating curve is a graph over the x_{n+1} -axis with bounded gradient, the curvature and all its derivatives remain bounded as well.

Proposition 5. *Under the assumptions of Proposition 4, there exist positive constants $c_3(n, c_0, R, v_0)$, c_4 and c_5 , such that the curvature of the evolving surfaces M_t is bounded*

$$\max_{t>0} |A|^2 \leq c_3(n, c_0, R, v_0) \left(\frac{c_4}{\sqrt{c_5}} + \frac{1}{c_5 t} \right), \quad \text{for } t \in (0, T_{\max}].$$

Remark. Actually $c_3(n, c_0, R, v_0)$ depends on $c_2(n, c_0, R, v_0)$ of Proposition 4, whereas c_5 is a technical constant, depending on the chosen testfunction in the proof, and c_4 depends on the dimension, the height estimates and the bounds on h and v , as from Lemma 2 and Proposition 4.

Proof. We proceed as in ([14], proof of Theorem 3.1) and calculate the evolution equation of the product $g = |A|^2 \varphi(v^2)$, where $\varphi(r) = \frac{r}{1-kr}$, $k > 0$, and $v = \langle \nu, \omega \rangle^{-1}$. The only difference being the volume constraint, which affects the evolution equations of $|A|^2$ and v by an additional $-2h\tilde{C}$ and $\frac{n-1}{u^2}v$, respectively, (compare Lemma 3), we end up with the inequality

$$\left(\frac{d}{dt} - \Delta \right) g \leq -2kg^2 - \frac{2k}{(1-kv^2)^2} |\nabla v|^2 g - 2\varphi v^{-3} \nabla \varphi \cdot \nabla g - 2h\tilde{C}(v^2) + \frac{2(n-1)}{u^2} v^2 \varphi'$$

which replaces (20) of [14].

For estimating the seconds-last term in our case, we use Young's inequality, and obtain

$$\begin{aligned} -2h\tilde{C}\varphi(v^2) &\leq 2h|A|^3\varphi(v^2) \\ &\leq \frac{3}{2}|A|^4\varphi^2(v^2) + \frac{1}{2}h^4\varphi^{-2}(v^2) \\ &= \frac{3}{2}g^2 + \frac{1}{2}h^4\varphi^{-2}(v^2). \end{aligned}$$

We now choose $k > \frac{3}{4}$ and deduce

$$\left(\frac{d}{dt} - \Delta \right) g \leq -c_5 g^2 - a \cdot \nabla g + c_4(n, c_0, c_1, c_2), \quad (1)$$

where $c_5 > 0$ and c_1 and c_2 denote the constants in the bounds on h and v in Lemma 2 and Proposition 4.

By Corollary 1.4 in [11], if g satisfies (1), then

$$g \leq \frac{c_4}{\sqrt{c_5}} + \frac{1}{c_5 t}$$

on M_t , $t \in (0, T]$. The result follows then from the definition of g as the product $|A|^2 \varphi(v^2)$ and the gradient estimates. \square

For the higher curvature derivatives we have under the assumptions of Proposition 4

Proposition 6. *For each $m \geq 1$ there is C_m such that*

$$|\nabla^m A|^2 \leq C_m$$

uniformly on M_t for $0 \leq t \leq T_{\max} \leq \infty$.

Proof. Having obtained uniform bounds on $|A|^2$ (Proposition 5) and h (Lemma 2) the proof is a repetition of that of Theorem 4.1 in [23].

Thus, we have long-time existence for the flow:

Corollary 7.

$$T_{\max} = \infty.$$

6. Convergence to surfaces of constant mean curvature

Having long-time existence for the flow it remains to show that it converges to a constant mean curvature surface as $t \rightarrow \infty$.

Proposition 8. *The mean curvature H of the evolving surfaces converges to its average*

$$\sup_{M_t} |H - h| \rightarrow 0 \quad \text{as } t \rightarrow \infty.$$

Proof. We note that

$$\frac{d}{dt}|M_t| = \int_M (H - h)^2 dg_t$$

and therefore

$$\int_0^\infty \int_M (H - h)^2 dg_t dt \leq |M_0|.$$

Given the uniform estimates on all curvature quantities and their derivatives (section 5), we have that

$$\int_M (H - h)^2 dg_t \quad \text{and} \quad \frac{d}{dt} \int_M (H - h)^2 dg_t$$

are uniformly bounded. Thus, we obtain

$$\int_M (H - h)^2 dg_t \rightarrow 0 \quad \text{for } t \rightarrow \infty.$$

In order to obtain the result we will now use a standard interpolation argument. We first estimate the supremum norm using the Sobolev inequality for $p > n$ with a constant uniform in time; we then use the a-priori estimates we obtained for the curvature and its derivatives and integration by parts to deduce

$$\begin{aligned} \sup_{M_t} |H - h| &\leq \left(\int_M |\nabla(H - h)|^p dg_t \right)^{1/p} \\ &\leq c \left\{ \left(\int_M (H - h)^2 dg_t \right)^{1/2} + \left(\int_{\partial M} (H - h)^2 d\sigma_t \right)^{1/2} \right\}, \end{aligned}$$

which proves the result. \square

Remark. Having long-time existence, the evolution will eventually converge to an extremum of the surface energy under a volume constraint. The above curvature estimates provide us then with a regularity result for such surfaces.

References

- [1] S. Altschuler, S. Angenent and Y. Giga, Mean curvature flow through singularities for surfaces of rotation, preprint.
- [2] M. Athanassenas, A variational problem for constant mean curvature surfaces with free boundary, *J. Reine Angew. Math.* **377** (1987), 97–107.
- [3] M. Athanassenas, A free boundary problem for capillary surfaces, *Manuscripta Math.* **76** (1992), 5–19.

- [4] R. Bartnik, Existence of maximal surfaces in asymptotically flat spacetimes, *Comm. Math. Phys.* **94** (1984), 155–175.
- [5] K. A. Brakke, *Math. Notes*, Princeton University Press, Princeton, N.Y. 1978.
- [6] X. Y. Chen, Y. Giga, S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations, *J. Diff. Geom.* **33** (1991), 749–786.
- [7] X. Y. Chen, Y. Giga and S. Goto, Analysis toward snow crystal growth, to appear in *Proceeding of functional Analysis and related topics*.
- [8] Ch. Delaunay, Sur la surface de révolution dont la courbure moyenne est constante, *Journ. Math. pures et appliquées* **6** (1841), 309–320.
- [9] G. Dziuk and B. Kawohl, On rotationally symmetric mean curvature flow, preprint Nr. 108, SFB 256, Bonn.
- [10] K. Ecker, Local techniques for mean curvature flow, *Proceedings of the Centre for Mathematics and its Applications, Australian National University* **26** (1991), 107–119.
- [11] K. Ecker, On mean curvature flow of spacelike hypersurfaces in asymptotically flat space-times, *J. Austral. Math. Soc. (series A)* **55** (1993), 41–59.
- [12] K. Ecker, An ϵ -regularity theorem for mean curvature flow of surfaces in 3-manifolds, to appear.
- [13] K. Ecker, G. Huisken, Mean curvature evolution of entire graphs, *Annals of Math.* **130** (1989), 453–471.
- [14] K. Ecker and G. Huisken, Interior estimates for hypersurfaces moving by mean curvature, *Invent. Math.* **105** (1991), 547–569.
- [15] L. S. Evans and J. Spuck, Motion of level sets by mean curvature I, *J. Diff. Geom.* **33** (1991), 635–681.
- [16] L. S. Evans and J. Spuck, Motion of level sets by mean curvature II, to appear.
- [17] M. Gage and R. S. Hamilton, The heat equation shrinking convex plane curves, *J. Diff. Geom.* **23** (1986), 69–96.
- [18] Y. Giga and S. Goto, Geometric evolutions of phase boundaries, preprint.
- [19] M. Grayson, The heat equation shrinks embedded plane curves to round points, *J. Diff. Geom.* **31** (1987), 285–314.
- [20] M. Grayson, A short note on the evolution of a surface by its mean curvature, *Duke Math. J.* **58** (1989), 555–558.
- [21] R. S. Hamilton, Three-manifolds with positive Ricci curvature, *J. Diff. Geom.* **17** (1982), 255–306.
- [22] G. Huisken, Flow by mean curvature of convex surfaces into spheres, *J. Diff. Geom.* **20** (1984), 237–266.
- [23] G. Huisken, The volume preserving mean curvature flow, *J. Reine Angew. Math.* **382** (1987), 35–48.
- [24] G. Huisken, Asymptotic behaviour for singularities of the mean curvature flow, *J. Diff. Geom.* **31** (1990), 285–299.
- [25] T. Ilmanen, Generalized flow of sets by mean curvature on a manifold, preprint.
- [26] T. Ilmanen, The level set flow on a manifold, preprint.
- [27] M. Simon, Mean curvature flow of rotationally symmetric surfaces, *B.Sc. Thesis, Dept. of Math. ANU, Canberra*.

Maria Athanassenas
 Department of Mathematics
 University of Melbourne
 Parkville, VIC 3052
 Australia
 e-mail: athanas@gizmo.maths.monash.edu.au

(Received: September 27, 1995)