**Zeitschrift:** Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

**Band:** 74 (1999)

Artikel: Acyclic covers

Autor: Schafer, James A.

**DOI:** https://doi.org/10.5169/seals-55781

# Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

## **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 03.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

## Commentarii Mathematici Helvetici

# Acyclic covers

James A. Schafer

**Abstract.** The following question, which is directly related to the Whitehead problem of sub-complexes of acyclic 2-complexes, is studied: If  $\mathfrak{P}$  is a class of groups, X is a 2-dimensional CW-complex and X' is an acyclic, infinite cyclic cover of X with  $\pi_1(X')$  in  $\mathfrak{P}$ , must X' be contractible? A positive answer is given if X is finite and  $\mathfrak{P}$  is the class of amenable groups.

Mathematics Subject Classification (1991). Primary 57M20; Secondary 20J05.

Keywords. 2-complexes, covering space, amenable.

In [12], J.H.C. Whitehead posed the following problem.

Whitehead's Aspherical Complex Question. If L is a connected subcomplex of an aspherical 2-dimensional CW-complex K, must L also be aspherical?

Although some early work was done on this question by Cockcroft [4] this paper is concerned with the approach originating with the following theorem of Adams [1]. For a history of results concerning this problem see [2] or [9].

**Theorem.** Let L be a connected subcomplex of an aspherical 2-complex K. Then there exists a characteristic subgroup N of  $\pi_1(L)$  such that the Galois cover  $L_N$  of L corresponding to N is acyclic.

#### Remarks.

- (1) It is obvious that if L is not aspherical then  $L_N$  cannot be contractible and hence  $N \neq 0$ .
- (2) The N in the above theorem is constructed as the intersection of all normal subgroups P of  $G = \ker \pi_1(L) \to \pi_1(K)$  such that G/P is conservative.
- (3) In [11] Strebel shows that the Galois covering of L corresponding to the perfect radical of G (i.e. the maximal perfect subgroup) is also acyclic.
- (4) Since the cellular chain complex of the cover  $L_N$  of K is acyclic, the group  $Q = \pi_1(L)/N$  and also G/N have cohomological dimension at most 2.

In the following all complexes are connected. The following exact sequence was

derived by Hopf. Let Y be a 2-dimensional CW-complex with fundamental group G, then

$$0 \to H_3(G) \to \pi_2(Y)_G \xrightarrow{h} H_2(Y) \to H_2(G) \to 0$$

is exact. This shows that the Galois cover  $Y' \to Y$  is acyclic if and only if  $P = \pi_1(Y')$  is superperfect (i.e.,  $H_1(P) = H_2(P) = 0$ ) and the Hurewicz map  $h: \pi_2(Y) \to H_2(Y')$  is zero. Hence the interest in the following result.

#### **Theorem 1.** The following are equivalent.

- (1) Let  $P \neq 1$  be a finite, superperfect normal subgroup of G and suppose Q = G/P contains an element of infinite order. Then if X is any [G, 2]-complex, the Hurewicz map  $h : \pi_2(X) \to H_2(X_P)$  is non-zero.
- (2) Let  $P \neq 1$  be a finite, superperfect normal subgroup of G and suppose Q = G/P is free and non-trivial. Then if X is any [G,2] complex the Hurewicz map  $h: \pi_2(X) \to H_2(X_P)$  is non-zero.
- (3) Suppose P is a perfect normal subgroup of G with Q = G/P free and non-trivial. Further suppose there exists a [G,2] complex X with  $H_2(X_P) = 0$ , then P = 1 or P is infinite.
- (4) Let  $X' \xrightarrow{p} X$  be a connected Galois cover with group  $\mathbb{Z}$ . Suppose X is 2-dimensional and X' is acyclic. Then either X' is contractible or  $\pi_1(X')$  is infinite.
- (5) Let  $X' \xrightarrow{p} X$  be a connected Galois cover whose group contains an element of infinite order. Suppose the dimension of X is 2 and X' is acyclic, then X' is contractible or  $\pi_1(X')$  is infinite.

*Proof.*  $(1) \Rightarrow (2)$ . Obvious since any non-trivial free group is torsion free.

- $(2) \Rightarrow (3)$ . From the exact sequence of Hopf, we see  $H_2(X_P) = 0$  implies  $H_2(P) = 0$  i.e. P is superperfect and  $h: \pi_2(X) \to H_2(X_P)$  is zero. If  $P \neq 1$  and finite this is impossible by (2).
- (3)  $\Rightarrow$  (4). If  $P = p_*(\pi_1(X'))$  then P is perfect, normal in  $G = \pi_1(X)$  and G/P is free of rank one. If X' is not contractible then  $P \neq 1$  and since  $H_2(X') = 0$ , P is infinite by (3).
- $(4) \Rightarrow (5)$ . Let X be a 2-complex,  $X' \to X$  a cover whose group  $\pi_1(X)/\pi_1(X')$  contains an element z of infinite order. Let  $H \subseteq \pi_1(X)$  be the inverse image of the subgroup generated by  $\{z\}$  under the natural projection and let  $X_H \to X$  be the Galois cover corresponding to H. Then  $X' \to X_H$  is a Galois cover with infinite cyclic group and by (4)  $\pi_1(X') = 1$  or is infinite.
- $(5) \Rightarrow (1)$ . Let X be a [G,2]-complex and  $X_P$  the Galois cover corresponding to P.  $H_1(X_P) = 0$  since P is perfect. Now  $H_2(P) = 0$  and if  $h: \pi_2(X) \to H_2(X_P)$  is also zero then  $H_2(X_P) = 0$  by the Hopf sequence. Hence  $X_P$  is acyclic and by (5) either  $X_P$  is contractible or P is infinite. But this contradicts the fact that P is finite and non-trivial. Hence h must be non-zero.

If one examines the above proof one sees that P being finite and non-trivial is

not essential. In fact we have

**Theorem 2.** Let  $\mathfrak{P}$  be a class of groups, e.g. finite, torsion, trivial, etc. The following are equivalent.

- (A) Let H be a superperfect normal subgroup of G and suppose G/H contains and element of infinite order. If H belongs to  $\mathfrak P$  and X is an arbitrary [G,2]-complex then the Hurewicz homomorphism  $h:\pi_2(X)\to H_2(X_H)$  is non-zero
- (B) If  $Y' \to Y$  is a Galois cover with group  $\mathbb{Z}$  where Y is two dimensional and Y' is acyclic then  $\pi_1(Y')$  does not belong to  $\mathfrak{P}$ .
- Proof.  $(A) \Rightarrow (B)$  Let  $Y' \to Y$  be a Galois cover with group  $\mathbb{Z}$ ,  $\dim(Y) = 2$  and Y' acyclic. Then  $H = \pi_1(Y')$  is perfect and by the Hopf sequence  $H_2(H) = 0$ , hence H is superperfect. If  $G = \pi_1(Y)$  then  $G/H \cong \mathbb{Z}$  contains an element of infinite order and so if H is in  $\mathfrak{P}$  then the Hurewicz map  $h : \pi_2(Y) \to H_2(Y')$  is non-zero. But this is ridiculous since  $H_2(Y') = 0$ .
- $(B)\Rightarrow (A)$  Let H be superperfect and normal in G and suppose G/H contains an element of infinite order. Let X be a [G,2]-complex. There exists a subgroup  $\bar{G}$ ,  $H\subseteq \bar{G}\subseteq G$  such that  $\bar{G}/H\cong \mathbb{Z}$ . If  $X_H,X_{\bar{G}}$  denote the covers of X corresponding to H,  $\bar{G}$  respectively, then  $X_H\to X_{\bar{G}}$  is a Galois cover with group  $\mathbb{Z}$ . If the Hurewicz map  $h:\pi_2(X)\to H_2(X_H)$  is zero then  $H_2(X_H)=0$  by Hopf. Hence  $X_H$  is acyclic and by (B)  $\pi_1(X_H)$  does not belong to  $\mathfrak{P}$ . Therefore the Hurewicz map must be zero.

**Corollary.** Let  $\mathfrak{P}$  be a class of groups. If there exists a subcomplex L of a 2-dimensional aspherical complex K which is not aspherical and such that the maximal perfect subgroup of  $\pi_1(L)$  belongs to  $\mathfrak{P}$  then there exists an acyclic, infinite cyclic non-contractible Galois cover  $X' \to X$  where X is 2-dimensional and  $\pi_1(X')$  belongs to  $\mathfrak{P}$ .

*Proof.* If there exists such a pair (K, L) then (A) is false for the class  $\mathfrak P$  by Strebel's result [11]. Hence (B) is false for this class, i.e. there exists a Galois cover  $X' \to X$  of 2-dimensional complexes, with group  $\mathbb Z$ , X' acyclic and  $\pi_1(X')$  in  $\mathfrak P$ .

Hence we see a positive answer to the following question would limit the maximal perfect subgroups that could appear in any example of a non-aspherical subcomplex of a two-dimensional apherical complex.

**Acyclic Cover Question.** Let  $\mathfrak{P}$  be a class of groups containing the trivial group. If X is a 2-dimensional CW-complex and X' is an acyclic, infinite cyclic Galois cover of X with  $\pi_1(X')$  in  $\mathfrak{P}$ , is X' contractible?

In [5], M.N. Dyer claimed to have proven part (2) of Theorem 1 (in fact, whenever Q has cohomological dimension 1 or 2) and hence to have answered the Acyclic Cover Question in the affirmative for the class of finite groups. However his proof is incorrect and not at all rectifible as he uses in an essential way the following lemma which is definitely false.

**Lemma([5], 3.1).** Let  $0 \to P \to G \xrightarrow{\pi} Q \to 0$  be an exact sequence of groups with P finite, then  $H^i(G,\mathbb{Z}G) \cong H^i(G,\mathbb{Z}Q) \cong H^i(Q,\mathbb{Q})$  for all i > 0. The first isomorphism is induced by  $\mathbb{Z}\pi : \mathbb{Z}G \to \mathbb{Z}Q$  and the second by  $\pi : G \to Q$ .

## Remarks.

- (1) The first error is minor and correctible. Dyer only needs these isomorphisms for i=1,2 and when P is also superperfect. Actually in view of Theorem 1 these isomorphisms are only needed for i=1 when P is perfect. In these cases the isomorphisms do exist.
- (2) The serious error lies in the second sentence. Although it is true the second isomorphism is induced by  $\pi$ , the first is induced by  $\alpha: \mathbb{Z}Q \to \mathbb{Z}G$  where  $\alpha(q) = \sum_{\pi(g)=q} g$  is the isomorphism of  $\mathbb{Z}Q$  and  $\mathbb{Z}G^P$ . Since  $\pi \cdot \alpha: \mathbb{Z}Q \to \mathbb{Z}Q$  is multiplication by |P| and by a theorem of Swan [10],  $H^1(Q,\mathbb{Z}Q) \cong H^1(G,\mathbb{Z}Q)$  is free abelian and non-zero for Q non-trivial and free, it follows  $\mathbb{Z}\pi: \mathbb{Z}G \to \mathbb{Z}Q$  can never be an isomorphism in this case.
- (3) Dyer uses that  $\mathbb{Z}\pi$  induces an isomorphism in step 3 of his proof to show a certain map is a split epimorphism which in turn leads to his contradiction.

There is some evidence for a positive answer to the Acyclic Cover Question for the class of finite groups. In fact if one one assumes that X is a finite, 2-dimensional CW-complex then there exists no infinite cyclic acyclic covers of X with finite fundamental group except for contractible ones. This is based on the following theorem of Eckmann [6] which in turn is an extension of a result of Cheegar and Gromov [3]. This result is a very sophisticated version of an theorem by Gottlieb [7] concerning centers and Euler characteristics. We need a definition.

**Definition.** A group A is called amenable if every action of A on a compact metric space has an A-invariant Borel measure.

### Remarks.

- (1) The class of amenable groups is closed under subgroups, quotient groups, extensions and increasing unions. It also contains all finite groups and all abelian groups and hence contains the class of elementary amenable groups, that is the class generated from finte and abelian groups by the operations of subgroups, quotient groups, extensions and increasing unions.
- (2) R. Grigorchuck [8] has produced examples of finitely generated amenable groups which are not elementary amenable. These are finitely generated

- subgroups of the group of based automorphisms of the based infinite triadic tree (3 edges at each vertex).
- (3) It is not difficult to show the free group on two generators,  $F_2$ , is not amenable and hence any group containing  $F_2$  is also non-amenable.

**Theorem [6].** Let X be a finite, connected complex of dimension m. Suppose  $\pi_i(X) = 0$  for 1 < i < m and  $\pi_1(X)$  is an infinite amenable group. Then  $(-1)^m \chi(X) \ge 0$  and  $\chi(X) = 0$  if and only if X is aspherical.

Using this result we obtain the following evidence for a positive answer to the Acyclic Cover Question for the class of finite groups.

**Theorem 3.** Let X be 2-dimensional and finite. Suppose X' is a Galois acyclic cover of X with group  $\mathbb{Z}$ . If  $\pi_1(X')$  is amenable then X' is a K(P,1) and hence if P is finite it must be trivial and X' contractible.

Proof. Since  $\pi_1(X')$  and  $\mathbb{Z}$  are amenable so is  $G = \pi_1(X)$  and it is clearly infinite since it maps onto  $\mathbb{Z}$ . Since X' is acyclic the spectral sequence of the Galois cover  $X' \to X$  shows X has the (integral) homology of a circle and hence of Euler chacteristic zero. By Eckmann's result X is aspherical and hence so is X'. If P is non-trivial and finite it has infinite cohomological dimension, so if P is finite it must be trivial.

#### References

- J. F. Adams, A new proof of a theorem of W.H.Cockcroft, J. London Math. Soc. 30 (1955), 482–488.
- [2] W. Bogley, J.H.C. Whitehead's asphericity question. In: Two-dimensional Homotopy and Combinatorial Group Theory (C. Hog-Angeloni, W. Metzler and A.J. Sieradski, eds), London Math. Society Lecture Note Series 197 Cambridge Univ. Press, 1993, 309-334.
- [3] J. Cheeger and M. Gromov, L<sub>2</sub>-cohomology and group cohomology, Topology 25 (1986), 189–215.
- [4] W. H. Cockcroft, On aspherical two-dimensional complexes, Proc. London Math. Soc. 4(3) (1954), 375–384.
- [5] M. N. Dyer, Groups with no infinite perfect subgroups and aspherical 2-complexes, Comm. Math. Helvetici 68 (1993), 333–339.
- [6] B. Eckmann, Amenable groups and Euler characteristic, Comm. Math. Helvetici 67 (1992), 383–393.
- [7] D. H. Gottlieb, A certain subgroup of the fundamental group, Amer. J. Math. 87 (1965), 840–856
- [8] R. Grigorchuk, Degrees of growth of finitely generated groups and the theory of invariant means, Math. USSR-Izv. 25 (1985), 259-300.
- [9] J. A. Schafer, The Whitehead Problem, Arhus University Preprint Series 13 (1994).
- [10] R. Swan, Groups of cohomological dimension one, Journal of Algebra 12 (1969), 585–601.
- [11] R. Strebel, Homological methods applied to the derived series of groups, Comment. Math.

Helv. 49 (1974), 302–322.
[12] J. H. C. Whitehead, On addding relations to homotopy groups, Annals of Math. 42 (1941), 409–428.

James A. Schafer Department of Mathematics University of Maryland College Park, MD 20742 USA

(Received: April 24, 1995)