

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 74 (1999)

Artikel: Preprojective algebras, differential operators and a Conze embedding for deformations of Kleinian singularities
Autor: Crawley-Boevey, William
DOI: <https://doi.org/10.5169/seals-55801>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Preprojective algebras, differential operators and a Conze embedding for deformations of Kleinian singularities

William Crawley-Boevey

Abstract. For any associative algebra A over a field K we define a family of algebras $\Pi^\lambda(A)$ for $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$. In case A is the path algebra of a quiver, one recovers the deformed preprojective algebra introduced by M. P. Holland and the author. In case A is the coordinate ring of a smooth curve, the family includes the ring of differential operators for A and the coordinate ring of the cotangent bundle for $\text{Spec } A$. In case A is quasi-free and $\Omega^1 A$ is a finitely generated A - A -bimodule we prove that $\Pi^\lambda(A)$ is well-behaved under localization. We use this to prove a Conze embedding for deformations of Kleinian singularities.

Mathematics Subject Classification (1991). Primary 16G20; Secondary 16S32.

Keywords. Preprojective algebra, quiver, quasi-free algebra, pseudoflat epimorphism, differential operator, Kleinian singularity, Conze embedding.

If K is an algebraically closed field of characteristic zero and Γ is a non-trivial finite subgroup of $\text{SL}_2(K)$ then the coordinate ring of the Kleinian singularity K^2/Γ has a family of deformations \mathcal{O}^λ where $\lambda \in Z(K\Gamma)$. They have been defined and studied in work of M. P. Holland and the author [5]. If λ has trace zero on the regular representation of Γ , then \mathcal{O}^λ is a commutative ring, and it occurs as the coordinate ring of a fibre of the semi-universal deformation of K^2/Γ . On the other hand, if λ has nonzero trace on the regular representation, then \mathcal{O}^λ is a non-commutative ring.

In this paper we construct an embedding $\psi_\lambda : \mathcal{O}^\lambda \rightarrow C_\nu$ where ν is the trace of λ on the regular representation of Γ , and $C_\nu = K\langle x, y \mid xy - yx = \nu \rangle$. This is an embedding of noetherian domains, and we show that it induces an isomorphism of quotient division rings. In the commutative case C_ν is a polynomial ring in two variables, so the embedding is a birational map from the affine plane to a deformation of the Kleinian singularity. In the noncommutative case C_ν is isomorphic to the first Weyl algebra, and the embedding is reminiscent of one constructed by N. Conze [3]. We therefore call ψ_λ a ‘Conze embedding’.

In the work of M. P. Holland and the author, the key idea for studying deformations of Kleinian singularities was to relate them to a new class of algebras which we introduced, the ‘deformed preprojective algebras’ associated to quivers of extended Dynkin type. In fact, in our earliest work we constructed Conze

embeddings. By dividing into cases according to the different types of extended Dynkin quivers, we constructed representations of the deformed preprojective algebras over C_ν . These representations induce maps $\mathcal{O}^\lambda \rightarrow C_\nu$, and we used computer calculations to prove that these maps are injective. Instead of publishing our work, we decided to wait for a better understanding of deformed preprojective algebras, and a natural proof of the existence of Conze embeddings. This paper is the result. Although M. P. Holland is not explicitly an author of this paper, he has contributed a great deal to it.

Let K be an arbitrary field. For any K -algebra A (associative, with 1), and any element $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ we define an algebra $\Pi^\lambda(A)$. It is equipped with a homomorphism $A \rightarrow \Pi^\lambda(A)$. Note that an element $\nu \in K$ determines the element $\nu \otimes [A] \in K \otimes_{\mathbb{Z}} K_0(A)$, and for simplicity we write $\Pi^\nu(A)$ rather than $\Pi^{\nu \otimes [A]}(A)$.

For finite-dimensional hereditary algebras this definition generalizes the preprojective algebras of Baer, Geigle and Lenzing [2], which we denote here by $\Pi_{\text{BGL}}(A)$.

Theorem 0.1. *If A is finite-dimensional and hereditary then $\Pi^0(A) \cong \Pi_{\text{BGL}}(A)$.*

Our next result relates the new definition to the original deformed preprojective algebras of [5], which we denote here by $\Pi_{\text{CBH}}^\lambda(Q)$, where Q is a quiver with vertex set I and $\lambda \in K^I$. We identify K^I with $K \otimes_{\mathbb{Z}} K_0(KQ)$, with $\lambda \in K^I$ corresponding to the element

$$\lambda = \sum_{i \in I} \lambda_i \otimes [KQe_i] \in K \otimes_{\mathbb{Z}} K_0(KQ)$$

(where e_i is the trivial path at vertex i).

Theorem 0.2. *If Q is a quiver and $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$ then $\Pi^\lambda(KQ) \cong \Pi_{\text{CBH}}^\lambda(Q)$.*

Theorems 0.1 and 0.2 together imply that $\Pi_{\text{BGL}}(KQ) \cong \Pi_{\text{CBH}}^0(Q)$. This has been known for some time—it was explained to the author by C. M. Ringel—and it was used implicitly in [5]. An alternative proof has recently been written up by Ringel [13].

Theorem 0.2 includes as a special case the fact that for a polynomial ring in one variable, $\Pi^\nu(K[x]) \cong C_\nu$. It is this isomorphism which leads to the appearance of C_ν in the Conze embedding. When K has characteristic zero, the algebra $\Pi^1(K[x])$ is the ring of differential operators for $K[x]$. This turns out to be no coincidence. Indeed we prove the following result.

Theorem 0.3. *If K is a field of characteristic zero and A is the coordinate ring of a smooth affine curve over K , then $\Pi^0(A)$ is the coordinate ring of the cotangent bundle of $\text{Spec } A$ and $\Pi^1(A)$ is the ring of differential operators for A .*

The first three theorems deal with examples of $\Pi^\lambda(A)$. We now turn to the functorial properties of $\Pi^\lambda(A)$. In Section 5 we prove the following result.

Theorem 0.4. *Suppose that e is an idempotent in an algebra A with $AeA = A$. If $\lambda \in K \otimes_{\mathbb{Z}} K_0(eAe)$, then $\Pi^\lambda(eAe) \cong e(\Pi^\mu(A))e$, where $\mu \in K \otimes_{\mathbb{Z}} K_0(A)$ corresponds to λ under the natural isomorphism $K_0(eAe) \cong K_0(A)$.*

It follows that the algebras $\Pi^\lambda(A)$ are well-behaved under passage to matrix rings and under Morita equivalence. For example if A and B are Morita-equivalent algebras, and $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ corresponds to $\mu \in K \otimes_{\mathbb{Z}} K_0(B)$, under the isomorphism $K_0(A) \cong K_0(B)$, then $\Pi^\lambda(A)$ and $\Pi^\mu(B)$ are Morita equivalent. In the next section we show that the algebras $\Pi^\lambda(A)$ are also well-behaved under direct products.

According to Cuntz and Quillen [6], an algebra A is *quasi-free* if the kernel $\Omega^1 A$ of the multiplication map $A \otimes A \rightarrow A$ is a projective A - A -bimodule. Such algebras are to be considered as coordinate rings of noncommutative manifolds. We say that A is *bimodule-finite* if $\Omega^1 A$ is a finitely generated bimodule. The following result is perhaps already known to experts.

Proposition 0.5. *An algebra A is bimodule-finite if and only if it has a finitely generated subalgebra C , such that the inclusion $C \rightarrow A$ is a ring epimorphism.*

It is easy to see that path algebras are both quasi-free and bimodule-finite. Now any quasi-free algebra is hereditary, and over the complex numbers Cuntz and Quillen observed that every finite-dimensional hereditary algebra is Morita equivalent to a path algebra, so is quasi-free. In the general case, however, the situation is slightly more complicated.

Proposition 0.6. *A finite-dimensional algebra A is quasi-free if and only if it is hereditary and $A/\text{rad } A$ is separable over K .*

Our real reason for working in an abstract setting is in order to prove a localization theorem. For any ring homomorphism $\theta : A \rightarrow B$, the functor $B \otimes_A -$ defines a homomorphism $\theta_* : K \otimes_{\mathbb{Z}} K_0(A) \rightarrow K \otimes_{\mathbb{Z}} K_0(B)$. Following [1, §5], a ring epimorphism $A \rightarrow B$ is said to be *pseudoflat* provided that $\text{Tor}_1^A(B, B) = 0$. See [15, Theorem 4.8] for a number of equivalent conditions.

Theorem 0.7. *If $\theta : A \rightarrow B$ is a pseudoflat epimorphism and $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ then there is a natural map $\Pi^\lambda(A) \rightarrow \Pi^{\theta_*(\lambda)}(B)$. If A is a quasi-free bimodule-finite algebra, then so is B , and the diagram*

$$\begin{array}{ccc} A & \longrightarrow & \Pi^\lambda(A) \\ \downarrow & & \downarrow \\ B & \longrightarrow & \Pi^{\theta_*(\lambda)}(B) \end{array}$$

is a pushout in the category of rings.

Now suppose that Q is an extended Dynkin quiver with vertex set I , and for simplicity suppose that K is an algebraically closed field. Let $\delta \in \mathbb{Z}^I$ be the minimal positive imaginary root for Q . By using universal localization one can easily construct a pseudoflat epimorphism $\theta : A \rightarrow M_N(K[x])$ such that the general representation of KQ of dimension vector δ is the restriction of a $M_N(K[x])$ -module. (It follows that $N = \sum_i \delta_i$.) If $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$, by the theorem there is an induced a pseudoflat epimorphism θ_λ from $\Pi^\lambda(KQ)$ to

$$\Pi^{\theta_*(\lambda)}(M_N(K[x])) \cong M_N(\Pi^\nu(K[x])) \cong M_N(C_\nu)$$

where in fact $\nu = \sum_i \lambda_i \delta_i$. By using the representation theory of Q we prove the following result.

Theorem 0.8. *If Q is an extended Dynkin quiver, K is an algebraically closed field and $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$ then $\Pi^\lambda(KQ)$ is a prime noetherian ring of Gelfand-Kirillov dimension 2. Moreover $\theta_\lambda : \Pi^\lambda(KQ) \rightarrow M_N(C_\nu)$ is injective, and it induces an isomorphism on simple artinian quotient rings.*

In an appendix we use the methods of this paper to study the variety of representations of $\Pi^0(KQ)$ of dimension δ .

We finally return to Kleinian singularities. Assume that K is algebraically closed of characteristic zero, and let Γ be a finite subgroup of $SL_2(K)$. Let Q be an orientation of the McKay quiver of Γ , and let 0 be an extending vertex. Recall that there is an isomorphism $\mathcal{O}^\lambda \cong e_0 \Pi^\lambda(KQ) e_0$, where $\lambda \in Z(K\Gamma)$ is identified with $\lambda \in K^I$ by letting λ_i be the trace of λ on the i th irreducible representation of Γ . Here is the result mentioned at the start of the introduction.

Theorem 0.9. *There is an embedding $\psi_\lambda : \mathcal{O}^\lambda \rightarrow C_\nu$ where ν is the trace of λ on the regular representation of Γ . Moreover ψ_λ induces an isomorphism on quotient division rings.*

1. Definition of $\Pi^\lambda(A)$

Let A be an algebra (associative, with 1) over a field K . Recall that A - A -bimodules are the same as A^e -modules, where $A^e = A \otimes A^{op}$. (Unadorned tensor products are always over the field K .) The *universal derivation bimodule*, or *bimodule of noncommutative differential 1-forms* is the kernel $\Omega^1 A$ of the multiplication map $A \otimes A \rightarrow A$. See for example [1] or [6].

If M is an A - A -bimodule, we write $\text{Der}(A, M)$ for the space of derivations from A to M . It is isomorphic to $\text{Hom}_{A^e}(\Omega^1 A, M)$, a homomorphism θ giving rise to the derivation d with $d(a) = \theta(a \otimes 1 - 1 \otimes a)$.

The space $\text{Der}(A, A \otimes A)$ becomes an A - A -bimodule via $adb = (r_b \otimes \ell_a)d$ where $\ell_a, r_b : A \rightarrow A$ denote left multiplication by a and right multiplication

by b respectively. We write Δ (or Δ_A) for the derivation $A \rightarrow A \otimes_K A$ with $\Delta(a) = a \otimes 1 - 1 \otimes a$.

If M is an A - A -bimodule we write $T_A M$ for the tensor algebra of M over A . For any $a \in A$ we define $\Pi^a(A) = T_A \text{Der}(A, A \otimes A) / (\Delta - a)$. We consider it as an A -ring, that is, as an algebra equipped with homomorphism $A \rightarrow \Pi^a(A)$. We have the following elementary results.

Lemma 1.1. *If u is a unit in the centre of A and $a \in A$, then $\Pi^a(A) \cong \Pi^{ua}(A)$.*

Proof. Multiplication on the left by u^{-1} induces an automorphism of the bimodule $\text{Der}(A, A \otimes A)$, and hence there is an algebra automorphism of $T_A \text{Der}(A, A \otimes A)$ which is the identity on A , but sends Δ to $u^{-1}\Delta$. Under this automorphism the ideal $(\Delta - a)$ is sent to $(u^{-1}\Delta - a) = (\Delta - ua)$. \square

Lemma 1.2. *Up to isomorphism $\Pi^a(A)$ depends only on the image of a in $H_0(A) = A/[A, A]$.*

Proof. Write $M = \text{Der}(A, A \otimes A)$. Given $b, c \in A$, it suffices to construct an automorphism of $T_A M$ sending $\Delta - a$ to $\Delta - a + [b, c]$. By the universal property of tensor algebras, the homomorphisms $\theta : T_A M \rightarrow T_A M$ which are the identity on A are in 1-1 correspondence with A - A -bimodule maps $M \rightarrow T_A M$. Taking the bimodule map sending d to $d + \sum_i y_i c x_i$, where x_i and y_i are defined by $d(b) = \sum_i x_i \otimes y_i$, it is clear that the resulting homomorphism θ is an automorphism, and that $\theta(\Delta - a) = \Delta - a + [b, c]$, as required. \square

Let $\text{tr} : K_0(A) \rightarrow H_0(A) = A/[A, A]$ be the trace map, sending the class of a projective module P to the image in $A/[A, A]$ of the trace of any idempotent $e \in M_n(A)$ with image isomorphic to P . This map extends to a linear map $K \otimes_{\mathbb{Z}} K_0(A) \rightarrow H_0(A)$ also denoted tr .

Definition 1.3. If $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ then

$$\Pi^\lambda(A) = \Pi^{a_\lambda}(A) = T_A \text{Der}(A, A \otimes A) / (\Delta - a_\lambda)$$

where a_λ is any lift of $\text{tr}(\lambda)$ to A .

The previous lemma shows that $\Pi^\lambda(A)$ does not depend on the choice of a_λ . If $a, b \in A$ then $a\Delta b$ is the derivation with $(a\Delta b)(x) = xb \otimes a - b \otimes ax$ for $x \in A$. Thus the A - A -sub-bimodule of $\text{Der}(A, A \otimes A)$ generated by Δ is the set of inner derivations. Therefore $\text{Der}(A, A \otimes A) / A\Delta A \cong H^1(A, A \otimes A)$, which has the following consequence.

Lemma 1.4. $\Pi^0 A \cong T_A H^1(A, A \otimes A)$.

Any tensor algebra $T_A M$ is naturally graded, with $(T_A M)_n$ being the n th

tensor power of M . Thus the lemma gives a grading of $\Pi^0(A)$. On the other hand the algebra $\Pi^a(A)$ is filtered

$$\Pi^a(A)_{\leq 0} \subseteq \Pi^a(A)_{\leq 1} \subseteq \Pi^a(A)_{\leq 2} \subseteq \dots,$$

where $\Pi^a(A)_{\leq n}$ is the image in $\Pi^a(A)$ of $\sum_{i=0}^n (T_A \operatorname{Der}(A, A \otimes A))_i$. As usual one can consider the associated graded algebra $\operatorname{gr} \Pi^a(A)$, and there is the following result.

Lemma 1.5. *There is a natural surjective graded algebra map $\Pi^0(A) \rightarrow \operatorname{gr} \Pi^a(A)$.*

Proof. An algebra homomorphism $\phi : \Pi^0(A) \rightarrow \operatorname{gr} \Pi^a(A)$ is determined by an algebra homomorphism $A \rightarrow \Pi^a(A)_{\leq 0}$, which we take to be the natural map, and an A - A -bimodule map $g : H^1(A, A \otimes A) \rightarrow (\operatorname{gr} \Pi^a(A))_1$. For ϕ to be surjective, we need g to be surjective, so it suffices to show that $(\operatorname{gr} \Pi^a(A))_1$ is naturally a quotient of $H^1(A, A \otimes A)$. Now by definition $(\operatorname{gr} \Pi^a(A))_1 = (A \oplus \operatorname{Der}(A, A \otimes A))/S$, where $S = A + I \cap (A \oplus \operatorname{Der}(A, A \otimes A))$ and I is the ideal $(\Delta - a)$ in $T_A \operatorname{Der}(A, A \otimes A)$. The result follows since S contains $A \oplus A \Delta A$. \square

2. Finite-dimensional hereditary algebras

Let D be the duality $\operatorname{Hom}_K(-, K)$.

Lemma 2.1. *For any algebra A there is a natural isomorphism $\operatorname{Ext}_A^1(DY, X) \cong H^1(A, X \otimes Y)$ for X and Y left A -modules, with Y finite-dimensional.*

Proof. Tensoring the defining sequence for $\Omega^1 A$ with DY one obtains an exact sequence

$$0 \rightarrow \Omega^1 A \otimes_A DY \rightarrow A \otimes DY \rightarrow DY \rightarrow 0.$$

Since $A \otimes DY$ is projective, if X is a left A -module one obtains

$$\dots \rightarrow \operatorname{Hom}_A(A \otimes DY, X) \rightarrow \operatorname{Hom}_A(\Omega^1 A \otimes_A DY, X) \rightarrow \operatorname{Ext}_A^1(DY, X) \rightarrow 0.$$

Now $\operatorname{Hom}_A(\Omega^1 A \otimes_A DY, X)$ can be identified with $\operatorname{Hom}_{A^e}(\Omega^1 A, X \otimes Y)$, so with $\operatorname{Der}(A, X \otimes Y)$. Also we can identify $\operatorname{Hom}_A(A \otimes DY, X)$ with $Y \otimes X$, and then the map $Y \otimes X \rightarrow \operatorname{Der}(A, X \otimes Y)$ sends an element of $Y \otimes X$ to the corresponding inner derivation. \square

Lemma 2.2. *If A is a finite-dimensional hereditary algebra then there is a natural isomorphism $\operatorname{Ext}_A^1(DA, X) \cong H^1(A, A \otimes A) \otimes_A X$ for X a left A -module.*

Proof. Since A is hereditary the functor $\text{Ext}^1(DA, -)$ is right exact. Since it also commutes with direct limits it is naturally isomorphic to the tensor product functor $\text{Ext}^1(DA, A) \otimes_A -$. \square

If A is a finite-dimensional hereditary algebra then the *inverse Auslander-Reiten translation* is the functor τ^- defined by $\tau^-(X) = \text{Ext}^1(DA, X)$. The algebra $\Pi_{\text{BGL}}(A)$ of [2, §3] is

$$\bigoplus_{n=0}^{\infty} \text{Hom}_A(A, \tau^{-n}(A)),$$

with the product defined by $uv = \tau^{-m}(u) \circ v$ for elements $u \in \text{Hom}_A(A, \tau^{-n}(A))$ and $v \in \text{Hom}_A(A, \tau^{-m}(A))$.

Theorem 2.3. *If A is finite-dimensional and hereditary then $\Pi^0(A) \cong \Pi_{\text{BGL}}(A)$.*

Proof. Combine Lemmas 1.4 and 2.2 with [2, Proposition 3.1]. \square

3. Deformed preprojective algebras

Let Q be a quiver with vertex set I and let KQ be its path algebra. Let $e_i \in KQ$ be the trivial path at vertex i . Recall that K^I is identified with $K \otimes_{\mathbb{Z}} K_0(KQ)$, with $\lambda \in K^I$ corresponding to the element

$$\lambda = \sum_{i \in I} \lambda_i \otimes [KQ e_i] \in K \otimes_{\mathbb{Z}} K_0(KQ).$$

Let \overline{Q} be the quiver obtained from Q by adjoining an arrow $a^* : j \rightarrow i$ for each arrow $a : i \rightarrow j$ in Q . The next result shows that $\Pi^\lambda(KQ)$ coincides with the deformed preprojective algebra $\Pi_{\text{CBH}}^\lambda(Q)$ of [5].

Theorem 3.1. *If $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$ then $\Pi^\lambda(KQ) \cong K\overline{Q}/J$ where J is the ideal generated by $\sum_{a \in Q} [a, a^*] - \sum_{i \in I} \lambda_i e_i$.*

Proof. There is a standard projective resolution of KQ as a KQ - KQ -bimodule

$$0 \rightarrow \bigoplus_{a: i \rightarrow j \text{ in } Q} KQ e_j \otimes e_i KQ \xrightarrow{f} \bigoplus_{i \in I} KQ e_i \otimes e_i KQ \xrightarrow{\text{mult}} KQ \rightarrow 0$$

where f sends $(e_j \otimes e_i)_a$ to $(e_j \otimes a)_j - (a \otimes e_i)_i$. Identifying

$$KQ \otimes KQ = \bigoplus_{r, s \in I} KQ e_r \otimes e_s KQ,$$

one obtains an exact sequence

$$0 \rightarrow \bigoplus_{a:i \rightarrow j} KQe_j \otimes e_i KQ \oplus \bigoplus_{r \neq s} KQe_r \otimes e_s KQ \xrightarrow{g} KQ \otimes KQ \xrightarrow{\text{mult}} KQ \rightarrow 0, \quad (\dagger)$$

where g sends $(e_j \otimes e_i)_a$ to $e_j \otimes a - a \otimes e_i$ and $(e_r \otimes e_s)_{rs}$ to $e_r \otimes e_s$.

If M is a KQ - KQ -bimodule, then $\text{Hom}_{KQ^e}(KQe_i \otimes e_j KQ, M) \cong e_i M e_j$, so if $M = KQ \otimes KQ$ the Hom space is isomorphic to $e_i KQ \otimes KQ e_j$. Now this tensor product is over K , so one can swap the order of the terms and write $KQe_j \otimes e_i KQ$. By doing this, one clearly sees the KQ - KQ -bimodule structure of this space.

Now the left hand term of the exact sequence (\dagger) is $\Omega^1 KQ$, and computing its homomorphisms to $KQ \otimes KQ$ one obtains

$$\text{Der}(KQ, KQ \otimes KQ) \cong \bigoplus_{a:i \rightarrow j} KQe_i \otimes e_j KQ \oplus \bigoplus_{r \neq s} KQe_s \otimes e_r KQ.$$

Thus $T_{KQ} \text{Der}(KQ, KQ \otimes KQ)$ is identified with the path algebra of the quiver \tilde{Q} obtained from \tilde{Q} by adding an arrow $c_{rs} : r \rightarrow s$ for each pair of vertices $r \neq s$. Also Δ corresponds to the map g , so to the element

$$\sum_{a:i \rightarrow j} (a \otimes e_j - e_i \otimes a)_a + \sum_{r \neq s} (e_s \otimes e_r)_{rs}.$$

Now with the identification of K^I and $K \otimes_{\mathbb{Z}} K_0(KQ)$, the element $\sum_i \lambda_i e_i$ is a lift to KQ of $\text{tr}(\lambda) \in KQ/[KQ, KQ]$. Thus $\Pi^\lambda(KQ) \cong K\tilde{Q}/(\xi)$ where

$$\xi = \sum_{a:i \rightarrow j} (aa^* - a^*a) + \sum_{r \neq s} c_{rs} - \sum_i \lambda_i e_i.$$

Clearly the ideal generated by ξ is also generated by the elements $\sum_i e_i \xi e_i$ with $i \in I$ and $e_s \xi e_r$ with $r \neq s$. These are $\sum_{a \in Q} [a, a^*] - \sum_{i=1}^n \lambda_i e_i$ and c_{rs} respectively. The result follows. \square

The path algebra of the quiver with one vertex and one loop is a polynomial ring $K[x]$, so we have the following special case.

Corollary 3.2. *If $\nu \in K$ then $\Pi^\nu(K[x]) \cong K\langle x, y \mid xy - yx = \nu \rangle$.*

4. Differential operators on curves

Throughout this section A is the coordinate ring of a smooth affine curve. Thus it is a commutative domain, finitely generated over K , of Krull dimension 1, and it is smooth over K , so for any commutative K -algebra C and any ideal I in

C with $I^2 = 0$, any K -algebra homomorphism $A \rightarrow C/I$ lifts to a K -algebra homomorphism $A \rightarrow C$. A suitable reference for smooth algebras is [10, §§25–30]. The following result is due to Schelter [14] (at least when K is algebraically closed).

Lemma 4.1. *$\Omega^1 A$ is a rank 1 projective A^e -module. Thus A is quasi-free.*

Proof. Since A is smooth over K , so is A^e , so for any maximal ideal \mathfrak{m} of A^e the localization $(A^e)_{\mathfrak{m}}$ is regular of dimension 2. Localizing the defining sequence for $\Omega^1 A$ at \mathfrak{m} gives an exact sequence

$$0 \rightarrow (\Omega^1 A)_{\mathfrak{m}} \rightarrow (A^e)_{\mathfrak{m}} \rightarrow A \otimes_{A^e} (A^e)_{\mathfrak{m}} \rightarrow 0,$$

and it suffices to prove that the ideal $(\Omega^1 A)_{\mathfrak{m}}$ of $(A^e)_{\mathfrak{m}}$ is projective. This is certainly true if \mathfrak{m} doesn't contain $\Omega^1 A$, for then some element of $\Omega^1 A$ is invertible in $(A^e)_{\mathfrak{m}}$, so $(\Omega^1 A)_{\mathfrak{m}} = (A^e)_{\mathfrak{m}}$. Thus suppose that \mathfrak{m} contains $\Omega^1 A$, so \mathfrak{m} is the inverse image under the multiplication map $A^e \rightarrow A$ of a maximal ideal \mathfrak{n} of A . This implies that $A \otimes_{A^e} (A^e)_{\mathfrak{m}} \cong A_{\mathfrak{n}}$. Now if a is a nonzero element of \mathfrak{n} then $1 \otimes a \in \mathfrak{m}$ and since $1 \otimes a$ acts as a non zero-divisor on $A_{\mathfrak{n}}$, the $(A^e)_{\mathfrak{m}}$ -module $A_{\mathfrak{n}}$ has depth at least 1. It follows that it has projective dimension at most 1, so that $(\Omega^1 A)_{\mathfrak{m}}$ is projective. \square

Since A^e is commutative, any Hom space between A - A -bimodules is naturally an A - A -bimodule. In particular $\text{Der}(A, M) = \text{Hom}_{A^e}(\Omega^1 A, M)$ is an A - A -bimodule for any M .

Lemma 4.2. *$\text{Der}(A, \Omega^1 A)$ is a free A - A -bimodule generated by Δ .*

Proof. It is the endomorphism ring of a rank 1 projective A^e -module. \square

Applying $\text{Hom}_{A^e}(\Omega^1 A, -)$ to the defining sequence for $\Omega^1 A$, we obtain a sequence

$$0 \rightarrow \text{Der}(A, \Omega^1 A) \rightarrow \text{Der}(A, A \otimes A) \rightarrow \text{Der}(A, A) \rightarrow 0$$

so by the previous lemma $H^1(A, A \otimes A)$ is naturally isomorphic to $\text{Der}(A, A)$. Now $\text{Der}(A, A)$ is a rank 1 projective A -module, so its tensor algebra over A coincides with its symmetric algebra, and we have the following result.

Theorem 4.3. *$\Pi^0(A)$ is isomorphic to the coordinate ring $K[\mathbb{T}^* \text{Spec } A]$ of the cotangent bundle of $\text{Spec } A$.*

We now investigate the relationship between $\Pi^1(A)$ and differential operators. Let $D(A)$ be the ring of differential operators for A . Elements of A are identified with the differential operators of order 0, with $a \in A$ corresponding to the homothety $a \text{Id}_A$. We write $D_{\leq 1}(A)$ for the set of differential operators of order ≤ 1 . It

becomes an A - A -bimodule by composition of maps, and there is an exact sequence of A - A -bimodules

$$0 \longrightarrow A \longrightarrow D_{\leq 1}(A) \xrightarrow{g} \operatorname{Der}(A, A) \longrightarrow 0$$

where g sends a differential operator d to $d - d(1)\operatorname{Id}_A$. Note that $\operatorname{Der}(A, A)$ is a subspace of $D_{\leq 1}(A)$ complementary to $A\operatorname{Id}_A$, but it is not a sub-bimodule.

Lemma 4.4. *The evaluation map $A \otimes \operatorname{Der}(A, A) \rightarrow A$ is surjective.*

Proof. If $a \in A$ and $h \in \operatorname{Hom}_{A^e}(\Omega^1 A, A)$ then since A is commutative, the assignment $d(x) = h(ax \otimes 1 - a \otimes x)$ defines a derivation $A \rightarrow A$. Now the assertion follows since firstly $\Omega^1 A$ is generated as a left A -module by elements of the form $x \otimes 1 - 1 \otimes x$, and secondly the natural map

$$\Omega^1 A \otimes_{A^e} \operatorname{Hom}_{A^e}(\Omega^1 A, A) \rightarrow A$$

is an isomorphism (since $\Omega^1 A$ is a rank 1 projective A^e -module). \square

Lemma 4.5. *$A\operatorname{Id}_A$ is a superfluous A - A -sub-bimodule of $D_{\leq 1}(A)$. That is, if M is an A - A -sub-bimodule of $D_{\leq 1}(A)$ and $M + A\operatorname{Id}_A = D_{\leq 1}(A)$, then $M = D_{\leq 1}(A)$.*

Proof. By assumption any derivation $d \in \operatorname{Der}(A, A)$ can be written as the sum of an element $m \in M$ and a homothety $a\operatorname{Id}_A$. Now if $x \in A$ then $d(x)y = d(xy) - xd(y)$ for $y \in A$, so

$$d(x)\operatorname{Id}_A = d \circ (x\operatorname{Id}_A) - (x\operatorname{Id}_A) \circ d = m \circ (x\operatorname{Id}_A) - (x\operatorname{Id}_A) \circ m \in M.$$

Thus $A\operatorname{Id}_A \subseteq M$ by the previous lemma. \square

Lemma 4.6. *There are surjective bimodule maps θ, ϕ giving rise to a commutative diagram with exact rows*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \operatorname{Der}(A, \Omega^1 A) & \longrightarrow & \operatorname{Der}(A, A \otimes A) & \longrightarrow & \operatorname{Der}(A, A) \longrightarrow 0 \\ & & \downarrow \phi & & \downarrow \theta & & \parallel \\ 0 & \longrightarrow & A & \longrightarrow & D_{\leq 1}(A) & \xrightarrow{g} & \operatorname{Der}(A, A) \longrightarrow 0 \end{array}$$

Proof. Since $\operatorname{Der}(A, A \otimes A)$ is a projective bimodule, the map from $\operatorname{Der}(A, A \otimes A)$ to $\operatorname{Der}(A, A)$ lifts to a map θ . Now θ is surjective since $A\operatorname{Id}_A$ is superfluous in $D_{\leq 1}(A)$. The result follows. \square

Theorem 4.7. *If K has characteristic zero and A is the coordinate ring of a smooth affine curve, then $\Pi^1(A)$ is isomorphic to the ring $D(A)$ of differential operators for A .*

Proof. Since the map ϕ in Lemma 4.6 is surjective and $\text{Der}(A, \Omega^1 A)$ is generated by Δ , the element $u = \phi(\Delta)$ must be a unit in A . Now θ induces a homomorphism $\psi : \Pi^u(A) \rightarrow D(A)$ of filtered rings, and there is a commutative square

$$\begin{array}{ccc} \Pi^0(A) & \longrightarrow & K[\mathbb{T}^* \text{Spec } A] \\ \downarrow & & \downarrow \\ \text{gr } \Pi^u(A) & \xrightarrow{\text{gr } \psi} & \text{gr } D(A) \end{array}$$

where the top map is the isomorphism of Theorem 4.3, the left hand vertical map is as in Lemma 1.5, and the right hand vertical map is the natural isomorphism resulting from the fact that A is smooth and K has characteristic zero. It follows that $\text{gr } \psi$ is an isomorphism, and then the result follows since $\Pi^1(A) \cong \Pi^u(A)$ by Lemma 1.1. \square

5. Morita equivalence

In this section A is an algebra and $e \in A$ is an idempotent with $AeA = A$, so that A is Morita equivalent to eAe . We define $f = 1 - e$.

Lemma 5.1. *If M is an A - A -bimodule, then the assignment sending d to the map $a \mapsto e d(a) e$ induces a surjective linear map $\text{Der}(A, M) \rightarrow \text{Der}(eAe, eMe)$ whose kernel is the set of inner derivations $a \mapsto am - ma$ with $m \in fMe \oplus eMf \oplus fMf$.*

Proof. Let $X = (Af \otimes eA) \oplus (Ae \otimes fA) \oplus (Af \otimes fA)$. Since $AeA = A$, it follows that $A \otimes A = \Omega^1 A + Ae \otimes eA$. Thus the projection $A \otimes A \rightarrow X$ induces an exact sequence

$$0 \rightarrow \Omega^1 A \cap Ae \otimes eA \rightarrow \Omega^1 A \rightarrow X \rightarrow 0.$$

This is split exact since X is a projective bimodule, so it induces an exact sequence

$$0 \rightarrow \text{Hom}_{Ae}(X, M) \xrightarrow{\theta} \text{Hom}_{Ae}(\Omega^1 A, M) \xrightarrow{\phi} \text{Hom}_{Ae}(\Omega^1(A) \cap Ae \otimes eA, M) \rightarrow 0.$$

Now $e(\Omega^1(A) \cap Ae \otimes eA)e = \Omega^1(eAe)$, so by Morita equivalence the last term can be identified with the space of eAe - eAe -bimodule homomorphisms from $\Omega^1(eAe)$ to eMe , and hence with $\text{Der}(eAe, eMe)$. Identifying also the middle term with $\text{Der}(A, M)$, the map ϕ is as stated. Now the left hand term is isomorphic to $fMe \oplus eMf \oplus fMf$, and the map θ sends an element m of this direct sum to the inner derivation $a \mapsto am - ma$, as required. \square

Lemma 5.2. *If J is the A - A -sub-bimodule of $\text{Der}(A, A \otimes A)$ generated by $e\Delta f$, $f\Delta e$ and $f\Delta f$, and $L = \text{Der}(A, A \otimes A)/J$, then there is an isomorphism of eAe - eAe -bimodules*

$$eLe \rightarrow \text{Der}(eAe, eAe \otimes eAe)$$

sending $e\overline{\Delta}e$ to Δ_{eAe} .

Proof. One can identify $e\operatorname{Der}(A, A \otimes A)e$ with $\operatorname{Der}(A, Ae \otimes eA)$, and then the lemma follows from the previous one. \square

There are inverse isomorphisms between $K_0(eAe)$ and $K_0(A)$ induced by the functors $P \mapsto Ae \otimes_{eAe} P$ and $Q \mapsto eQ$ on projective modules. This enables one to identify $K \otimes_{\mathbb{Z}} K_0(eAe)$ and $K \otimes_{\mathbb{Z}} K_0(A)$.

Theorem 5.3. *If $\lambda \in K \otimes_{\mathbb{Z}} K_0(eAe)$ then $\Pi^\lambda(eAe) \cong e(\Pi^\mu(A))e$, where μ is the corresponding element of $K \otimes_{\mathbb{Z}} K_0(A)$.*

Proof. It suffices to show that $\Pi^a(eAe) \cong e(\Pi^a A)e$ for $a \in eAe$, as there is a commutative square

$$\begin{array}{ccc} K \otimes_{\mathbb{Z}} K_0(eAe) & \longrightarrow & K \otimes_{\mathbb{Z}} K_0(A) \\ \operatorname{tr} \downarrow & & \operatorname{tr} \downarrow \\ eAe/[eAe, eAe] & \xrightarrow{i} & A/[A, A] \end{array}$$

where i is induced by the inclusion of eAe in A , and the top map is the bijection mentioned above. Now

$$\begin{aligned} \Pi^a(A) &= (T_A \operatorname{Der}(A, A \otimes A))/(\Delta - a) \\ &= (T_A \operatorname{Der}(A, A \otimes A))/(\Delta f, f\Delta e, f\Delta f, e\Delta e - a) \\ &\cong T_A L/(e\overline{\Delta}e - a) \end{aligned}$$

where L is as in the previous lemma. Thus

$$\begin{aligned} e\Pi^a(A)e &\cong e[T_A L/(e\overline{\Delta}e - a)]e \cong e(T_A L)e/(e\overline{\Delta}e - a) \\ &\cong T_{eAe}(eLe)/(e\overline{\Delta}e - a) \cong \Pi^a(eAe) \end{aligned}$$

by the previous lemma. \square

If A and B are Morita-equivalent algebras, then there is an equivalence from the category of left A -modules to the category of left B -modules. This induces a bijection from $K \otimes_{\mathbb{Z}} K_0(A)$ to $K \otimes_{\mathbb{Z}} K_0(B)$. In particular this applies to the algebras A and $M_n(A)$.

Corollary 5.4. *If $\lambda \in K \otimes_{\mathbb{Z}} K_0(M_n(A))$ then $\Pi^\lambda(M_n(A)) \cong M_n(\Pi^\mu(A))$ where μ is the corresponding element of $K \otimes_{\mathbb{Z}} K_0(A)$.*

Proof. The matrix units e_{ij} ($1 \leq i, j \leq n$) for $M_n(A)$ induce matrix units in $\Pi^\lambda(M_n(A))$. Thus $\Pi^\lambda(M_n(A)) \cong M_n(R)$ where $R = e_{11}\Pi^\lambda(M_n(A))e_{11}$, and then $R \cong \Pi^\mu(A)$ by the theorem. \square

Corollary 5.5. *If A and B are Morita-equivalent algebras, and $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ corresponds to $\mu \in K \otimes_{\mathbb{Z}} K_0(B)$, then $\Pi^\lambda(A)$ and $\Pi^\mu(B)$ are Morita equivalent.*

Proof. It suffices to prove this in two cases, when $B = M_n(A)$, and when $B = eAe$ with e an idempotent with $AeA = A$; see for example [11, Proposition 3.5.6]. The first corollary deals with the case of a matrix ring. The theorem deals with the other case, for e is an idempotent in $\Pi^\lambda(A)$ with $\Pi^\lambda(A)e\Pi^\lambda(A) = \Pi^\lambda(A)$, so that $\Pi^\lambda(A)$ is Morita equivalent to $e\Pi^\lambda(A)e \cong \Pi^\mu(eAe)$. \square

6. Products

In this section we suppose that A decomposes as a direct product of two algebras, $A = B \times C$. We identify B and C as subsets of A with $A = B \oplus C$ and $BC = CB = 0$. This leads to a decomposition $1 = e + f$ with $e \in B$ and $f \in C$ idempotents. There is a natural isomorphism $K_0(A) \cong K_0(B) \times K_0(C)$, and if $\lambda \in K \otimes_{\mathbb{Z}} K_0(B)$ and $\mu \in K \otimes_{\mathbb{Z}} K_0(C)$ we write (λ, μ) for the corresponding element of $K \otimes_{\mathbb{Z}} K_0(A)$. We prove that $\Pi^{(\lambda, \mu)}(A) \cong \Pi^\lambda(B) \times \Pi^\mu(C)$.

Lemma 6.1. *If I is the A - A -sub-bimodule of $\text{Der}(A, A \otimes A)$ generated by $e\Delta f$ and $f\Delta e$ then there is an isomorphism*

$$\text{Der}(A, A \otimes A)/I \cong \text{Der}(B, B \otimes B) \oplus \text{Der}(C, C \otimes C)$$

with $\overline{e\Delta_A e}$ corresponding to Δ_B and $\overline{f\Delta_A f}$ corresponding to Δ_C

Proof. Clearly $\text{Der}(B, B \otimes B) \oplus \text{Der}(C, C \otimes C)$ can be identified with

$$D = \{d \in \text{Der}(A, A \otimes A) \mid d(e) = 0\},$$

so it suffices to prove that D and I are complementary. Now I consists of all inner derivations of the form $s(a) = a(p + q) - (p + q)a$ with $p \in eA \otimes Af$ and $q \in fA \otimes Ae$, and since $s(e) = q - p$, it follows that $D \cap I = 0$. On the other hand, if $d : A \rightarrow A \otimes A$ is any derivation then the fact that e is idempotent implies that $d(e) \in eA \otimes Af \oplus fA \otimes Ae$. Thus there is an inner derivation $d' \in I$ with $(d + d')(e) = 0$. It follows that $D + I = \text{Der}(A, A \otimes A)$. \square

Theorem 6.2. *If $A = B \times C$, $\lambda \in K \otimes_{\mathbb{Z}} K_0(B)$, $\mu \in K \otimes_{\mathbb{Z}} K_0(C)$, and (λ, μ) is the corresponding element of $K \otimes_{\mathbb{Z}} K_0(A)$, then $\Pi^{(\lambda, \mu)}(A) \cong \Pi^\lambda(B) \times \Pi^\mu(C)$.*

Proof. It suffices to prove that if $a \in A$ then $\Pi^a(A) \cong \Pi^u(B) \times \Pi^v(C)$ where $u = eae$ and $v = faf$. Now

$$\begin{aligned} \Pi^a(A) &= T_A \text{Der}(A, A \otimes A) / (\Delta - u - v) \\ &= T_A \text{Der}(A, A \otimes A) / (e\Delta f, f\Delta e, e\Delta e - u, f\Delta f - v) \\ &\cong T_A [\text{Der}(B, B \otimes B) \oplus \text{Der}(C, C \otimes C)] / (\Delta_{eAe} - u, \Delta_{fAf} - v) \end{aligned}$$

by the previous lemma. Now this last tensor algebra is isomorphic to

$$T_B \operatorname{Der}(B, B \otimes B) \times T_C \operatorname{Der}(C, C \otimes C),$$

and the result follows. \square

7. Bimodule-finite algebras

Lemma 7.1. *If $A \rightarrow B$ is a ring epimorphism then there is an exact sequence of B - B -bimodules $0 \rightarrow \operatorname{Tor}_1^A(B, B) \rightarrow B \otimes_A \Omega^1 A \otimes_A B \rightarrow \Omega^1 B \rightarrow 0$. Thus if A is bimodule-finite then so is B .*

Proof. Take the defining exact sequence for $\Omega^1 A$ and tensor each side with B . Now use the fact that multiplication induces an isomorphism $B \otimes_A B \rightarrow B$. \square

There is the following obvious consequence.

Lemma 7.2. *If $A \rightarrow B$ is a pseudoflat ring epimorphism then $\Omega^1 B$ is isomorphic to $B \otimes_A \Omega^1 A \otimes_A B$. Thus if A is quasi-free then so is B .*

Proposition 7.3. *An algebra A is bimodule-finite if and only if it has a finitely generated subalgebra C , such that the inclusion $C \rightarrow A$ is a ring epimorphism.*

Proof. If there is such a subalgebra C then $\Omega^1 C$ is known to be finitely generated (see for example [1]), and the short exact sequence of Lemma 7.1 shows that $\Omega^1 A$ is finitely generated. For the converse, choose generators of $\Omega^1 A$, and choose a finitely generated subalgebra C sufficiently large so that the generators all lie in $C \otimes C \subseteq A \otimes A$. Now there is a commutative diagram with exact rows

$$\begin{array}{ccccccc} A \otimes_C \Omega^1 C \otimes_C A & \longrightarrow & A \otimes A & \longrightarrow & A \otimes_C A & \longrightarrow & 0 \\ & & f \downarrow & & \parallel & & g \downarrow \\ 0 & \longrightarrow & \Omega^1 A & \longrightarrow & A \otimes A & \longrightarrow & A \longrightarrow 0. \end{array}$$

By construction f is surjective, and it follows that g is injective. In other words the inclusion $C \rightarrow A$ is a ring epimorphism. \square

8. Finite-dimensional quasi-free algebras

Lemma 8.1. *Let M be an A - A -bimodule which is flat as a right A -module. If $M \otimes_A X$ is flat for all left A -modules X , then $\mathrm{Tor}_1^{A^e}(X \otimes Y, M) = 0$ for all left A -modules X and right A -modules Y .*

Proof. Fix an exact sequence $0 \rightarrow L \rightarrow P \rightarrow M \rightarrow 0$ with P a projective A - A -bimodule. Tensoring first with X and then with Y , the hypotheses imply that the tensor product sequence

$$0 \rightarrow Y \otimes_A L \otimes_A X \rightarrow Y \otimes_A P \otimes_A X \rightarrow Y \otimes_A M \otimes_A X \rightarrow 0$$

is exact. But this sequence is identified with the sequence

$$0 \rightarrow (X \otimes Y) \otimes_{A^e} L \rightarrow (X \otimes Y) \otimes_{A^e} P \rightarrow (X \otimes Y) \otimes_{A^e} M \rightarrow 0$$

so $\mathrm{Tor}_1^{A^e}(X \otimes Y, M) = 0$. \square

Lemma 8.2. *Suppose that A is a finite-dimensional algebra, and that $A/\mathrm{rad} A$ is separable over K . Let M be an A - A -bimodule which is projective as a right A -module. If $M \otimes_A X$ is projective for any left A -module X , then M is a projective bimodule.*

Proof. Since $\mathrm{rad} A$ is nilpotent, every simple A - A -bimodule occurs as a composition factor of $(A/\mathrm{rad} A) \otimes (A/\mathrm{rad} A)$, and the separability hypothesis implies that $(A/\mathrm{rad} A) \otimes (A/\mathrm{rad} A)$ is semisimple. The previous lemma now shows that $\mathrm{Tor}_1^{A^e}(S, M) = 0$ for all simple A^e -modules S . Since A^e is finite-dimensional, it follows that M is a projective A^e -module. \square

Lemma 8.3. *If A and B are finite-dimensional self-injective algebras, then so is $A \otimes B$.*

Proof. If D is duality with the field, then $D(A)$ is a projective A -module and $D(B)$ is a projective B -module. Now the isomorphism $D(A \otimes B) \cong D(A) \otimes D(B)$ shows that $D(A \otimes B)$ is a projective $A \otimes B$ -module, so $A \otimes B$ is self-injective. \square

Lemma 8.4. *If A is a finite-dimensional self-injective algebra then every A -module is either projective, or has infinite projective dimension.*

Proof. Looking at the last two terms in the minimal projective resolution of a module of finite projective dimension, there must be an injection of one projective into another which is not split. This is impossible if all projective modules are injective. \square

Proposition 8.5. *A finite-dimensional algebra A is quasi-free if and only if it is hereditary and $A/\text{rad } A$ is separable over K .*

Proof. If A is hereditary, the bimodule $\Omega^1 A$ satisfies the hypotheses of Lemma 8.2. Thus if $A/\text{rad } A$ is separable over K then $\Omega^1 A$ is a projective bimodule, so A is quasi-free.

Now suppose that A is quasi-free. Certainly this implies that A is hereditary. Let S be a simple A -module, and let B be the corresponding simple factor of $A/\text{rad } A$. Since A is hereditary, its Gabriel quiver has no oriented cycles (since any nonzero map between indecomposable projectives must be injective). It follows that $\text{Ext}_A^1(S, S) = 0$. Now the projection $A \rightarrow B$ is a pseudoflat epimorphism by [15, Theorem 4.8], since the restriction to A of any B -module is isomorphic to a direct sum of copies of S . Thus B is quasi-free, so B has projective dimension ≤ 1 as a B^e -module. Now B^e is self-injective by Lemma 8.3, so actually B is a projective module. Thus B is separable. Repeating for each simple A -module it follows that $A/\text{rad } A$ is separable. \square

9. Localization

Lemma 9.1. *If $\theta : A \rightarrow B$ is a pseudoflat epimorphism and M is a B - B -bimodule, then restriction induces a bijection $\text{Der}(B, M) \rightarrow \text{Der}(A, M)$.*

Proof. We have isomorphisms

$$\text{Hom}_{B^e}(\Omega^1 B, M) \cong \text{Hom}_{B^e}(B \otimes_A \Omega^1 A \otimes_A B, M) \cong \text{Hom}_{A^e}(\Omega^1 A, M)$$

using Lemma 7.2. \square

Lemma 9.2. *If $\theta : A \rightarrow B$ is a pseudoflat epimorphism and $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$ then there is a natural homomorphism $\Pi^\lambda(A) \rightarrow \Pi^{\theta_*(\lambda)}(B)$. Moreover the diagram*

$$\begin{array}{ccc} A & \longrightarrow & \Pi^\lambda(A) \\ \downarrow & & \downarrow \\ B & \longrightarrow & \Pi^{\theta_*(\lambda)}(B) \end{array}$$

commutes.

Proof. The question of naturality is slightly delicate, since the definition of $\Pi^\lambda(A)$ depends on the choice of a lift of $\text{tr}(\lambda)$ to A , and for the homomorphism from $\Pi^\lambda(A)$ to $\Pi^{\theta_*(\lambda)}(B)$ one should choose compatible lifts. The map θ induces a map

$\bar{\theta} : A/[A, A] \rightarrow B/[B, B]$, and the square

$$\begin{array}{ccc} K \otimes_{\mathbb{Z}} K_0(A) & \xrightarrow{\theta_*} & K \otimes_{\mathbb{Z}} K_0(B) \\ \text{tr} \downarrow & & \text{tr} \downarrow \\ A/[A, A] & \xrightarrow{\bar{\theta}} & B/[B, B] \end{array}$$

commutes. If a is a lift to A of $\text{tr}(\lambda)$ we use $\theta(a)$ as a lift of $\theta_*(\lambda)$. Thus it suffices to construct a natural map $\Pi^a(A) \rightarrow \Pi^{\theta(a)}(B)$. Now there is a natural map

$$\text{Der}(A, A \otimes A) \rightarrow \text{Der}(A, B \otimes B) \cong \text{Der}(B, B \otimes B)$$

which is in fact a homomorphism of A - A -bimodules, and it sends Δ_A to Δ_B . Combining this with the algebra map $A \rightarrow B$ one obtains an algebra map

$$T_A \text{Der}(A, A \otimes A) \rightarrow T_B \text{Der}(B, B \otimes B).$$

This map sends Δ_A to Δ_B and a to $\theta(a)$. The result follows. \square

Theorem 9.3. *Suppose that $\theta : A \rightarrow B$ is a pseudoflat epimorphism and that $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$. If A is a quasi-free bimodule-finite algebra, then so is B , and the diagram*

$$\begin{array}{ccc} A & \longrightarrow & \Pi^\lambda(A) \\ \downarrow & & \downarrow \\ B & \longrightarrow & \Pi^{\theta_*(\lambda)}(B) \end{array}$$

is a pushout in the category of rings.

Proof. We have $B \otimes_A \text{Hom}_{A^e}(\Omega^1 A, A \otimes A) \otimes_A B \cong \text{Hom}_{A^e}(\Omega^1 A, B \otimes B)$ since $\Omega^1 A$ is a finitely generated projective A - A -bimodule. Thus the induced map

$$B \otimes_A \text{Der}(A, A \otimes A) \otimes_A B \rightarrow \text{Der}(B, B \otimes B)$$

is an isomorphism of B - B -bimodules, sending $1 \otimes \Delta_A \otimes 1$ to Δ_B . Let a be a lift of $\text{tr}(\lambda)$ to A . Since $\Pi^a(A) \cong (T_A \text{Der}(A, A \otimes A))/(\Delta_A - a)$, the pushout of $A \rightarrow B$ and $A \rightarrow \Pi^a(A)$ is isomorphic to

$$T_B(B \otimes_A \text{Der}(A, A \otimes A) \otimes_A B)/(1 \otimes \Delta_A \otimes 1 - \theta(a)),$$

and this is isomorphic to $\Pi^{\theta_*(\lambda)}(B)$. \square

Corollary 9.4. *Under the hypotheses of the theorem, the map $\Pi^\lambda(A) \rightarrow \Pi^{\theta_*(\lambda)}(B)$ is a pseudoflat epimorphism.*

Proof. Use [1, Proposition 5.2]. \square

We shall also need one further observation.

Corollary 9.5. *Under the hypotheses of the theorem, the map $\Pi^\lambda(A) \rightarrow \Pi^{\theta_*(\lambda)}(B)$ preserves filtrations, that is it sends $\Pi^\lambda(A)_{\leq n}$ into $\Pi^{\theta_*(\lambda)}(B)_{\leq n}$ for all n . Moreover, it induces a commutative square*

$$\begin{array}{ccc} \Pi^0(A) & \longrightarrow & \Pi^0(B) \\ \downarrow & & \downarrow \\ \text{gr } \Pi^\lambda(A) & \longrightarrow & \text{gr } \Pi^{\theta_*(\lambda)}(B). \end{array}$$

One of the main examples of a pseudoflat epimorphism is given by universal localization, see [1]. Here we consider just the special case arising from perpendicular categories. If A is hereditary and \mathcal{X} is a collection of finitely presented left A -modules, then the *perpendicular category* \mathcal{X}^\perp is the category of modules M with $\text{Hom}_A(X, M) = \text{Ext}^1(X, M) = 0$ for all $X \in \mathcal{X}$. Considering the universal localization of A with respect to projective presentations of the modules in \mathcal{X} , one obtains a pseudoflat epimorphism $A \rightarrow A_\mathcal{X}$ with the property that restriction induces an equivalence between the category of left $A_\mathcal{X}$ -modules and \mathcal{X}^\perp . The theorem now implies the following.

Corollary 9.6. *Suppose that A is quasi-free and bimodule-finite and $\lambda \in K \otimes_{\mathbb{Z}} K_0(A)$. If \mathcal{X} is a collection of finitely presented left A -modules and $\theta : A \rightarrow A_\mathcal{X}$ is the corresponding universal localization, then restriction via the natural map induces an equivalence from the category of $\Pi^{\theta_*(\lambda)}(A_\mathcal{X})$ -modules to the category of $\Pi^\lambda(A)$ -modules whose restriction to A is in \mathcal{X}^\perp .*

10. Module varieties

Let K be an algebraically closed field, let A be a finitely generated K -algebra and let e_i ($i \in I$) be a complete set of orthogonal idempotents, so $e_i e_j = 0$ for $i \neq j$ and $\sum_i e_i = 1$. If $\alpha \in \mathbb{N}^I$, we write $\text{Rep}(A, \alpha)$ for the variety of A -module structures on $K^\alpha = \bigoplus_i K^{\alpha_i}$ under which each e_i acts as projection onto the i th summand. Thus $\text{Rep}(A, \alpha)$ consists of all algebra maps $A \rightarrow \text{End}_K(K^\alpha)$ sending e_i to the projection onto K^{α_i} . Elements of $\text{Rep}(A, \alpha)$ are representations of A of dimension vector α . The group $\text{GL}(\alpha) = \prod_i \text{GL}(\alpha_i)$ acts naturally on this variety, and the orbits correspond to isomorphism classes of representations. The stabilizer of a representation X can be identified with the automorphism group of X . It follows that the orbit of X has dimension $\sum_i \alpha_i^2 - \dim \text{End}(X)$.

Lemma 10.1. *If $\beta \leq \alpha$ then the set of elements of $\text{Rep}(A, \alpha)$ such that the corresponding representation has a subrepresentation of dimension vector β , is closed.*

Proof. It is the image of a closed set under the projection $\text{Rep}(A, \alpha) \times P \rightarrow \text{Rep}(A, \alpha)$, where P is the product of Grassmannians of subspaces of dimension β_i in K^{α_i} . Now use the fact that P is a projective variety. See [16, Lemma 3.1]. \square

We write $\text{Rep}(A, \alpha)_s$ for the $\text{GL}(\alpha)$ -stable subset consisting of simple A -modules. By the lemma it is open.

Lemma 10.2. *If A is a finitely generated K -algebra of GK dimension d , then for any α we have $\dim \text{Rep}(A, \alpha)_s \leq \sum_i \alpha_i^2 + d - 1$.*

Proof. Passing to the quotient of A by the intersection of the annihilators of all simple representations of dimension vector α , we may suppose that A is a semiprime PI ring, satisfying the identities of $N \times N$ matrices, where $N = \sum_i \alpha_i$. By [11, Theorem 13.4.4], A has only finitely many minimal prime ideals, so we may assume that it is prime. By [11, Proposition 13.7.4] there is a central localization A_c which is an Azumaya algebra. Now each simple A -module is either an A_c -module, or an $A/(c)$ -module, so by an induction we reduce to the case when A is an Azumaya algebra, say with centre Z .

Now there is a natural map $f : \text{Rep}(A, \alpha)_s \rightarrow \text{maxspec } Z$ sending a simple A -module to its central character. For, Z is finitely generated by [11, Lemma 13.9.10], and if z_1, \dots, z_r are generators, they identify $\text{maxspec } Z$ with a closed subset of K^r . Now if $N = \sum_i \alpha_i$ then $\text{End}_K(K^\alpha) \cong M_N(K)$, and if $\theta : A \rightarrow M_N(K)$ is an element of $\text{Rep}(A, \alpha)_s$ then the map $z \mapsto \theta(z)_{11}$ is the central character of θ . Thus we can define f by sending θ to $(\theta(z_1)_{11}, \dots, \theta(z_r)_{11}) \in K^r$.

Now each fibre of f meets only finitely many $\text{GL}(\alpha)$ -orbits, and each orbit has dimension $\sum_i \alpha_i^2 - 1$. The result follows since Z has Krull dimension at most d . \square

Now let Q be a quiver with vertex set I and let $\alpha \in \mathbb{N}^I$. Clearly $\text{Rep}(KQ, \alpha)$ can be identified with the affine space

$$\bigoplus_{a:i \rightarrow j} \text{Hom}_K(K^{\alpha_i}, K^{\alpha_j}).$$

Also, by Section 3 and [5], one can identify $\text{Rep}(\Pi^0(KQ), \alpha)$ with the fibre over 0 of the moment map

$$\mu : \text{Rep}(K\overline{Q}, \alpha) \rightarrow \text{End}(\alpha)_0, \quad x \mapsto \sum_{a \in Q} [x_a, x_{a^*}],$$

where $\text{End}(\alpha) = \prod_i \text{End}(K^{\alpha_i})$ and $\text{End}(\alpha)_0 = \{(\theta_i) \in \text{End}(\alpha) \mid \sum_i \text{tr}(\theta_i) = 0\}$. Recall that a module M is called a *brick* if $\text{End}(M) = K$.

Lemma 10.3. *The map μ is smooth at a point $x \in \text{Rep}(K\overline{Q}, \alpha)$ if and only if the corresponding module is a brick.*

Proof. Identifying $\text{Rep}(K\overline{Q}, \alpha)$ and $\text{End}(\alpha)_0$ with their tangent spaces at x and $\mu(x)$ respectively, μ induces the map

$$d\mu_x : \text{Rep}(K\overline{Q}, \alpha) \rightarrow \text{End}(\alpha)_0, \quad y \mapsto \sum_{a \in Q} ([x_a, y_{a^*}] + [y_a, x_{a^*}]).$$

Now if D is duality with the field, there is a trace pairing which identifies the vector spaces $D(\text{Rep}(K\overline{Q}, \alpha)) \cong \text{Rep}(K\overline{Q}, \alpha)$ (with arrows a and a^* being interchanged), and identifies $D(\text{End}(\alpha)_0) \cong \text{End}(\alpha)/K$. Then $D(d\mu_x)$ is the map

$$\begin{aligned} \text{End}(\alpha)/K &\rightarrow \text{Rep}(K\overline{Q}, \alpha), \\ (\theta_i) &\mapsto \sum_{a:i \rightarrow j \in Q} ((\theta_j x_a - x_a \theta_i)_{a^*} + (x_{a^*} \theta_j - \theta_i x_{a^*})_a), \end{aligned}$$

so if X is the $\Pi^0(KQ)$ -module corresponding to x , then $\text{Ker } D(d\mu_x) \cong \text{End}(X)/K$. Now X is a brick if and only if $D(d\mu_x)$ is injective, so if and only if μ is smooth at x . \square

Now let Q be an extended Dynkin quiver and let δ be the minimal positive imaginary root for Q .

Lemma 10.4. *The restriction map $\pi : \text{Rep}(\Pi^0(KQ), \delta) \rightarrow \text{Rep}(KQ, \delta)$ is onto, and the fibre over a point $x \in \text{Rep}(KQ, \delta)$ is irreducible of dimension $\dim \text{End}(X)$, where X is the KQ -module corresponding to x .*

Proof. The fibre over x is isomorphic to the fibre $c^{-1}(0)$ in [5, Lemma 4.2], so it is isomorphic to $D \text{Ext}^1(X, X)$. Now since X has dimension δ the Ringel form implies that this has dimension $\dim \text{End}(X)$. \square

Lemma 10.5. *The variety $\text{Rep}(\Pi^0(KQ), \delta)$ is irreducible and Cohen-Macaulay of dimension $1 + \sum_i \delta_i^2$, and the general element is a simple $\Pi^0(KQ)$ -module. Moreover, if $\text{Rep}(\Pi^0(KQ), \delta)$ is considered as a scheme using the natural scheme structure on the fibre $\mu^{-1}(0)$, then it is reduced.*

Proof. Equip $\text{Rep}(\Pi^0(KQ), \delta)$ with the scheme structure. The argument of [5, Lemma 8.3] shows that $\text{Rep}(\Pi^0(KQ), \delta)$ is irreducible of dimension $1 + \sum_i \delta_i^2$, hence a complete intersection, so Cohen-Macaulay.

Supposing that the general element is not simple, it follows from the irreducibility and Lemma 10.1 that there is $0 < \beta < \delta$ such that every $\Pi^0(KQ)$ -module of dimension δ has a subrepresentation of dimension β . Now any KQ -module of

dimension δ can be considered trivially as a $\Pi^0(KQ)$ -module, so has a submodule of dimension β . Similarly any KQ^{op} -module of dimension δ can be considered as a $\Pi^0(KQ)$ -module, so has a submodule of dimension β . Dualizing, this implies that any KQ -module of dimension δ has a submodule of dimension $\delta - \beta$. This is impossible by [16, Theorem 3.4].

Now μ is smooth at the general point of $\text{Rep}(\Pi^0(KQ), \delta)$ by Lemma 10.3, so $\text{Rep}(\Pi^0(KQ), \delta)$ is generically smooth, hence generically reduced. With the Cohen-Macaulay property, this implies that it is reduced. \square

11. A Conze embedding

Let Q be an extended Dynkin quiver with vertex set I and let $\delta \in \mathbb{Z}^I$ be the minimal positive imaginary root for Q . Let K be an algebraically closed field.

Lemma 11.1. *There is a pseudoflat epimorphism $\theta : KQ \rightarrow M_N(K[x])$ such that the general representation of KQ of dimension δ is the restriction of a $M_N(K[x])$ -module.*

Proof. If Q is of type \tilde{A}_n , oriented as a cycle, then \mathcal{X} should consist of n of the $n+1$ one-dimensional simple modules. If Q has no oriented cycles then \mathcal{X} should contain all the regular simple modules in one tube in the Auslander-Reiten quiver of KQ , and all but one regular simple module in every other tube. Localizing at a set which contains all but one regular simple in each tube, one obtains by [4, Theorem 4.2] a tame hereditary algebra with two simple modules. Since the base field is algebraically closed, this algebra is Morita equivalent to the Kronecker algebra. Now localizing at one further regular simple module, one obtains an algebra Morita equivalent to $K[x]$. \square

Henceforth we suppose that $\theta : KQ \rightarrow M_N(K[x])$ is a pseudoflat epimorphism such that the general representation of KQ of dimension δ is the restriction of a $M_N(K[x])$ -module. If $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$ there is a corresponding element $\lambda \in K^I$. See Section 3.

Lemma 11.2. *We have $N = \sum_i \delta_i$, and if P is the indecomposable projective module for $M_N(K[x])$, then $\theta_*(\lambda) = \sum_i \lambda_i \delta_i \otimes [P]$ for $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$.*

Proof. The general representation of KQ of dimension δ is a brick, so it must be the restriction of a simple $M_N(K[x])$ -module. The result follows. \square

Recall from the introduction that θ induces a map θ_λ from $\Pi^\lambda(KQ)$ to $M_N(C_\nu)$ for some ν . The previous lemma and Corollary 5.4 show that $\nu = \sum_i \lambda_i \delta_i$.

Lemma 11.3. *The map θ_0 is injective.*

Proof. We factorize θ_0 as a product

$$\Pi^0(KQ) \xrightarrow{f} \Pi^0(KQ)/\text{Ker}(\theta_0) \xrightarrow{g} M_N(C_0).$$

Both of these maps are ring epimorphisms, so we obtain injective morphisms

$$\text{Rep}(M_N(C_\nu), \delta) \rightarrow \text{Rep}(\Pi^0(KQ)/\text{Ker}(\theta_0), \delta) \rightarrow \text{Rep}(\Pi^0(KQ), \delta).$$

The image of the composition of these is the open set of representations whose restriction to KQ is in \mathcal{X}^\perp . Now Lemma 10.5 implies that $\text{Rep}(\Pi^0(KQ)/\text{Ker}(\theta_0), \delta)_s$ has dimension $1 + \sum_i \delta_i^2$. Thus $\Pi^0(KQ)/\text{Ker}(\theta_0)$ has GK dimension at least 2 by Lemma 10.2. Now $\Pi^0(KQ)$ is prime of GK dimension 2 by [2], and it follows that $\text{Ker}(\theta_0) = 0$, as claimed. \square

Theorem 11.4. *The map θ_λ is injective, the natural map $\Pi^0(KQ) \rightarrow \text{gr } \Pi^\lambda(KQ)$ is an isomorphism, and $\Pi^\lambda(KQ)$ is prime of GK dimension 2.*

Proof. By Corollary 9.5 there is a commutative square

$$\begin{array}{ccc} \Pi^0(KQ) & \xrightarrow{\theta_0} & M_N(C_0) \\ \downarrow & & \downarrow \\ \text{gr } \Pi^\lambda(KQ) & \xrightarrow{\text{gr } \theta_\lambda} & \text{gr } M_N(C_\nu). \end{array}$$

The associated graded algebra for the first Weyl algebra is the polynomial ring in two variables, so the map $M_N(C_0) \rightarrow \text{gr } M_N(C_\nu)$ is an isomorphism. Now the map $\Pi^0(KQ) \rightarrow \text{gr } \Pi^\lambda(KQ)$ is surjective, so the fact that θ_0 is injective implies that $\Pi^0(KQ) \rightarrow \text{gr } \Pi^\lambda(KQ)$ is an isomorphism and $\text{gr } \theta_\lambda$ is injective. It follows that θ_λ is injective. Finally, $\Pi^\lambda(KQ)$ is prime of GK dimension 2 by [2]. \square

Lemma 11.5. *The ring $\Pi^\lambda(KQ)$ has Goldie rank at least N .*

Proof. Let M be the restriction to $\Pi^\lambda(KQ)$ of the simple module for its quotient ring. Letting $E = \text{End}(M)^{op}$ we consider M as a $\Pi^\lambda(KQ)$ - E -bimodule. One can consider M as a representation of the quiver \overline{Q} by right E -vector spaces and E -linear maps, satisfying the usual relations for the deformed preprojective algebra. Now these vector spaces are finite dimensional over E , and lead to a dimension vector $\alpha \in \mathbb{N}^I$, and the Goldie rank of $\Pi^\lambda(KQ)$ is $\sum_i \alpha_i$.

If i is a loopfree vertex and $\lambda_i \neq 0$ then the reflection functor of [5, Theorem 5.1] evidently defines an equivalence from the category of $\Pi^\lambda(KQ)$ - E -bimodules to the category of $\Pi^{\lambda'}(KQ)$ - E -bimodules, for some λ' , which acts as s_i on dimension vectors.

By applying a sequence of reflection functors to M we pass to a $\Pi^{\lambda'}(KQ)$ - E -bimodule M' of dimension vector α' (for a new λ'), and we choose the sequence to ensure that α' is minimal. This implies that for any vertex i , either $\lambda'_i = 0$ or $(\alpha', \epsilon_i) \leq 0$.

If i is a vertex with $\lambda'_i = 0$, the 1-dimensional simple module S_i at vertex i has projective resolution

$$0 \rightarrow \Pi^{\lambda}(KQ)e_i \rightarrow \bigoplus_{a:i \rightarrow j \text{ in } \overline{Q}} \Pi^{\lambda}(KQ)e_j \rightarrow \Pi^{\lambda}(KQ)e_i \rightarrow S_i \rightarrow 0.$$

Now M is an injective module over $\Pi^{\lambda}(KQ)$, so M' is injective over $\Pi^{\lambda'}(KQ)$, so applying $\dim_E \operatorname{Hom}(-, M')$ we deduce that $(\alpha', \epsilon_i) = 0$.

Thus α' is in the fundamental region for Q , so is a multiple of the vector δ . Now δ is unchanged by the reflections s_i , so α must have been a multiple of δ . The result follows. \square

Let D_{ν} be the quotient division algebra for C_{ν} .

Theorem 11.6. *The map θ_{λ} induces an isomorphism from the simple artinian quotient ring of $\Pi^{\lambda}(KQ)$ to $M_N(D_{\nu})$.*

Proof. First observe that if $k < N$ and D and E are division rings, then there can be no homomorphism $M_N(E) \rightarrow M_k(D)$, for if S is the module obtained by inducing the simple $M_N(E)$ -module up to $M_k(D)$, then $S^N \cong M_k(D)$ is semisimple of length k .

By [11, Proposition 3.1.16], the quotient ring of $\Pi^{\lambda}(KQ)$ embeds in $M_k(D_{\nu})$ for some $k \leq N$. By the discussion above we must have $k = N$, and inspecting the proof of the cited result we see that $M_N(D_{\nu})$ is simple as a $\Pi^{\lambda}(KQ)$ - $M_N(D_{\nu})$ -bimodule, and then that it is torsion free over $\Pi^{\lambda}(KQ)$. This means that θ_{λ} does induce a map from the quotient ring of $\Pi^{\lambda}(KQ)$ to $M_N(D_{\nu})$. Moreover this map is an isomorphism since it is a ring epimorphism. \square

We now apply this to Kleinian singularities. Let K be an algebraically closed field of characteristic zero, and let Γ be a finite subgroup of $\operatorname{SL}_2(K)$. Let Q be an orientation of the McKay quiver of Γ . Recall that there is an isomorphism $\mathcal{O}^{\lambda} \cong e_0 \Pi^{\lambda}(KQ) e_0$, where $\lambda \in Z(K\Gamma)$ is identified with $\boldsymbol{\lambda} \in K^I$ by letting λ_i be the trace of λ on the i th irreducible representation of Γ , and hence with $\lambda \in K \otimes_{\mathbb{Z}} K_0(KQ)$.

Corollary 11.7. *There is an embedding $\psi_{\lambda} : \mathcal{O}^{\lambda} \rightarrow C_{\nu}$ where ν is the trace of λ on the regular representation of Γ . Moreover ψ_{λ} induces an isomorphism on quotient division rings.*

Proof. It suffices to observe that if e is an idempotent in a prime Goldie ring R

with simple artinian quotient ring $Q(R)$, then eRe is prime Goldie with quotient ring $eQ(R)e$. \square

Appendix

In this appendix we use the methods of the paper to prove the normality of a certain variety. This result is used in the article [9] by M. P. Holland. If K is an algebraically closed field and Q is an extended Dynkin quiver with minimal imaginary root δ , then the variety $\text{Rep}(\Pi^0(KQ), \delta)$ need not be normal (see below). Here we prove the normality of the open subvariety $\text{Rep}(\Pi^0(KQ), \delta)_{ss}$ of semistable $\Pi^0(KQ)$ -modules, where the semistable modules are defined as follows. If M is a KQ - or a $\Pi^0(KQ)$ -module of dimension vector α , then its *defect* is defined by the formula $d(M) = \langle \delta, \alpha \rangle$. One says that M is *semistable* if $d(M) = 0$ and $d(N) \leq 0$ for all submodules $N \subseteq M$. It is well known that the semistable KQ -modules are exactly the regular modules.

Lemma 12.1. *A $\Pi^0(KQ)$ -module is semistable if and only if it is semistable as a KQ -module.*

Proof. Certainly if M is semistable as a KQ -module then it is as a $\Pi^0(KQ)$ -module, for any $\Pi^0(KQ)$ -submodule N is also a KQ -submodule, so $d(N) \leq 0$.

Now suppose that M is semistable as a $\Pi^0(KQ)$ -module. To show that it is semistable over KQ , it suffices to show that all indecomposable KQ -submodules of M are preprojective or regular. For a contradiction, let N be an indecomposable preinjective submodule. Now by Theorem 2.3,

$$\Pi^0(KQ) \otimes_{KQ} N \cong N \oplus \tau^- N \oplus \tau^{-2}(N) \oplus \dots,$$

and since N is preinjective this sum terminates, so it is a finite-dimensional preinjective KQ -module. Now the $\Pi^0(KQ)$ -submodule \overline{N} of M generated by N is a quotient of this sum, so it is preinjective as a KQ -module. Thus $d(\overline{N}) > 0$, contrary to the assumption. \square

Lemma 12.2. *If Q is an extended Dynkin quiver then any semistable KQ -module M of dimension δ can be extended to a $\Pi^0(KQ)$ -module which is a brick.*

Proof. Certainly this is true if Q has type \tilde{A}_n , so that $\delta_i = 1$ for all vertices i . One considers the KQ -module M as a representation of Q in which the vector space at each vertex is 1-dimensional. Now one extends this to a representation of \overline{Q} by letting a^* be a non-zero map if and only if a is zero. Clearly this defines a $\Pi^0(KQ)$ -module, and it is easy to see that it is a brick.

To deal with other quivers, we first formulate the assertion in a Morita-invariant way, and then we use universal localization to reduce to type \tilde{A}_n .

Observe that a KQ -module M is semistable of dimension δ if and only if it is regular, and its regular composition factors are exactly the regular simples for some tube in the Auslander-Reiten quiver of KQ , each with multiplicity one. For simplicity we call this property (*).

In view of the Morita equivalence property for $\Pi^0(A)$, the lemma may be formulated as the following claim: if A is a tame hereditary algebra and M is an A -module with property (*), then M can be extended to a $\Pi^0(A)$ -module which is a brick. We prove this claim by induction on the number of simple modules for A . Let M be a module with property (*).

If there are no inhomogeneous tubes in the Auslander-Reiten quiver for A , except possibly the one containing M , then A is of type \tilde{A}_n , and we have checked the claim at the start of the proof.

If there is such an inhomogeneous tube, choose a regular simple module X contained in this tube, and let $A \rightarrow B$ be the corresponding universal localization. Then B is a tame hereditary algebra with one fewer simple module than A by [4, Theorem 4.2]. Now M is in the perpendicular category to X , so it is the restriction of a B -module M' , and clearly M' has property (*). By induction M' can be extended to a $\Pi^0(B)$ -module M'' which is a brick, and then since $\Pi^0(A) \rightarrow \Pi^0(B)$ is a ring epimorphism, the restriction of M'' to $\Pi^0(A)$ is a brick. The claim follows. \square

Theorem 12.3. *If Q is an extended Dynkin quiver then $\text{Rep}(\Pi^0(KQ), \delta)_{ss}$ is a normal variety.*

Proof. It is Cohen-Macaulay by Lemma 10.5, so it suffices to prove that its singular locus S has dimension at most $-1 + \sum_i \delta_i^2$. Consider the projection

$$\pi : \text{Rep}(\Pi^0(KQ), \delta)_{ss} \rightarrow \text{Rep}(KQ, \delta)_{ss}.$$

Now the general element r of $\text{Rep}(KQ, \delta)_{ss}$ is a brick, so every element x of the fibre $\pi^{-1}(r)$ is a brick. Thus by Lemma 10.3 the map μ is smooth at x , so $\text{Rep}(\Pi^0(KQ), \delta)_{ss}$ is smooth at x . Besides the bricks, there are only finitely many $\text{GL}(\delta)$ -orbits of non-bricks in $\text{Rep}(KQ, \delta)_{ss}$. If \mathcal{O}_X is one of these orbits, then

$$\dim \pi^{-1}(\mathcal{O}_X) = \dim \text{End}(X) + \dim \mathcal{O}_X = \dim \text{GL}(\delta) = \sum_i \delta_i^2.$$

Now the general element of $\pi^{-1}(\mathcal{O}_X)$ is a brick by the previous lemma, and so $\pi^{-1}(\mathcal{O}_X) \cap S$ has dimension at most $-1 + \sum_i \delta_i^2$, as required. \square

Finally we justify our claim that $\text{Rep}(\Pi^0(KQ), \delta)$ need not be normal. By Lemma 10.5 the natural scheme structure on $\text{Rep}(\Pi^0(KQ), \delta)$ is reduced, and hence in the notation of Section 10, the tangent space at a point $x \in \text{Rep}(\Pi^0(KQ), \delta)$ can be identified with $\text{Ker}(d\mu_x)$. It follows that $\text{Rep}(\Pi^0(KQ), \delta)$ is smooth at x if and only if x is a brick.

Now Ringel [12] has pointed out that if Q is extended Dynkin, not of type \tilde{A}_n , then there is a KQ -module X of dimension δ which is not the restriction of any brick for $\Pi^0(KQ)$. For example, let I be an indecomposable injective KQ -module of defect ≥ 2 , say of dimension vector α . Then $\delta - \alpha$ is a positive root, so is the dimension vector of an indecomposable P , necessarily preprojective. Let $X = P \oplus I$. The condition on defect implies that $\text{Hom}(X, I)$ has dimension at least 2, so $\text{Hom}(P, I) \neq 0$, and hence X has a non-zero endomorphism ϕ which kills I and has image contained in I . By Lemmas 1.4 and 2.2, a $\Pi^0(KQ)$ -module structure on X is determined by a map $\tau^-X \rightarrow X$. Now since $\tau^-I = 0$ and $\text{Hom}(\tau^-P, P) = 0$, it is easy to see that ϕ is a $\Pi^0(KQ)$ -endomorphism.

Now the inverse image of the $\text{GL}(\delta)$ -orbit of X under the projection

$$\text{Rep}(\Pi^0(KQ), \delta) \rightarrow \text{Rep}(KQ, \delta)$$

has codimension 1 by the argument of Theorem 12.3. Since all points of the inverse image are singular, $\text{Rep}(\Pi^0(KQ), \delta)$ cannot be normal.

References

- [1] G. M. Bergman and W. Dicks, Universal derivations and universal ring constructions, *Pacific J. Math.* **79** (1978), 293–337.
- [2] D. Baer, W. Geigle and H. Lenzing, The preprojective algebra of a tame hereditary algebra, *Commun. Algebra* **15** (1987), 425–457.
- [3] N. Conze, Algèbres d'opérateurs différentiels et quotients des algèbres enveloppantes, *Bull. Soc. math. France* **102** (1974), 379–415.
- [4] W. Crawley-Boevey, Regular modules for tame hereditary algebras, *Proc. London Math. Soc.* **62** (1991), 490–508.
- [5] W. Crawley-Boevey and M. P. Holland, Noncommutative deformations of Kleinian singularities, *Duke Math. J.* **92** (1998), 605–635.
- [6] J. Cuntz and D. Quillen, Algebra extensions and nonsingularity, *J. Amer. Math. Soc.* **8** (1995), 251–289.
- [7] V. Dlab and C. M. Ringel, The preprojective algebra of a modulated graph. In: *Representation theory II, Proc. Ottawa 1979*, eds V. Dlab and P. Gabriel, Lec. Notes in Math. **832**, Springer, Berlin 1980, pp. 216–231.
- [8] I. M. Gelfand and V. A. Ponomarev, Model algebras and representations of graphs, *Funkcional Anal. i Prilozhen* **13** (1979), 1–12. English translation: *Func. Anal. Appl.* **13** (1979), 157–166.
- [9] M. P. Holland, Quantization of the Marsden-Weinstein reduction for extended Dynkin quivers, to appear in *Ann. Sci. École Norm. Sup.*
- [10] H. Matsumura, *Commutative Ring Theory*, Cambridge Univ. Press 1986.
- [11] J. C. McConnell and J. C. Robson, *Noncommutative noetherian rings*, Wiley 1987.
- [12] C. M. Ringel, private communication, 30 July 1997.
- [13] C. M. Ringel, The preprojective algebra of a quiver, in: *Algebras and modules II*, Proc. Geiranger 1996, *Canad. Math. Soc. Conf. Proc.* **24**, Amer. Math. Soc. 1998, 467–480.
- [14] W. F. Schelter, Smooth Algebras, *J. Algebra* **103** (1986), 677–685.
- [15] A. Schofield, *Representations of rings over skew fields*, London Math. Soc. Lec. Note Ser. **92**, Cambridge Univ. Press 1985.
- [16] A. Schofield, General representations of quivers, *Proc. London Math. Soc.* **65** (1992), 46–64.

William Crawley-Boevey
Department of Pure Mathematics
University of Leeds
Leeds LS2 9JT
UK
e-mail: w.crawley-boevey@leeds.ac.uk

(Received: March 4, 1998)