

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 77 (2002)

Artikel: Marked length rigidity for symmetric spaces
Autor: Dal'Bo, Françoise / Kim, Inkang
DOI: <https://doi.org/10.5169/seals-57933>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Marked length rigidity for symmetric spaces

Françoise Dal'Bo and In Kang Kim¹

Abstract. We give conditions under which a homomorphism between two Zariski dense subgroups of connected semisimple Lie groups G and G' without compact factors and with trivial center can be extended to a continuous isomorphism between G and G' . In particular we prove the marked length rigidity and the marked translation vector rigidity. This last result was motivated by a Margulis's question.

Mathematics Subject Classification (2000). 22E40, 22F30.

Keywords. Symmetric spaces, length spectrum.

Introduction

Let G, G' be connected semisimple Lie groups without compact factors and with trivial center. The motivation of this paper is to give conditions under which a homomorphism between two Zariski dense subgroups of G and G' can be extended to a continuous isomorphism between G and G' . Much study of lattices has been done, yet the study of general co-infinite volume groups is relatively less carried out. Fix a closed Weyl chamber \mathcal{A}^+ included in the Lie algebra of G . The translation vector $v(g)$ of $g \in G$, is, by definition, the unique $a \in \mathcal{A}^+$ such that e^a is conjugate to the hyperbolic part of the Jordan decomposition of g (see section 1). The Euclidean norm of $v(g)$ is denoted $\ell(g)$ and is called the length of g . If X is a symmetric space associated to G , one has: $\ell(g) = \inf_{x \in X} d(x, g(x))$. In the particular case where $G = PSL(n, \mathbb{R})$ and \mathcal{A}^+ is the set of diagonal matrices $\text{diag}(a_1, \dots, a_n)$ with $a_1 \geq \dots \geq a_n$, one has: $v(g) = \text{diag}(\text{Log} |\lambda_1|, \dots, \text{Log} |\lambda_n|)$ where λ_i is the i^{th} complex eigenvalue of g . Let $\Gamma \subset G$, the limit cone, $\mathcal{L}(\Gamma)$, associated to Γ is, by definition, the smallest closed cone in \mathcal{A}^+ containing all $v(\gamma)$ for $\gamma \in \Gamma$. An important result due to Y. Benoist [1] says that the interior of $\mathcal{L}(\Gamma)$ is not empty, if Γ is a Zariski dense group. The originality of this paper is to explore this property to obtain strong rigidity results in a short and elementary way.

Let us give the main results.

¹Partially supported by the KOSEF interdisciplinary grant 1999-2-101-5.

Theorem A. *Let $\Gamma \subset G, \Gamma' \subset G'$ be Zariski dense subgroups. If φ is a surjective homomorphism between Γ and Γ' such that $\ell(\gamma) = \ell(\varphi(\gamma))$ for any $\gamma \in \Gamma$ then φ can be extended to a continuous isomorphism between G and G' .*

Following the way of A. Parreau [15], we give applications of Theorem A to the space of representations of an abstract group into G .

Theorem A is already known for symmetric spaces of rank 1 ([4], [11]) and their products ([12]). For simple Lie groups it is shown in ([6]). Along this line, Besson, Courtois, Gallot and Hamenstädt ([2], [9]) showed that, if M is a negatively curved locally symmetric compact manifold and N is an arbitrary negatively curved manifold which has the same marked length spectrum with M , then they are isometric. Actually it is conjectured that two negatively curved compact manifolds with the same marked length spectrum are isometric. This conjecture is proved in dimension 2 ([14]).

The following theorem gives a positive answer to a Margulis's question raised during the rigidity conference at Paris in June 1998.

Theorem B. *Suppose $G = G'$ and rank $G \geq 2$. Let Γ, Γ' be Zariski dense subgroups of G . If φ is a surjective homomorphism between Γ and Γ' such that for all $\gamma \in \Gamma$ there exists $k(\gamma) \in \mathbb{R}^*$ such that $v(\varphi(\gamma)) = k(\gamma)v(\gamma)$, then φ can be extended to a continuous automorphism of G .*

We first study the simple case where G and G' are simple. Using a criterion of conjugacy proved in [6] we give a family of conditions (including conditions of Theorems A and B) under which a surjective homomorphism between Zariski dense subgroups can be extended.

1. Benoist's theorem for limit cone

An element g of a real reductive connected linear group can be uniquely written $g = ehu$ where e is elliptic (all the eigenvalues have modulus 1), u is unipotent ($u - \text{Id}$ is nilpotent), h is hyperbolic (all the eigenvalues are real positive), and all three commute. This decomposition is called the Jordan decomposition of g . If $G = \text{KAN}$ is any Iwasawa decomposition of a connected semisimple Lie group G , then e is conjugate to an element in K , h is conjugate to an element in A and u is conjugate to an element in N ([1], [7]). Fix a closed Weyl chamber \mathcal{A}^+ in the Lie algebra of G , there exists a unique $a \in \mathcal{A}^+$, called the translation vector of g and denoted $v(g)$, such that h is conjugate to e^a . Geometrically, if X is a symmetric space associated to G , then $\|v(g)\| = \ell(g)$ where $\ell(g) = \inf_{x \in X} d(x, g(x))$ (see [15] for an interpretation of $v(g)$). Let Γ be a subgroup of G , one defines the limit cone of Γ , denoted $\mathcal{L}(\Gamma)$, as the smallest closed cone in \mathcal{A}^+ containing $v(\Gamma)$. If $G = \text{PSL}(2, \mathbb{R}) \times \text{PSL}(2, \mathbb{R})$ and $\mathcal{A}^+ = \{(r_1 M, r_2 M) / r_1, r_2 \in \mathbb{R}^+\}$ where $M = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$, then $\mathcal{L}(\Gamma)$ is the closure

of $\{(r\ell(\gamma_1)M, r\ell(\gamma_2)M)/r \in \mathbb{R}^+, (\gamma_1, \gamma_2) \in \Gamma\}$ where $\ell(\gamma_i) = 0$ if γ_i is elliptic or parabolic and $\ell(\gamma_i) > 0$ is the displacement of γ_i if γ_i is hyperbolic. The following result, due to Y. Benoist, plays a key role in this paper.

Theorem 1.1 [1]. *If Γ is a Zariski dense subgroup of G then $\mathcal{L}(\Gamma)$ is convex and has nonempty interior.*

In the particular case where Γ is a Zariski dense subgroup of $\mathrm{PSL}(2, \mathbb{R}) \times \mathrm{PSL}(2, \mathbb{R})$ associated to the diagonal action of two isomorphic Fuchsian groups $\Gamma_1 \xrightarrow{\varphi} \Gamma_2$, this theorem says that $\left\{ \frac{\ell(\gamma_1)}{\ell(\varphi(\gamma_1))}, \gamma_1 \in \Gamma_1 \right\}$ is an interval $[a, b] \subset [0, \infty]$ with $a \neq b$. This property was already remarked in the context of rank 1 semisimple groups by M. Burger [4] (see also [5]).

2. Rigidity results for simple groups

In this section one supposes that G and G' are connected, noncompact, **simple** Lie groups with trivial center. Let $\varphi : \Gamma \rightarrow \Gamma'$ be a homomorphism between two subgroups of G and G' . One defines the graph group $\Gamma_\varphi \subset G \times G'$ by $\Gamma_\varphi = \{(\gamma, \varphi(\gamma))/\gamma \in \Gamma\}$. The following result is proved in [6].

Criterion of conjugacy 2.1 [6]. *Let φ be a surjective homomorphism between two Zariski dense subgroups Γ, Γ' included in connected non compact simple Lie groups, G and G' , with trivial center. The following properties are equivalent:*

- 1) φ can be extended to a continuous isomorphism between G and G'
- 2) Γ_φ is not Zariski dense in $G \times G'$.

This criterion is false if G and G' are not simple. Take for example $G = \mathrm{PSL}(2, \mathbb{R})$ and $G' = G \times G$. Denote \mathcal{A}^+ the closed Weyl chamber of G defined by $\mathcal{A}^+ = \{rM/r \in \mathbb{R}^+\}$ where $M = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Let $\varphi : \Gamma_1 \rightarrow \Gamma_2$ be an isomorphism between non conjugate and non elementary Fuchsian groups. The groups Γ_1 and $\Gamma_{1\varphi}$ are Zariski dense subgroups respectively of G and G' . Consider the isomorphism $\Psi : \Gamma_1 \rightarrow \Gamma_{1\varphi}$ defined by $\Psi(\gamma) = (\gamma, \varphi(\gamma))$. The limit cone of the graph group associated to Ψ is included in $\{(rM, rM, sM)/r, s \in \mathbb{R}^+\} \subset \mathcal{A}^+ \times \mathcal{A}^+$ and hence has empty interior. According to Benoist's theorem (section 1), $\Gamma_{1\Psi}$ is not Zariski dense. On the other hand Ψ cannot be extended.

One deduces from the previous criterion the following corollary.

Corollary 2.2. *Let Ad be the adjoint representation. If there exists an algebraic relation satisfied by all $(\mathrm{Ad}(\gamma)), \mathrm{Ad}(\varphi(\gamma))$ with $\gamma \in \Gamma$, then φ can be extended to a continuous isomorphism between G and G' .*

In the case where $G = \mathrm{PSL}(n, \mathbb{R})$, $G' = \mathrm{PSL}(n', \mathbb{R})$ and φ preserves the trace, Corollary 2.2 is proved in [16].

Remark that the condition $\ell(\gamma) = \ell(\varphi(\gamma))$ for each $\gamma \in \Gamma$ is not in general an algebraic condition. But in this case, since $\|v(\gamma)\| = \|v(\varphi(\gamma))\|$ for $\gamma \in \Gamma$, the limit cone of the graph group has empty interior. Applying Benoist's theorem, one concludes that Γ_φ is not Zariski dense and hence that φ can be extended. More generally, one has the following result.

Corollary 2.3. *If the interior of $\mathcal{L}(\Gamma_\varphi)$ is empty then φ can be extended to a continuous isomorphism between G and G' .*

Let us give three different conditions under which Γ_φ is not Zariski dense and hence φ can be extended:

- 1) $\ell(\gamma) = \ell(\varphi(\gamma))$ for any $\gamma \in \Gamma$.
- 2) $v(\gamma)$ and $v(\varphi(\gamma))$ are colinear for any $\gamma \in \Gamma$.
- 3) The largest modulus of the complex eigenvalue of $\mathrm{Ad}(\gamma)$ equals the largest one of $\mathrm{Ad}(\varphi(\gamma))$ for any $\gamma \in \Gamma$.

Conditions 1) and 2) correspond to Theorems A and B when G and G' are simple. Contrary to the conditions 1) and 2), if φ satisfies condition 3) and G and G' are not simple, φ cannot be necessarily extended. For example, fix two isomorphic Schottky groups $\rho: \Gamma \rightarrow \Gamma'$ in $\mathrm{PSL}(2, \mathbb{R})$. Suppose that $\ell(\gamma) > \ell(\rho(\gamma))$ for each $\gamma \in \Gamma$ (see [5] for the construction of such groups). Consider the isomorphism $\varphi: \Gamma \rightarrow \Gamma_\rho$ defined by $\varphi(\gamma) = (\gamma, \rho(\gamma))$. The groups Γ, Γ_ρ are Zariski dense respectively in $\mathrm{PSL}(2, \mathbb{R})$ and $\mathrm{PSL}(2, \mathbb{R}) \times \mathrm{PSL}(2, \mathbb{R})$ and the condition 3) is satisfied but φ cannot be extended.

3. Proofs of Theorems A and B

In this section G and G' denote connected semisimple groups with trivial center and without compact factor. Such a group can be decomposed into a product of connected noncompact simple groups with trivial center.

Lemma 3.1. *Let Γ, Γ' be Zariski dense subgroups of G and G' . Suppose that φ is a surjective homomorphism between Γ and Γ' and set $\Gamma_\varphi = \{(\gamma, \varphi(\gamma)) / \gamma \in \Gamma\}$. The projections of the identity component of the Zariski closure of Γ_φ into G and G' are surjective.*

Proof. The Lie algebra \mathcal{G} of G can be decomposed into a direct sum of simple ideals $\mathcal{G} = \mathcal{F}_1 + \cdots + \mathcal{F}_n$. Moreover each ideal of \mathcal{G} is a direct sum of certain \mathcal{F}_i ([10] corollary II.6.3). Let G_i be the connected Lie subgroup in G associated to \mathcal{F}_i . Since G has trivial center, $G = G_1 \times \cdots \times G_n$. Let H be the identity component of the Zariski closure of Γ_φ . Denote p the projection of H into G and T_p its tangent map at identity. The image, \mathcal{F} , of the Lie algebra of H by T_p is a non trivial

subalgebra of \mathcal{G} normalized by Γ . Since Γ is Zariski dense, \mathcal{F} is an ideal and hence $\mathcal{F} = \mathcal{F}_{i_1} + \cdots + \mathcal{F}_{i_k}$, $k \leq n$. This implies that $p(H) = G_{i_1} \times \cdots \times G_{i_k}$. Since the index of H in the Zariski closure of Γ_φ is finite and Γ is Zariski dense, $p(H)$ is also Zariski dense. This proves that $k = n$ and thus that p is surjective. Since φ is surjective, the same argument holds for the projection of H into G' . \square

Proof of Theorem A. Denote H the identity component of the Zariski closure of Γ_φ and \mathcal{H} its Lie algebra. We want to prove that the projection p (resp. p') of H into G (resp. G') is injective. Let us first show that \mathcal{H} is semisimple. Consider its solvable radical $\mathcal{R} \subset \mathcal{H}$. The image of \mathcal{R} by the tangent map Tp of p at identity is normalized by Γ . Since Γ is Zariski dense in G , $Tp(\mathcal{R})$ is a solvable ideal. The semi simplicity of G implies that $Tp(\mathcal{R})$ is trivial. Since φ is surjective, the same argument holds for p' . This shows that \mathcal{R} is trivial. Fix a Cartan decomposition $\mathcal{H} = \mathcal{P}'' + \mathcal{T}''$ of \mathcal{H} , since $G \times G'$ is semisimple, there exists a Cartan decomposition $\mathcal{P} + \mathcal{T}$ of the Lie algebra of $G \times G'$ such that $\mathcal{P}'' \subset \mathcal{P}$ and $\mathcal{T}'' \subset \mathcal{T}$ ([10] VI exercise 8(i)). Choose a Weyl chamber $\mathcal{W} \subset \mathcal{P}''$ since $\mathcal{P}'' \subset \mathcal{P}$ one has $\mathcal{W} \subset \mathcal{A} \times \mathcal{A}'$ where \mathcal{A} and \mathcal{A}' are Cartan subalgebras of the Lie algebra $\mathcal{G}, \mathcal{G}'$ of G and G' . Let us analyze $\text{Ker } p$. This group is normalized by Γ' because φ is surjective. Since Γ' is Zariski dense and the center of G' is trivial, either $\text{Ker } p = \{\text{Id}\}$ or $\text{Ker } p$ is a normal non trivial Lie subgroup of G' . In the last case, denote \mathcal{I} the Lie algebra of the identity component of $\text{Ker } p$. One has $\mathcal{I} = \mathcal{I}'_1 + \cdots + \mathcal{I}'_p$ where \mathcal{I}'_j are noncompact simple ideals of \mathcal{G}' such that $\mathcal{G}' = \mathcal{I}'_1 + \cdots + \mathcal{I}'_k$ with $k \geq p$ ([10] corollary II.6.3). It follows that \mathcal{W} contains an element $a = (0, \omega) \in \mathcal{A} \times \mathcal{A}'$ with $\|\omega\| \neq 0$. Since $\Gamma_\varphi \cap H$ is Zariski dense in H , according to Benoist's theorem, the interior of its limit cone, $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$, relatively to \mathcal{W} , is not empty. Moreover $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ is included in $S = \{(u, u') \in \mathcal{A} \times \mathcal{A}' / \|u\| = \|u'\|\}$ because φ preserves the translation length and $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ is included in the image of the limit cone of $\Gamma_\varphi \cap H$ relatively to $\mathcal{A}^+ \times \mathcal{A}'^+$ by the Weyl group. Let $b = (u, u')$ an element of the interior of $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H) \subset \mathcal{W}$. One can suppose $\|u\| = \|u'\| = 1$. Since the interior of $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ in \mathcal{W} is not empty, the intersection of the plane generated by a and b with $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ contains an open disc. There is a contradiction with the fact that the intersection of this plane with S is the curve $\{\alpha a + \beta b / 2\alpha\beta \langle u', \omega \rangle + \alpha^2 \|\omega\|^2 = 0\}$. In conclusion p is injective. The same argument holds for p' , because φ is surjective. Applying the lemma 3.1, one obtains that p and p' are bijective. Consider now the projections q (resp. q') of the Zariski closure $\overline{\Gamma_\varphi}^Z$ of Γ_φ into G (resp. G'). The maps q and q' are surjective. Let us prove that they are injective. Take $g \in \text{Ker } q$, for any $h \in H$ one has $q(ghg^{-1}h^{-1}) = \text{Id}$. Since H is normalized by $\overline{\Gamma_\varphi}^Z$ and p is injective, $gh = hg$. Using the fact that p' is surjective one obtains $p'(g)g' = g'p'(g)$ for any $g' \in G'$. Because the center of G' is trivial, $g = \text{Id}$. The same argument also holds for p' . Consider the map $f = p' \circ p^{-1}$, it is a continuous isomorphism between G and G' whose restriction to Γ coincides with φ . \square

Proof of Theorem B. The proof is similar to the previous one. Let us just adapt the end of the proof of Theorem A, when we suppose that $\text{Ker } p$ is nontrivial. Under this assumption one obtains that \mathcal{W} contains an element $a = (0, \omega) \in \mathcal{A} \times \mathcal{A}$ with $\omega \neq 0$. Since $v(\gamma) = k(\gamma)v(\varphi(\gamma))$ for each $\gamma \in \Gamma$, the limit cone $\mathcal{L}^{\mathcal{A}^+ \times \mathcal{A}^+}(\Gamma_\varphi \cap H)$ is included in $T = \{(u, u') \in \mathcal{A}^+ \times \mathcal{A}^+ / u \text{ and } u' \text{ are colinear}\}$ and hence $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ is included in $\bigcup_{g \in \text{Weyl}} gT$ where Weyl is the Weyl group of $\mathcal{A} \times \mathcal{A}$. The interior of $\mathcal{L}^{\mathcal{W}}(\Gamma_\varphi \cap H)$ in \mathcal{W} is not empty according to Benoist's theorem. It follows that for some $g \in \text{Weyl}$, the interior I of $g(T)$ is not empty in \mathcal{W} . Let $b = (u, u') \in I$. Since $\text{rank } G \geq 2$ one can assume that u' is not colinear to u . The intersection of the plane P generated by a and b with I contains an open disc. There is a contradiction with the fact that the intersection of T with $g^{-1}(P)$ is a line. \square

4. Applications of Theorem A to the space of representations

Fix a connected semisimple Lie group G without compact factor and with trivial center, and a symmetric space X associated to G . A subgroup of G is said parabolic if it fix a point of the geometric boundary, ∂X , of X .

Proposition 4.1. *Let Γ be a nonparabolic subgroup of G and H the identity component of its identity component. If $H \neq G$ then H fix a totally geodesic submanifold $Y \subsetneq X$.*

Proof. We thank P. Eberlein for helpful arguments.

The group H is reductive or parabolic ([3] corollaire 3.3). The last case cannot happens because H is normalized by Γ which does not fix any point in ∂X . Let $H = ST$ be the Levi decomposition of H where S is a connected semisimple group and T is a torus, corresponding to the identity component of the center of H . If $T \neq \text{Id}$ there exists a flat totally geodesic submanifold $T \subset X$ such that T leaves F invariant and F/T is compact ([8]). Let C be the union of all totally geodesic submanifolds which are parallel to F . Then C is invariant under H and is isometric to $F \times N$ for some closed convex subset N of X ([7] proposition 1.6.7). The set C is a totally geodesic submanifold possible with boundary. Let Y be a complete totally geodesic submanifold of X with $\dim Y = \dim C$. Since H leaves C invariant and C contains an open subset of Y , the group H leaves Y invariant. Remark that $Y \neq X$, because Y contains an Euclidean factor. If $T = \{\text{Id}\}$ then H is semisimple, and there exists $x \in X$ such that Hx is a totally geodesic submanifold ([13] lemma 7.21). By the assumption $H \neq G$ hence $Hx \neq X$. \square

Let Γ be an abstract group and $\rho : \Gamma \rightarrow G$ be a faithful representation. One always supposes that the Zariski closure, H_ρ , of $\rho(\Gamma)$ is connected and that the representation ρ is nonparabolic (i.e. $\rho(\Gamma)$ is nonparabolic). In this case H_ρ is reductive (proof of proposition 4.1). Let $H_\rho = ST$ be the Levi decomposition

of H_ρ . The representation ρ is noncompact if S is a semisimple group without compact factor and with trivial center. Under this assumption H_ρ stabilizes a totally geodesic submanifold of X isometric to $N \times F$ where N is a symmetric space on which S acts transitively and F is a flat on which T acts by translation with compact quotient (proof of the proposition 4.1). Two faithful, nonparabolic and noncompact representations ρ and ρ' of Γ are equivalent if there exists an isometry f between $N \times F$ and $N' \times F'$ such that $f \circ \rho(\gamma) = \rho'(\gamma) \circ f$ for any $\gamma \in \Gamma$. If F and F' are empty, then ρ and ρ' are equivalent if and only if $\rho' \circ \rho^{-1}$ can be extended to a continuous isomorphism between S and S' ([7] proposition 3.9.11). Denote R_{fnpc}/\sim the space of faithful nonparabolic, noncompact representations of Γ into G , up to the equivalence relation. The following result is an application of Theorem A to the context of representations.

Proposition 4.2. *The map $L: R_{fnpc}/\sim \rightarrow \mathbb{R}^\Gamma$ defined by $L([\rho])(\gamma) = \ell(\rho(\gamma))$ is injective.*

Proof. Let $\rho_1, \rho_2 \in R_{fnpc}$. Suppose $L(\rho_1) = L(\rho_2)$. For $i = 1, 2$ set $\Gamma_i = \rho_i(\Gamma)$, $H_i = H\rho_i$ and $T_i = S_i T_i$.

a) Suppose $S_1 = S_2 = \{e\}$, then T_i acts by translation on the flat $(F_i, \langle \cdot \rangle_i)$ and F_i/T_i is compact. Let us identify $\rho_i(\gamma)$ with its translation vector. Choose a basis, $\rho_1(\gamma_1), \dots, \rho_1(\gamma_n)$ of F_1 , such a basis exists because Γ_1 is Zariski dense in T_1 . For $\gamma \in \Gamma$, write $\rho_1(\gamma) = \sum_{i=1}^n a_i \rho_1(\gamma_i)$ and $\rho_2(\gamma) = \sum_{i=1}^n b_i \rho_2(\gamma_i) + \omega$ where ω is orthogonal to each $\rho_2(\gamma_i)$. Since $\|\rho_1(\gamma)\| = \|\rho_2(\gamma)\|$, one has $\langle \rho_1(\gamma), \rho_1(\gamma') \rangle_1 = \langle \rho_2(\gamma), \rho_2(\gamma') \rangle_2$ for any $\gamma, \gamma' \in \Gamma$. Put $c_{ij} = \langle \rho_1(\gamma_i), \rho_1(\gamma_j) \rangle_1 = \langle \rho_2(\gamma_i), \rho_2(\gamma_j) \rangle_2$. One has $\langle \rho_1(\gamma), \rho_1(\gamma_j) \rangle_1 = \sum_{i=1}^n a_i c_{ij}$ and $\langle \rho_2(\gamma), \rho_2(\gamma_j) \rangle_2 = \sum_{i=1}^n b_i c_{ij}$ hence $\sum_{i=1}^n (a_i - b_i) c_{ij} = 0$ for any $1 \leq j \leq n$. This proves that $a_i = b_i$. Moreover $\|\rho_1(\gamma)\| = \|\rho_2(\gamma)\|$ hence $\omega = 0$. One thus obtains $\rho_2(\gamma) = \sum_{i=1}^n a_i \rho_2(\gamma_i)$ and $\dim F_2 = n$ because Γ_2 is Zariski dense in T_2 . The linear map $f: F_1 \rightarrow F_2$ defined by $f(\rho_1(\gamma_i)) = \rho_2(\gamma_i)$ is an isometry satisfying $f \circ \rho_1(\gamma) = \rho_2(\gamma) \circ f$, hence $[\rho_1] = [\rho_2]$.

b) Suppose $S_1 \neq \{e\}$, then $S_2 \neq \{e\}$. Decompose S_i into a product of noncompact simple factors with trivial center $S_i = S_{i1} \times \dots \times S_{ik_i}$ and denote p_{is} the projection of S_i into S_{is} . Since Γ_i is Zariski dense in $S_i \times T_i$ then $p_{is}(\Gamma)$ is Zariski dense in S_{is} . Set $D = [\Gamma, \Gamma]$ and $D_i = \rho_i(D)$. The group D_i is normalized by Γ_i and is included in S_i , hence one can suppose that the Zariski closure of D_i equals $S_{i1} \times \dots \times S_{in_i}$ with $n_i \leq k_i$. Moreover $n_i = k_i$ because $p_{is}(D_i)$ is normalized by $p_{is}(\Gamma)$ which is Zariski dense in S_{is} and the center of S_{is} is trivial. In conclusion D_i is Zariski dense in S_i . By assumption $\ell(\rho_1(d)) = \ell(\rho_2(d))$ for any $d \in D$. One deduces from Theorem A that the restriction of $\rho_2 \circ \rho_1^{-1}$ to D_1 can be extended to a continuous isomorphism φ between S_1 and S_2 . Up to φ , one can suppose $S_1 = S_2$ and $\rho_1(d) = \rho_2(d)$ for any $d \in D$. Let $\gamma \in \Gamma$, since $\rho_1(\gamma d \gamma^{-1}) = \rho_2(\gamma d \gamma^{-1})$ and $\rho_1(d) = \rho_2(d)$, the projection of $\rho_2^{-1}(\gamma) \rho_1(\gamma)$ into S_1 commutes with all $\rho_1(d)$. Since D_1 is Zariski dense and the center of S_1 is trivial, the projection of $\rho_2^{-1}(\gamma) \rho_1(\gamma)$ into S_1 is trivial. Consider now the projection p_i of

Γ_i into T_i . One has $\ell(p_1 \circ \rho_1(\gamma)) = \ell(p_2 \circ \rho_2(\gamma))$, moreover $p_i(\Gamma_i)$ is Zariski dense in T_i . Using arguments developed in a), one obtains the existence of a isometry $f : F_1 \rightarrow F_2$ such that $f \circ (p_1 \circ \rho_1(\gamma)) = p_2 \circ \rho_2(\gamma) \circ f$, hence $[\rho_1] = [\rho_2]$. \square

The following part is inspired by the section 5 of A. Parreau's thesis ([15]). Let us consider the particular case where Γ is an infinite group of finite type. Fix a finite set, S , of generators. One associates to a representation $\rho : \Gamma \rightarrow G$ its minimal displacement, $\lambda(\rho) = \inf_{x \in X} (\sup_{s \in S} d(x, \rho(s)(x)))$. If $\lambda(\rho) = 0$ there exists a sequence $(x_n)_{n \geq 1}$ in X such that $\lim_n d(x_n, \rho(s)(x_n)) = 0$ for any $s \in S$. Up to a subsequence one can suppose that $(x_n)_{n \geq 1}$ converges in $X \cup \partial X$. If $\lim_n x_n = x \in X$ then $\rho(s)(x) = x$ for any $s \in S$ and hence $\rho(\Gamma)$ belongs to a compact subgroup. Otherwise $\lim_n x_n = \xi \in \partial X$ and $\rho(s)(\xi) = \xi$ for any $s \in S$. In this case ρ is parabolic. In conclusion, if $\rho \in R_{fnpnc}$ then $\lambda(\rho) > 0$. Let us consider the map $\frac{V}{\lambda} : R_{fnpnc} / \sim \rightarrow \mathbb{R}^\Gamma$ defined by $L([\rho])(\gamma) = \frac{\ell(\rho(\gamma))}{\lambda(\rho)}$. This map is continuous ([15] propositions V.2.3 and V.3.8) and its image is included in a compact set ([15] proposition V.4.1). One deduces from these properties and from the proposition 4.2 the following result.

Corollary 4.3. *The map $\frac{L}{\lambda} : R_{fnpnc} / \sim \rightarrow \mathbb{R}^\Gamma$ is injective, continuous and its image is included in a compact set.*

Acknowledgments. The authors deeply thank Y. Benoist and P. Eberlein for numerous discussions. The second author wants to thank G. Besson for his hospitality and inspiring discussions during his visit to Institut Fourier.

References

- [1] Y. Benoist, Propriétés asymptotiques des groupes linéaires, *GAF* **7** (1997), 1–47.
- [2] G. Besson, G. Courtois and S. Gallot, Entropies et rigidités des espaces localement symétriques de courbure strictement négative, *GAF* **5** (1997), 731–799
- [3] A. Borel and J. Tits, Eléments unipotents et sous-groupes paraboliques de groupes réductifs. *Inv. math.* **12** (1971), 95–104.
- [4] M. Burger, Intersection, the Manhattan curve and Patterson–Sullivan theory in Rank 2, *International Mathematics Research Notices* **7** (1993), 217–225.
- [5] F. Dal'Bo, Famille de groupes agissant sur le produit de deux variétés de Hadamard, *Séminaire de théorie spectrale et géométrie de Grenoble* (1997).
- [6] F. Dal'Bo and I. Kim, A criterion of conjugacy for Zariski dense subgroups, *CRAS* t. 330, série I (2000), 647–650.
- [7] P. Eberlein, Geometry of nonpositively curved manifolds, *University of Chicago Press*, 1996.
- [8] D. Gromoll and J. Wolf, Some relations between the metric structure and the algebraic structure of the fundamental group in manifolds of nonpositive curvature, *Bull. Amer. Math. Soc.* **77** (1971), 545–552.

- [9] U. Hamenstadt, Cocycles, symplectic structures and intersection, *GAF* **9** (1) (1999), 90–140.
- [10] S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Academic Press, 1978.
- [11] I. Kim, Marked length rigidity of rank one symmetric spaces, *Topology*, **40** n°6 (2001), 1295–1323.
- [12] I. Kim, Ergodic theory and rigidity on the symmetric space of noncompact type, *Ergodic theory and dynamical systems*, **21** n° 1 (2001), 93–114.
- [13] G. Margulis, Discrete subgroups of semisimple Lie group, Springer-Verlag, 1991.
- [14] J.-P. Otal, Le spectre marqué des surfaces à courbure négative, *Ann. of Math. (2)* **131** (1990), 151–162.
- [15] A. Parreau, Dégénérescences de sous-groupes discrets de groupes de Lie semisimples et actions de groupes sur des espaces affines, Thèse, Orsay, 2000.
- [16] J. H. Sampson, Sous-groupes conjugués d'un groupe linéaire, *Ann. Inst. Fourier* **26** (2) (1976), 1–6.

Françoise Dal'Bo
Université de Rennes 1
Institut de Recherche Mathématique de Rennes
Campus de Beaulieu
35042 Rennes cedex
France
e-mail: dalbo@univ-rennes1.fr

Inkang KIM
Department of Mathematics
Seoul National University
151-741 Seoul
Korea
e-mail: inkang@math.snu.ac.kr

(Received: September 18, 2001)



To access this journal online:
<http://www.birkhauser.ch>
