Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	78 (2003)
Artikel:	Classification of graded Hecke algebras for complex reflection groups
Autor:	Ram, Arun / Shepler, Anne V.
DOI:	https://doi.org/10.5169/seals-58758

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Comment. Math. Helv. 78 (2003) 308–334 0010-2571/03/020308-27

© 2003 Birkhäuser Verlag, Basel

Commentarii Mathematici Helvetici

Classification of graded Hecke algebras for complex reflection groups

Arun Ram and Anne V. Shepler

Abstract. The graded Hecke algebra for a finite Weyl group is intimately related to the geometry of the Springer correspondence. A construction of Drinfeld produces an analogue of a graded Hecke algebra for any finite subgroup of GL(V). This paper classifies all the algebras obtained by applying Drinfeld's construction to complex reflection groups. By giving explicit (though non-trivial) isomorphisms, we show that the graded Hecke algebras for finite real reflection groups constructed by Lusztig are all isomorphic to algebras obtained by Drinfeld's construction. The classification shows that there exist algebras obtained from Drinfeld's construction which are not graded Hecke algebras as defined by Lusztig for real as well as complex reflection groups.

Mathematics Subject Classification (2000). 20C, 20F, 52B.

Keywords. Reflection group, Coxeter group, Weyl group, affine Hecke algebra, Iwahori–Hecke algebra, representation theory, graded Hecke algebra.

0. Introduction

This paper is motivated by a general effort to generalize the theory of Weyl groups and their relation to groups of Lie type to the setting of complex reflection groups. One natural question is whether there are affine Hecke algebras corresponding to complex reflection groups. If they exist then it might be possible to use these algebras to build an analogue of the Springer correspondence for complex reflection groups.

A priori, one knows how to construct affine Hecke algebras corresponding only to Weyl groups since both a finite real reflection group W and a W-invariant lattice (the existence of which forces W to be a Weyl group) are needed in the construction. Our search for analogues of graded Hecke algebras for complex reflection groups was motivated by Lusztig's results [Lu2] showing that the geometric

Research of the first author supported in part by the National Security Agency and by EPSRC Grant GR K99015 at the Newton Institute for Mathematical Sciences. Research of the second author supported in part by National Science Foundation grant DMS-9971099.

Graded Hecke algebras

information contained in the affine Hecke algebra can be recovered from the corresponding graded Hecke algebra. Lusztig [Lu] defines the graded Hecke algebra for a finite Weyl group W with reflection representation V. Let $t_g, g \in W$, be a basis for the group algebra $\mathbb{C}W$ of W and let $k_\alpha \in \mathbb{C}$ be "parameters" indexed by the roots in the root system of W such that k_α depends only on the length of the root α . Then the graded Hecke algebra $H_{\rm gr}$ depending on the parameters k_α is the (unique) algebra structure on $S(V) \otimes \mathbb{C}W$ such that

- (a) the symmetric algebra of $V, S(V) = S(V) \otimes 1$, is a subalgebra of $H_{\rm gr}$,
- (b) the group algebra $\mathbb{C}W = 1 \otimes \mathbb{C}W = \text{span}{\{1 \otimes t_g \mid g \in W\}}$ is a subalgebra of H_{gr} , and
- (c) $t_{s_i}v = (s_iv)t_{s_i} k_{\alpha_i}\langle v, \alpha_i^{\vee} \rangle$ for all $v \in V$ and simple reflections s_i in the simple roots α_i .

This definition applies to all finite real reflection groups W since the simple roots and simple reflections are well defined. Unfortunately, the need for simple reflections in the construction makes it unclear how to define analogues for complex reflection groups.

For finite real reflection groups, the graded Hecke algebra H_{gr} is a "semidirect product" of the polynomial ring S(V) and the group algebra $\mathbb{C}W$. Drinfeld [Dr] defines a *different* type of semidirect product of S(V) and $\mathbb{C}W$, and Drinfeld's construction applies to all finite subgroups G of GL(V). In this paper, we

- (1) classify all the algebras obtained by applying Drinfeld's construction to finite complex reflection groups G,
- (2) show that every graded Hecke algebra $H_{\rm gr}$ (as defined by Lusztig) for a finite real reflection group is isomorphic to an algebra obtained by Drinfeld's construction by giving explicit isomorphisms between these algebras.

The results from (2) show how Drinfeld's construction is a true generalization of Lusztig's construction of graded Hecke algebras, something which is not obvious. Our classification in (1) gives a complete solution to the problem of finding all graded Hecke algebras for finite reflection groups.

A consequence of our classification is that there exist graded Hecke algebras for finite real reflection groups which are not obtained with Lusztig's construction. In this sense, Drinfeld's construction is a strict generalization of the algebras $H_{\rm gr}$. These new algebras, and the algebras corresponding to complex reflection groups that are not real reflection groups, deserve further study and probably have interesting representation theories.

For us, one surprising result of our classification is that no nontrivial graded Hecke algebra structures exist for many complex reflection groups. In some sense, this is disappointing, as we would have liked to find nontrivial and interesting structures for *each* complex reflection group.

It might be that we have not yet hit upon the "right" definition of graded Hecke algebras. For example, we show that there do not exist nontrivial graded Hecke algebra structures, according to Drinfeld's definition, for any of the complex reflection groups $G(r, 1, n) = (\mathbb{Z}/r\mathbb{Z}) \wr S_n$ when r > 2 and n > 3. On the other A. Ram and A. V. Shepler

hand, in the last section of this paper we are able to construct algebras that "look" like they ought to be graded Hecke algebras corresponding to these groups. Is it possible that there is a "better" definition of graded Hecke algebras which applies to complex reflection groups and which includes the algebras that we introduce in Section 5 as examples?

Acknowledgements. We thank C. Kriloff for numerous stimulating conversations during our work on this paper. A. Ram thanks the Newton Institute for the Mathematical Sciences at Cambridge University for hospitality and support (EPSRC Grant No. GR K99015) during Spring 2001 when the writing of this paper was completed.

1. Graded Hecke algebras

In this section, we define the graded Hecke algebra following Drinfeld [Dr]. Our main result in this section is Theorem 1.9c, which determines exactly how many degrees of freedom one has in defining a graded Hecke algebra.

Let V be an n dimensional vector space over \mathbb{C} and let G be a finite subgroup of GL(V). The group algebra of G is

$$\mathbb{C}G = \mathbb{C}\operatorname{-span}\{t_g \mid g \in G\}, \quad \text{with} \quad t_g t_h = t_{gh}.$$

Let $a_g: V \times V \longrightarrow \mathbb{C}$ be skew symmetric bilinear forms indexed by the elements of G and let A be the associative algebra generated by V and $\mathbb{C}G$ with the additional relations

$$t_h v t_{h^{-1}} = h v$$
 and $[v, w] = \sum_{g \in G} a_g(v, w) t_g$, for $h \in G$ and $v, w \in V$,
(1.1)

where [v, w] = vw - wv. These relations allow every element $a \in A$ to be written in the form

$$a = \sum_{g \in G} p_g t_g, \qquad p_g \in S(V), \tag{1.2}$$

where S(V) is the symmetric algebra of V. More precisely, one must fix a section of the canonical surjection $T(V) \to S(V)$ from the tensor algebra of V to S(V) and take the elements p_g to be in the image of this section.

The structure of A depends on the choices of the "parameters" $a_g(v, w) \in \mathbb{C}$. Our goal is to determine when the algebra A will be a "semidirect product" of S(V) and $\mathbb{C}G$. This idea motivates the following definition [Dr].

The algebra A is a graded Hecke algebra for G if

$$A \cong S(V) \otimes \mathbb{C}G$$
 as a vector space,

or, equivalently, if the expression in (1.2) is unique for each $a \in A$. A general element $a \in A$ is a linear combination of products of elements t_g and u_i , where $\{u_1, u_2, \ldots, u_n\}$ is a basis of V. There are two straightening operations needed to put a in the form (1.2):

(a) moving t_h 's to the right, and (b) putting $u_i u_j$ pairs in the correct order.

These two straightening operations correspond to the two identities in (1.1). Note that the "correct order" of $u_i u_j$ is determined by the choice of the section of the projection $T(V) \to S(V)$. Let v_1, v_2, v_3 be arbitrary elements of V and let $h \in G$. Applying the straightening operations to $t_h v_1 v_2$ gives

$$\begin{aligned} t_h v_1 v_2 &= t_h [v_1, v_2] + t_h v_2 v_1 & (\text{rearrange } v_1 \text{ and } v_2) \\ &= t_h [v_1, v_2] + (h v_2) (h v_1) t_h & (\text{move } t_h \text{ to the right}), \end{aligned}$$

and applying the straightening operations in a different order gives

$$\begin{aligned} t_h v_1 v_2 &= (hv_1)(hv_2)t_h \\ &= [hv_1, hv_2]t_h + (hv_2)(hv_1)t_h. \end{aligned} \qquad (\text{move } t_h \text{ to the right}) \end{aligned}$$

Setting these two expressions equal gives the relation

$$t_h[v_1, v_2]t_{h^{-1}} = [hv_1, hv_2], \quad \text{for all } h \in G, \, v_1, v_2 \in V.$$
(1.3)

Similarly, applying the straightening operations to $v_1v_2v_3$ gives

$v_1v_2v_3=[v_1,v_2]v_3+v_2v_1v_3$	(moving v_1 past v_2)
$= [v_1, v_2] v_3 + v_2 [v_1, v_3] + v_2 v_3 v_1 \\$	(moving v_1 past v_3)
$= [v_1, v_2]v_3 + v_2[v_1, v_3] + [v_2, v_3]v_1 + v_3v_2v_1$	(straightening v_2 and v_3),

and applying the straightening operations in a different order gives

$v_1v_2v_3=v_1[v_2,v_3]+v_1v_3v_2$	(moving v_2 past v_3)
$= v_1[v_2,v_3] + [v_1,v_3]v_2 + v_3v_1v_2 \\$	(moving v_1 past v_3)
$= v_1[v_2, v_3] + [v_1, v_3]v_2 + v_3[v_1, v_2] + v_3v_2v_1$	(straightening v_1 and v_2).

These are equal if

 $[v_1, [v_2, v_3]] + [v_2, [v_3, v_1]] + [v_3, [v_1, v_2]] = 0, \qquad \text{for all } v_1, v_2, v_3 \in V. \tag{1.4}$

Conversely, the identities (1.3) and (1.4) are exactly what is needed to guarantee that any order of application of the straightening operations (a) and (b) will produce the same normal form (1.2) for the element *a*. Thus we have

Lemma 1.5. Let A be an algebra defined as in (1.1). Then A is a graded Hecke algebra if and only if the identities (1.3) and (1.4) hold in A.

Using (1.1), the relations (1.3) and (1.4) can be rewritten in terms of the bilinear forms $a_g: V \times V \to \mathbb{C}$ as

$$a_g(v_1, v_2) = a_{hgh^{-1}}(hv_1, hv_2)$$
 and (1.6)

$$a_g(v_3, v_1)(gv_2 - v_2) + a_g(v_2, v_3)(gv_1 - v_1) + a_g(v_1, v_2)(gv_3 - v_3) = 0$$
(1.7)

for $v_1, v_2, v_3 \in V$ and $g, h \in G$.

Let $\langle \ ,\ \rangle:V\times V\to \mathbb{C}$ be a G-invariant nondegenerate Hermitian form on V. For each $g\in G,$ set

$$V^{g} = \{ v \in V \mid gv = v \},$$

$$(V^{g})^{\perp} = \{ v \in V \mid \langle v, w \rangle = 0 \text{ for all } w \in V^{g} \}, \text{ and }$$

$$\ker a_{g} = \{ v \in V \mid a_{g}(v, w) = 0 \text{ for all } w \in V \}.$$

Lemma 1.8. Let G be a finite subgroup of GL(V) and let $g \in G$.

(a) $(V^g)^{\perp} = \{ v - gv \mid v \in V \}.$

(b) Suppose $g \neq 1$. If $\operatorname{codim}(V^g) = 2$ and $a: V \times V \to \mathbb{C}$ is a skew symmetric bilinear form such that ker $a = V^g$, then a satisfies (1.7).

Let A be a graded Hecke algebra for G defined by skew symmetric bilinear forms $a_g: V \times V \to \mathbb{C}$.

(c) If $g \neq 1$ then ker $a_g \supseteq V^g$.

(d) If $g \neq 1$ and $a_g \neq 0$ then ker $a_g = V^g$ and $\operatorname{codim}(V^g) = 2$.

(e) If $g \neq 1$ and $a_g \neq 0$ then, for all $h \in G$,

$$a_{h^{-1}gh}(b_1, b_2) = \det(h^{\perp})a_g(b_1, b_2),$$

where $\{b_1, b_2\}$ is a basis of $(V^g)^{\perp}$ and $h^{\perp}: (V^g)^{\perp} \to (V^g)^{\perp}$ is the composition of h restricted to $(V^g)^{\perp}$ with the canonical projection $V \to V/V^g$.

Proof. (a) Consider the map $\phi: V \to V$ given by $\phi(v) = v - gv$. Then ker $\phi = V^g$ and im $\phi \subseteq (V^g)^{\perp}$ since, if $v \in V, w \in V^g$, then

$$\langle v - gv, w
angle = \langle v, w
angle - \langle gv, w
angle = \langle v, w
angle - \langle gv, gw
angle = \langle v, w
angle - \langle v, w
angle.$$

Since dim(im ϕ) = codim(ker ϕ) = codim(V^g) it follows that im $\phi = (V^g)^{\perp}$.

(b) Let $v_1, v_2, v_3 \in V$. If any $v_i \in V^g$, then (1.7) holds trivially for the skew symmetric form a. So assume each $v_i \notin V^g$ and write each v_i as $v_i^+ + v_i^-$ where $v_i^+ \in V^g$ and $v_i^- \in (V^g)^{\perp}$. Then

$$a(v_i,v_j)=a(v_i^-,v_j^-) \qquad ext{and} \qquad v_i-gv_i=v_i^--gv_i^-.$$

Since $\dim(V^g)^{\perp} = 2$, at least one of the v_i^- is a linear combination of the other two. Say $v_1^- = c_2 v_2^- + c_3 v_3^-$ with $c_2, c_3 \in \mathbb{C}$. Substituting $v_i - gv_i = v_i^- - gv_i^-$ and $v_1^- = c_2 v_2^- + c_3 v_3^-$ then yields

$$\begin{split} &a(v_3,v_1)(gv_2-v_2)+a(v_2,v_3)(gv_1-v_1)+a(v_1,v_2)(gv_3-v_3)\\ &=a(v_3^-,v_1^-)(gv_2^--v_2^-)+a(v_2^-,v_3^-)(gv_1^--v_1^-)+a(v_1^-,v_2^-)(gv_3^--v_3^-)=0, \end{split}$$

Graded Hecke algebras

and so (1.7) holds.

(c) Let $v_3 \in V^g$ and $v_2 \in V$.

If $v_2 \in V^g$, then $a_g(v_2, v_3)(gv_1 - v_1) = 0$ for all $v_1 \in V$ by (1.7). Since $V^g \neq V$, there is some v_1 such that $gv_1 \neq v_1$ and so $a_g(v_2, v_3) = 0$.

If
$$v_2 \notin V^g$$
, let $v_1 = \sum_{k=1}^{r-1} g^k v_2$, where r is the order of g. By (1.6), $a_g(v_3, g^k v_2) = a_g(g^{-k}v_3, v_2) = a_g(v_3, v_2)$, for any k, and so

$$\begin{split} 0 &= a_g(v_3,v_1)(gv_2-v_2) + a_g(v_2,v_3)(gv_1-v_1) \\ &= (r-1)a_g(v_3,v_2)(gv_2-v_2) + a_g(v_3,v_2)(gv_2-v_2) = ra_g(v_3,v_2)(gv_2-v_2). \end{split}$$

Thus $a_g(v_3, v_2) = 0$. Hence, for all $v_2 \in V$, $a_g(v_3, v_2) = 0$ and so $V^g \subseteq \ker a_g$.

(d) By (c), $\operatorname{codim}(V^g) \geq \operatorname{codim}(\ker a_g)$. Since $a_g \neq 0$, there exist $v, w \in V$ with $a_g(v, w) \neq 0$ and so $\operatorname{codim}(\ker a_g) \geq 2$. Let $v_1 - gv_1$ and $v_2 - gv_2$ be linearly independent elements of $(V^g)^{\perp}$. Then (1.7) implies that any element $v_3 - gv_3 \in (V^g)^{\perp}$ is a linear combination of $v_1 - gv_1$ and $v_2 - gv_2$, and so

$$2 \ge \dim((V^g)^{\perp}) = \operatorname{codim}(V^g) \ge \operatorname{codim}(\ker a_g) \ge 2.$$

Thus $V^g = \ker a_g$ and $\operatorname{codim}(V^g) = 2$.

(e) Write $hb_1 = h_{11}b_1 + h_{21}b_2 + (hb_1)^g$ and $hb_2 = h_{12}b_1 + h_{22}b_2 + (hb_2)^g$ with $h_{ij} \in \mathbb{C}$ and $(hb_i)^g \in V^g$. Then

$$\begin{split} a_{h^{-1}gh}(b_1,b_2) &= a_g(hb_1,hb_2) \\ &= a_g(h_{11}b_1 + h_{21}b_2 + (hb_1)^g,h_{12}b_1 + h_{22}b_2 + (hb_2)^g) \\ &= (h_{11}h_{22} - h_{21}h_{12})a_g(b_1,b_2) = \det(h^{\perp})a_g(b_1,b_2) \end{split}$$

since a_g is skew symmetric and $V^g \subseteq \ker a_g$.

The following theorem is a slightly strengthened version of statements (given without proof) in [Dr].

Theorem 1.9. Let G be a finite subgroup of GL(V) and let $Z_G(g) = \{h \in G \mid hg = gh\}$ denote the centralizer of an element g in G.

(a) If A is a graded Hecke algebra for G, then the values of $a_{h^{-1}gh}$ are determined by the values of a_g via the equation

$$a_{h^{-1}ah}(v_1, v_2) = a_g(hv_1, hv_2), \quad \text{for all } g, h \in G, v_1, v_2 \in V.$$

- (b) For $g \neq 1$, there is a graded Hecke algebra A with $a_g \neq 0$ if and only if
 - $\ker a_g = V^g$, $\operatorname{codim}(V^g) = 2$, and $\det(h^{\perp}) = 1$, for all $h \in Z_G(g)$,

A. Ram and A. V. Shepler

where h^{\perp} is h restricted to the space $(V^g)^{\perp}$. In this case, a_g is determined by its value $a_g(b_1, b_2)$ on a basis $\{b_1, b_2\}$ of $(V^g)^{\perp}$.

 (c) Let d be the number of conjugacy classes of g ∈ G such that codim(V^g) = 2 and det(h[⊥]) = 1 for all h ∈ Z_G(g), where h[⊥] is h restricted to the space (V^g)[⊥]. The sets {a_g}_{g∈G} corresponding to graded Hecke algebras A form a vector space of dimension d + dim((Λ² V)^G).

Proof. (a) is simply a restatement of (1.6).

(b) \implies : If A is a graded Hecke algebra and $a_g \neq 0$ then by Lemma 1.8d, $\operatorname{codim}(V^g) = 2$ and $\ker a_g = V^g$. So a_g is determined by its value $a_g(b_1, b_2)$ on a basis b_1, b_2 of $(V^g)^{\perp}$. Suppose $h \in Z_G(g)$. Then, by Lemma 1.8e,

$$a_g(b_1, b_2) = a_{hgh^{-1}}(hb_1, hb_2) = a_g(hb_1, hb_2) = \det(h^{\perp}) \ a_g(b_1, b_2),$$

and so det $(h^{\perp}) = 1$. Note that $h(V^g) = V^g$ and $h(V^g)^{\perp} = (V^g)^{\perp}$ since, for each $v \in V^g$, h(v) = hg(v) = gh(v).

 \Leftarrow : If $\operatorname{codim}(V^g) = 2$ then, up to constant multiples, there is a unique skew symmetric form on V which is nondegenerate on $(V^g)^{\perp}$ and which has ker $a_g = V^g$. Fix such a form and then define forms $a_h, h \in G$, by

$$a_h(v_1, v_2) = \begin{cases} a_g(kv_1, kv_2) & \text{if } h = k^{-1}gk, \\ 0 & \text{otherwise,} \end{cases}$$
(1.10)

for $v_1, v_2 \in V$. Let a_1 be any *G*-invariant skew symmetric form on *V*. Then this collection $\{a_g\}_{g\in G}$ of skew symmetric bilinear forms satisfies (1.6) by definition and (1.7) by Lemma 1.8b. Thus (by Lemma 1.5), it determines a graded Hecke algebra *A* via (1.1).

(c) From (a) and (b) it follows that the sets $\{a_g\}_{g\in G}$, running over all graded Hecke algebras A for G, form a vector space. Since each of the collections $\{a_g\}_{g\neq 1}$ constructed by (1.10) has its support on a single conjugacy class, these collections form a basis of the vector space of sets $\{a_g\}_{g\neq 1}$. The only condition on the form a_1 is that it satisfies (1.6), which means that it is a G-invariant element of $(\bigwedge^2 V)^*$.

The following consequence of Theorem 1.9 will be useful for completing the classification of graded Hecke algebras for complex reflection groups.

Corollary 1.11. Assume that G contains $h = \xi \cdot 1$ for some $\xi \in \mathbb{C} \setminus \{\pm 1\}$. If A is a graded Hecke algebra for G, then $a_q = 0$ for all $g \neq 1$.

Proof. If $h = \xi \cdot 1 \in G$, then $h \in Z_G(g)$ for every $g \in G$ and $\det(h^{\perp}) = \xi^2$ if $\operatorname{codim}(V^g) = 2$. The statement then follows from Theorem 1.9b.

2. The classification for reflection groups

A reflection is an element of GL(V) that has exactly one eigenvalue not equal to 1. The reflecting hyperplane of a reflection is the (n-1)-dimensional subspace which is fixed pointwise. A complex reflection group G is a finite subgroup of GL(V)generated by reflections. The group G is *irreducible* if V cannot be written in the form $V = V_1 \oplus V_2$ where V_1 and V_2 are G-invariant subspaces. The group G is a real reflection group if $V = \mathbb{C} \otimes_{\mathbb{R}} V_{\mathbb{R}}$ for a real vector space $V_{\mathbb{R}}$ and $G \subseteq GL(V_{\mathbb{R}})$.

The following facts about reflection groups are well known.

Lemma 2.1. Let G be an irreducible reflection group.

- (a) [ST, Theorem 5.3] The number of elements $g \in G$ such that $\operatorname{codim}(V^g) = 2$ is $\sum_{i < j} m_i m_j$ where m_1, \ldots, m_n are the exponents of G.
- (b) [Ca, Lemma 2] If G is a real reflection group and $g \in G$ with $\operatorname{codim}(V^g) = 2$, then g is the product of two reflections.
- (c) [OT, Theorem 6.27] For any $g \in G$, the space V^g is the intersection of reflecting hyperplanes.

Remark. The statement of Lemma 2.1b does not hold for complex reflection groups. Consider the exceptional complex reflection group G_4 of rank 2, in the notation of Shephard and Todd [ST]. All the reflections have order 3 and $-1 \in G_4$. Suppose -1 = rs for two reflections r and s. If s has eigenvalues 1 and ω , where ω is a primitive cube root of unity, then $r^{-1} = -s$ has eigenvalues -1 and $-\omega$, a contradiction to the assumption that r is a reflection. Thus $-1 \in G_4$ is not a product of two reflections.

Lemma 2.2. Let $G \subseteq GL(V)$ be a complex reflection group. Let A be a graded Hecke algebra for G and let $g \in G$. Let $V^G = \{v \in V \mid gv = v \text{ for all } g \in G\}$ be the invariants in V.

- (a) If g = 1 and dim $V^G \leq 1$, then $a_g = 0$.
- (b) If the order of g is 2, then $a_g = 0$.

Proof. (a) Let $\langle , \rangle : V \times V \to \mathbb{C}$ be a nondegenerate *G*-invariant Hermitian form on V and write $V = V^G \oplus (V^G)^{\perp}$ where $(V^G)^{\perp} = \{v \in V \mid \langle v, w \rangle = 0 \text{ for all } w \in V^G\}$. Since dim $(V^G) \leq 1$ and a_1 is skew symmetric, a_1 restricted to V^G is 0. There is a basis $\alpha_1, \ldots, \alpha_k$ of $(V^G)^{\perp}$ and constants $\xi_1, \ldots, \xi_k \in \mathbb{C}, \ \xi_i \neq 1$, such that the reflections s_1, \ldots, s_k given by

$$s_i v = v + (\xi_i - 1) \frac{\langle v, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle} \alpha_i, \quad \text{for } v \in V,$$

are in G. Equation (1.6) implies that, for any $v \in V$,

$$a_1(\alpha_i, v) = a_1(s_i\alpha_i, s_iv) = a_1\left(\xi_i\alpha_i, v + (\xi_i - 1)\frac{\langle v, \alpha_i \rangle}{\langle \alpha_i, \alpha_i \rangle}\alpha_i\right) = \xi_i a_1(\alpha_i, v),$$

since $a_1(\alpha_i, \alpha_i) = 0$ (as a_1 is skew symmetric). Since $\xi_i \neq 1$, $a_1(\alpha_i, v) = 0$ for $1 \leq i \leq k$. Thus ker $a_1 = V$.

(b) Since $g^2 = 1$, all eigenvalues of g are ± 1 . If $\operatorname{codim}(V^g) \neq 2$, then $a_g = 0$ by Theorem 1.9b. If $\operatorname{codim}(V^g) = 2$, then

$$g = \mathrm{id}_{V^g} \oplus (-\mathrm{id}_{(V^g)^\perp})$$

as a linear transformation on V. By [St1, Theorem 1.5], [Bou V, §5 Ex. 8], the stabilizer, $\operatorname{Stab}(V^g)$, of V^g is a reflection subgroup of G and so there is a reflection $s \in \operatorname{Stab}(V^g)$ that is the identity on V^g . So $s \in Z_G(g)$ and $\det(s) = \det(s^{\perp}) \neq 1$, where s^{\perp} is s restricted to $(V^g)^{\perp}$. Thus, by Theorem 1.9b, $a_g = 0$.

2A. Real reflection groups

If $G \subseteq GL(V)$ is a real reflection group then $V = \mathbb{C} \otimes_{\mathbb{R}} V_{\mathbb{R}}$ and $G \subseteq GL(V_{\mathbb{R}})$, where $V_{\mathbb{R}}$ is a real vector space. We shall assume that G is irreducible.

Let us recall some basic facts about real reflection groups which can be found in [Hu] or [Bou]. The action of G on $V_{\mathbb{R}}$ has fundamental chambers wC indexed by $w \in G$. The *roots* for G are vectors $\alpha \in V_{\mathbb{R}}$ such that the reflections in G are the reflections s_{α} in the hyperplanes

$$H_{\alpha} = \{ v \in V_{\mathbb{R}} \mid \langle v, \alpha \rangle = 0 \}.$$

For each fundamental chamber C, the reflections s_1, s_2, \ldots, s_n in the hyperplanes $H_{\alpha_1}, H_{\alpha_2}, \ldots, H_{\alpha_n}$ that bound C form a set of simple reflections for G. The simple reflections obtained from a different choice of fundamental chamber wC are $ws_1w^{-1}, \ldots, ws_nw^{-1}$.

Theorem 2.3. Let $G \subseteq GL(V_{\mathbb{R}})$ be a real reflection group. Let s_1, \ldots, s_n be a set of simple reflections in G and let m_{ij} be the order of $s_i s_j$. Then $g \in G$ satisfies $g^2 \neq 1$, $\operatorname{codim}(V^g) = 2$, and $\det(h^{\perp}) = 1$ for all $h \in Z_G(g)$ (the conditions in Theorem 1.9c) if and only if g is conjugate to

$$(s_i s_j)^k$$
, with $0 < k < \frac{m_{ij}}{2}$,

for some $1 \leq i, j \leq n$.

Proof. \Longrightarrow : Let α and β be two roots such that $V^g = H_\alpha \cap H_\beta$ (see Lemma 2.1c). Then $H_\alpha \cap H_\beta$ has nontrivial intersection with some fundamental chamber C for W, and we may assume that H_α and H_β are walls of the chamber C (since C is a cone in \mathbb{R}^n). Since choosing simple reflections with respect to a different chamber wC corresponds to conjugation by w, we may assume that the reflections in the hyperplanes H_α and H_β are simple reflections and $\alpha = \alpha_1$ and $\beta = \alpha_2$.

Graded Hecke algebras

The element g is an element of the stabilizer $\operatorname{Stab}(V^g)$, which is a reflection group by [St1, Theorem 1.5]. Since $\operatorname{codim}(V^g) = 2$, $\operatorname{Stab}(V^g)$ is a rank two real reflection group, and therefore a dihedral group. This dihedral group is generated by the two simple reflections s_1 and s_2 in the hyperplanes H_{α_1} and H_{α_2} (restricted to $(V^g)^{\perp}$) and all reflections have determinant -1. Let g^{\perp} be the element g restricted to $(V^g)^{\perp}$. Since $g \in Z_G(g)$, $\det(g^{\perp}) = 1$, and so g must be a product of an even number of reflections. Thus $g = (s_1s_2)^k$ or $g = (s_2s_1)^k$, for some $0 < k \le m/2$, where m is the order of s_1s_2 . Since $g^2 \ne 1$, $k \ne m/2$, and so g is conjugate to $(s_1s_2)^k$ for some 0 < k < m/2.

 $\begin{array}{l} \Leftarrow : \text{Assume that } g = (s_i s_j)^k \text{ for some } 0 < k < m_{ij}/2. \text{ Then } V^g = H_{\alpha_i} \cap H_{\alpha_j} \\ \text{and so } \operatorname{codim}(V^g) = 2. \text{ Since } g \text{ is a product of an even number of reflections,} \\ \det(g^{\perp}) = 1. \text{ The only elements of } O(V_{\mathbb{R}}) \cong O_2(\mathbb{R}) \text{ that are diagonalizable in} \\ GL(V_{\mathbb{R}}) \cong GL_2(\mathbb{R}) \text{ are } \pm 1 \text{ and elements with determinant } -1. \text{ Thus, the eigenvectors of the element } g^{\perp} \text{ (which has distinct eigenvalues since it is not } \pm 1) \text{ do} \\ \text{not lie in } V_{\mathbb{R}}, \text{ only in } V = \mathbb{C} \otimes_{\mathbb{R}} \mathbb{R}. \text{ Let } h \in Z_G(g) \text{ and let } h^{\perp} \in O(V_{\mathbb{R}}) \cong O_2(\mathbb{R}) \\ \text{denote } h \text{ restricted to } (V^g)^{\perp}. \text{ Since } h^{\perp} \text{ commutes with } g^{\perp} \text{ and } g^{\perp} \text{ has distinct eigenvalues, } g^{\perp} \text{ and } h^{\perp} \text{ have the same eigenvectors. Hence, } \det h^{\perp} = 1. \end{array}$

Using Theorem 2.3 and Theorem 1.9b, we can read off the graded Hecke algebras for the irreducible real reflection groups from the Dynkin diagrams. For each irreducible real reflection group, label a set of simple reflections s_1, \ldots, s_n using the Dynkin diagrams below. If nodes i and j and nodes j and k are connected by single edges, then $s_i s_j$ is conjugate to $s_j s_k$ via the element $s_i s_j s_k$.

The following table gives representatives of the conjugacy classes of $g \in G$ that may have $a_g \neq 0$ for some graded Hecke algebra A. We assume that the reflection group G is acting on its irreducible reflection representation V. When G is the symmetric group S_n acting on an *n*-dimensional vector space V by permutation matrices, then dim $(V^G) = 1$ and, by Lemma 2.2a and Theorem 2.3, $a_g \neq 0$ for some graded Hecke algebra A only if g is conjugate to the three cycle $(1, 2, 3) = s_1 s_2$ (this example is analyzed in Section 3).

Type	$\begin{array}{c} \text{Representative} g \\ \text{with} a_g \neq 0 \end{array}$
A_{n-1}	$s_{1}s_{2}$
B_n	$s_1s_2, \ s_2s_3$
D_n	$s_{2}s_{3}$
E_{6}, E_{7}, E_{8}	s_1s_4
F_4	$s_1s_2, \ s_2s_3, \ s_3s_4$
H_3, H_4	$s_1s_2, \ (s_1s_2)^2, \ s_2s_3$
$I_2(m)$	$(s_1 s_2)^k, 0 < k < m/2$

Table 1. Graded Hecke algebras for real reflection groups



FIGURE 1. Coxeter-Dynkin diagrams for real reflection groups

2B. Complex reflection groups G(r, p, n)

The irreducible complex reflection groups were classified by Shephard and Todd [ST]. There is one infinite family denoted G(r, p, n) and a list of exceptional complex reflection groups denoted G_4, \ldots, G_{35} . In this subsection, we classify the graded Hecke algebras for the groups G(r, p, n).

Let r, p and n be positive integers with p dividing r and let $\xi = e^{2\pi i/r}$. Let S_n be the symmetric group of $n \times n$ matrices and let

$$\xi_j = \operatorname{diag}(1, 1, \ldots, 1, \xi, 1, \ldots, 1),$$

where ξ appears in the *j*th entry. Then

$$G(r, p, n) = \{\xi_1^{\lambda_1} \cdots \xi_n^{\lambda_n} w \mid w \in S_n, \ 0 \le \lambda_i \le r-1, \ \lambda_1 + \cdots + \lambda_n = 0 \bmod p\}.$$

For $\lambda = (\lambda_1, \ldots, \lambda_n) \in (\mathbb{Z}/r\mathbb{Z})^n$, let $\xi^{\lambda} = \xi_1^{\lambda_1} \cdots \xi_n^{\lambda_n}$. Then the multiplication in G(r, p, n) is described by the relations

$$\xi^{\lambda}\xi^{\mu} = \xi^{\lambda+\mu}$$
 and $w\xi^{\lambda} = \xi^{w\lambda}w$, for $\lambda, \mu \in (\mathbb{Z}/r\mathbb{Z})^n$, $w \in S_n$,

where S_n acts on $(\mathbb{Z}/r\mathbb{Z})^n$ by permuting the factors. Let v_i be the column vector with 1 in the *i*th entry and all other entries 0. The group G(r, p, n) acts on $V := \mathbb{C}^n$ with orthonormal basis $\{v_1, \ldots, v_n\}$ as a complex reflection group.

Graded Hecke algebras

Every real reflection group is a complex reflection group and several of these are special cases of the groups G(r, p, n). In particular,

(a) G(1,1,n) is the symmetric group S_n ,

- (b) G(2,1,n) is the Weyl group WB_n of type B_n ,
- (c) G(2,2,n) is the Weyl group WD_n of type D_n , and
- (d) G(r, r, 2) is the dihedral group $I_2(r)$ of order 2r.

The reflections in G(r, p, n) are

$$\begin{aligned} \xi_i^{kp}, & 1 \le i \le n, \ 0 \le k \le (r/p) - 1, \ \text{and} \\ \xi_i^k \xi_j^{-k}(i,j), & 1 \le i < j \le n, \ 0 \le k \le r - 1, \end{aligned}$$

where (i, j) is the transposition in S_n that switches i and j.

Conjugacy in G(r, p, n). Each element of G(r, p, n) is conjugate by elements of S_n to a disjoint product of cycles of the form

$$\xi_i^{\lambda_i}\cdots\xi_k^{\lambda_k}(i,i+1,\ldots,k).$$

By conjugating this cycle by $\xi_i^{-c}\xi_{i+1}^{\lambda_i}\xi_{i+2}^{\lambda_i+\lambda_{i+1}}\cdots\xi_k^{\lambda_i+\dots+\lambda_{k-1}} \in G(r,r,n)$, we have

 $\xi_i^{-c}\xi_k^{c+\lambda_i+\dots+\lambda_k}(i,\dots,k), \quad \text{where } c = (k-i)\lambda_i + (k-i-1)\lambda_{i+1} + \dots + \lambda_{k-1}.$

If $i_1, i_2, \ldots, i_{\ell}$ denote the minimal indices of the cycles and c_1, \ldots, c_{ℓ} are the numbers c for the various cycles, then after conjugating by $\xi_{i_1}^{c_1} \cdots \xi_{i_{\ell-1}}^{c_{\ell-1}} \xi_{i_{\ell}}^{-(c_1+\cdots+c_{\ell-1})} \in G(r, r, n)$, each cycle becomes

$$\xi_k^{\lambda_i+\dots+\lambda_k}(i,\dots,k) \qquad \text{except the last, which is} \qquad \xi_{i_\ell}^{-a}\xi_n^b(i_\ell,\dots,n),$$

where $a = c_1 + \cdots + c_{\ell}$ and $b = a + \lambda_{i_{\ell}} + \cdots + \lambda_n$. If $k = n - i_{\ell} + 1$ is the length of the last cycle, then conjugating the last cycle by $\xi_{i_{\ell}}^{k-1}\xi_{i_{\ell}+1}^{-1}\cdots\xi_n^{-1} \in G(r, r, n)$ gives

$$\xi_{i_{\ell}}^{-a+k}\xi_{n}^{b-k}(i_{\ell},\ldots,n).$$

If we conjugate the last cycle by $\xi_{i_{\ell}}^{p} \in G(r, p, n)$, we have

$$\xi_{i_{\ell}}^{-a+p}\xi_{n}^{b-p}(i_{\ell},\ldots,n).$$

In summary, any element g of G(r, p, n) is conjugate to a product of disjoint cycles where each cycle is of the form

$$\xi_k^a(i, i+1, \dots, k), \qquad 0 \le a \le r-1,$$
(2.4a)

except possibly the last cycle, which is of the form

$$\xi_{i_{\ell}}^{a}\xi_{n}^{b}(i_{\ell}, i_{\ell}+1, \dots, n), \quad \text{with } 0 \le a \le \gcd(p, k) - 1, \quad (2.4b)$$

where $k = n - i_{\ell} + 1$ is the length of the last cycle.

Centralizers in G(r, p, n). Let $Z_{G(r, p, n)}(g) = \{h \in G(r, p, n) \mid hg = gh\}$ denote the centralizer of $g \in G(r, p, n)$. Since G(r, p, n) is a subgroup of G(r, 1, n),

$$Z_{G(r,p,n)}(g) = Z_{G(r,1,n)}(g) \cap G(r,p,n),$$

for any element $g \in G(r, p, n)$. Suppose that g is an element of G(r, 1, n) which is a product of disjoint cycles of the form $\xi_k^a(i, \ldots, k)$ and that $h \in G(r, 1, n)$ commutes with g. Conjugation by h effects some combination of the following operations on the cycles of g:

(a) permuting cycles of the same type, $\xi_k^a(i, \ldots, k)$ and $\xi_m^b(j, \ldots, m)$ with b = a and k - i = m - j,

- (b) conjugating a single cycle $\xi_k^a(i, \ldots, k)$ by powers of itself, and
- (c) conjugating a single cycle $\xi_k^a(i, \ldots, k)$ by $\xi_i^b \cdots \xi_k^b$, for any $0 \le b \le r-1$.

Furthermore, the elements of G(r, 1, n) which commute with g are determined by how they "rearrange" the cycles of g and a count (see [Mac, p. 170]) of the number of such operations shows that if $g \in G(r, 1, n)$ and $m_{a,k}$ is the number of cycles of type $\xi_{i+k}^a(i, i+1, \ldots, i+k)$ for g, then

$$\operatorname{Card}(Z_{G(r,1,n)}(g)) = \prod_{a,k} (m_{a,k}! \cdot k^{m_{a,k}} \cdot r^{m_{a,k}}).$$
(2.5)

Determining the graded Hecke algebras for G(r, p, n). It follows from Lemma 1.8a that if $g = \xi_i^{a+b} \xi_k^{-a}(i, \ldots, k)$, then $(V^g)^{\perp}$ has basis

$$\{v_k - v_{k-1}, v_{k-1} - v_{k-2}, \dots, v_{i+1} - \xi^a v_i\}$$
 if $b = 0$, and $\{v_i, \dots, v_k\}$ if $b \neq 0$.

Thus, if $g \in G(r, p, n)$ and $\operatorname{codim}(V^g) = 2$, then g is conjugate to one of the following elements:

$b = \xi_1^a \xi_3^{-a}(1,2,3),$	$0 \le a \le \gcd(p,3) - 1,$
$c = \xi_1^{a+\ell} \xi_2^{-a}(1,2),$	$\ell \neq 0$ (so $r \neq 1$),
$d = \xi_1^{\ell_1} \xi_2^{\ell_2},$	$\ell_1 \neq 0, \ \ell_2 \neq 0 \ (\text{so } r \neq 1),$
$e=(1,2)\xi_3^\ell,$	$\ell eq 0,$
$f = (1,2)\xi_3^a \xi_4^{-a}(3,4).$	

It is interesting to note that these elements are also representatives of the conjugacy classes of elements in G(r, p, n) which can be written as a product of two reflections.

We determine conditions on the above elements and on r, p, and n to give nontrivial graded Hecke algebras:

Graded Hecke algebras

(z) The center of G(r, p, n) is

$$Z(G(r,p,n)) = \{\xi_1^\ell \cdots \xi_n^\ell \mid n\ell = 0 \bmod p\}.$$

Since $\xi_1^p \cdots \xi_n^p \in Z(G(r, p, n))$, it follows that p = r or p = r/2 whenever

- $Z(G(r, p, n)) \subseteq \{\pm 1\} = \{\xi_1^0 \cdots \xi_n^0, \xi_1^{r/2} \cdots \xi_n^{r/2}\}.$ (b1) If $n \ge 4$, the element $\xi_1 \xi_2 \xi_3 \xi_4^{-3} \in Z_G(b)$ and has determinant ξ^2 on $(V^b)^{\perp} =$ span- $\{v_3 v_2, v_2 \xi^a v_1\}.$
- (b2) If n = 3 and $p = 0 \mod 3$, the element $\xi_1^{p/3} \xi_2^{p/3} \xi_3^{p/3} \in Z_G(b)$ and has determinant $\xi^{2p/3}$ on $(V^b)^{\perp}$. (c1) If $n \ge 3$, the element $\xi_1 \xi_2 \xi_3^{-2} \in Z_G(c)$ and has determinant ξ^2 on $(V^c)^{\perp} =$
- span- $\{v_1, v_2\}$.
- (c2) If n = 2, p = r/2 and p is odd, the element $\xi_1^{p/4} \xi_2^{p/4} \in Z_G(c)$ and has determinant $\xi^{r/2}$ on $(V^c)^{\perp}$.
- (d1) If $n \geq 3$, the element $\xi_1 \xi_3^{-1} \in Z_G(d)$ and has determinant ξ on $(V^d)^{\perp} =$ span- $\{v_1, v_2\}$.
- (d2) If p = r/2, the element $\xi_1^{r/2} \in Z_G(d)$ and has determinant $\xi^{r/2}$ on $(V^d)^{\perp}$. (ef) The elements e and f have order 2.

Thus, it follows from Corollary 1.11, Theorem 1.9b, and Lemma 2.2b that if A is a graded Hecke algebra for G(r, p, n), then

$a_b = 0$	unless	(i) $r = 1$, or (ii) $r = 2$, or (iii) $n = 3$ and $p \neq 0 \mod 3$,
$a_c = 0$	unless	(i) $r = 2$ and $p = 1$, or (ii) $n = 2$ and $p = r/2$,
$a_d = 0$	unless	$p = r, n = 2 \text{ and } p \neq 0 \mod 2,$
$a_e = 0$	always,	and
$a_f=0$	always.	

In the remaining cases, one uses the description of $Z_G(g)$ given just before (2.5) to check that all elements of $Z_G(g)$ have determinant 1 on $(V^g)^{\perp}$. Note that n=3and $p \neq 0 \mod 3$ imply that $a_b = 0$ for the elements $b = \xi_1^a \xi_3^{-a}(1,2,3)$.

We arrive at the following enumeration of the nontrivial graded Hecke algebras for complex reflection groups. (The tensor product algebra $S(V) \otimes \mathbb{C}G$ always exists and corresponds to the case when all of the skew symmetric forms a_a are zero). The table below gives representatives of the conjugacy classes of $g \in G$ that may have $a_g \neq 0$ for some graded Hecke algebra A.

Group	$\begin{array}{c} \text{Representative } g \\ \text{with } a_g \neq 0 \end{array}$
$G(1,1,n)=S_n$	(1,2,3)
$G(2,1,n)=WB_n, \ \ n\geq 3$	$\xi_1(1,2), (1,2,3)$
$G(2,2,n)=WD_n, \ n\geq 3$	(1,2,3)
$G(r,r,2)=I_2(r)$	$\xi_1^k \xi_2^{r-k}, \ 0 < k < r/2$
G(r, r/2, 2), r/2 odd	$\xi_2^{r/2}(1,2)$
$G(r,r,3), \ \ r eq 0 mod 3$	(1,2,3)
$G(r, r/2, 3), \ r/2 \neq 0 \bmod 3, \ r \neq 2$	(1,2,3)

Table 2. Graded Hecke algebras for the groups G(r, p, n)

2C. Exceptional complex reflection groups

The irreducible exceptional complex reflection groups G are denoted G_4, \ldots, G_{35} in the classification of Shephard and Todd. From Table VII in [ST], one sees that the center of G is ± 1 only in the cases G_4 , G_{12} , G_{24} and G_{33} . By Schur's lemma, the center of an irreducible complex reflection group consists of multiples of the identity. Thus, by Corollary 1.11, the only exceptional complex reflection groups that could have a nontrivial graded Hecke algebra (i.e., with some $a_g \neq 0$) are G_4 , G_{12} , G_{24} and G_{33} (we exclude the real groups). We determine the graded Hecke algebras for these groups using Theorem 1.9b and Lemma 2.2.

The rank 2 group G_4 has order 24 and seven conjugacy classes. The following data concerning these conjugacy classes are obtained from the computer software GAP [S+] using the package CHEVIE [G+]. In the following table, ω is a primitive cube root of unity and C(g) denotes the conjugacy class of g.

	Conjug	acy cla	lss repr	esentat	ives for G	4	
$\operatorname{Order}(g)$	1	4	3	6	6	3	2
$\det(g)$	1	1	ω	ω	ω^{-1}	ω^{-1}	1
$ \mathcal{C}(g) $	1	6	4	4	4	4	1
$ Z_G(g) $	24	4	6	6	6	6	24

The elements with determinant 1 and order more than 2 in G_4 all have order 4. If g is an element of order 4, then $|Z_G(g)| = 4$ and every element of $Z_G(g)$ has determinant 1 since $Z_G(g)$ is generated by g. Hence, by Theorem 1.9b and Lemma 2.2, a_g can be nonzero for a graded Hecke algebra for G_4 exactly when g has order

4. Thus, the dimension of the space of parameters for graded Hecke algebras of G_4 is 1.

The rank 2 group G_{12} has order 48. The computer software GAP provides the following information about the conjugacy classes of G_{12} .

Conjugacy class representatives for G_{12}								
$\operatorname{Order}(g)$	1	2	8	6	8	2	3	4
$\det(g)$	1	-1	-1	1	-1	1	1	1
$ \mathcal{C}(g) $	1	12	6	8	6	1	8	6
$ Z_G(g) $	48	4	8	6	8	48	6	8

If g is an element in G_{12} with order more than 2 and determinant 1, then g has order 3, 4, or 6. Let h be any element of order 8. Then h has determinant -1and commutes with h^2 of order 4. Hence, by Theorem 1.9b, if g has order 4, then $a_g = 0$. Let g_6 be a representative from the class of elements of order 6. Since $|Z_G(g_6)| = 6$, $Z_G(g_6)$ is generated by g_6 and hence every element of $Z_G(g_6)$ has determinant 1. We can choose g_6^2 as a representative for the conjugacy class of elements of order 3. As g_6 and g_6^2 commute, $\langle g_6 \rangle \subset Z_G(g_6^2)$. But $|\langle g_6 \rangle| =$ $6 = |Z_G(g_3)|$, so $Z_G(g_6^2)$ is generated by g_6 and every element of $Z_G(g_3)$ has determinant 1. Thus, a_g can be nonzero for a graded Hecke algebra A for G_{12} exactly when g has order 3 or 6. Thus, the dimension of the space of parameters of graded Hecke algebras for G_{12} is 2.

The rank 3 group G_{24} has order 336. Note that $-1 \in G_{24}$ since $Z(G) = \{\pm 1\}$. Up to *G*-orbits, there are two codimension 2 subspaces, *L* and *M*, that are equal to V^g for some $g \in G_{24}$ (see [OT, App. C, Table C.5]). Furthermore, $\operatorname{Stab}(L) \cong A_2$ and $\operatorname{Stab}(M) \cong B_2$. We need only consider elements of order 3 in $\operatorname{Stab}(L) \cong A_2$ and of order 4 in $\operatorname{Stab}(M) \cong B_2$ (as the rest have order 1 or 2). In G_{24} , there is only one conjugacy class of elements of order 3 and only one conjugacy class of elements of order 4 and determinant 1. The table below (obtained using GAP) records information about these classes.

Certain classes of G_{24}				
$\operatorname{Order}(g)$	3	4		
$\det(g)$	1	1		
$ \mathcal{C}(g) $	56	42		
$ Z_G(g) $	6	8		

If g has order 3, $Z_G(g)$ must contain 1, g, g^2 , and -1, and hence $Z_G(g)$ is generated by these elements since $|Z_G(g)| = 6$. Thus all elements of $Z_G(g)$ have determinant 1

A. Ram and A. V. Shepler

on $(V^g)^{\perp}$. If g has order 4 and determinant 1, then $Z_G(g)$ must contain $1, g, g^2, g^3$, and -1, elements which all have determinant 1 on $(V^g)^{\perp}$. Since $|Z_G(g)| = 8$, these elements generate $Z_G(g)$ and so every element of $Z_G(g)$ has determinant 1 on $(V^g)^{\perp}$. Hence, a_g can be nonzero for a graded Hecke algebra of G_{24} exactly when g has order 3 or g has order 4 and determinant 1. Thus, the dimension of the space of parameters for graded Hecke algebras for G_{24} is 2.

The group G_{33} is the only exceptional complex reflection group of rank 5. It has order 72 · 6! and degrees 4, 6, 10, 12, 18. There are 45 reflecting hyperplanes and the corresponding reflections all have order 2. Up to *G*-orbits, there are two codimension 2 subspaces, *L* and *M*, that are equal to V^g for some $g \in G_{33}$ (see [OT, App. C, Table C.14]). Furthermore, $\operatorname{Stab}(L) \cong A_1 \times A_1$ and $\operatorname{Stab}(M) \cong A_2$. We need not consider the case where $V^g = L$ since then *g* has order 2 and hence $a_g = 0$ for any graded Hecke algebra by Proposition 2.2b.

We use a presentation for G_{33} in six coordinates from [ST]: Let $V = \mathbb{C}^6$ with standard coordinate functions x_i and consider the group generated by order 2 reflections about the hyperplanes $H_1 = \{x_2 - x_3 = 0\}$, $H_2 = \{x_3 - x_4 = 0\}$, $H_3 = \{x_1 - x_2 = 0\}$, $H_4 = \{x_1 - \omega x_2 = 0\}$, $H_5 = \{x_1 + x_2 + x_3 + x_4 + x_5 + x_6 = 0\}$, where ω is a primitive cube root of unity. The fixed point space of this (reducible) group is $Y = H_1 \cap \cdots \cap H_5 = \{(0, 0, 0, 0, x, -x) \mid x \in \mathbb{C}\}$, and G_{33} is just the restriction to Y^{\perp} . Let s_i be the order 2 reflection about H_i . Let $g = s_1 s_3$. Then $V^g = H_1 \cap H_3$ and $\operatorname{Stab}(V^g) \cong A_2$. Let $h = (s_1 s_3 s_4)^2$, the diagonal matrix with diagonal $\{\omega, \omega, \omega, 1, 1, 1\}$. Then h acts as ω times the identity on $(V^g)^{\perp}$ as $(V^g)^{\perp} \subseteq \mathbb{C}$ -span $\{x_1, x_2, x_3\}$. Hence, h commutes with g. But $(V^g)^{\perp}$ has dimension 2 and h has determinant $\omega^2 \neq 1$ on $(V^g)^{\perp}$. Thus, by Theorem 1.9b and Lemma 2.2, $a_g = 0$ for any graded Hecke algebra. The same argument applied to Y^{\perp} shows that G_{33} has no nontrivial graded Hecke algebras. In summary, the dimension of the space of parameters for graded Hecke algebras for G_{33} is zero.

Group	$g ext{ with } a_g eq 0$
G_4	$\operatorname{Order}(g)=4$
G_{12}	$\operatorname{Order}(g_1) = 3$ and $\operatorname{Order}(g_2) = 6$
G_{24}	$\operatorname{Order}(g_1) = 3 \text{ and } \operatorname{Order}(g_2) = 4, \det(g_2) = 1$

Table 3. Graded Hecke algebras for exceptional complex reflection groups

3. The graded Hecke algebras $H_{\rm gr}$

In [Lu], Lusztig gives a definition of graded Hecke algebras for real reflection groups which is different from the definition in Section 1, which applies to more general groups. It is not obvious that Lusztig's algebras are examples of the graded Hecke

Graded Hecke algebras

algebras defined in Section 1. In this section, we show explicitly how the definition of Section 1 includes Lusztig's algebras.

Let W be a finite real reflection group acting on V and let R be the root system of W. Let $\alpha_1, \ldots, \alpha_n$ be a choice of simple roots in R and let s_1, \ldots, s_n be the corresponding simple reflections in W. Let s_{α} be the reflection in the root α so that, for $v \in V$,

$$s_{\alpha}v = v - \langle v, \alpha^{\vee} \rangle \alpha, \qquad ext{where } \alpha^{\vee} = 2 \alpha / \langle \alpha, \alpha \rangle.$$

Let $R^+ = \{\alpha > 0\}$ denote the set of positive roots in R.

Let k_{α} be fixed complex numbers indexed by the roots $\alpha \in R$ satisfying

$$k_{w\alpha} = k_{\alpha}, \quad \text{for all } w \in W, \ \alpha \in R.$$
 (3.1)

This amounts to a choice of either one or two "parameters", depending on whether all roots in R are the same length or not. As in Section 1, let $\mathbb{C}W = \mathbb{C}$ -span $\{t_g \mid g \in W\}$, with $t_g t_h = t_{gh}$, and let S(V) be the symmetric algebra of V. Lusztig [Lu] defines the "graded Hecke algebra" with parameters $\{k_\alpha\}$ to be the unique algebra structure $H_{\rm gr}$ on the vector space $S(V) \otimes \mathbb{C}W$ such that

$$S(V) = S(V) \otimes 1 \text{ is a subalgebra of } H_{\text{gr}}, \tag{3.2a}$$

$$\mathbb{C}W = 1 \otimes \mathbb{C}W \text{ is a subalgebra of } H_{\text{gr}}, \tag{3.2b}$$

and

$$t_{s_i}v = (s_iv)t_{s_i} - k_{\alpha_i}\langle v, \alpha_i^{\vee} \rangle,$$

for all $v \in V$ and simple reflections s_i in the simple roots α_i .

(3.2c)

We shall show that every algebra $H_{\rm gr}$ as defined by (3.2a–c) is a graded Hecke algebra A for a specific set of skew symmetric bilinear forms a_g .

Let $k_{\alpha} \in \mathbb{C}$ as in (3.1). Use the notation

$$h = \frac{1}{2} \sum_{\alpha > 0} k_{\alpha} \alpha^{\vee} t_{s_{\alpha}}, \quad \text{so that} \quad \langle v, h \rangle = \frac{1}{2} \sum_{\alpha > 0} k_{\alpha} \langle v, \alpha^{\vee} \rangle t_{s_{\alpha}} \quad (3.3)$$

for $v \in V$. The element h should be viewed as an element of $V \otimes \mathbb{C}W$, and $\langle v, h \rangle \in \mathbb{C}W$. With this notation, let A be the algebra (as in Section 1) generated by V and $\mathbb{C}W$ with relations

$$t_g v = (gv)t_g$$
 and $[v, w] = -[\langle v, h \rangle, \langle w, h \rangle],$ for $v, w \in V, g \in W.$

(3.4)

Note that A is defined by the bilinear forms

$$a_g(v,w) = \frac{1}{4} \sum_{\substack{\alpha,\beta > 0 \\ g = s_\alpha s_\beta}} k_\alpha k_\beta \left(\langle v, \beta^{\vee} \rangle \langle w, \alpha^{\vee} \rangle - \langle v, \alpha^{\vee} \rangle \langle w, \beta^{\vee} \rangle \right).$$

The following theorem shows that the algebra A satisfies the defining conditions (3.2a–c) of the algebra $H_{\rm gr}$.

Theorem 3.5. Let W be a finite real reflection group and let A be the algebra defined by (3.4).

(a) As vector spaces, A ≈ S(V) ⊗ CW (and hence, A is a graded Hecke algebra).
(b) If ṽ = v - ⟨v, h⟩ for v ∈ V, then

$$[\tilde{v}, \tilde{w}] = 0$$
 and $t_{s_i} \tilde{v} = (\widetilde{s_i v}) t_{s_i} - k_{\alpha_i} \langle v, \alpha_i^{\lor} \rangle,$

for all $v, w \in V$ and simple reflections s_i in W.

Proof. First note that if $u, v \in V$ then

$$[u, \langle v, h \rangle] = \frac{1}{2} \sum_{\alpha > 0} k_{\alpha} \langle v, \alpha^{\vee} \rangle \langle u, \alpha^{\vee} \rangle \alpha t_{s_{\alpha}} = [v, \langle u, h \rangle]. \tag{\ast}$$

Thus, for $u, v, w \in V$,

$$\begin{split} [u, [v, w]] + [w, [u, v]] &+ [v, [w, u]] \\ &= [u, [\langle w, h \rangle, \langle v, h \rangle]] + [w, [\langle v, h \rangle, \langle u, h \rangle]] + [v, [\langle u, h \rangle, \langle w, h \rangle]] \\ &= [[u, \langle w, h \rangle], \langle v, h \rangle] + [\langle w, h \rangle, [u, \langle v, h \rangle]] + [[w, \langle v, h \rangle], \langle u, h \rangle] \\ &+ [\langle v, h \rangle, [w, \langle u, h \rangle]] + [[v, \langle u, h \rangle], \langle w, h \rangle] + [\langle u, h \rangle], [v, \langle w, h \rangle]] \\ &= [[w, \langle u, h \rangle], \langle v, h \rangle] + [\langle w, h \rangle, [v, \langle u, h \rangle]] + [[v, \langle w, h \rangle], \langle u, h \rangle] \\ &+ [\langle v, h \rangle, [w, \langle u, h \rangle]] + [[v, \langle u, h \rangle], \langle w, h \rangle] + [\langle u, h \rangle, [v, \langle w, h \rangle]] \\ &= 0. \end{split}$$

For $v \in V$, $h \in W$, and s_i a simple reflection,

$$t_{s_{i}}\langle v,h\rangle t_{s_{i}} = \frac{1}{2} \sum_{\alpha>0} k_{\alpha} \langle v,\alpha^{\vee} \rangle t_{s_{s_{i}\alpha}} = \left(\frac{1}{2} \sum_{\alpha>0} k_{\alpha} \langle v,s_{i}\alpha^{\vee} \rangle t_{s_{\alpha}}\right) + k_{\alpha_{i}} \langle v,\alpha_{i}^{\vee} \rangle t_{s_{i}}$$
$$= \left(\frac{1}{2} \sum_{\alpha>0} k_{\alpha} \langle s_{i}v,\alpha^{\vee} \rangle t_{s_{\alpha}}\right) + k_{\alpha_{i}} \langle v,\alpha_{i}^{\vee} \rangle t_{s_{i}} = \langle s_{i}v,h \rangle + k_{\alpha_{i}} \langle v,\alpha_{i}^{\vee} \rangle t_{s_{i}}.$$
(3.7)

326

(3.6)

Using this equality, we obtain

$$\begin{split} t_{s_{i}}[v,w]t_{s_{i}} &= -t_{s_{i}}[\langle v,h\rangle,\langle w,h\rangle]t_{s_{i}} \\ &= -[\langle s_{i}v,h\rangle + k_{\alpha_{i}}\langle v,\alpha_{i}^{\vee}\rangle t_{s_{i}},\langle s_{i}w,h\rangle + k_{\alpha_{i}}\langle w,\alpha_{i}^{\vee}\rangle t_{s_{i}}] \\ &= [s_{i}v,s_{i}w] - k_{\alpha_{i}}\langle v,\alpha_{i}^{\vee}\rangle [t_{s_{i}},\langle s_{i}w,h\rangle] - k_{\alpha_{i}}\langle w,\alpha_{i}^{\vee}\rangle [\langle s_{i}v,h\rangle,t_{s_{i}}] \\ &= [s_{i}v,s_{i}w] - k_{\alpha_{i}}\langle v,\alpha_{i}^{\vee}\rangle (t_{s_{i}}\langle s_{i}v,h\rangle t_{s_{i}} - \langle s_{i}v,h\rangle)t_{s_{i}} \\ &\quad + k_{\alpha_{i}}\langle w,\alpha_{i}^{\vee}\rangle (t_{s_{i}}\langle s_{i}v,h\rangle t_{s_{i}} - \langle s_{i}v,h\rangle)t_{s_{i}} \\ &= [s_{i}v,s_{i}w] - k_{\alpha_{i}}\langle v,\alpha_{i}^{\vee}\rangle (\langle w,h\rangle t_{s_{i}} + k_{\alpha_{i}}\langle s_{i}v,\alpha_{i}^{\vee}\rangle - \langle s_{i}w,h\rangle t_{s_{i}}) \\ &\quad + k_{\alpha_{i}}\langle w,\alpha_{i}^{\vee}\rangle (\langle v,h\rangle t_{s_{i}} + k_{\alpha_{i}}\langle s_{i}v,\alpha_{i}^{\vee}\rangle - \langle s_{i}v,h\rangle t_{s_{i}}) \\ &= [s_{i}v,s_{i}w] - k_{\alpha_{i}}\langle v,\alpha_{i}^{\vee}\rangle \langle w,\alpha_{i}^{\vee}\rangle \langle \alpha_{i},h\rangle t_{s_{i}} - k_{\alpha_{i}}^{2}\langle v,\alpha_{i}^{\vee}\rangle \langle w,s_{i}\alpha_{i}^{\vee}\rangle \\ &\quad + k_{\alpha_{i}}\langle w,\alpha_{i}^{\vee}\rangle \langle v,\alpha_{i}^{\vee}\rangle \langle \alpha_{i},h\rangle t_{s_{i}} + k_{\alpha_{i}}^{2}\langle w,\alpha_{i}^{\vee}\rangle \langle v,s_{i}\alpha_{i}^{\vee}\rangle \\ &= [s_{i}v,s_{i}w]. \end{split}$$

$$(3.8)$$

The two identities (3.6) and (3.8), as in (1.3) and (1.4), show that the algebra A is isomorphic to $S(V) \otimes \mathbb{C}W$.

(b) This can now be proved by direct computation. If $v, w \in V$ then

$$[\tilde{v},\tilde{w}] = [v - \langle v,h\rangle, w - \langle w,h\rangle] = [v,w] + [\langle v,h\rangle, \langle w,h\rangle] - [v,\langle w,h\rangle] + [w,\langle v,h\rangle] = 0,$$

by equation (3.4) and equation (*) in the proof of Theorem 3.5. If $v \in V$ and s_i is a simple reflection then, by (3.7),

$$t_{s_i}\tilde{v}t_{s_i} = t_{s_i}vt_{s_i} - t_{s_i}\langle v,h\rangle \\ t_{s_i} = s_iv - \langle s_iv,h\rangle - k_{\alpha_i}\langle v,\alpha_i^\vee\rangle \\ t_{s_i} = \widetilde{s_iv} - k_{\alpha_i}\langle v,\alpha_i^\vee\rangle$$

Theorem 3.5b shows that if A is the graded Hecke algebra defined by (3.4), then the elements \tilde{v} , for $v \in V$, generate a subalgebra of A isomorphic to S(V)and these elements together with the t_{s_i} satisfy the relations of (3.2c). Since part (a) of Theorem 3.5 shows that A is isomorphic to $S(V) \otimes \mathbb{C}W$ as a vector space, it follows that A satisfies the conditions (3.2a–c), relations which uniquely define the graded Hecke algebra A. Thus, Lusztig's algebras are special cases of the graded Hecke algebras defined in Section 1. Furthermore, by comparing the dimensions of the parameter spaces, we see that there are graded Hecke algebras that are not isomorphic to algebras defined by Lusztig for the Coxeter groups F_4 , H_3 , H_4 , and $I_2(m)$.

4. Examples

4A. The symmetric group $G(1, 1, n) = S_n$

Let V be an n dimensional vector space with orthonormal basis v_1, \ldots, v_n and let S_n act on V by permuting the v_i . Let A be a graded Hecke algebra for S_n . Any element which is a product of two reflections is conjugate to (1, 2, 3) or (1, 2)(3, 4). The element (1, 2)(3, 4) has order 2 and so, in the algebra A,

$$[v_i, v_j] = \sum_{k \neq i, j} (a_{(i,j,k)}(v_i, v_j)t_{(i,j,k)} + a_{(j,i,k)}(v_i, v_j)t_{(j,i,k)}),$$

since v_i or v_j is in $V^g = \ker a_g$ for all other three cycles g. Since, by (1.6), $a_{(j,i,k)}(v_i, v_j) = a_{(i,j,k)}(v_j, v_i) = -a_{(i,j,k)}(v_i, v_j)$, the graded Hecke algebra A is defined by the relations

$$[v_i, v_j] = \beta \sum_{k \neq i,j} (t_{(i,j,k)} - t_{(j,i,k)}) \quad \text{and} \quad t_w v_i = v_{w(i)} t_w, \tag{4.1}$$

where $w \in S_n$, $1 \leq i, j \leq n$, $i \neq j$, and $\beta = a_{(1,2,3)}(v_1, v_2)$. Let $k \in \mathbb{C}$. Then, with h as in (3.3),

$$\langle v_i, h \rangle = \frac{1}{2} \sum_{\ell < m} k \langle v_i, v_\ell - v_m \rangle t_{(\ell,m)}$$

$$= \frac{k}{2} \left(\sum_{i < \ell} t_{(i,\ell)} - \sum_{i > \ell} t_{(\ell,i)} \right) = \frac{k}{2} \sum_{i \neq \ell} \operatorname{sgn}(\ell - i) t_{(i,\ell)}.$$
(4.2)

If $f \in \mathbb{C}S_n$, let $f|_{t_g}$ denote the coefficient of t_g in f. Let A be the graded Hecke algebra defined by the relations in (4.1) with

$$\beta = a_{(i,j,\ell)}(v_i, v_j) = [\langle v_i, h \rangle, \langle v_j, h \rangle] \big|_{t_{(i,j,\ell)}} = (k^2/4)(t_{(i,\ell)}t_{(j,\ell)} + t_{(i,j)}t_{(i,\ell)} - t_{(j,\ell)}t_{(i,j)}) \big|_{t_{(i,j,\ell)}} = k^2/4.$$
(4.3)

If $\tilde{v}_i = v_i - \langle v_i, h \rangle$ and s_i is the simple reflection (i, i+1) then, by Theorem 3.5,

$$\tilde{v}_i \tilde{v}_j = \tilde{v}_j \tilde{v}_i, \qquad t_{s_i} \tilde{v}_i = \tilde{v}_{i+1} t_{s_i} + k, \qquad t_{s_i} \tilde{v}_{i+1} = \tilde{v}_i t_{s_i} - k, \quad \text{and} \\
t_{s_i} \tilde{v}_i = \tilde{v}_i t_{s_i}, \qquad \text{for } |i-j| > 1,$$
(4.4)

and the algebra A is the graded Hecke algebra $H_{\rm gr}$ for S_n which is defined in Section 3. When k=1, the map

is a surjective algebra homomorphism.

Graded Hecke algebras

4B. The hyperoctahedral group $G(2,1,n) = WB_n$

We use the notation from Section 2B so that the group G(2, 1, n) is acting by orthogonal matrices on the *n* dimensional vector space *V* with orthonormal basis $\{v_1, \ldots, v_n\}$. In this case, ξ_i denotes the diagonal matrix with all ones on the diagonal except for -1 in the (i, i)th entry.

Let A be a graded Hecke algebra for G(2,1,n). If $\beta_1 = a_{(i,j,k)}(v_i,v_j)$ and $\beta_2 = a_{\xi_1(1,2)}(v_1,v_2)$, then, in the algebra A,

$$[v_{i}, v_{j}] = \beta_{2}(t_{\xi_{1}(1,2)} - t_{\xi_{2}(1,2)}) + \beta_{1} \sum_{\ell \neq i,j} \begin{pmatrix} t_{(i,j,\ell)} - t_{\xi_{i}\xi_{\ell}(i,j,\ell)} - t_{\xi_{i}\xi_{j}(i,j,\ell)} + t_{\xi_{j}\xi_{\ell}(i,j,\ell)} \\ + t_{\xi_{i}\xi_{j}(j,i,\ell)} + t_{\xi_{j}\xi_{\ell}(j,i,\ell)} - t_{\xi_{i}\xi_{\ell}(j,i,\ell)} - t_{(j,i,\ell)} \end{pmatrix}.$$

$$(4.6)$$

Let $k_s, k_\ell \in \mathbb{C}$. Then, with h as in (3.3),

$$\langle v_i, h \rangle = \frac{k_s}{2} \sum_{\ell} \langle v_i, 2v_{\ell} \rangle t_{\xi_{\ell}} - \frac{k_{\ell}}{2} \sum_{\ell < m} \langle v_i, v_{\ell} - v_m \rangle t_{(\ell,m)} - \frac{k_{\ell}}{2} \sum_{\ell < m} \langle v_i, v_{\ell} + v_m \rangle t_{\xi_{\ell}(\ell,m)}$$
$$= k_s t_{\xi_i} - \frac{k_{\ell}}{2} \left(\sum_{i < \ell} (t_{(i,\ell)} + t_{\xi_i \xi_{\ell}(i,\ell)}) + \sum_{i > \ell} (-t_{(i,\ell)} + t_{\xi_i \xi_{\ell}(i,\ell)}) \right).$$
(4.7)

If $f \in \mathbb{C}G(2,1,n)$, let $f|_{t_g}$ denote the coefficient of t_g in f. With notation as in (4.6), let A be the graded Hecke algebra for G(2,1,n) with

$$\begin{split} \beta_1 &= a_{(i,j,\ell)}(v_i, v_j) = \left[\langle v_i, h \rangle, \langle v_j, h \rangle \right] \Big|_{t_{(i,j,\ell)}} \\ &= \left(k_{\ell}^2 / 4 \right) (t_{(i,\ell)} t_{(j,\ell)} + t_{(i,j)} t_{(i,\ell)} - t_{(j,\ell)} t_{(i,j)}) \Big|_{t_{(i,j,\ell)}} = k_{\ell}^2 / 4, \quad \text{and} \\ \beta_2 &= \left[\langle v_i, h \rangle, \langle v_j, h \rangle \right] \Big|_{t_{\xi_i(i,j)}} \\ &= \left(1/2 \right) k_s k_{\ell} (-t_{\xi_i} t_{(i,j)} + t_{(i,j)} t_{\xi_j} - t_{\xi_j} t_{\xi_i \xi_j(i,j)} - t_{\xi_i \xi_j(i,j)} t_{\xi_i}) \Big|_{t_{\xi_i(i,j)}} = -k_s k_{\ell}. \end{split}$$

If $\tilde{v}_i = v_i - \langle v_i, h \rangle$, then, by Theorem 3.5, the \tilde{v}_i commute and the algebra A is the algebra $H_{\rm gr}$ for WB_n defined in Section 3.

4C. The type D_n Weyl group $G(2,2,n) = WD_n$

We shall use the notation from Section 2B so that the group G(2, 2, n) is acting by orthogonal matrices on the *n* dimensional vector space *V* with orthonormal basis $\{v_1, \ldots, v_n\}$. This is an index 2 subgroup of G(2, 1, n), and our notation is the same as used above for WB_n .

Let A be a graded Hecke algebra for G(2,2,n). If $\beta = a_{(i,j,k)}(v_i,v_j)$ then, in the algebra A,

$$[v_i, v_j] = \beta \sum_{\ell \neq i, j} \begin{pmatrix} t_{(i,j,k)} - t_{\xi_i \xi_\ell(i,j,\ell)} - t_{\xi_i \xi_j(i,j,\ell)} + t_{\xi_j \xi_\ell(i,j,\ell)} \\ + t_{\xi_i \xi_j(j,i,\ell)} + t_{\xi_j \xi_\ell(j,i,\ell)} - t_{\xi_i \xi_\ell(j,i,\ell)} - t_{(j,i,\ell)} \end{pmatrix}.$$
(4.8)

Let $k \in \mathbb{C}$. Then, with h as in (3.3),

$$\langle v_i, h \rangle = \frac{k}{2} \left(\sum_{i < \ell} (t_{(i,\ell)} + t_{\xi_i \xi_\ell(i,\ell)}) + \sum_{i > \ell} (-t_{(i,\ell)} + t_{\xi_i \xi_\ell(i,\ell)}) \right).$$
(4.9)

If $f \in \mathbb{C}G(2,2,n)$, let $f|_{t_g}$ denote the coefficient of t_g in f. With notation as in (4.8), let A be the graded Hecke algebra for G(2,2,n) with

$$\beta = a_{(i,j,\ell)}(v_i, v_j) = \left[\langle v_i, h \rangle, \langle v_j, h \rangle \right] \Big|_{t_{(i,j,\ell)}}$$
$$= (k^2/4)(t_{(i,\ell)}t_{(j,\ell)} + t_{(i,j)}t_{(i,\ell)} - t_{(j,\ell)}t_{(i,j)}) \Big|_{t_{(i,j,\ell)}} = k^2/4.$$

If $\tilde{v}_i = v_i - \langle v_i, h \rangle$, then, by Theorem 3.5, the \tilde{v}_i commute and the algebra A is the algebra $H_{\rm gr}$ for WD_n defined in Section 3.

4D. The dihedral group $I_2(r) = G(r, r, 2)$ of order 2r

We shall use the notation for G(r, r, 2) from Section 2B so that the group G(r, r, 2) is acting by unitary matrices on the 2 dimensional vector space V with orthonormal basis $\{v_1, v_2\}$. The group G(r, r, 2) is realized as a real reflection group by using the basis

$$\varepsilon_1 = \frac{1}{\sqrt{2}}(v_1 + v_2), \qquad \varepsilon_2 = \frac{-1}{i\sqrt{2}}(v_1 - v_2).$$

This basis is also orthonormal and, with respect to this basis, G(r, r, 2) acts by the matrices

$$\begin{pmatrix} \cos(2\pi m/r) & \mp \sin(2\pi m/r) \\ \sin(2\pi m/r) & \pm \cos(2\pi m/r) \end{pmatrix}, \qquad 0 \le m \le r-1.$$

Let A be a graded Hecke algebra for G(r, r, 2). The conjugacy classes of elements which are products of two reflections are $\{\xi_1^k \xi_2^{-k}, \xi_1^{-k} \xi_2^k\}, 0 < k < r/2$. Then, in the algebra A,

$$[\varepsilon_1, \varepsilon_2] = \sum_{0 < k < r/2} \beta_k (t_{\xi_1^k \xi_2^{-k}} - t_{\xi_1^{-k} \xi_2^k}), \quad \text{where} \quad \beta_k = a_{\xi_1^k \xi_2^{-k}} (\varepsilon_1, \varepsilon_2).$$
(4.10)

When r is even, there are two conjugacy classes of reflections

$$\{\xi_1^{2k}\xi_2^{-2k}(1,2) \mid 0 \le k < r/2\} \qquad \text{and} \qquad \{\xi_1^{2k+1}\xi_2^{-(2k+1)}(1,2) \mid 0 \le k < r/2\}.$$

The reflection $\xi_1^m \xi_2^{-m}(12)$ is the reflection in the line perpendicular to the vector

$$\alpha_m = \sin(-2\pi m/2r)\varepsilon_1 + \cos(-2\pi m/2r)\varepsilon_2,$$

Graded Hecke algebras

and the vectors α_m can be taken as a root system for G(r, r, 2). With h as in (3.3) and $k_s, k_\ell \in \mathbb{C}$,

$$\langle \varepsilon_{1}, h \rangle = \sum_{0 \le k < r/2} \left(k_{s} \sin(-2k \, 2\pi/2r) t_{\xi_{1}^{2k} \xi_{2}^{-2k}(1,2)} + k_{\ell} \sin(-(2k+1) \, 2\pi/2r) t_{\xi_{1}^{2k+1} \xi_{2}^{-(2k+1)}(1,2)} \right),$$

$$\langle \varepsilon_{2}, h \rangle = \sum_{0 \le k < r/2} \left(k_{s} \cos(-2k \, 2\pi/2r) t_{\xi_{1}^{2k} \xi_{2}^{-2k}(1,2)} + k_{\ell} \cos(-(2k+1) \, 2\pi/2r) t_{\xi_{1}^{2k+1} \xi_{2}^{-(2k+1)}(1,2)} \right).$$

$$(4.11)$$

If $f \in \mathbb{C}G(r, r, 2)$, let $f|_{t_g}$ denote the coefficient of t_g in f. With notation as in 4.10, let A be the graded Hecke algebra for G(r, r, 2) with

$$\beta_{k} = a_{\xi_{1}^{k}\xi_{2}^{-k}}(\varepsilon_{1}, \varepsilon_{2}) = \left[\langle \varepsilon_{1}, h \rangle, \langle \varepsilon_{2}, h \rangle \right] \Big|_{t_{\xi_{1}^{k}\xi_{2}^{-k}}}$$

$$= \begin{cases} \sin(k \, 2\pi/2r)rk_{s}k_{\ell} & \text{if } k \text{ is odd} \\\\ \sin(k \, 2\pi/2r)\frac{r}{2}(k_{s}^{2} + k_{\ell}^{2}) & \text{if } k \text{ is even.} \end{cases}$$

$$(4.12)$$

If $\tilde{\varepsilon}_i = \varepsilon_i - \langle \varepsilon_i, h \rangle$, then by Theorem 3.5, the $\tilde{\varepsilon}_i$ commute and the algebra A is the algebra $H_{\rm gr}$ for $I_2(r)$ defined in Section 3.

When r is odd, all aspects of the calculation in (4.11) and (4.12) are the same as for the case r even except that there is only one conjugacy class of reflections, $\{\xi_1^k \xi_2^{-k}(1,2) \mid 0 \le k \le r-1\}$, and so $k_s = k_{\ell}$.

4E. The group G(r, r/2, 2), r/2 odd

We use the notation from Section 2B, or from above for the group G(r, r, 2). In this case, the group is not a real reflection group, hence G(r, r/2, 2) acts by unitary matrices but not by orthogonal matrices.

Let A be a graded Hecke algebra for G(r, r/2, 2). The only conjugacy class for which a_g can be nonzero is $\{t_{\xi_1^k \xi_2^{r/2-k}(1,2)} \mid 0 \leq k < r\}$. Thus, in the algebra A,

$$[v_1, v_2] = \beta \sum_k (t_{\xi_1^{2k} \xi_2^{r/2 - 2k}(1, 2)} - t_{\xi_1^{r/2 - 2k} \xi_2^{2k}(1, 2)}), \qquad \text{where} \quad \beta = a_{\xi_2^{r/2}(1, 2)}(v_1, v_2).$$

5. A different graded Hecke algebra for G(r, 1, n)

The classification of graded Hecke algebras for complex reflection groups in Section 2 shows that there do not exist graded Hecke algebras $A \cong S(V) \otimes \mathbb{C}G$ for the groups G = G(r, 1, n), r > 2, n > 3. In this section, we define a different "semidirect product" of the symmetric algebra S(V) and the group algebra $\mathbb{C}G$ for the groups G(r, 1, n). These algebras are not graded Hecke algebras in the sense of Section 1, but they do have a structure similar to what we would expect from experience with graded Hecke algebras for real reflection groups. Is it possible that there is a general definition of graded Hecke algebras, different from that given in Section 1, which includes the algebras defined below as examples for the groups G(r, 1, n)?

We shall use the notation for the groups G(r, 1, n) as in Section 2B so that the group G(r, 1, n) is acting by monomial matrices on a vector space V of dimension n with orthonormal basis $\{v_1, \ldots, v_n\}$. Let s_i denote the permutation $(i, i + 1) \in G(r, 1, n)$.

Define $H^*_{r,1,n}$ to be the algebra generated by the group algebra $\mathbb{C}G(r,1,n)$ and V with relations

$$\begin{aligned} v_i v_j &= v_j v_i, & \text{for all } 1 \le i, j \le n, \\ t_{\xi_i} v_j &= v_j t_{\xi_i}, & \text{for all } 1 \le i, j \le n, \\ t_{s_i} v_k &= v_k t_{s_i}, & \text{if } k \notin \{i, i+1\}, \end{aligned}$$
(5.1)

$$t_{s_i} v_{i+1} = v_i t_{s_i} + \sum_{\ell=0}^{r-1} t_{\xi_i^{\ell} \xi_{i+1}^{-\ell}}, \quad \text{for } 1 \le i \le n-1$$

The following proposition establishes an "evaluation homomorphism" for the algebras $H_{r,1,n}^*$ which is a generalization of the homomorphism in (4.5).

Proposition 5.2. Define elements \bar{v}_k in the group algebra $\mathbb{C}G(r, 1, n)$ by setting $\bar{v}_1 = 0$ and

$$\bar{v}_k = \frac{1}{r} \sum_{i < k} \sum_{0 \le \ell \le r-1} t_{\xi_i^{\ell} \xi_k^{-\ell}(i,k)}, \quad \text{for } 2 \le k \le n.$$

Then there is a surjective algebra homomorphism

$$\begin{array}{cccc} H^*_{r,1,n} & \longrightarrow & \mathbb{C}G(r,1,n) \\ t_g & \longmapsto & t_g \\ v_k & \longmapsto & \bar{v}_k \end{array}$$

Proof. We must check that the defining relations (5.1) of $H_{r,1,n}^*$ hold with the v_k replaced by the \bar{v}_k .

Graded Hecke algebras

For each $1 \leq k \leq n$, let

$$ar{z}_k = ar{v}_1 + \dots + ar{v}_k = rac{1}{r}\sum_{\substack{1 \leq i < j \leq k \ 0 \leq \ell < r-1}} t_{\xi_i^\ell \xi_j^{-\ell}(i,j)}.$$

Then, for each $k, \bar{z}_k \in Z(\mathbb{C}G(r, 1, k))$ since it is the sum of the elements of the conjugacy class of reflections $t_{\xi_i^\ell \xi_j^{-\ell}(i,j)}$ in G(r, 1, k). So \bar{z}_k commutes with $\bar{z}_1, \ldots, \bar{z}_k$ and therefore $\bar{z}_1, \ldots, \bar{z}_n$ commute. Since $\bar{v}_k = \bar{z}_k - \bar{z}_{k-1}$, it follows that $\bar{v}_1, \ldots, \bar{v}_n$ also commute.

If m > k then t_{ξ_m} clearly commutes with \bar{z}_k . If $m \leq k$ then t_{ξ_m} commutes with \bar{z}_k since $\bar{z}_k \in Z(G(r, 1, k))$. So t_{ξ_m} commutes with $\bar{z}_1, \ldots, \bar{z}_n$ and hence with $\bar{v}_1, \ldots, \bar{v}_n$.

Since

$$\begin{aligned} t_{s_k} \bar{v}_k t_{s_k} &= t_{s_k} \left(\sum_{\substack{i < k \\ 0 \le \ell \le r-1}} t_{\xi_i^{\ell} \xi_k^{-\ell}(i,k)} \right) t_{s_k} &= \sum_{\substack{i < k \\ 0 \le \ell \le r-1}} t_{\xi_i^{\ell} \xi_{k+1}^{-\ell}(i,k+1)} \\ &= \sum_{\substack{i < k+1 \\ 0 \le \ell \le r-1}} t_{\xi_i^{\ell} \xi_{k+1}^{-\ell}(i,k+1)} - \sum_{\substack{0 \le \ell \le r-1}} t_{\xi_k^{\ell} \xi_{k+1}^{-\ell}(k,k+1)} \\ &= \bar{v}_{k+1} - \sum_{\substack{0 \le \ell \le r-1}} t_{\xi_k^{\ell} \xi_{k+1}^{-\ell}} t_{s_k}, \end{aligned}$$

it follows that

$$\bar{v}_k t_{s_k} = t_{s_k} \bar{v}_{k+1} - \sum_{0 \le \ell \le r-1} t_{s_k} t_{\xi_k^\ell \xi_{k+1}^{-\ell}} t_{s_k} = t_{s_k} \bar{v}_{k+1} - \sum_{0 \le \ell \le r-1} t_{\xi_k^{-\ell} \xi_{k+1}^\ell}$$
$$= t_{s_k} \bar{v}_{k+1} - \sum_{0 \le \ell \le r-1} t_{\xi_k^\ell \xi_{k+1}^{-\ell}}.$$

6. References

[Bou] N. Bourbaki, Groupes et algebres de Lie, Chapt. IV-VI, Masson, Paris, 1981.

- [Ca] R. W. Carter, Conjugacy classes in the Weyl group, Compositio Math. 25 (1972), 1–59.
- [Dr] V. G. Drinfeld, Degenerate affine Hecke algebras and Yangians, Funct. Anal. Appl. 20 (1986), 58–60.

A. Ram and A. V. Shepler

- [G+] M. Geck, G. Hiss, F. Lübeck, G. Malle and G. Pfeiffer, CHEVIE A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras, AAECC 7 (1996), 175–210.
- [Hu] J. E. Humphreys, Reflection groups and Coxeter groups, Cambridge Studies in Advanced Mathematics 29, Cambridge University Press, Cambridge, 1990.
- [Lu] G. Lusztig, Affine Hecke algebras and their graded version, J. Amer. Math. Soc. 2 (1989), 599–635.
- [Lu2] G. Lusztig, Cuspidal local systems and graded Hecke algebras I, Inst. Hautes Études Sci. Publ. Math. 67 (1988), 145–202.
- [Mac] I. G. Macdonald, Symmetric functions and Hall polynomials, Second edition, Oxford Mathematical Monographs, Oxford University Press, New York, 1995.
- [OS1] P. Orlik and L. Solomon, Coxeter arrangements, Proc. Symp. Pure Math. 40 (1983), 269–291.
- [OS2] P. Orlik and L. Solomon, Arrangements defined by unitary reflection groups, Math. Ann. 261 (1982), 339–357.
- [OT] P. Orlik and H. Terao, Arrangements of hyperplanes, Springer-Verlag, Berlin-Heidelberg, 1992.
- [S+] M. Schönert et al., GAP Groups, Algorithms, and Programming, Lehrstuhl D für Mathematik, Rheinisch Westfälische Technische Hochschule, Aachen, Germany, fifth edition, 1995.
- [ST] G. C. Shephard and J. A. Todd, Finite Unitary Reflection Groups, Canad. J. Math. 6 (1954), 274–304.
- [St1] R. Steinberg, Differential equations invariant under finite reflection groups, Trans. Amer. Math. Soc. 112 (1964), 392–400.
- [St2] R. Steinberg, Endomorphisms of linear algebraic groups, Mem. Amer. Math. Soc. 80 (1968), 1–108.

Arun Ram Department of Mathematics University of Wisconsin Madison, WI 53706 USA e-mail: ram@math.wisc.edu Anne V. Shepler Department of Mathematics University of North Texas Denton, TX 76203 USA e-mail: ashepler@unt.edu

(Received: July 25, 2001)



To access this journal online: http://www.birkhauser.ch CMH