

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 78 (2003)

Artikel: Hyperbolic Coxeter groups of large dimension
Autor: Januszkiewicz, Tadeusz / wittkowski, Jacek
DOI: <https://doi.org/10.5169/seals-58770>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Hyperbolic Coxeter groups of large dimension

Tadeusz Januszkiewicz and Jacek Świątkowski

Abstract. We construct examples of Gromov hyperbolic Coxeter groups of arbitrarily large dimension. We also extend Vinberg's theorem to show that if a Gromov hyperbolic Coxeter group is a virtual Poincaré duality group of dimension n , then $n \leq 61$.

Coxeter groups acting on their associated complexes have been extremely useful source of examples and insight into nonpositively curved spaces over last several years. Negatively curved (or Gromov hyperbolic) Coxeter groups were much more elusive. In particular their existence in high dimensions was in doubt.

In 1987 Gabor Moussong [M] conjectured that there is a universal bound on the virtual cohomological dimension of any Gromov hyperbolic Coxeter group. This question was also raised by Misha Gromov [G] (who thought that perhaps any construction of high dimensional negatively curved spaces requires nontrivial number theory in the guise of arithmetic groups in an essential way), and by Mladen Bestvina [B2].

In the present paper we show that high dimensional Gromov hyperbolic Coxeter groups do exist, and we construct them by geometric or group theoretic but not arithmetic means.

Mathematics Subject Classification (2000). 20F55, 20F65, 20F67, 51F15, 57P10.

Keywords. Coxeter group, Gromov hyperbolic group, Poincaré duality, dimension.

Theorem 1. *For any n there exists a right angled Coxeter group (W, S) of virtual cohomological dimension $vc(W) = n$ which does not contain $Z \oplus Z$ and thus is Gromov hyperbolic.*

It is known by work of Mike Davis [D2] and Moussong [M] that this theorem reduces to the existence of a simplicial complex L_n which is (i) flag, (ii) contains no empty square and (iii) satisfies certain homological condition (which is implied by say requiring that L_n is an oriented pseudomanifold of dimension $n-1$). Such a complex gives rise to the Coxeter group (W, S) with generators indexed by vertices of L_n and relations $(x_i x_j)^2 = 1$ iff vertices i, j are connected by an edge. Then (i) implies that the nerve of (W, S) is L_n , (ii) implies hyperbolicity and (iii) implies that $vc(W) = n$.

Both authors were supported by a KBN grant 5 P03A 035 20.

We construct simplicial complexes required for the proof of Theorem 1 using complexes of groups technique of [BH]. The resulting spaces are interesting on their own right and we study them further in a forthcoming paper. Let us mention here only that the L_n we construct turn out to be $K(\pi, 1)$ spaces.

Gromov hyperbolic Coxeter groups of large dimension are very different from fundamental groups of closed aspherical manifolds. For example we have

Theorem 2. *If (W, S) is a virtual Poincaré duality Coxeter group of dimension n which does not contain $Z \oplus Z$ as a subgroup (equivalently, which is Gromov hyperbolic) then $n \leq 61$. Moreover, if (W, S) is right angled, the bound improves to $n \leq 4$ and is sharp.*

This theorem is an extension of a result of Ernest Borisovich Vinberg [V] (which motivated Moussong's question) which states that if there is a cocompact reflection group action on the Lobačevski space of dimension n then $n \leq 29$. Vinberg presents also a significantly easier proof of a weaker bound $n \leq 61$, which we follow closely. It was observed already by Moussong [M] that Vinberg's result uses only Gromov hyperbolicity. The new ingredient we use to obtain the present form of the Theorem is a recent result of Davis on virtual Poincaré duality Coxeter groups [D1].

The right angled case of Vinberg theorem has the following corollary dealing with arbitrary cubical complexes.

Corollary. *If M is a manifold of dimension greater than 4, then it does not admit a negatively curved piecewise hyperbolic cubical metric.*

The content of the paper is as follows.

Section 1 contains a necessary background on Coxeter groups.

Section 2 deals with the nonexistence results. We prove Theorem 2 following closely Vinberg's paper, discussing only points which are specific to our situation. We do not claim much originality here. The reader should be ready to consult [V]. We also establish a slightly stronger version of the Corollary.

In Section 3 we recall the simplex of groups construction in the form we need. We focus on the properties of (multi-) simplicial complexes arising as developments (coverings) of simplices of groups.

In Section 4 we define (and establish properties of) special complexes of groups which we call retractible. This is done by enriching the structure of a complex of groups with additional maps: retractions. We found functorial language very convenient here.

Section 5 contains the discussion of a further restriction on the complexes of groups. It is a secondary form of retractibility and we call it *extra retractibility*. The main result is Proposition 4.12 stating that extra retractibility implies the no-square condition on the associated complexes L .

Section 6 contains a construction of extra retractible complexes of groups in arbitrary dimension, thus concluding the proof of Theorem 1.

The final three short sections contain variations and applications of the main result. In Section 7 we show how to construct many $CAT(-1)$ cubical complexes by a hyperbolization procedure which uses complexes L_n constructed in the previous section.

Section 8 shows how to obtain non right-angled Gromov hyperbolic Coxeter groups from right-angled ones without changing dimension.

Finally, Section 9 addresses the issue of complexity. The complexes L_n we construct are strikingly large. We establish a lower bound on the complexity of simplicial flag no-square pseudomanifold of dimension n in general. While this is much smaller than the complexity of our examples, it is still fairly large.

We also want to call readers attention to [MS], where $CAT(-1)$ pseudomanifolds of large dimensions are constructed by a different (but related) method.

Acknowledgements. We thank Jan Dymara for useful discussions and the referee for careful reading of the manuscript.

1. Background on Coxeter groups

We present the necessary material concisely, since all of it is well documented in the literature, see for example [D2].

1.1. Generalities

Let $M = (m_{st})_{s,t \in S}$ be a symmetric matrix with diagonal entries equal 1 and off diagonal entries in $\{2, 3, \dots, \infty\}$. We associate with M a *Coxeter system* (W, S) , where W is a group, S is the set of involutions $s^2 = 1$ generating W and the nontrivial relations are $(st)^{m_{st}} = 1$. If $m_{st} = \infty$, there is no relation between s and t implied.

Let A be a finite graph, that is a symmetric, anti-reflexive relation on the vertex set S . Define a matrix

$$M^A = (m_{st}^A)_{s,t \in S} = \begin{cases} 1 & \text{if } s = t, \\ 2 & \text{if } s, t \text{ are connected by an edge in } A, \\ \infty & \text{otherwise.} \end{cases}$$

The Coxeter system (W^A, S) thus obtained is called the *right angled* Coxeter system associated with the graph A . When using the usual notation of Coxeter-Dynkin diagrams for Coxeter systems, one obtains A as the dual to the Coxeter-Dynkin diagram of M^A , which has an edge marked ∞ for any pair of non-commuting generators.

1.2. The nerve and the Davis complex

Subgroups generated by subsets of S are called *standard parabolic subgroups*. The parabolic subgroup generated by $T \subset S$ is denoted W_T . Finite standard parabolics (excluding the trivial group) ordered by inclusion form a partially ordered set. Its geometric realization [BH, p. 370] is called the *nerve* and denoted $N(W, S)$. A coarser simplicial structure on the nerve will be useful: its vertices are elements of S while a subset $T \subset S$ spans a simplex iff T generates a finite subgroup of W .

For right angled systems the nerve has a simple description in terms of the graph A . Namely form a simplicial complex A^f for which A is the one skeleton, and whose higher dimensional simplices correspond to complete subgraphs in A . Thus one completes A by filling in (inductively) a simplex if we see its boundary in the part of the complex already constructed. The result is the coarse simplicial structure of the nerve. The nerve $N(W, S)$ is the barycentric subdivision of A^f .

A simplicial complex $K = (K_{(1)})^f$ obtained as above from its one skeleton $K_{(1)}$ is called a *flag complex*.

There is a related poset, consisting of left cosets gW_T of all finite parabolics (this time including the trivial one), ordered again by inclusion. Its geometric realization is called the *Davis complex* of (W, S) and is denoted by $|W|$. It is a truncated version of the Coxeter complex of (W, S) . Clearly W acts on $|W|$ on the left (with a strict fundamental domain, [BH, p. 372]). The quotient of $|W|$ by W is the cone over the nerve $N(W, S)$.

1.3. Nonpositive and negative curvature

Since the simplicial cone over the barycentric subdivision of any simplicial complex carries a natural cubical structure, the Davis complex of a Coxeter group can be considered as a cubical complex. Thus it carries a geodesic metric, obtained by declaring each k -cube to be isometric with $[0, 1]^k$.

A fundamental observation, due to Gromov, is that a cubical complex is locally $CAT(0)$ (that is it is nonpositively curved in the comparison sense of Alexandrov and Toponogov) iff the link of any vertex is a flag complex. Moussong [M] amplified this observation, and showed what is the right choice of metrics on cubes yielding a $CAT(0)$ metric for an arbitrary Coxeter group. Moreover, he showed that if there is no “obvious” obstruction then this metric can be modified to a (W -invariant) $CAT(-1)$ metric on $|W|$. The obvious obstruction arises when either (W, S) contains an affine Coxeter system of rank at least 2 as a standard parabolic subgroup, or it contains a standard parabolic which is a product of two infinite Coxeter systems. Thus a right angled Coxeter group acts isometrically and cocompactly on a $CAT(-1)$ space iff it does not contain a product of two infinite dihedral groups as a standard parabolic. This condition is equivalent to the following combinatorial fact: *If a, b, c, d are vertices of A forming a square, ie. there are edges ab, bc, cd, da in A , then at least one of the edges ac, bd is in A .*

We say that a flag simplicial complex A^f as above contains no empty square

or that it is a *no-square complex*.

1.4. The virtual cohomological dimension

The virtual cohomological dimension of a Coxeter group, denoted $vc(W)$, has been computed by Bestvina [B1], in the form of an inductive algorithm giving the answer. A clearer formulation is due to Davis [D1], who gave a formula for cohomology of W with coefficients in the group ring (which gives the dimension). Namely

$$vc(W, S) = \max\{n : H^n(D, D^T; \mathbb{Z}) \neq 0\} = 1 + \max\{n : \tilde{H}^n(D^T; \mathbb{Z}) \neq 0\}$$

Here the maximum is taken over all $T \subset S$ such that W_T is finite, D denotes the cone over the nerve $N(W, S)$, and D^T is the union of closed simplices of $N(W, S)$ which contain as a vertex a singleton $s \in S - T$. The maximum over the empty set is -1 .

If T spans a finite parabolic subgroup of W , D^T is homotopy equivalent to the nerve of the parabolic subgroup spanned by $S - T$. The homotopy equivalence is obtained by deformation retraction of D^T onto the full subcomplex in the nerve $N(W, S)$ (with respect to the coarse simplicial structure on this nerve) spanned by the subset $S - T$ of vertices.

Thus

$$vc(W, S) = 1 + \max\{n : \tilde{H}^n(N(W_{S-T}, S-T); \mathbb{Z}) \neq 0\}.$$

Notice that if (W, S) is a right angled Coxeter group, whose nerve is a closed orientable pseudomanifold of dimension n (that is if each simplex of dimension $(n-1)$ is contained in exactly two simplices of dimension n , and the manifold obtained by removing simplices of codimension 2 is orientable), then $vc(W) = n+1$.

The discussion above shows that to construct a hyperbolic Coxeter group of dimension $n+1$ it is sufficient to construct a simplicial flag complex with no empty squares, which is an orientable pseudomanifold of dimension n . This will be done in Sections 3–6.

2. An extension of Vinberg's theorem

2.1. Coxeter groups and Poincaré duality

A group Γ is a virtual Poincaré duality group of dimension n , if for some finite index subgroup Γ_0 , the cohomology of $B\Gamma_0$ satisfies the Poincaré duality [D1].

A simplicial complex N is a *generalized homology sphere* (GHS) of dimension k if it has the same homology as S^k and if the link of each simplex in N is a

generalized homology sphere of appropriate dimension. In particular, each simplex is contained in a simplex of dimension k , and N is a pseudomanifold.

Let (W, S) be a Coxeter system such that W is a virtual Poincaré duality group of dimension n . It is a theorem of M. W. Davis [D1] that (W, S) decomposes into a product of Coxeter systems $(W, S) = (W_1, S_1) \times (W_2, S_2)$, so that W_2 is a finite group and the nerve of (W_1, S_1) is a generalized homology sphere of dimension $n - 1$.

2.2. Combinatorics of triangulations of generalized homology spheres

Let N be a finite simplicial complex of dimension $(n - 1)$ and let f_i denote the number of $(i - 1)$ -dimensional simplices in N ($f_0 = 1$). Define numbers h_i by the polynomial equation

$$\sum_{i=0}^n h_i t^i = \sum_{i=0}^n f_{i-1} (t-1)^{n-i}.$$

A theorem of Richard Stanley [S] asserts that if N is a generalized homology sphere of dimension $(n - 1)$ then

$$h_i = h_{n-i} \quad \text{and} \quad h_i \geq 0.$$

V. V. Nikulin (see [V, Prop. 6.1]) showed that these restrictions on h_i imply that

$$\binom{n}{2} \frac{f_{n-1}}{f_{n-3}} < \begin{cases} \frac{4(n-1)}{n-2} & \text{for } n \text{ even} \\ \frac{4n}{n-1} & \text{for } n \text{ odd.} \end{cases}$$

The quantity on the left hand side is the average number of $(n - 1)$ -simplices containing given $(n - 3)$ -simplex. If $n \geq 5$ the inequality implies that there exists an $(n - 3)$ -simplex in N contained in three or four $(n - 1)$ -simplices.

Suppose $A = \sigma_{n-3}$ is contained in three $(n - 1)$ -simplices. Since N is a pseudomanifold, any two of these three simplices intersect along an $(n - 2)$ -simplex (otherwise we would have more than three top dimensional simplices containing A). In other words, the simplex A contained in three $(n - 1)$ -simplices produces a subcomplex $Y_3(A) \subset N$ on $(n - 2) + 3 = n + 1$ vertices. The three additional vertices are called “black” vertices.

Similarly, if $A = \sigma_{n-3}$ is contained in four $(n - 1)$ -simplices $\sigma^0, \sigma^1, \sigma^2, \sigma^3$, the argument above implies that (up to renumbering) simplices σ^i, σ^{i+1} intersect along an $(n - 1)$ -simplex (taking indices mod 4). Thus we have a subcomplex $Y_4(A) \subset N$. Four vertices of $Y_4(A)$ outside A are again called “black”, and we see that they form a square.

Proof of the right angled case of Theorem 2. Let (W, S) be a right angled Coxeter system whose nerve $N = N(W, S)$ is a generalized homology sphere of dimension $(n - 1) \geq 4$. Recall that (by the construction of the nerve) N is a flag complex.

If N contains Y_3 , it contains the one skeleton of n dimensional simplex, hence it is not of dimension $(n-1)$.

If N contains Y_4 , then it contains an empty square, since any additional edge between black vertices in Y_4 would produce a n -simplex. Thus the group W contains $Z \oplus Z$ generated by the black vertices in Y_4 and is not hyperbolic.

To see that the bound is sharp, one uses the 120 cell. \square

Proof of the general case of Theorem 2. Vinberg considers weights $w(A, B)$ associated to flags $f = (A \subset B)$ of a codimension 2 and top dimensional simplex in a GHS of dimension $n-1$. If the following conditions are met with some constant c , for all simplices A, B :

$$w(B) = \sum_{A|A \subset B} w(A, B) \leq cn \quad (1)$$

$$w(A) = \sum_{B|B \supset A} w(A, B) \geq 5 - |A|, \quad (2)$$

then $n < 8c + 6$ (Prop 6.2 of [V]). Here $|A|$ denotes the number of top dimensional simplices containing A .

Choose the weights as follows: look at the two “black” (that is not contained in A) vertices of B and compute their distance in the Dynkin diagram of (W, S) ; if the distance is at most 7, the weight is 1, otherwise it is 0.

In [V, Lemma 6.1] condition (1) is checked with $c = 7$. It is a fact about any finite Coxeter system, thus holds in the present generality.

Condition (2) has to be checked only for A such that $|A| = 3, 4$.

If $|A| = 3$, first note that the subgroup W_Y generated by the vertices of $Y_3(A)$ is infinite (otherwise N would contain an n -simplex). Take a minimal infinite parabolic subgroup $L < W_Y$ containing all black vertices and denote the generating set inherited from W_Y by l . Removing a black vertex from l yields a finite subgroup by construction of $Y_3(A)$. Removing a white vertex yields a finite subgroup by minimality of L . The subsystem (L, l) is irreducible since otherwise it would contain an infinite factor. Thus L is either affine or hyperbolic since its associated quadratic form contains a positive definite subspace of codimension one. Since L has at least 3 generators, it cannot be affine: an irreducible affine Coxeter group on $k+1$ generators contains a subgroup isomorphic to Z^k . Thus L is one of the Lanner groups (i.e. cocompact reflection groups on a Lobačevskii space, whose fundamental domain is a simplex), and Vinberg showed that $w(A) \geq 2 = 5 - |A|$.

If $|A| = 4$, an argument similar to the one above shows that a minimal infinite parabolic subgroup of $Y_4(A)$ containing two opposite black vertices, and not containing the other pair of black vertices, is a Lanner group, and hence Vinberg’s argument estimating the sum of weights $w(A) \geq 1 = 5 - |A|$ goes through.

Since $c = 7$, the bound $n < 8c + 6 = 62$ follows and the proof is complete. \square

2.3. Examples and comments

There are some issues related to Theorem 2 which still should be addressed. First, there is a question if the bound proved is sharp. Vinberg provides a better bound $n \leq 29$ for cocompact reflection groups acting on Lobačevski hyperbolic space; we did not try to adapt his argument to the present context.

It is quite conceivable that the maximum dimension of a Gromov hyperbolic Poincaré duality group is not attained on a Lobačevski hyperbolic group. Anyway there are many more Gromov hyperbolic Poincaré Duality Coxeter groups than Lobačevski hyperbolic.

Example 2.3.1. Let W be a cocompact right angled Coxeter group on H^4 with fundamental domain (and the quotient) the simple polytope P . Thus faces of P correspond to generators of W , and the relations are $s^2 = 1$ and $(st)^2 = 1$ iff faces corresponding to s, t intersect. Let m_{st} be the Coxeter matrix of (W, S)

We change the orbifold structure on P , by choosing a face of codimension 2 (thus two faces of codimension 1, s_0 and t_0 , containing it), and declaring the dihedral angle along that face to be $\frac{\pi}{n}$. In other words we take the Coxeter system (W_n, S) which has the same system of generators as W , and the Coxeter matrix is

$$n_{s,t} = \begin{cases} m_{s,t} & \text{if } (s,t) \neq (s_0, t_0), \\ n & \text{otherwise.} \end{cases}$$

Coxeter systems (W, S) and (W_n, S) have isomorphic nerves, thus the Davis complex of W_n is homeomorphic to R^4 . The fundamental domain for W_n , denoted P_n , is combinatorially isomorphic to P .

Using Moussong's theorem we see that W_n is Gromov hyperbolic (an elementary smoothing argument shows that there exists a W_n invariant smooth Riemannian metric of negative curvature on $|W_n|$). We claim that W_n does not act discretely cocompactly by isometries on H^4 .

Suppose it does. Since the argument becomes simpler for $n = 2k$, we assume that. First, since the ideal boundaries of $|W_n|$ and H^4 are W_n -equivariantly homeomorphic (uniqueness of the boundary of a hyperbolic group), W_n acts on H^4 as a reflection group and H^4 is equivariantly homeomorphic to $|W_n|$.

Consider a generator s and the action of the centralizer of s on the $Fix(s) = H^3$; we easily see that $Z_{W_n}(s)$ is isomorphic to $Z_W(s)$ (this is where the condition $n = 2k$ is useful), and their actions on H^3 are isometric.

Thus the polytopes P, P_n are combinatorially equivalent, and their corresponding faces are isometric. This implies the fundamental domains, hence the Coxeter systems (W, S) and (W_n, S) are isomorphic, a contradiction.

This example essentially comes from [GT], where the smoothing to negatively curved metric is described. The change of the orbifold structure of the quotient is analogous to taking a ramified cover. The argument we used is an elementary

substitute to Mostow rigidity. One can produce many more examples of this kind by changing dihedral angles along the set of disjoint codimension 2 faces of P . These variations seem to give an enormous supply of negatively curved 4 dimensional reflection groups.

Example 2.3.2. Higher dimensional examples are harder to construct. The largest known dimension of a cocompact Lobačevski hyperbolic Coxeter group is eight and it has been found in [Bu]. On the other hand we have the following easy example of an eight dimensional Gromov hyperbolic Coxeter group.

Consider a Coxeter group W with 10 generators, whose Dynkin diagram is an interval with labels $(5, 3, 3, 3, 3, 3, 3, 3, 5)$ on the edges. One readily checks with Moussong theorem that it is Gromov hyperbolic. The Davis complex is homeomorphic to R^8 (if instead of the 3 in the middle one puts 2, ie. if one cuts the middle edge, the resulting group is the product of two 4-dimensional hyperbolic Lanner groups, and the two groups have the same poset of finite parabolic subgroups).

One checks with an argument similar to that in Example 2.3.1 that W is not Lobačevski hyperbolic.

2.4. Nonexistence of cubical $CAT(-1)$ metrics

Proof of the Corollary. A piecewise hyperbolic metric on a cubical complex is a geodesic metric obtained by declaring each cube to be isometric to the symmetric cube in the Lobačevskii space of curvature -1 , of some given edge length.

It is a theorem of Gromov, (a more general result is proved in [M]) that for a cubical complex X , there exists an edge length of the cube such that the cubical metric is locally $CAT(-1)$ iff the link of any vertex in the complex is a flag complex which satisfies the no-square condition. By the discussion in Section 1 existence of such a metric on a manifold of dimension $n \geq 5$ implies the existence of a $CAT(-1)$ right-angled Coxeter group, for which the Davis complex is an n -manifold. This contradicts Theorem 2. \square

Observe that we indeed proved a stronger result:

Proposition. 2.4.1. *Let X be a $CAT(-1)$ cubical complex and let c be a cube of codimension $(k+1)$ (so that the link of c has dimension k). If $k \geq 4$, then the link of c is not a generalized homology sphere.* \square

Proposition 2.4.1 shows that not every link can arise in a $CAT(-1)$ cubical complex. Simple arguments with subdivision [Dr] show there is no such obstruction for links of dimension ≤ 2 . It is unclear to us what happens for links of dimension 3. It would also be interesting to obtain further restrictions on links in higher dimensions.

Remark. One should not become too enthusiastic about the Corollary. It is very

likely that many manifolds with locally $CAT(0)$ cubical metrics have Gromov hyperbolic fundamental groups and carry (noncubical) $CAT(-1)$ metrics. Examples of this kind are known to us in low dimensions.

3. Simple complexes of groups

We will construct complexes required for the proof of Theorem 1 as normal covers of certain simple complexes of groups. Such complexes are necessarily fairly large, and we view their symmetry groups as an efficient bookkeeping device. Recall first quickly the formalism of simple complexes of groups [BH, II.12].

Let (\mathcal{Q}, \leq) be a partially ordered set; consider it as a small category with an object for each element in \mathcal{Q} and a unique morphism $S \rightarrow T$ for any pair of elements with $S \leq T$. A simple complex of groups over \mathcal{Q} , denoted $G(\mathcal{Q})$, is a functor from \mathcal{Q} to the category of groups and injective maps. A complex is developable if all the groups G_S inject into the direct limit $\widehat{G(\mathcal{Q})}$.

For a developable complex together with a homomorphism $\phi : \widehat{G(\mathcal{Q})} \rightarrow G$, injective on local groups G_S , one performs The Basic Construction ([BH, II.12.18]). It gives a space $D(\mathcal{Q}, \phi)$ called the development of $G(\mathcal{Q})$ associated to ϕ . The group G acts on $D(\mathcal{Q}, \phi)$ with a strict fundamental domain $|\mathcal{Q}|$, which is isomorphic to the geometric realization of the poset \mathcal{Q} . The isotropy subgroups of points in the fundamental domain are the local groups G_S .

3.1. Developable simplices of groups

For a finite set S , let Δ_S be the simplex dual to S , i.e. the simplex of dimension $|S|-1$, with codimension one faces labeled by elements of S and denoted $\sigma_s : s \in S$. For a nonempty subset $T \subset S$ we put $\sigma_T = \sigma_T^S := \cap_{s \in T} \sigma_s$. We also use the convention that $\sigma_\emptyset = \sigma_\emptyset^S := \Delta_S$. For vertices σ_{S-s} we use shorter notation v_s . In this notation, the vertex v_s is opposite to the face σ_s , so that $\Delta_S = v_s * \sigma_s$. Moreover, any face σ_T of Δ_S is spanned by the vertices $v_s : s \in S - T$. For a point $p \in \Delta_S$, denote by $S(p)$ the (unique) subset of S such that $p \in \text{int } \sigma_{S(p)}$.

Let $\mathcal{Q} = P(S)$ be the poset of all subsets of S and let F be a simple complex of groups over \mathcal{Q} . In other words, to any $T \subset S$ assign a group F_T , so that if $U \subset T$, then $F_U \subset F_T$. Groups of the form $F_{\{s\}}$ will be called *singleton groups*.

Restricting F to the subposet $\mathcal{Q} \setminus \{S\}$ we obtain a simplex of groups $F(\Delta_S)$ equipped with the homomorphism $\phi_S : F(\Delta_S) \rightarrow F_S$ induced by inclusions $F_T \rightarrow F_S$.

Define an equivalence relation \sim on $\Delta_S \times F_S$ by

$$(p, \gamma) \sim (q, \eta) \quad \text{iff} \quad p = q \quad \text{and} \quad \gamma^{-1}\eta \in F_{S(p)};$$

denote by $[p, \eta]$ the equivalence class of (p, η) , let

$$L_S = \Delta_S \times F_S / \sim.$$

This is the Basic Construction of [BH, II.12] applied to the pair $(F(\Delta_S), \phi_S)$, except that we insist on using a coarser simplicial structure than [BH] (who use the barycentric subdivisions of our simplices).

Proposition 3.2.

- (1) L_S is a pure¹($|S| - 1$)-dimensional multisimplicial² complex.
- (2) The group F_S acts on L_S by $\gamma[p, \eta] = [p, \gamma\eta]$, and the quotient map is equal to the map induced by the projection onto the first factor. The stabilizer of the point $[p, 1]$ is $F_{S(p)}$.
- (3) L_S is finite iff the index $[F_S : F_\emptyset]$ is finite. L_S is locally finite iff all indices $[F_R : F_\emptyset]$ for proper subsets $R \subset S$ are finite.
- (4) L_S is a closed pseudomanifold iff $[F_{\{s\}} : F_\emptyset] = 2$ for all singleton groups $F_{\{s\}}$.
- (5) L_S is gallery connected³ iff the singleton subgroups $F_{\{s\}}$ generate F_S .
- (6) L_S is an orientable pseudomanifold iff in addition to (4) there is a homomorphism $\rho : F_S \rightarrow \mathbb{Z}_2$ whose restriction $\rho_s : F_{\{s\}} \rightarrow \mathbb{Z}_2$ has kernel equal to F_\emptyset for each $s \in S$.
- (7) L_S is a simplicial complex iff $F_R \cap F_T = F_{R \cap T}$ for all subsets of S .

Proof. Assertions (1) and (2) are obvious. The first part of (3) follows by observing that top dimensional simplices in L_S are indexed by cosets F_S/F_\emptyset . Similarly, if σ is a simplex in L_S whose image under the quotient map $L_S \rightarrow \Delta_S$ is σ_T , then the set of top dimensional simplices containing σ is indexed by the cosets F_T/F_\emptyset . This implies the second part of (3). The same argument applied to simplices σ of codimension 1 proves (4).

Assertion (5) follows by observing that simplices $[\Delta_S, g_1]$ and $[\Delta_S, g_2]$ share a codimension 1 face iff $g_2 = g_1g$ for some g in one of the singleton groups $F_{\{s\}}$.

To prove (6), suppose that L_S is a pseudomanifold. Pulling back a fixed orientation of Δ_S by the quotient map $L_S \rightarrow \Delta_S$, we equipped top dimensional simplices of L_S with orientations which are opposite for any pair of adjacent (i.e. sharing a codimension 1 face) simplices. Homomorphism ρ as in (6) induces the map from top dimensional simplices to \mathbb{Z}_2 with a similar property: its values at any pair of adjacent simplices are distinct. Modify the induced orientations by reversing them on simplices assigned with the nonzero value in \mathbb{Z}_2 . This makes the orientations consistent and proves one of the implications in (6). The converse follows by a similar argument.

¹ X is pure n -dimensional if any $x \in X$ is in a closure of an n -dimensional simplex.

² In a multisimplicial complex there may be more than one simplex with a given vertex set, but the complex is still a sum of embedded closed simplices intersecting along sums of their faces.

³ A pure complex X is gallery connected if for any two top dimensional simplices Δ_1, Δ_2 there is a sequence of top dimensional simplices $\Delta_1 = y_1, y_2 \dots y_k = \Delta_2$, so that y_i has a common face of codimension 1 with y_{i+1} .

To prove (7), observe that k -dimensional simplices in L_S are indexed by cosets of the form F_S/F_{S-K} , where K is a k -element subset of S . Assignment of the set of its vertices to a k -simplex gF_{S-K} can be viewed as a map $F_S/F_{S-K} \rightarrow \prod_{s \in K} F_S/F_{S-\{s\}}$ mapping gF_{S-K} to $gF_{S-\{s\}}$. Clearly this map is equivariant with respect to the left action of F_S . A simplex is uniquely determined by the set of its vertices iff the map $F_S/F_{S-K} \rightarrow \prod_{s \in K} F_S/F_{S-\{s\}}$ is injective. This happens iff $F_{S-K} = \bigcap_{s \in K} F_{S-\{s\}}$. The last equality is satisfied for all nonempty subsets K iff $F_R \cap F_T = F_{R \cap T}$ for all subsets. \square

4. Retractability

We start by enriching the category $P(S)$ with additional morphisms. Consider an element $1 \notin S$, called a *basepoint*, and put $T_* := T \cup \{1\}$ for each $T \in P(S)$. As the set of morphisms from R to T declare the set of all maps $m : R_* \rightarrow T_*$ with the property that $m(s) \in \{1, s\}$. Denote the new category by $P_r(S)$.

Observe that for each morphism $m : R \rightarrow T$ we have $m(1) = 1$ and $m(s) = 1$ for all $s \in R - T$.

For any pair R, T there is a distinguished morphism $m_{RT} : R_* \rightarrow T_*$, given by

$$m_{RT}(s) = \begin{cases} s & \text{if } s \in R \cap T, \\ 1 & \text{otherwise.} \end{cases}$$

Morphisms m_{RT} with $R \subset T$ and $T \subset R$ respectively, will be called *inclusions* and *retractions*. Since for any morphism $m : R \rightarrow T$ we have $m = m_{m(R)-\{1\}, T} \circ m_{R, m(R)-\{1\}}$, it follows that any morphism is a composition of an inclusion and a retraction. In particular, for any R, T we have $m_{RT} = m_{T \cap R, R} \circ m_{T, T \cap R}$.

Definition 4.1. A *retractible complex of groups* over the poset $P(S)$ is a covariant functor F from the category $P_r(S)$ to the category of groups.

We still denote by F_T the group assigned by a functor to a subset $T \subset S$. For any subsets $R, T \subset S$, the homomorphism induced by the morphism m_{RT} is denoted by $h_{RT} : F_R \rightarrow F_T$.

Lemma 4.2.

- (1) *Homomorphisms induced by inclusions $R \subset T$ are injective.*
- (2) *Homomorphisms h_{TR} with $R \subset T$ (i.e. homomorphisms induced by retractions) map the subgroup $F_R \subset F_T$ identically to itself.*
- (3) *If $R, U \subset T$, then $h_{TR}|_{F_U} = h_{U \cap R, R} \circ h_{U, U \cap R}$, and hence $h_{TR}(F_U) = F_{U \cap R}$. In particular, $h_{TR}(F_{\{s\}}) = F_{\emptyset}$ for $s \in T - R$.*
- (4) *If $R, T \subset S$, then $F_R \cap F_T = F_{R \cap T}$ as subgroups of F_S .*

Proof. If $R \subset T$ then by functoriality we have $h_{TR} \circ h_{RT} = id_{F_R}$, which proves both (1) and (2). Property (3) follows from (1) and (2) by noting that $h_{TR} \circ$

$h_{UT} = h_{U \cap R, R} \circ h_{U, U \cap R}$ (which follows by functoriality from a similar identity for morphisms in the category $P_r(S)$).

To see (4), note that $h_{SR}|_{F_R} = id_{F_R}$ and $h_{SR}(F_T) = F_{T \cap R}$. Thus, if $x \in F_R \cap F_T$, then $x = h_{SR}(x) \in F_{T \cap R}$, and hence $F_R \cap F_T \subset F_{R \cap T}$. The opposite inclusion is obvious. \square

If F is a retractible complex of groups on $P(S)$, the associated complex L_S has additional properties.

Proposition 4.3. *Let F be a retractible complex of groups on the poset $P(S)$ and let L_S be the associated development resulting from the basic construction in 3.1.*

- (1) *L_S is a simplicial complex.*
- (2) *If $v \in L_S$ is a vertex that projects to v_s under the quotient map $L_S \rightarrow \Delta_S$, then the star $st(v, L_S)$ of this vertex in L_S is a full subcomplex isomorphic to the simplicial cone over the development $L_{S-\{s\}}$ (associated to the complex of groups F restricted to $P(S-\{s\})$).*
- (3) *L_S is flag.*
- (4) *If for each $s \in S$ we have $[F_{\{s\}} : F_{\emptyset}] = 2$ (in particular if $F_{\emptyset} = \{1\}$ and $F_{\{s\}} = \mathbb{Z}_2$) then L_S is an orientable pseudomanifold.*

Proof. Property (1) is an immediate consequence of Proposition 3.2(7) and Lemma 4.2 (4).

To prove (2) we use the complex D_s introduced in Section 5, just before Lemma 5.1. Observe that the star of $v = [v_s, g]$ in L_S is equal to the translate gD_s of D_s . By Lemma 5.2, this star is then isomorphic to the simplicial cone over $L_{S-\{s\}}$. To prove that this star is full, it is sufficient to prove that D_s is full in L_S . Let τ be the simplex of L_S with all vertices in D_s . By Lemma 5.1, the complex D_s is a strict fundamental domain for the action of a subgroup $H_s^S \subset F_S$ on L_S . Thus the image of the simplex τ under the quotient map $L_S \rightarrow D_s$ (associated to the action of H_s^S) has the same vertices as τ , and hence is equal to τ . It follows that τ is contained in D_s , which finishes the proof of (2).

Please note that proofs of Lemmas 5.1 and 5.2 which we used are independent of results in Section 4.

A simplicial complex X with full stars of vertices is flag iff all stars of vertices in X are flag. Moreover, a cone over a flag complex is again flag. Thus one can use (2) inductively to conclude L_S is flag. To start induction we note that 0-dimensional complexes are flag. This proves (3).

The hypothesis (4) yields a homomorphism $\rho_s : F_{\{s\}} \rightarrow \mathbb{Z}_2$ as in Proposition 3.2 (6) for each $s \in S$. A homomorphism $\rho : F_S \rightarrow \mathbb{Z}_2$ given by the formula

$$\rho(g) := \sum_{s \in S} \rho_s \circ h_{S, \{s\}}(g),$$

is well defined and extends all homomorphisms ρ_s for $s \in S$, which can be easily verified using Lemma 4.2. Applying Proposition 3.2(6) finishes the proof. \square

Definition 4.4. A retractible complex of groups F over the poset $P(S)$ is *reduced* if $F_\emptyset = \{1\}$.

Observe that to any retractible complex of groups F there is associated a reduced complex F^0 over the same poset, with groups $F_T^0 := \ker(h_{T,\emptyset} : F_T \rightarrow F_\emptyset)$ and with the induced homomorphisms equal to the restrictions of the corresponding homomorphisms for F . The development L_S^0 associated to F^0 is isomorphic to L_S , and the isomorphism is given by $[p, g] \rightarrow [p, h_{S,\emptyset}(g)]$.

Definition 4.5. A retractible complex of groups F over the poset $P(S)$ is *minimal* if each of the groups F_T is generated by the union of the singleton subgroups $F_{\{s\}} : s \in T$.

To any retractible complex of groups F there is associated a minimal complex F^{\min} over the same poset, with groups $F_T^{\min} := \langle \cup_{s \in T} F_{\{s\}} \rangle$ and with the induced homomorphisms equal to the restrictions of the corresponding homomorphisms for F . The development L_S^{\min} associated to F^{\min} is easily seen to be the gallery connected component of the simplex $[\Delta_S, 1]$ in L_S . The whole L_S is isomorphic to the disjoint union of F_S/F_S^{\min} copies of L_S^{\min} , glued along some faces of codimension at least 2.

4.6. Examples of retractible complexes of groups

First examples of functors on $P_r(S)$ are given by direct product and free product of singleton groups. More specifically assign to any $s \in S$ a group G_s ; then put $F_T^{\text{direct}} := \prod_{s \in T} G_s$ and $F_T^{\text{free}} := *_{s \in T} G_s$.

For another example, suppose that S is a vertex set of a graph A . With $T \subset S$ we associate the direct limit of the poset of groups $\lim \text{dir } \prod_{s \in K} G_s$, where $K \subset T$ spans a complete subgraph of A . This yields the graph product functor over $P_r(S)$.

Finally, suppose that we are given an infinite Coxeter matrix $(m_{s,t})_{s,t \in S}$, with even off diagonal elements (∞ is declared an even number). Then assigning to $T \subset S$ the Coxeter group with the matrix $(m_{s,t})_{s,t \in T}$ we get a functor on $P_r(S)$. The restriction to even entries is necessary to have homomorphisms induced by retractions.

The examples described above are all reduced and minimal. Note that any reduced and minimal functor F on $P_r(S)$ is sandwiched between the free and the direct product of its singleton groups. More precisely, there are natural transformations yielding surjective homomorphisms $F_T^{\text{free}} \rightarrow F_T \rightarrow F_T^{\text{direct}}$ for each $T \subset S$.

4.7. Examples of developments

We discuss developments L_S associated to functors from 4.6.

For $F = F^{\text{direct}}$, the complex L_S is the simplicial join $*_{s \in S} F_{\{s\}}$. In particular,

if all the singleton groups are Z_2 , F_T^{direct} is Z_2^T and the complex L_S is isomorphic to the standard hyperoctahedral triangulation of the sphere $S^{|S|-1}$. In the general case, when the singleton groups are nontrivial, L_S is a spherical Tits building with hyperoctahedral apartments.

If $F = F^{\text{free}}$ and the singleton groups are Z_2 , the complex L_S is isomorphic to a thickening of a regular S -valent tree with $(|S| - 1)$ -dimensional simplices.

More generally, if F is a graph product functor based on a graph A with the vertex set S , the space L_S is the Tits (simplicial) building associated with the right angled Weyl group given by the matrix M^A . The thickness of the building at the face corresponding to s is equal $|F_{\{s\}}|$.

Finally, if F comes from an even Coxeter matrix M , L_S is the Coxeter complex of the Coxeter system (F_S, S) given by the matrix M . Here we identify the elements $s \in S$ with the generators in the singleton subgroups $F_{\{s\}} = Z_2$.

5. Extra retractability and no-square condition

We say that a simplicial complex X contains *no empty square* (or satisfies the *no-square* condition) if for any vertices a, b, c, d in X connected by edges ab, bc, cd and da (thus forming a square $abcd$ in X) at least one of the edges ac or bd is in X .

The hyperoctahedral triangulation of $S^{|S|-1}$ mentioned in 4.7 shows that the space L_S associated to an arbitrary retractible complex of groups may contain empty squares. Thus when trying to construct no-square flag complexes, we need further restrictions on the complex of groups. To formulate them a preliminary discussion is necessary.

We will often deal with sets of form $S \setminus \{s\}$, and we use a simplified notation $S - s$ for them. We also denote by s^g the conjugation gsg^{-1} , and apply the same convention $H^g = gHg^{-1}$ for conjugation of subgroups.

Let F be a retractible complex of groups over $P(S)$. For a subset $T \subset S$ and for any $s \in T$, denote by H_s^T the kernel of the homomorphism $h_{T, T-s} : F_T \rightarrow F_{T-s}$. We focus attention on the groups H_s^S and their actions on the complex L_S . Put

$$D_s = \bigcup_{g \in F_{S-s}} [\Delta_S, g] \subset L_S.$$

Lemma 5.1. *The subcomplex D_s is a strict fundamental domain for the action of H_s^S on L_S , i.e. the restriction of the quotient map $q_s : L_S \rightarrow H_s^S \setminus L_S$ to D_s is a bijection.*

Proof. The map $j_s : H_s^S \setminus L_S \rightarrow D_s$ defined by

$$j_s(H_s^S \cdot [p, \gamma]) = [p, h_{S, S-s}(\gamma)]$$

is the inverse of $q_s|_{D_s}$. □

Lemma 5.2. *The complex D_s is isomorphic to the simplicial cone $C(L_{S-s})$, with vertex v_s as cone vertex.*

Proof. Since we have $\Delta_S = v_s * \sigma_s$, we write

$$D_s = \Delta_S \times F_{S-s} / \sim = (v_s * \sigma_s) \times F_{S-s} / \sim \cong v_s * (\Delta_{S-s} \times F_{S-s} / \sim_s) = v_s * L_{S-s},$$

where \sim_s above denotes the equivalence relation used to define L_{S-s} . The isomorphism in the middle comes from comparing the equivalence relation \sim restricted to $\sigma_s \times F_{S-s}$ and the equivalence relation \sim_s . We omit the straightforward details. \square

We will view the complex D_s as a stratified space: its top closed stratum is D_s and remaining closed strata are the simplices not containing the center v_s . The boundary ∂D_s of D_s is the union of all nonmaximal strata $\cup_{g \in F_{S-s}} [\sigma_s, g]$. By Lemma 5.2, the boundary is isomorphic to the complex L_{S-s} . For a point $p \in D_s$ denote by $\sigma(p)$ the unique minimal stratum (face) of D_s with $p \in \sigma(p)$. Lemma 5.1 implies the following.

Lemma 5.3. *For each face σ of D_s let Γ_σ be the stabilizing subgroup of the group H_s^S at the face σ . We then have*

$$L_S = D_s \times H_s^S / \sim,$$

where $(p, \gamma) \sim (q, \eta)$ iff $p = q$ and $\gamma^{-1}\eta \in \Gamma_{\sigma(p)}$.

Now we will take a closer look at the subgroups Γ_σ .

Lemma 5.4.

- (1) *Each face $\sigma \subset \partial D_s$ has a form $\sigma = [\sigma_T, g]$ for some subset $T \subset S$ containing s and for some $g \in F_{S-s}$.*
- (2) *If $\sigma = [\sigma_T, g]$ for some $s \in T \subset S$ and some $g \in F_{S-s}$, then $\Gamma_\sigma = (H_s^T)^g \subset H_s^S$.*

Proof. Only the second assertion requires a proof. Observe that $\text{Stab}(F_S, [\sigma_T, 1]) = F_T$, and hence $\text{Stab}(F_S, [\sigma_T, g]) = (F_T)^g$. Using the fact that the subgroup H_s^S is normal in F_S , we have $\Gamma_\sigma = \text{Stab}(H_s^S, [\sigma_T, g]) = (F_T)^g \cap H_s^S = (F_T \cap H_s^S)^g = (F_T \cap \ker h_{S, S-s})^g = (\ker h_{T, T-s})^g = (H_s^T)^g$. \square

From now on all retractible complexes of groups will be reduced and minimal. Under this restriction we obtain nice descriptions of various groups in terms of generators.

Lemma 5.5. *If F is reduced and minimal then the subgroup H_s^T is generated by the union of the conjugates $(F_{\{s\}})^g : g \in F_{T-s}$.*

Proof. By minimality the subgroups F_{T-s} and $F_{\{s\}}$ generate F_T . Since F is reduced, we have $F_{\{s\}} \subset H_s^S$. The lemma follows by combining these two facts.

Next lemma describes the subgroups Γ_σ in terms of generators.

Lemma 5.6. *Let $\sigma = [\sigma_T, \gamma] \subset \partial D_s$. Then the subgroup $\Gamma_\sigma \subset H_s^S$ is generated by the union of the conjugates $(F_{\{s\}})^g : g \in \gamma \cdot F_{T-s}$.*

Proof. Applying Lemmas 5.4 and 5.5 we have

$$\begin{aligned} \Gamma_\sigma &= (H_s^T)^\gamma = \langle h^\gamma : h \in \bigcup_{g \in F_{T-s}} (F_{\{s\}})^g \rangle = \langle \bigcup_{g \in F_{T-s}} (F_{\{s\}})^{\gamma g} \rangle \\ &= \langle \bigcup_{g \in \gamma \cdot F_{T-s}} (F_{\{s\}})^g \rangle. \end{aligned}$$

□

Observe that the top dimensional simplices of ∂D_s have the form $[\sigma_s, g] : g \in F_{S-s}$ and that they correspond bijectively to the elements of F_{S-s} . Put $\sigma_g := [\sigma_s, g]$. The following reformulation of Lemma 5.6 will be useful later in this section.

Lemma 5.7. *Let σ be a simplex of ∂D_s . Then the group Γ_σ is generated by the union of the conjugates $(F_{\{s\}})^g : \sigma \subset \sigma_g$.*

Proof. The inclusion $\sigma = [\sigma_T, \gamma] \subset [\sigma_s, g] = \sigma_g$ holds iff $\gamma^{-1}g \in F_{T-s}$ iff $g \in \gamma \cdot F_{T-s}$. □

Definition 5.8. A reduced, minimal, retractible complex of groups F over the poset $P(S)$ is *extra retractible* if for each $s \in S$ and for any subset $T \subset S$ containing s , there is a (uniquely determined) homomorphism $r_{s,T}^S : H_s^S \rightarrow H_s^T$ such that:

- (1) $r_{s,T}^S(h) = h$ if $h \in (F_{\{s\}})^g$ and $g \in F_{T-s}$;
- (2) $r_{s,T}^S(h) = 1$ if $h \in (F_{\{s\}})^g$ and $g \in F_{S-s} - F_{T-s}$.

Note that condition (1) above implies that $r_{s,T}^S(h) = h$ for each $h \in H_s^T$ and thus the homomorphism $r_{s,T}^S$ is a retraction. This explains why we call the property from Definition 5.8 extra retractibility.

Lemma 5.9. *Let F be an extra retractible complex of groups over the poset $P(S)$. Then for each simplex $\sigma \in \partial D_s$ there is a (uniquely determined) homomorphism $r_\sigma : H_s^S \rightarrow \Gamma_\sigma$ such that*

- (1) $r_\sigma(h) = h$ if $h \in (F_{\{s\}})^g$ and $\sigma \subset \sigma_g$;
- (2) $r_\sigma(h) = 1$ if $h \in (F_{\{s\}})^g$ and σ is not contained in σ_g .

Proof. Let $\sigma = [\sigma_T, \gamma]$. In view of Lemmas 5.6 and 5.7, the homomorphism defined by $r_\sigma(h) := [r_{s,T}^S(h^{\gamma^{-1}})]^\gamma$ is as required. To see this, observe that if h is as in (1)

then $h = k^{\gamma g}$ for some $k \in F_{\{s\}}$ and for some $g \in F_{T-s}$. Thus

$$r_\sigma(h) = r_\sigma(k^{\gamma g}) = [r_{s,T}^S((k^{\gamma g})^{\gamma^{-1}})]^\gamma = [r_{s,T}^S(k^g)]^\gamma = (k^g)^\gamma = k^{\gamma g} = h.$$

Similarly, one shows that $r_\sigma(h) = 1$ if h is as in (2). \square

Remark.

- (1) Existence of homomorphisms r_{σ_g} as in Lemma 5.9 implies that for distinct $g_1, g_2 \in F_{S-s}$ we have $(F_{\{s\}})^{g_1} \cap (F_{\{s\}})^{g_2} = \{1\}$.
- (2) Remark (1) above implies that the complex of groups F^{direct} (for which $F_T^{\text{direct}} = (Z_2)^T$) is not extra retractible, since the groups in this complex are abelian.
- (3) Slightly less clear, but still true, is the fact that the complex of groups F^{free} is extra retractible. However, the groups F_T in this complex are in general infinite and hence useless for our purpose. In Section 6 we construct a functor F which is extra retractible and for which all the groups F_T are finite.

Remark (1) above can be generalized as follows.

Lemma 5.10. *Let σ and τ be two simplices of ∂D_s that are not contained in any common simplex of ∂D_s . Then $\Gamma_\sigma \cap \Gamma_\tau = \{1\}$.*

Proof. The groups Γ_σ and Γ_τ are generated by the unions of the families of conjugates $(F_{\{s\}})^g : \sigma \subset \sigma_g$ and $(F_{\{s\}})^g : \tau \subset \sigma_g$ respectively. Due to our assumption on σ and τ , these two families are disjoint. It follows that the homomorphism r_σ as in Lemma 5.9 maps Γ_σ identically to itself, and it maps all of Γ_τ to 1. This implies that $\Gamma_\sigma \cap \Gamma_\tau = \{1\}$. \square

We now turn to describing the complex L_S as tessellated by the isomorphic copies of D_s . We take as tiles the translates $h \cdot D_s : h \in H_s^S$ and denote the whole tessellation by $H_s^S \cdot D_s$. Using the description $L_S = D_s \times H_s^S / \sim$ from Lemma 5.3, we can identify each tile $h \cdot D_s$ with the set $[D_s, h]$. It is then clear that the tiles cover all of L_S and that they do not overlap, i.e. the intersection of any two tiles is contained in the boundary of each of them. Assuming extra retractibility, we have the following stronger

Lemma 5.11. *Suppose that F is an extra retractible complex of groups over the poset $P(S)$. Then the intersection of any two distinct tiles in the tessellation $H_s^S \cdot D_s$ is either empty or it is a single simplex contained in the boundary of each of the tiles.*

Proof. Let h_1, h_2 be two distinct elements of the group H_s^S and $[D_s, h_1], [D_s, h_2]$ the corresponding two distinct tiles. If $x \in [D_s, h_1] \cap [D_s, h_2]$ then $x = [p, h_1] = [q, h_2]$

and $(p, h_1) \sim (q, h_2)$. It follows that $p = q$ and $h_1^{-1}h_2 \in \Gamma_{\sigma(p)}$. Thus, identifying the intersection of the tiles with a subset in D_s , we have

$$[D_s, h_1] \cap [D_s, h_2] = \cup\{\sigma \in \partial D_s : h_1^{-1}h_2 \in \Gamma_\sigma\}.$$

Now, using homomorphisms r_σ and the argument as in the proof of Proposition 4.2. (3), we see that $\Gamma_\sigma = \cap\{\Gamma_v : v \text{ is a vertex of } \sigma\}$. This implies that if all vertices of a simplex $\sigma \in \partial D_s$ are in the intersection of the two tiles, so is the whole σ , and thus the intersection is a full subcomplex in ∂D_s . On the other hand, all the vertices contained in the intersection have to belong to a single simplex of ∂D_s , since otherwise they span a full subcomplex in ∂D_s containing some two simplices σ and τ not contained in a single simplex of ∂D_s . Applying Lemma 5.10, we then get $h_1^{-1}h_2 \in \Gamma_\sigma \cap \Gamma_\tau = \{1\}$, and hence $h_1 = h_2$. This contradicts our initial assumption proving that the intersection is either empty or a single simplex of ∂D_s . \square

Proposition 5.12. *Suppose that F is an extra retractible complex of groups on the poset $P(S)$. Then the complex L_S satisfies no-square condition.*

Proof. Suppose there is a square in L_S with vertices a, b, c, d and with edges ab, bc, cd and da . By translating with an element of F_S we may (and will) assume that ab is an edge of the simplex $[\Delta_S, 1]$ in L_S . Let s, t be the elements of S such that $[v_s, 1] = a$ and $[v_t, 1] = b$. Consider the subgroup $H_s^S \subset F_S$ and the tessellation $H_s^S \cdot D_s$ of L_S . The vertex a is then the unique interior vertex of the tile D_s , and we will refer to it as centre of D_s . Consider next the subgroup $H_s^{S-t} \subset H_s^S$ and the homomorphism $r_{s, S-t}^S : H_s^S \rightarrow H_s^{S-t}$. Define the subgroup $H_{s,t}^S := \ker r_{s, S-t}^S \subset H_s^S$ and note that its restricted action on L_S clearly preserves the tessellation $H_s^S \cdot D_s$. Put $D_{s,t} = \cup\{h \cdot D_s : h \in H_s^{S-t}\}$.

Claim 1. *$D_{s,t}$ is a strict fundamental domain for the action of $H_{s,t}^S$ on L_S .*

Proof of Claim 1. Viewing L_S as $H_s^S \times D_s / \sim$ (see Lemma 5.3), define a map $j_{s,t} : H_{s,t}^S \setminus L_S \rightarrow D_{s,t}$ by the formula

$$j_{s,t}(H_{s,t}^S \cdot [p, \gamma]) = [p, r_{s, S-t}^S(\gamma)].$$

This map is easily checked to be an inverse of the quotient map $L_S \rightarrow H_{s,t}^S \setminus L_S$ restricted to $D_{s,t}$, hence the claim.

Claim 2. *$D_{s,t}$ is the union of all tiles of the tessellation $H_s^S \cdot D_s$ that contain the vertex $b = [v_t, 1]$.*

Proof of Claim 2. Using again the description $L_S = H_s^S \times D_s / \sim$, observe that a tile $h \cdot D_s = [D_s, h]$ contains the vertex $[v_t, 1]$ iff $h \in \Gamma_{v_t} = \Gamma_{\sigma_{S-t}} = H_s^{S-t}$ (see Lemma 5.4). This implies the claim.

By Claim 2 and the fact that a is the centre of the tile $D_s \subset D_{s,t}$, the edges ab , bc and da of the square $abcd$ are contained in $D_{s,t}$. By Claim 1, the vertices c and d are connected by an edge in $D_{s,t}$, namely by the image of the edge cd under the quotient map $L_S \rightarrow H_{s,t}^S \setminus L_S = D_{s,t}$. This edge is in fact equal to cd , since two vertices span at most one edge in L_S .

Consider now the following two cases.

Case 1: $c \in D_s$. Since D_s is a simplicial cone centered at a (see Lemma 4.2), there is an edge ac in $D_s \subset L_S$. Thus the square $abcd$ is not empty in this case.

Case 2: $c \notin D_s$. Let $g \cdot D_s \subset D_{s,t}$ be a tile containing the edge cd . By our assumption on c we have $g \cdot D_s \neq D_s$. Next, since a is the centre of D_s , we have $d \in D_s$ and consequently $d \in D_s \cap g \cdot D_s$. By Lemma 5.11, the intersection $D_s \cap g \cdot D_s$ is a simplex, and by Claim 2 this simplex contains the vertex b . It follows that there is an edge bd in L_S , and again the square $abcd$ is not empty. \square

6. Construction and the proof of Theorem 1

To prove Theorem 1 it is sufficient to construct for any natural number n a flag simplicial complex L_n of dimension n which is a closed orientable pseudomanifold and contains no empty square. According to previous sections, such L_n comes as the development L_S of an extra retractible complex of groups F on the poset $P(S)$, if $|S| = n + 1$, all singleton groups are equal to Z_2 and the group F_S is finite (Propositions 3.2, 4.3 and 5.12). We will construct such complexes of groups for any n . This will be done inductively, by extending the complexes of groups constructed in earlier steps to bigger posets.

Let $(P_f(A), \leq)$ be the poset of all finite countably infinite sets A . View $P_f(A)$ as the category with inclusions and retractions in the obvious analogy with the category $P_r(S)$ defined at the beginning of Section 4. As before, the induced groups are denoted by F_S , the groups $F_{\{s\}}$ are called singleton groups, and the homomorphisms induced by morphisms m_{RT} are denoted h_{RT} .

For any morphism F from the category $P_f(A)$ to the category of groups, and for any finite subset $S \subset A$, the restriction of F to $P_r(S)$, denoted $F(S)$, is a retractible complex of groups. Thus we will call F a retractible complex of groups on $P_f(A)$. We say that F is reduced (minimal, extra retractible), if for any $S \in P_f(A)$ the restriction $F(S)$ is reduced (minimal, extra retractible, respectively). To prove Theorem 1, it is sufficient to construct an extra retractible complex of groups F on $P_f(A)$, with all singleton groups equal to Z_2 and all groups F_S finite. We will call such a complex of groups (or functor) *wonderful*.

For a natural number n , denote by $P_n(A)$ the full subcategory of $P_f(A)$ whose objects are all sets $S \subset A$ with cardinality $|S| \leq n$. It means that $P_n(A)$ contains all morphisms from the category $P_f(A)$ between the relevant objects. We adapt the notion of a wonderful complex of groups to functors on the categories $P_n(A)$ in the obvious way. A wonderful functor F on $P_f(A)$ will be constructed by a

sequence of extensions from a wonderful functor on $P_n(A)$ to a wonderful functor on $P_{n+1}(A)$.

The construction. For the first step of the induction, define F on $P_1(A)$ in the only possible way, putting $F_{\{s\}} = Z_2$ for each $s \in A$. Observe that this functor is clearly wonderful. Next, assume that F defined on $P_n(A)$ is wonderful. For each $S \subset A$ with $|S| = n+1$, denote by $F(\Delta_S)$ the simplex of groups (over the simplex Δ_S) with local groups F_T at faces σ_T of Δ_S . Declare \mathcal{F}_S to be the direct limit of this simplex of groups.

Lemma 6.1. *Local groups F_T in the simplex of groups $F(\Delta_S)$ inject into the direct limit \mathcal{F}_S . In other words, simplex of groups $F(\Delta_S)$ is developable.*

Proof. For a fixed local group F_T consider the morphism from $F(\Delta_S)$ to F_T given by the family of homomorphisms $h_{RT} : F_R \rightarrow F_T$ (indexed by proper subsets $R \subset S$). This simple morphism induces a homomorphism $\mu_{S,T} : \mathcal{F}_S \rightarrow F_T$ whose composition $F_T \rightarrow \mathcal{F}_S \rightarrow F_T$ with the map $F_T \rightarrow \mathcal{F}_S$ (associated to the direct limit) is clearly the identity. Thus the map $F_T \rightarrow \mathcal{F}_S$ is injective. \square

Observe that any subset $S \subset A$ with $|S| = n+1$ identifies with the generating set for the group \mathcal{F}_S , where an element $s \in S$ represents the generator in the singleton subgroup $F_{\{s\}} \subset \mathcal{F}_S$ (isomorphic to Z_2). The fact that these generators are all distinct follows from the existence of retractions $\mu_{S,\{s\}} : \mathcal{F}_S \rightarrow F_{\{s\}}$ (described in the proof of Lemma 6.1). This observation is extended by the following

Proposition 6.2. *The functor on the category $P_{n+1}(A)$ with groups F_T for $|T| \leq n$, and with groups \mathcal{F}_S for $|S| = n+1$, is extra retractible.*

Remark. The functor as in Proposition 6.2 satisfies all properties of a wonderful functor except that the groups \mathcal{F}_S are in general not finite. Finite groups F_S giving a wonderful functor on $P_{n+1}(A)$ will be defined later as certain quotients of the groups \mathcal{F}_S .

Proof of Proposition 6.2. We have first to check the existence of all homomorphisms prescribed by functoriality. Those ones which are induced by morphisms from the subcategory $P_n(A)$ are simply the homomorphisms given by the functor F . The remaining homomorphisms are obtained by composing the inclusions $F_T \rightarrow \mathcal{F}_S$ and the retractions $\mu_{S,T} : \mathcal{F}_S \rightarrow F_T$ defined in the proof of Lemma 6.1. This is so, because any morphism $m : S \rightarrow T$ in $P_{n+1}(A)$ is of the form $m = m_{m(S)-\{1\},T} \circ m_{S,m(S)-\{1\}}$.

Since minimality and reducedness of the considered functor are obvious, we turn to proving extra retractibility. Let $S \subset A$ be a set with $|S| = n + 1$. Denote by \mathcal{H}_s^S the kernel of the homomorphism $\mu_{S-s}^S : \mathcal{F}_S \rightarrow \mathcal{F}_{S-s}$. According to Lemma 5.5, \mathcal{H}_s^S is generated by the set $\{s^g : g \in \mathcal{F}_{S-s}\}$, and the generating sets for groups H_s^T with $T \subset S$ have similar form. It is sufficient to prove that for any $T \subset S$ containing s there is a homomorphism $r_{s,T}^S : \mathcal{H}_s^S \rightarrow H_s^T$ such that

- (1) $r_{s,T}^S(h) = h$ for each $h \in \{s^g : g \in \mathcal{F}_{T-s}\}$;
- (2) $r_{s,T}^S(h) = 1$ for each $h \in \{s^g : g \in \mathcal{F}_{S-s} - \mathcal{F}_{T-s}\}$.

This requires some preparatory discussion.

Let $\mathcal{L}_S = \mathcal{F}_S \times \Delta_S / \sim$ be the development of the complex of groups $F(\Delta_S)$ associated to the direct limit \mathcal{F}_S . It follows from general theory [BH II.12] that \mathcal{L}_S is connected and simply connected. Consider the complex of groups $\mathcal{H}_s^S \setminus \mathcal{L}_S$ associated to the action of \mathcal{H}_s^S on \mathcal{L}_S . By lemma 5.1, $\mathcal{H}_s^S \setminus \mathcal{L}_S$ is a simple complex of groups, and since \mathcal{L}_S is connected and simply connected, the group \mathcal{H}_s^S is the direct limit of this complex. To prove the existence of the homomorphism $\mathcal{H}_s^S \rightarrow H_s^T$ as required, it is sufficient to construct the appropriate simple morphism $\mathcal{H}_s^S \setminus \mathcal{L}_S \rightarrow H_s^T$, i.e. the consistent family of homomorphisms from the local groups of $\mathcal{H}_s^S \setminus \mathcal{L}_S$ to H_s^T .

According to Lemma 5.4, local groups of $\mathcal{H}_s^S \setminus \mathcal{L}_S$ have the form $(H_s^R)^g$ for the appropriate R and g . The required homomorphisms to H_s^T are then the compositions

$$(H_s^R)^g \rightarrow (H_s^{R \cap T})^g \rightarrow (H_s^T)^g \rightarrow H_s^T,$$

where the first map is the conjugated retraction $r_{s,R \cap T}^R$ (provided by extra retractibility in the inductive hypothesis), the second map is the conjugated inclusion, and the third map is the identity if $g \in \mathcal{F}_{T-s}$ while it is the trivial map to $1 \in H_s^T$ otherwise. Conditions (1) and (2) for the induced homomorphism can be checked directly, by tracing the images of the generators under all involved maps. We omit the details. \square

We will now improve the functor described in Proposition 6.2 by replacing groups \mathcal{F}_S with their appropriate finite quotients. To do this, we need certain normal subgroups of finite index in those groups. For any distinct elements $s, t \in S$, denote by $\mathcal{H}_{s,t}^S$ the kernel of the retraction $r_{s,S-t}^S : \mathcal{H}_s^S \rightarrow H_s^{S-t}$. Observe that, since the group H_s^{S-t} is finite, the group $\mathcal{H}_{s,t}^S$ has finite index in \mathcal{H}_s^S , and hence also in \mathcal{F}_S . Put $K_S := \bigcap_{s \neq t} \mathcal{H}_{s,t}^S$ and normalize it to the normal subgroup $N_S := \bigcap_{g \in \mathcal{F}_S} (K_S)^g$ which still has finite index in \mathcal{F}_S .

Lemma 6.3. *The normal subgroup N_S is contained in the kernel of any retraction $\mathcal{H}_s^S \rightarrow H_s^T$ and any homomorphism $\mathcal{F}_S \rightarrow \mathcal{F}_R$ induced by the functor from Proposition 6.2.*

Proof. By definition, N_S is contained in the kernel $\mathcal{H}_{s,t}^S$ of any map $\mathcal{H}_s^S \rightarrow H_s^{S-t}$. Now, since $T \subset S - t$ for some t , and since the map $\mathcal{H}_s^S \rightarrow H_s^T$ is equal to the composition $\mathcal{H}_s^S \rightarrow H_s^{S-t} \rightarrow H_s^T$ of the retractions, the kernel of $\mathcal{H}_s^S \rightarrow H_s^T$ contains the kernel of $\mathcal{H}_s^S \rightarrow H_s^{S-t}$, and hence contains N_S .

The part of the lemma concerning morphisms $\mathcal{F}_S \rightarrow F_R$ has very similar proof which we omit. \square

The next proposition completes the inductive construction of a wonderful functor F on $P_f(A)$, thus proving Theorem 2.

Proposition 6.4. *The functor on $P_{n+1}(A)$ obtained from a wonderful functor F on $P_n(A)$ by putting $F_S := \mathcal{F}_S/N_S$ for any $S \subset A$ with $|S| = n + 1$ is wonderful.*

Proof. Since by Lemma 6.3, the kernels of all homomorphisms from \mathcal{F}_S to F_R (induced by the functor as in Proposition 6.2) contain N_S , they induce homomorphisms from \mathcal{F}_S to F_R which turn the improved functor into a reduced and minimal retractible complex of groups on $P_{n+1}(A)$. Similarly, the kernels of retractions $\mathcal{H}_s^S \rightarrow H_s^T$ all contain N_S , thus inducing homomorphisms $\mathcal{H}_s^S/N_S \rightarrow H_s^T$. By observing that these homomorphisms are the desired retractions $H_s^S \rightarrow H_s^T$ we get extra retractibility. Since the rest of the desired properties are obvious, the proposition follows. \square

7. Hyperbolization

Thus far we produced essentially a single $CAT(-1)$ pseudomanifold, and the reader may think that such spaces are isolated/exotic. We now show how to produce many $CAT(-1)$ cubical complexes, using hyperbolization procedure.

We start with cubical complexes which are locally $CAT(0)$, and convert them to locally $CAT(-1)$ complexes. There are several hyperbolization procedures described in [DJ1], which, starting with any cubical complex, produce a locally $CAT(0)$ one. Putting these together we end up with a hyperbolization procedure which converts any cubical complex to a locally $CAT(-1)$ one.

Let us recall [BH, II.5] that a piecewise constant curvature κ space has curvature bounded above by κ , or satisfies the $CAT(\kappa)$ inequality locally if and only if all its links, considered as piecewise constant curvature 1 metric spaces, are large: the length of the shortest geodesic loop in any link is $\geq 2\pi$. In particular a piecewise flat metric on a cubical complex is $CAT(0)$ iff the links are flag.

For any minimal, reduced, retractible complex of groups over $P(S)$ we have, as discussed in 4.6, a surjective map $A_S : \mathcal{F}_S \rightarrow F_S^{\text{direct}}$. Assume that the singleton groups are all isomorphic to Z_2 . Then this last group is isomorphic with $Z_2^{|S|}$, and A_S coincides with the abelianization map. Since the complex L_S^{direct} associated with $Z_2^{|S|}$ is the hyperoctahedral triangulation of the sphere, we have the associated

map $a_S : L_S \rightarrow S^{|S|-1}$ of complexes. One can interpret this map as taking the quotient of L_S by the kernel of A_S .

Recall that the cone over the barycentric subdivision of any simplicial complex X has a natural structure of a cubical complex. We will call it the cubical cone over X and denote it by ccX . An alternative description is as follows. For any simplex $\sigma \subset X$ we take a cube c_σ of dimension $\dim \sigma + 1$ with a distinguished vertex 0, identify the link of 0 in c_σ with σ . Now identify cubical faces containing 0 following the identifications on X . The cubes of this cubical structure are called small cubes in ccX .

Since the coning is obviously functorial, we form the cubical cone over the map a_S , to obtain a map of cubical complexes

$$cc(a_S) : ccL_S \rightarrow ccS^{|S|-1} = C^S,$$

with C^S , the cube $[-1, 1]^S$, decomposed into $2^{|S|}$ images of the small cube $[0, 1]^S$ under sign changing symmetries of the large cube.

One can view ccL_S as a ramified covering of C^S .

We use the map $cc(a_S)$ as an input in a fiber product hyperbolization for cubical complexes as in [DJ1], and accordingly change the notation slightly. Now $C_h^S = ccL_S$ denotes the hyperbolized cube. For $C^T, T \subset S$, a face of dimension $|T|$ in C^S is obtained by fixing coordinates not in T to be $+1$ or -1 . We denote by C_h^T a connected component of the preimage of C^T by $cc(a_S)$. It is, in fact, a cubical cone ccL_T .

Let X be a locally $CAT(0)$ cubical complex.

Suppose there exists a folding map $f_X : X \rightarrow C^S$ to the $|S|$ -cube C^S ; the C^S is considered here as a single cube (i.e. it is not subdivided). This means that there is a map of cubical complexes which is injective on each cube in X to the $|S|$ -cube (thus the complex is of dimension at most $|S|$).

Let X_h be the fiber product of f_X and $cc(a_S)$, that is $X \times C_h^S \supset X_h = \{(x, h) | f_X(x) = cc(a_S)(h)\}$.

The intuitive content of this construction is that we replace a cube in X with a copy of C_h preserving the gluing pattern of X . The fiber product construction allows us to say it concisely.

The projection $X_h \rightarrow C_h$ restricted to the preimage of a top dimensional cube in X under the projection $X_h \rightarrow X$ is an isomorphism, thus it endows X_h with the geodesic metric, which on each C_h is the Moussong metric.

For the reader's convenience we recall briefly Moussong's construction. First we need the notion of hyperbolic cubes. In the disc model of H^n take coordinate hyperplanes $x_i = 0$. Take a point (t, \dots, t) , and all of its $2^n - 1$ orthogonal projections to the intersections of coordinate hyperplanes. The convex hull of these 2^n points is a cube. The solid angle at the vertex $(0, \dots, 0)$ is the spherical simplex with all dihedral angles $\frac{\pi}{2}$ (all-right simplex); the opposite solid angle is a fairly small regular spherical simplex, depending on t . The small angle is placed at the tip of the cubical cone, while the all-right angle is placed at the vertex corresponding to the barycenter of the $(n - 1)$ simplex.

We need to describe links of points in X_h . For this we need some notation. Let $C(x)$ denote the cube of X containing x in its interior. Denote by $Lk(C(x) \subset X)$ the normal link of any point c in the interior of $C(x)$ (which depends on $C(x)$ only) and by $Lk(a \in A)$ the link of a in A . Then we have

Proposition 7.1.

(1) *Let $(x, h) \in X_h$. Then we have an isometry of links*

$$Lk((x, h) \in X_h) = Lk(C(x) \subset X) * Lk(h \in C(x)_h).$$

*Here $X * Y$ denotes the spherical join of piecewise spherical complexes X, Y (see [BH p. 63]).*

(2) *If X is locally $CAT(0)$ and F_S is an extra retractible complex of groups, then the complex X_h with its Moussong metric is locally $CAT(-1)$.*

Proof. (1) is straightforward from the definition of the hyperbolization and the description of the metric.

(2) follows from (1) and the following three facts: the join of large links is large, the assumption that X has large links and the fact that the Moussong metric has large links.

Remarks.

- (1) The most notable defect of this hyperbolization is that it does not preserve the property of X being a manifold for $\dim X \geq 3$. In dimension 3 this can be fixed by an explicit local resolution of singularities. In higher dimension such a resolution, preserving $CAT(-1)$ property is unknown to us.
- (2) It is not really necessary to use cubical complexes which admit a folding map. If F_S has a better functorial property, that is if a permutation of S induces an automorphism $F_S \rightarrow F_S$, respecting the subgroup structure of F_S , then the complex L_S is symmetric, and we can apply the hyperbolization to complexes without a folding map. The complex constructed in Section 6 has such a symmetry.

8. Non right-angled groups

Let (m_{st}) be a right angled Coxeter matrix with the set of indices S . Let (m_{st}^f) be any Coxeter matrix obtained from (m_{st}) by the following modification:

$$(m_{st}^f) = \begin{cases} m_{st} & \text{if } m_{st} \neq \infty \\ f_{st} & \text{otherwise} \end{cases}$$

with $f_{st} \in \{5, 6, \dots, \infty\}$.

Proposition 8.1. *Suppose the Coxeter group (W, S) associated to (m_{st}) is Gromov hyperbolic. Then*

- (1) *The group (W^f, S) associated to any of the matrices m^f is Gromov hyperbolic.*
- (2) *If the nerve $N(W, S)$ is connected, then $N(W^f, S)$ has the homotopy type of the wedge of $N(W, S)$ and a certain number of circles.*
- (3) *If $vcd(W) \geq 2$, then $vcd(W) = vcd(W^f)$.*

Proof. To prove (1) we use Moussong's criterion for hyperbolicity: a Coxeter group is Gromov hyperbolic if none of its parabolic subgroups is affine of rank ≥ 2 or splits into a product of two infinite parabolic subgroups.

If $T \subset S$, then hyperbolicity of (W, S) implies that the Dynkin diagram of W restricted to T has at most one connected component with more than one point.

If the nontrivial connected component contains more than two points, the subgroup of (W^f, S) spanned by T is neither finite nor affine (as one sees by examining well known lists of finite and affine irreducible Coxeter groups) and does not split as a product of infinite groups.

On the other hand, if the nontrivial component has two points, or if the Dynkin diagram has no edges at all, the group spanned by T in W^f is either finite or contains an infinite dihedral subgroup of finite index. This proves (1).

To prove (2), consider the coarse simplicial structures on the nerves. Clearly $N(W, S)$ is a subcomplex of $N(W^f, S)$. Let τ be a simplex in $N(W^f, S)$ not in $N(W, S)$ and let $T \subset S$ denote the set of vertices of τ . From the proof of Assertion (1) we know that the Dynkin diagram of W^f restricted to T has one connected component with two vertices say t, s with $m_{st} = \infty$ and f_{st} finite (while the other components of the Dynkin diagram are points). This implies that τ (in fact the edge ts) is contained in the unique maximal simplex of $N(W^f, S)$, whose vertices are t, s and all elements of S commuting with both t, s . Thus either the edge ts is itself a maximal simplex in $N(W^f, S)$ or its maximal simplex deformation retracts through its interior onto the union of its faces not containing ts , which is a subcomplex of $N(W, S)$. These deformation retractions can be performed independently and together they give a collapse of $N(W^f, S)$ onto the union of $N(W, S)$ and certain number of 1-simplices. This proves (2).

For the proof of (3) we need the following

Lemma 8.2. *Let K be a finite flag simplicial complex, having the property that for any simplex σ in K (including the empty simplex), $H^i(K_\sigma; \mathbb{Z}) = 0$ for $i > k$, where K_σ is the full subcomplex of K spanned by the vertices not in σ . Then for any full subcomplex L of K we have $H^i(L; \mathbb{Z}) = 0$ for $i > k$.*

Proof. Let V be the set of vertices of K , and let L be spanned by $V - A \subset V$. We proceed by induction on cardinality of A . Note that for $|A| = 0, 1$ the conclusion follows from the assumptions.

Now let $|A| \geq 2$. If A spans a simplex in K , we are done. Otherwise there are

two vertices in A , say v, w , not connected by an edge in K . Consider subcomplexes $L_v, L_w, L_{v,w}$ of K spanned by $V - (A - \{v\})$, $V - (A - \{w\})$, $V - (A - \{v, w\})$ respectively. Then $L_v \cap L_w = L$, and $L_v \cup L_w = L_{v,w}$. Now we have a piece of the Mayer–Vietoris sequence

$$H^i(L_v; \mathbb{Z}) \oplus H^i(L_w; \mathbb{Z}) \rightarrow H^i(L; \mathbb{Z}) \rightarrow H^{i+1}(L_{v,w}; \mathbb{Z}).$$

For $i > k$ groups on both left and right are zero by inductive assumption, hence the Lemma.

Now the proof of (3) in Proposition 8.1 proceeds as follows. By Assertion (2), the nerve $N(W_{S-T}^f, S-T)$ has the homotopy type of the wedge of $N(W_{S-T}, S-T)$ and certain number of circles. By the discussion in Section 1.4 and the Lemma, $H^i(N(W_{S-T}, S-T); \mathbb{Z}) = 0$ for $i > vcd(W)$ and any T . Thus if $vcd(W) \geq 2$, $H^i(N(W_{S-T}^f, S-T); \mathbb{Z}) = 0$ for $i > vcd(W)$ and any T , and (3) follows.

9. Complexity issue

The complexes L_S we construct have strikingly large complexity measured in terms of the number of either vertices or top dimensional simplices. Let S_n, V_n denote respectively the number of simplices of dimension n and of vertices in L_S , for $|S| = n + 1$. Clearly $S_n = |F_S|$.

Since the link of any vertex v in L_S is isomorphic to L_{S-s} , v is contained in S_{n-1} simplices. Thus by double counting

$$V_n = \frac{(n+1)S_n}{S_{n-1}}.$$

Since under the abelianization map H_s^S surjects onto $Z_2^{S_{n-1}}$, we have

$$S_n \geq S_{n-1} 2^{S_{n-1}} \quad \text{and} \quad V_n \geq (n+1) 2^{S_{n-1}}.$$

Actual values of S_n are probably much larger than predicted by these inequalities, making the growth of S_n ridiculously fast. It may be desirable to improve this aspect of the construction.

On the other hand we would like to point out that flag no-square simplicial pseudomanifolds cannot be too simple, that is the number of their top simplices grows with the dimension significantly faster than exponentially.

Proposition 9.1. *Let v_n, s_n denote the minimal number of vertices and n -simplices respectively in a simplicial, flag, no-square pseudomanifold of dimension n . Then*

$$s_{n+1} > \frac{s_n^2}{n+2} \quad \text{and} \quad v_{n+1} > s_n.$$

Proof. Let X be a simplicial, flag, no-square pseudomanifold of dimension $n + 1$. Then the link of any simplex σ in X is also a simplicial, flag, no-square pseudomanifold of appropriate dimension. To prove this notice that since X is flag, the join $\sigma * \text{link}(\sigma)$ embeds in X as the set of all top dimensional simplices containing σ . Thus a square in $\text{link}(\sigma)$ is a square in X , and hence it has one of its diagonals in X . Since X is flag, the diagonal is in the link.

By double counting, the number of $(n + 1)$ -simplices in X is equal to $\frac{1}{n+2} \sum_v |\text{lk}(v)|$, where v runs over all vertices of X , and $|\text{lk}(v)|$ denotes the number of n -simplices in the link of v .

Since links in X are flag, no-square pseudomanifolds, we obtain the estimate

$$s_{n+1} \geq \frac{v_{n+1} s_n}{n+2}.$$

Now fix a vertex $x \in X$. Each n -simplex σ in the link of x determines the unique $(n+1)$ -simplex in X which contains σ and is not contained in the star of x . Hence it determines a vertex of X : the vertex of σ not in the link of x . The vertices so obtained are all distinct by the no-square condition. Counting vertices in the star of x , and vertices obtained above we get

$$v_{n+1} \geq 1 + v_n + s_n.$$

Putting the two displayed inequalities together finishes the proof. \square

References

- [B1] M. Bestvina, Virtual cohomological dimension of Coxeter groups, in: G. Niblo and M. Roller eds., *Geometric Group Theory I*, 19–23, LMS Lect. notes 181, Cambridge University Press, 1993.
- [B2] M. Bestvina, <http://www.math.utah.edu/~bestvina/>.
- [BH] M. Bridson and A. Haefliger, *Metric spaces of Nonpositive Curvature*, Grundlehren der Math. Wiss. 319, Springer, 1999.
- [Bu] W. O. Bugaenko, Arithmetic crystallographic groups generated by reflections and reflective hyperbolic lattices, *Lie groups, their discrete subgroups, and invariant theory*, 33–55, Adv. Sov. Math. 8, AMS, Providence, 1992.
- [D1] M. W. Davis, The cohomology of a Coxeter group with group ring coefficients, *Duke Math. J.* **91** (1998) no. 2, 297–314; *Duke Math. J.* **95** (1998) no. 3, 635.
- [D2] M. W. Davis, Nonpositive curvature and reflection groups, in: R. Daverman and R. Shor eds., *Handbook of Geometric Topology*, Elsevier, 2001.
- [DJ1] M. W. Davis and T. Januszkiewicz, Hyperbolization of polyhedra, *J. Diff. Geom.* **34** (1991), 347–388.
- [Dr] A. N. Dranishnikov, Boundaries of hyperbolic Coxeter groups, *Topology and Applications* **110** (2001), 29–38.
- [G] M. L. Gromov, Asymptotic invariants of infinite groups, in: G. Niblo and M. Roller eds., *Geometric Group Theory II*, LMS Lect. notes 182, CUP, 1993.
- [GT] M. L. Gromov and W. P. Thurston, Pinching constants for hyperbolic manifolds, *Inv. Math.* **89** (1987), 1–12.

[MS] L. Mosher and M. Sageev, Non manifold hyperbolic groups of high cohomological dimension. <http://newark.rutgers.edu/~mosher>

[M] G. Moussong, Hyperbolic Coxeter group, Ohio State University PhD thesis, 1987.

[S] R. Stanley, *Combinatorics and commutative algebra*, Birkhäuser, 1983.

[V] E. B. Vinberg, Hyperbolic reflection groups, *Usp. Mat. Nauk* **40** (1) (1985), 29–66.

T. Januszkiewicz and J. Świątkowski
Instytut Matematyczny UWr
pl. Grunwaldzki 2/4
PL-50-384 Wrocław
Poland

(Received: January 2, 2002)



To access this journal online:
<http://www.birkhauser.ch>