

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 78 (2003)

Artikel: Structure of foliations of codimension greater than one
Autor: Marzougui, Habib / Salhi, Ezzedine
DOI: <https://doi.org/10.5169/seals-58777>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Structure of foliations of codimension greater than one

Habib Marzougui and Ezzeddine Salhi

Abstract. We study the structure of a foliation of high codimension which admits a transverse foliation. We introduce four families of open saturated sets. These open sets have a simple characterization and allow us to establish a structure theorem as in codimension 1.

Mathematics Subject Classification (2000). 57R30.

Keywords. Foliation, leaves, regular open set, class, local minimal set.

1. Introduction

This paper concerns the study of the global structure of foliations of codimension greater than one in a closed manifold of finite dimension. In codimension 1, if the foliation is of class C^2 , its topological structure is fairly well understood (cf. [2], chap.4). When the foliation is of class C^0 , many phenomena occur which cannot occur in class C^2 ; such foliations were studied for instance in [1], [3], [4]. In codimension $q \geq 2$, there are many particular foliations which have been studied; for example, Epstein studied foliations with all leaves compact; Molino studied the Riemannian foliations case; Blumenthal studied the case of transversely homogeneous foliations (cf. [2], chap. 3)... It is natural to try to understand foliations in a closed manifold of codimension $q \geq 2$: The purpose here is to develop in this general situation a setting of a structure theorem. However, there are many differences between codimension 1 and codimension ≥ 2 . But very little is known about them, in particular about their global structure. In order to generalize the case of codimension 1, we relate them to the notion of regular open set introduced by Salhi in [3], [4], and some related concepts.

We refer to section 5 for the statement of the main result of this paper.

The plan of this paper is as follows. In section 2 we give some preliminaries. In sections 3 and 4, we introduce a family of open sets and technical lemmas which will make up the main theorem. In section 5, we prove the main theorem.

Research supported by the unit of research Algebra–Topology–Geometry: 99UR15-15.

2. Preliminaries

In this section we recall some basic definitions and notations, and introduce notions which will be useful in the sequel. Let M be a closed C^∞ manifold of dimension n , and \mathcal{F} a codimension q , $q \geq 1$ and class C^r , $r \geq 1$, foliation on M . A subset of M which is a union of leaves is called an \mathcal{F} -saturated set. An \mathcal{F} -saturated set E is called a *local minimal set* of \mathcal{F} if there exists an open \mathcal{F} -saturated set U such that E is a minimal set of $\mathcal{F} \mid U$; that is for every leaf $L \subset E$, we have $\overline{L} \cap U = E$. Two distinct local minimal sets of \mathcal{F} are disjoint.

We call the class of a leaf L of \mathcal{F} the union $cl(L)$ of all leaves G of \mathcal{F} such that $\overline{G} = \overline{L}$. If L is proper, $cl(L) = L$. If L is contained in a local minimal set E then $cl(L) = E$. An open \mathcal{F} -saturated set U is called an *open local minimal set* of \mathcal{F} if all leaves of $\mathcal{F} \mid U$ are dense in U . It is equivalent to the following: for every leaf L of $\mathcal{F} \mid U$, $cl(L) = U$. In this case, we have also $U \subset \overline{L}$.

For more information about this terminology, see [3], [4], or see also ([2], Chap. 4, section 4).

Here we give some trivial lemmas which are useful in the sequel.

Lemma 1. *If U is an open connected set of M and V is a nonempty open set of M strictly contained in U then $Fr(V) \cap U \neq \emptyset$.*

Indeed, V is not closed in U since U is connected and distinct from V . Therefore $Fr(V) \cap U = (\overline{V} - V) \cap U \neq \emptyset$.

Lemma 2. *Let L be a leaf of \mathcal{F} and $X \subset \overline{L}$ be a closed nonempty \mathcal{F} -saturated set of M with $X \neq \overline{L}$. If U is the connected component of $M - X$ containing L then $X = Fr(U)$.*

Proof. Since U is closed in $M - X$, $Fr(U) = \overline{U} - U \subset X$. On the other hand, we have $X \subset \overline{L} \subset \overline{U}$ and $X \subset M - U$. Hence $X \subset \overline{U} - U$ and then $X = Fr(U)$. \square

The Lemma 2 is also true if we replace the leaf L by any closed connected nonempty \mathcal{F} -saturated subset of M .

Corollary 1. *If G and L are leaves of \mathcal{F} such that $\overline{G} \subset \overline{L}$ and $\overline{G} \neq \overline{L}$ then $\overline{G} = Fr(U)$ where U is the connected component of $M - \overline{G}$ containing L .*

In codimension $q \geq 2$, there does not necessarily exist a foliation transverse to \mathcal{F} of dimension q . In order to give some analogies as in codimension 1, we suppose in all the sections below, that *there exists a foliation Γ transverse to \mathcal{F} of dimension q* .

3. Regular and foliated fiber open sets

In this section, we adapt to our purpose some viewpoints due to E. Salhi in [3], [4] (see also [2], chap. 4, section 4).

Let U denote a nonempty open connected \mathcal{F} -saturated set distinct from M . If T is a transverse leaf, we denote by $\text{Sat}(T)$ its saturated set for the leaves of \mathcal{F} .

Definition 1. U satisfies:

- Property (a): if for every $x \in U$, the transverse leaf T_x of Γ passing through x meets the complementary $M - U$ of U .
- Property (c): if U contains a transverse leaf and for every transverse leaf $T \subset U$, we have $\text{Sat}(T) = U$.

Definition 2. U is called *regular* if for every transverse leaf T of Γ meeting U , every leaf of $\mathcal{F} \mid U$ meets T (i.e. $U \subset \text{Sat}(T)$).

Definition 3. U is called a *foliated fiber* if it is regular and satisfies Property (a).

Note that if $U = M$, we say that M is regular (or a foliated fiber) if for every transverse leaf T we have: $M = \text{Sat}(T)$.

Remark 1. If \mathcal{F} is of codimension $q = 1$:

- U is a foliated fiber open set if and only if it satisfies the Property (a).
- U is regular if and only if U satisfies one of the Properties (a) or (c). (cf. [3], proposition 3).

Remark 2. If \mathcal{F} is of codimension $q \geq 2$, a regular open set U satisfies always Properties (a) or (c). The converse is not true in general:

– There exists an open set U which satisfies Property (a) but it is not regular as can be shown by the following example:

Example 1. Let $\mathcal{F} = \mathcal{F}_1 \times \mathcal{F}_2$ be the product foliation of $T^3 = T^2 \times S^1$ where \mathcal{F}_1 is the discrete foliation of the circle S^1 (i.e. each leaf is a single-point), and \mathcal{F}_2 is the foliation of the torus T^2 having a compact leaf L_0 and such that $(T^2 - L_0)$ is a Reeb component. \mathcal{F} is of codimension $q = 2$. Let Γ_1 (resp. Γ_2) be the foliation transverse to \mathcal{F}_1 (resp. \mathcal{F}_2). Since S^1 is the unique transverse leaf of Γ_1 , then any open set $U_1 \subset S^1$ distinct from S^1 is a foliated fiber. Also, since Γ_2 has a transverse leaf T_2 such that $\text{Sat}(T_2) = T^2 - L_0$, then T^2 is not regular. It follows that the open set $U = T^2 \times U_1$ in T^3 satisfies Property (a) but is not regular.

– There exists an open set U which satisfies Property (c) but it is not regular as can be pointed out to us by the referee.

At first, we prepare some properties which are useful in the sequel:

Proposition 1. *Let U_1 and U_2 be open, nonempty, connected \mathcal{F} -saturated sets of M such that $U_1 \subset U_2$. Then we have the following:*

- (i) *If U_2 is regular and $U_1 \neq U_2$ then U_1 is foliated fiber.*
- (ii) *If U_2 is an open local minimal set of \mathcal{F} then $U_1 = U_2$.*

Proof. (i) First, U_1 is regular since it is contained in a regular set U_2 . Let T_1 be a transverse leaf meeting U_1 . We have $U_2 \subset \text{Sat}(T_1)$. Then by Lemma 1, $\text{Fr}(U_1) \cap \text{Sat}(T_1) \neq \emptyset$. Therefore T_1 meets $M - U_1$, and U satisfies Property (a).

(ii) This follows from the fact that: For every leaf L of $\mathcal{F} \mid U_2$, $\text{cl}(L) = U_2$ and then $\text{cl}(L) = U_1$. \square

We introduce four families of \mathcal{F} -saturated open sets:

- \mathcal{A} the union of all open sets which satisfy Property (a)
- \mathcal{D} the union of all open local minimal sets of \mathcal{F}
- \mathcal{R} the union of all regular open sets.
- \mathcal{C} the union of all open sets which satisfy Property (c).

Note that:

- $\mathcal{A} \cap \mathcal{R}$ is the union of all foliated fiber open sets
- a regular non foliated fiber open set satisfies Property (c).

Corollary 2. *Let U be an \mathcal{F} -saturated open connected set of M . Then:*

- (i) *If U is an open local minimal set of \mathcal{F} then U is regular. In particular, we have $\mathcal{D} \subset \mathcal{R} \subset \mathcal{A} \cup \mathcal{C}$.*
- (ii) *U is an open local minimal set of \mathcal{F} if and only if $U \subset \mathcal{D}$.*

Proof. (i) Let U be an open local minimal set of \mathcal{F} . If T is a transverse leaf of Γ which meets U and T_U be a connected component of $T \cap U$, then $\text{Sat}(T_U) \subset U$ is an open, nonempty, connected \mathcal{F} -saturated set of M . It follows by (Proposition 1, (ii)) that $\text{Sat}(T_U) = U$. Thus $U \subset \text{Sat}(T)$ and U is regular.

(ii) If U is an open local minimal set of \mathcal{F} we have $U \subset \mathcal{D}$ by definition of \mathcal{D} . Suppose now that $U \subset \mathcal{D}$. Since two distinct open local minimal sets of \mathcal{F} are disjoint and U is connected, there exists an open local minimal set V of \mathcal{F} such that $U \subset V$. Therefore, by (Proposition 1, (ii)) we have $U = V$. \square

Proposition 2. *If U_1 is an open set which satisfies Property (c) and U_2 is regular such that $U_1 \cap U_2 \neq \emptyset$, then $U_2 \subset U_1$.*

Proof. By hypothesis of U_1 , there exists a transverse leaf T_1 such that $\text{Sat}(T_1) = U_1$. Since $U_1 \cap U_2 \neq \emptyset$, T_1 meets every leaf in $U_1 \cap U_2$. Therefore $U_2 \subset \text{Sat}(T_1) = U_1$, as desired. \square

Corollary 3. *If U_1 is an open set which satisfies Property (c) and U_2 is regular non foliated fiber such that $U_1 \cap U_2 \neq \emptyset$ then $U_1 = U_2$. In particular, every regular non foliated fiber open set is a connected component of \mathcal{R} .*

Corollary 4. *If U satisfies Property (c) then $\text{Fr}(U) \subset M - \mathcal{R}$.*

Indeed, otherwise there exists a regular set U_2 such that $\text{Fr}(U) \cap U_2 \neq \emptyset$. Then $U \cap U_2 \neq \emptyset$ and, we have $U_2 \subset U$ (Proposition 2), a contradiction.

Proposition 3. *Every connected component of \mathcal{R} is regular*

Proof. Suppose on the contrary, there exists a connected component U of \mathcal{R} which is not regular, that is, there exists a transverse leaf T meeting U such that $U \not\subset \text{Sat}(T)$. So, $U \cap \text{Sat}(T)$ is nonempty and strictly contained in U . By Lemma 1, $\text{Fr}(\text{Sat}(T)) \cap U \neq \emptyset$. Let L be a leaf in $\text{Fr}(\text{Sat}(T)) \cap U$. There exists a regular open set V such that $L \subset V$. It follows that $\text{Fr}(\text{Sat}(T)) \cap V \neq \emptyset$. Therefore, $V \subset \text{Sat}(T)$ and then $L \subset \text{Sat}(T)$, which is impossible. The proof is complete. \square

Corollary 5. *Let U be an \mathcal{F} -saturated open connected set of M . Then U is regular if and only if $U \subset \mathcal{R}$.*

Indeed, if $U \subset \mathcal{R}$ then there exists a regular open set V such that $U \subset V$ (Proposition 3). Therefore, U is regular (Proposition 1, i)).

Remark 3. If U satisfies Property (a) then $U \subset \mathcal{A}$. The converse is not true in general.

4. Technical lemmas

Hereafter, we prepare some lemmas which play an important role in the proof of the main theorem.

Lemma 3. *If P is a union of transverse leaves of Γ and $U = \text{Sat}(P)$, then there exists a minimal set E of \mathcal{F} | U and U is compact by saturation (i.e. quasi-compact for the topology defined by the saturated open subsets of M).*

Proof. This lemma follows from Zorn's lemma since we have $\overline{P} \subset U$. \square

Lemma 4. *Let P_n ($n \geq 1$) be a union of transverse leaves of Γ . If $\text{Sat}(P_{n+1}) \subset \text{Sat}(P_n)$ then $(\text{Sat}(P_n))_{n \geq 1}$ is stationary.*

Proof. Suppose that $(\text{Sat}(P_n))_{n \geq 1}$ is infinite. So, we may suppose, for every $n \geq 1$ that $\text{Sat}(P_{n+1})$ is strictly contained in $\text{Sat}(P_n)$. Let $L_n \subset (\text{Sat}(P_n) - \text{Sat}(P_{n+1}))$, be a leaf of \mathcal{F} . Then L_n meets a transverse leaf $T_n \subset P_n$; let $x_n \in T_n \cap L_n$. One can assume, passing to a subsequence if necessary, that $(x_n)_{n \geq 1}$ converges to a limit x . Let W_x be a bidistinguished open set which contains x . Then, there exists an integer N such that for $n \geq N$, $x_n \in W_x$. Therefore, the leaf L_n of \mathcal{F}

passing through x_n meets a transverse leaf $T_{n+1} \subset P_{n+1}$; which is impossible. \square

Lemma 5. *Every \mathcal{F} -saturated open set U has only a finite number of connected components which do not satisfy Property (a).*

Proof. Suppose on the contrary, there exist infinitely many transverse leaves $(T_n)_{n \geq 1}$ in U such that $(\text{Sat}(T_n))_{n \geq 1}$ are pairwise disjoint. Let $x_n \in T_n$ for every n . One can assume that $(x_n)_{n \geq 1}$ converges to a limit x . By taking a bidistinguished open set W_x which contains x , there exists an integer N such that for $n \geq N$, $x_n \in W_x$. Therefore, the leaf L_x of \mathcal{F} passing through x meets all T_n for $n \geq N$; a contradiction. \square

Lemma 6. *If $(U_n)_{n \geq 1}$ is a sequence of open sets which do not satisfy Property (a) and with $U_{n+1} \subset U_n$ then there exists an open set $V \subset \bigcap_{n \geq 1} U_n$.*

Proof. By assumption, for every n , there exists a transverse leaf $T_n \subset U_n$. Denote by P_n the union of all transverse leaves contained in U_n , and we let $V_n = \text{Sat}(P_n)$. It follows that for every $n \geq 1$, $V_{n+1} \subset V_n$ and $V_n \subset U_n$. By Lemma 4, $(V_n)_{n \geq 1}$ is stationary; so there exists an integer N such that for $n \geq N$, $V_n = V_N \subset U_n$. Hence, $V_N \subset \bigcap_{n \geq 1} U_n$, as desired. \square

Lemma 7. *Let U be an open set which does not satisfy Property (a). Then U contains an open set which satisfies Property (c).*

Proof. Suppose the contrary. Then in particular U does not satisfy Property (c). There exists a transverse leaf T_1 such that $\text{Sat}(T_1)$ is strictly contained in U . Since $\text{Sat}(T_1)$ does not satisfy Property (c), there exists a transverse leaf T_2 such that $\text{Sat}(T_2)$ is strictly contained in $\text{Sat}(T_1)$. In this way, we construct a sequence $(T_n)_{n \geq 1}$ of transverse leaves such that $\text{Sat}(T_{n+1})$ is strictly contained in $\text{Sat}(T_n)$, which contradicts Lemma 4. \square

Proposition 4. *If U is an open set which satisfies Property (c) then U has one of the following properties:*

- (i) *U is an open local minimal set of \mathcal{F} .*
- (ii) *There exists a minimal set E of $\mathcal{F} \mid U$ with empty interior and for every minimal set E of $\mathcal{F} \mid U$, each connected component of $U - E$ satisfies Property (a).*

Proof. By the hypothesis of U , there exists a transverse leaf T of Γ such that $\text{Sat}(T) = U$. Then, by Lemma 3 there exists a minimal set E of $\mathcal{F} \mid U$. Now, assume that (i) does not hold, that is U is distinct from E , and denote by V a connected component of $U - E$. We claim that V satisfies Property (a). If a transverse leaf T_1 meets V , then one of the following holds:

- (i) T_1 meets $M - U$; in this case, T_1 meets $M - V$ and V satisfies Property (a).

(ii) T_1 is contained in U ; in this case $\text{Sat}(T_1) = U$. Therefore, T_1 meets $M - V$, and then V satisfies Property (a). \square

Note that a regular non foliated fiber open set satisfies Proposition 4.

Corollary 6. *Let U be an open set which satisfies Property (c). If U contains two distinct minimal sets E_1 and E_2 then $U \subset \mathcal{A}$.*

Indeed, by Proposition 4, we have $U - E_1 \subset \mathcal{A}$ and $U - E_2 \subset \mathcal{A}$. So, $U \subset \mathcal{A}$.

Corollary 7. *If all leaves of \mathcal{F} are proper then every regular non foliated fiber open set U contains a leaf closed in U .*

Proof. This follows from Proposition 4 and from the fact that a local minimal set in this case is a proper leaf. \square

Remark 4. If \mathcal{F} is of codimension $q = 1$ and U satisfies one of the properties (i) or (ii) of Proposition 4 then U is regular. If \mathcal{F} is of codimension $q \geq 2$, and if U satisfies Property (i) then U is regular by Corollary 2, (i).

Proposition 5. *There exist finitely many \mathcal{F} -saturated connected open sets V_1, V_2, \dots, V_p which satisfy Property (c) such that:*

- (i) $\mathcal{C} = V_1 \cup V_2 \cup \dots \cup V_p$.
- (ii) *Each regular non foliated fiber open set is one of the V_i , $1 \leq i \leq p$.*

Proof. (i) For each $x \in \mathcal{C}$ there exists an open set $V_x \subset \mathcal{C}$ which satisfies Property (c). So, there exists a transverse leaf $T_{\phi(x)}$ in V_x with $V_x = \text{Sat}(T_{\phi(x)})$ and $\mathcal{C} = \bigcup_{x \in \mathcal{C}} V_x$. Since $\mathcal{C} = \text{Sat}(\bigcup_{x \in \mathcal{C}} T_{\phi(x)})$ is compact by saturation (Lemma 3) then there exist finitely many transverse leaves $T_{\phi(x_1)}, T_{\phi(x_2)}, \dots, T_{\phi(x_p)}$ such that $\mathcal{C} = \bigcup_{1 \leq i \leq p} \text{Sat}(T_{\phi(x_i)}) = \bigcup_{1 \leq i \leq p} V_{x_i}$, as desired. Property (ii) follows from the fact that: if R is a regular non foliated fiber open set then $R \cap V_i \neq \emptyset$ for some i , $1 \leq i \leq p$. It follows that $R = V_i$ by Corollary 3. \square

5. Main result

Theorem 1. 1. *The union $\mathcal{A} \cup \mathcal{D}$ is open and dense in M and we have:*

- $\mathcal{A} \cup \mathcal{D} \subset \mathcal{A} \cup \mathcal{R} \subset \mathcal{A} \cup \mathcal{C}$.
- 2. $M - \mathcal{A} \cup \mathcal{C} = \bigcup_{1 \leq i \leq p} \text{Fr}(C_i)$ where $(C_i)_{1 \leq i \leq p}$ are the finite connected components of $\mathcal{A} \cup \mathcal{C}$ which do not satisfy Property (a).
- 3. $\mathcal{C} - \mathcal{A}$ is a union of finite local minimal sets of \mathcal{F} .

Proof. Assertion (1): Suppose the contrary; that is there exists a nonempty, open,

connected \mathcal{F} -saturated set $U \subset M - \overline{\mathcal{A} \cup \mathcal{D}}$. Since U does not satisfy Property (a), there exists by Lemma 7, a transverse leaf T_1 such that $\text{Sat}(T_1) \subset U$ and $\text{Sat}(T_1)$ satisfies Property (c). By Lemma 3, there exists a minimal set E in $\text{Sat}(T_1)$. Since $U \subset M - \mathcal{A} \cup \mathcal{D}$ we have $E \neq \text{Sat}(T_1)$ and $\text{Sat}(T_1) - E$ satisfies Property (a) by Proposition 4, a contradiction.

Assertion (2): First, we deduce from assertion (1) that $\mathcal{A} \cup \mathcal{C}$ is dense in M . Now, let $(C_i)_{1 \leq i \leq p}$ be the family of connected components of $\mathcal{A} \cup \mathcal{C}$ which do not satisfy Property (a). (We know that such connected components are finite by Lemma 5). Now, since the C_i are closed in $\mathcal{A} \cup \mathcal{C}$ then, $\bigcup_{1 \leq i \leq p} \text{Fr}(C_i) \subset M - \mathcal{A} \cup \mathcal{C}$. To prove the other inclusion $M - \mathcal{A} \cup \mathcal{C} \subset \bigcup_{1 \leq i \leq p} \text{Fr}(C_i)$, let $x \in M - \mathcal{A} \cup \mathcal{C}$. Take $U_n = \text{Sat}(B_n)$ where B_n is the open ball centered at x with radius $\frac{1}{n}$. Thus, U_n is an open \mathcal{F} -saturated set which does not satisfy the Property (a). By Lemma 7, U_n contains an open set V_n which satisfies Property (c). Thus, we can suppose that for every n , V_n is contained in one of the $(C_i)_{1 \leq i \leq p}$, say C_1 , and $x \notin C_1$. Take $x_n \in V_n \cap B_n \subset C_1$. Then the sequence $(x_n)_{n \geq 1}$ converges to x and we have $x \in \text{Fr}(C_1)$, this completes the proof of Assertion (2).

Assertion (3): Let $x \in \mathcal{C} - \mathcal{A}$ and L_x be a leaf passing through x . From Proposition 5, $x \in V_i$ for some i , $1 \leq i \leq p$. We will show that $\text{cl}(L_x)$ is the only minimal set of $\mathcal{F} \mid V_i$: If V_i is not an open local minimal set of \mathcal{F} , then by Proposition 4 and Corollary 6, there exists a unique minimal set E_i in V_i such that each connected component of $V_i - E_i$ satisfies Property (a). Since $x \notin \mathcal{A}$, so $x \in E_i$ and then $\text{cl}(L_x) = E_i$. \square

Corollary 8 ([3]). *If \mathcal{F} is of codimension $q = 1$, then*

1. *\mathcal{R} is open and dense in M .*
2. *There exist a finite number of half-proper leaves L_1, L_2, \dots, L_p of \mathcal{F} such that $M - \mathcal{R} = \overline{L_1 \cup L_2 \cup \dots \cup L_p}$*
3. *There exist finitely many classes which are contained in regular non foliated product sets.*

Proof. Since \mathcal{F} is of codimension $q = 1$, we have $\mathcal{A} \cup \mathcal{C} = \mathcal{R}$. Hence, Assertion (1) follows from Theorem 1, (1). Assertion (2) follows from Theorem 1, (2) since the connected components $(R_i)_{1 \leq i \leq p}$ of \mathcal{R} which do not satisfy Property (a) are exactly the regular non foliated product open sets and since $\text{Fr}(R_i)$ are the closure of finitely many leaves and these leaves are half-proper and attracting in a proper side (cf. [3]). Assertion (3) follows from Theorem 1, (3). \square

Acknowledgements. We wish to express our thanks to the referee for reading carefully the paper and for providing us with helpful comments.

References

- [1] P. R. Dippolito, Codimension one foliations of closed manifolds, *Annals of Math.* **107** (1978), 403–453.
- [2] C. Godbillon, *Feuilletages. Études géométriques*, Progress in Mathematics, Vol. 98, Birkhäuser Verlag, 1991.
- [3] E. Salhi, Sur un théorème de structure des feuilletages de codimension 1, *C. R. Acad. Sc. Paris, série I* **300** (1985), 635–638.
- [4] E. Salhi, Niveau des feuilles, *C. R. Acad. Sc. Paris, série I* **301** (1985), 219–222.

Habib Marzougui
Department of Mathematics
Faculty of Sciences of Bizerte
Zarzouna 7021
Tunisia
e-mail: habib.marzouki@fsb.rnu.tn

Ezzeddine Salhi
Department of Mathematics
Faculty of Sciences of Sfax
B.P. 802
3018 Sfax
Tunisia
e-mail: Ezzeddine.Salhi@fss.rnu.tn

(Received: November 30, 2000; revised version: March 20, 2001)



To access this journal online:

<http://www.birkhauser.ch>