

Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	78 (2003)
Artikel:	An infinitesimal trace formula for the Laplace operator on compact Riemann surfaces
Autor:	Golse, François / Lochak, Pierre
DOI:	https://doi.org/10.5169/seals-58778

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

An infinitesimal trace formula for the Laplace operator on compact Riemann surfaces

François Golse and Pierre Lochak

Abstract. We derive an infinitesimal (or variational) version of the Selberg trace formula for compact Riemann surfaces, which gives information on the behaviour of the eigenvalues of the Laplace–Beltrami operator as the surface varies over the appropriate moduli space.

Mathematics Subject Classification (2000). 11F72, 32G34 35P05.

Keywords. Riemann surfaces, Selberg trace formula, moduli spaces of curves.

1. Introduction

In this note we derive a variational form of the Selberg trace formula for compact Riemann surfaces, as the surface varies over the appropriate moduli space. We use the real analytic theory of moduli spaces, taking advantage in particular of Wolpert’s variational formula ([6]) which lies at the heart of his study of the symplectic geometry of the moduli spaces.

Let X be a compact Riemann surface of genus $g > 1$, equipped with its Poincaré metric of constant curvature -1 . Let Δ_X be the attending Laplace operator on X and $\{\lambda_n\}_{n \geq 0}$ be its spectrum, that is the sequence of the (positive) eigenvalues of $-\Delta_X$ labelled in increasing order and with multiplicities. For any $n \in \mathbf{Z}_+$ we define as usual the auxiliary spectral numbers ρ_n via the equalities $\lambda_n = \frac{1}{4} + \rho_n^2$; true this defines $\rho_n \in \mathbf{C}$ only up to sign but we will consider only even functions of the ρ_n ’s. We let \mathcal{M}_g denote the moduli space of Riemann surfaces of genus g , viewed as a complex orbifold (see below for more detail).

A lot of work has been devoted to analysing the variation of the spectra of Riemann surfaces. However the information which has been collected to-date mostly concerns the behavior of the low lying eigenvalues as the surface degenerates. More precisely \mathcal{M}_g can be compactified into $\overline{\mathcal{M}}_g$ where the divisor at infinity $\mathcal{D} = \overline{\mathcal{M}}_g \setminus \mathcal{M}_g$ classifies the Riemann surfaces with nodes – or stable curves as they are called by algebraic geometers. For any point in \mathcal{D} , the corresponding surface can be thought of as a graph with surfaces of lower genera sitting at the vertices,

such that the total arithmetic genus is g . Let now X_ε be a family of surfaces such that as ε tends to 0, X_ε tends to a surface represented by a graph with k vertices ($2 \leq k \leq 2g - 2$); then as ε tends to 0, the first k eigenvalues (counting 0) of X_ε tend to 0 and the $(k + 1)$ -st one is uniformly bounded away from 0. In this prototypical situation tools have been developed which make it possible to prove the assertions summarized above and also to study more accurately the behavior of the $k - 1$ nonzero but vanishing eigenvalues. We refer to ([1]) for a short history of the development of this line of thought, references, as well as a recent and precise result of this type.

Very little seems to be known about the variation of arbitrary (in particular non vanishing) eigenvalues of arbitrary (in particular non degenerating) surfaces. We refer to ([8]) for results in the case of degenerating surfaces but non vanishing eigenvalues. The formula derived in this note can be used, like the original trace formula, in the vicinity of any surface, not necessarily near the boundary of \mathcal{M}_g . However, again like the trace formula, it gives global, not individual information on the (variation of the) spectrum. It may be that using well-adapted test functions, in particular functions which are peaked around a given point, one can extract more local information. We postpone to the concluding section some more speculative remarks about possible applications and amplifications and presently recall some additional background material.

Considering again a compact Riemann surface X of genus $g > 1$, it can be uniformized as $X = \mathcal{H}/\Gamma$, where \mathcal{H} denotes the Poincaré upper-half plane and $\Gamma \subset PSL(2, \mathbf{R})$ is a hyperbolic Fuchsian group of the first kind. Let γ be a closed geodesic on X of length $\ell(\gamma)$; one can also regard γ as an element of Γ up to conjugacy and the length is given in matrix term *via* the formula: $\text{trace}(\gamma) = 2 \cosh(\ell(\gamma)/2)$. Lastly we denote by \mathcal{P} the set of oriented primitive closed geodesics, *i.e.* the set of non divisible conjugacy classes in Γ . For any $\gamma \in \Gamma$, there are unique $p \in \mathcal{P}$ and $m \in \mathbf{Z}^*$ such that $\gamma = p^m$; note that we do not distinguish conjugacy classes and representatives in the notation, when no confusion is liable to arise from this abuse.

Selberg's trace formula deals with even "test" functions $h \equiv h(\rho)$, with Fourier transform $\hat{h}(\tau)$ defined by:

$$\hat{h}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} h(\rho) \cos(\tau\rho) d\rho.$$

The even test function h has to satisfy certain regularity and growth conditions which we state in the following

Assumptions

- (i) h can be continued into an analytic function over a strip $|Im(\rho)| < c$ for some $c > \frac{1}{2}$ and it satisfies the estimates $h(\rho) = O(|\rho|^{-\beta})$ as well as $h'(\rho) = O(|\rho|^{-\beta-1})$ for some $\beta > 4$, as ρ goes to infinity inside the strip;

(ii) *The Fourier transform \hat{h} satisfies $\hat{h}(\tau) = O(e^{-c|\tau|})$ as τ goes to infinity on \mathbf{R} .*

Under the above assumptions h satisfies the Selberg trace formula (cf. e.g. [3], section 2.6), to wit:

$$\begin{aligned} \sum_{n \geq 0} h(\rho_n) &= (2g-2) \int_{-\infty}^{+\infty} \rho \tanh(\pi\rho) h(\rho) d\rho \\ &\quad + \sum_{\gamma \in \mathcal{P}} \sum_{m=1}^{\infty} \frac{\ell(\gamma)}{2 \sinh(m\ell(\gamma)/2)} \hat{h}(m\ell(\gamma)). \end{aligned} \tag{1}$$

The trace formula (1) depends on the surface X , and the aim of this note is so to speak to “take the derivative of (1) w.r.t. X ”. To this end we introduce the Teichmüller space \mathcal{T}_g associated with (compact) surfaces of genus g ; the moduli space is then given as the quotient $\mathcal{M}_g = \mathcal{T}_g/\Gamma_g$, where Γ_g denotes the mapping class group (or Teichmüller modular group) of genus g , that is the orbifold fundamental group of \mathcal{M}_g . We recall but a few definitions in order to fix notation and refer the reader to e.g. [2], [4] or [5] for more information on these objects. So let S denote “the” differentiable, closed, orientable and compact surface of genus g . Given a Riemann surface X of genus g , a marking of X is given by a diffeomorphism $\phi : S \rightarrow X$, considered up to isotopy. Two marked Riemann surfaces (X_1, ϕ_1) and (X_2, ϕ_2) are equivalent if there exists a *conformal* diffeomorphism $h : X_1 \rightarrow X_2$ such that $\phi_2 = h \circ \phi_1$, again up to isotopy. A point of \mathcal{T}_g is an equivalence class $\mathbf{t} = [(X, \phi)]$ of marked Riemann surfaces. We denote by $\pi : \mathcal{T}_g \rightarrow \mathcal{M}_g$ the canonical projection which consists in forgetting the marking: $\pi(\mathbf{t}) = X$.

Any isotopy class α of closed curves on S determines a function ℓ_α on \mathcal{T}_g , such that $\ell_\alpha(\mathbf{t}) = \ell_X(\phi(\alpha))$ is by definition the length of the unique geodesic on X which is isotopic to $\phi(\alpha)$. If moreover α is simple, meaning that there exists a simple representative of α (a regular curve in the class of α without self-intersection), it also determines a vector field τ_α on \mathcal{T}_g , namely the infinitesimal Fenchel–Nielsen twist along α (see [4], [5] or [7]). For any $\mathbf{t} \in \mathcal{T}_g$, one can explicitly find $6(g-1)$ simple closed curves α_i , $1 \leq i \leq 6(g-1)$, such that the corresponding $\tau_{\alpha_i}(\mathbf{t})$ span the tangent space of \mathcal{T}_g at \mathbf{t} (see [7], Corollary 3.5). Recall also that \mathcal{M}_g is smooth at $X = \pi(\mathbf{t})$ if X has no nontrivial automorphism, in which case the images $\pi_*(\tau_{\alpha_i})$ span the tangent space of \mathcal{M}_g at X . Lastly we mention that the differential of ℓ_α is conjugate to τ_α for the symplectic structure defined by the Weil–Petersson two-form; in other words ℓ_α is nothing else but Hamilton’s function for the vector field τ_α (see section 8.3 of [4] or [7]).

From now on we do not notationally distinguish α from the geodesic which is isotopic to $\phi(\alpha)$ on some marked Riemann surface. An important ingredient in what follows is

Lemma 1 (S. Wolpert, [6]). *The variation of the geodesic length along a twist is*

given by:

$$\tau_\alpha(\ell(\gamma)) = C(\alpha, \gamma), \quad \text{where} \quad C(\alpha, \gamma) = \sum_{p \in \alpha \cap \gamma} \widehat{\cos(\alpha, \gamma)}_p. \quad (2)$$

Here $\widehat{(\alpha, \gamma)}_p$ denotes the angle of the two intersecting geodesics at p , with the following sign convention (see [6]): lift p to $\tilde{p} \in \mathcal{H}$ and assume w.l.o.g. that α lifts to the vertical axis; then $\widehat{(\alpha, \gamma)}_p$ is defined to be the angle between this axis (pointing upward) and the half-line tangent to γ pointing towards the right half-plane. This convention makes the expression $\widehat{\cos(\alpha, \gamma)}_p$ skew symmetric in (α, γ) , thus also the sum $C(\alpha, \gamma)$.

2. The infinitesimal trace formula

The main result of this note is embodied in the following infinitesimal trace formula:

Theorem 2. *Let h be an even function satisfying assumptions (i) and (ii) of section 1, and let α be a simple loop on S . Then the following identity holds:*

$$\sum_{n \geq 0} \frac{h'(\rho_n)}{2\rho_n} \tau_\alpha^\pm(\lambda_n) = \sum_{\gamma \in \mathcal{P}} C(\alpha, \gamma) \left(\sum_{m=1}^{\infty} \phi'_h(m\ell(\gamma)) \right), \quad (3)$$

where ϕ_h denotes the map $x \mapsto x/2(\sinh(x/2))^{-1}\hat{h}(x)$ and τ_α^+ (resp. τ_α^-) denotes the right (resp. left) derivative along the vector field τ_α .

The pieces of notation $C(\alpha, \gamma)$ and ϕ_h have been introduced for convenience only. Explicitly we have:

$$\phi'_h(x) = \frac{x/2}{\sinh(x/2)} \hat{h}'(x) + \frac{1 - x/2 \coth(x/2)}{2 \sinh(x/2)} \hat{h}(x).$$

Note furthermore that the l.h.s. of (3) can also be written in the form $\sum_{j \geq 0} h'(\rho_j) \tau_\alpha(\rho_j)$, which is more natural, provided this expression is correctly interpreted; see the remark at the end of section 3 below.

Formula (3), which can be considered as “the derivative of (1) along α ”, can be formally deduced from (1) by means of (2). The next two sections are devoted to justifying this derivation by showing the absolute convergence of the l.h.s. (the spectral side) and then of the r.h.s. (the geometric side) of (3).

3. The spectral side

Let \mathbf{t}_0 be a point of \mathcal{T}_g and α a simple closed loop on S . We denote by $\mathbf{t}_\theta \in \mathcal{T}_g$ the point which is obtained from \mathbf{t}_0 after twisting along α by the angle θ (see *e.g.* [4], section 3.2 and pp. 219–221). Let $X_\theta = \pi(\mathbf{t}_\theta)$; g_θ denotes the Poincaré metric on X_θ and Δ_θ the Laplace operator on X_θ for the metric g_θ . The increasing sequence of the eigenvalues of $-\Delta_\theta$ counted with multiplicities reads $0 = \lambda_0(\theta) \leq \lambda_1(\theta) \leq \dots \leq \lambda_n(\theta) \leq \dots$

Now for any even function h satisfying the assumptions of section 1, there is a unique function H_h which is analytic in a neighborhood of the real half-line $]-\frac{1}{2}, +\infty[$ and satisfies $H_h(z^2) = h(z)$ for $z \in \mathbf{C}$ such that $z^2 \in]-\frac{1}{2}, +\infty[$. In this section we show that under the assumptions (i)–(ii) of section 1, the map $\theta \mapsto \sum_{n \geq 0} H_h(\lambda_n(\theta) - \frac{1}{4})$ has left and right derivatives at $\theta = 0$ and that the following holds:

$$\tau_\alpha^\pm \left(\sum_{n \geq 0} H_h(\lambda_n(\theta) - \frac{1}{4}) \right) = \sum_{n \geq 0} H'_h(\lambda_n(\theta) - \frac{1}{4}) \tau_\alpha^\pm(\lambda_n). \quad (4)$$

We first prove:

Lemma 3. *The map $\theta \mapsto (I - \Delta_\theta)^{-1}$ is analytic over \mathbf{R} , with values in the space of bounded operators on L^2 , endowed with the topology given by the operator norm.*

Proof. The last two formulas on p. 133 of [6] explicitly describe a one-parameter family of Beltrami coefficients which define the points $\mathbf{t}_\theta \in \mathcal{T}_g$. The third formula on p. 134 of that paper gives explicit formula for the corresponding hyperbolic line element ds . This formula shows in particular that the dependence of the metric tensor ds^2 upon θ is real analytic. Using the expression of the Laplace operator in terms of the metric, one finds that $(I - \Delta_\theta)(I - \Delta_0)^{-1}$ is a linear combination of the four operators $(I - \Delta_0)^{-1}$, $\partial_z(I - \Delta_0)^{-1}$, $\partial_{\bar{z}}(I - \Delta_0)^{-1}$ and $\partial_z \partial_{\bar{z}}(I - \Delta_0)^{-1}$ with coefficients that are real-analytic functions of θ . Since the four operators above are bounded on $L^2(S)$, this shows that the map $\theta \mapsto (I - \Delta_\theta)(I - \Delta_0)^{-1}$ is real analytic over \mathbf{R} with values in the algebra of bounded operators on $L^2(S)$, equipped with the operator norm. Furthermore, for each $\theta \in \mathbf{R}$ the bounded operator $(I - \Delta_\theta)(I - \Delta_0)^{-1}$ is an invertible element of that algebra. Thus the map $\theta \mapsto [(I - \Delta_\theta)(I - \Delta_0)^{-1}]^{-1}$ is also real analytic on \mathbf{R} , which finishes the proof of the lemma. \square

As the operator Δ_θ is self-adjoint on $L^2(X_\theta)$ for any $\theta \in \mathbf{R}$, lemma 3 combined with the Kato–Rellich theorem (*cf.* [9], section XII.13) implies that for any $n \in \mathbf{Z}_+$, λ_n is Lipschitz continuous on \mathbf{R} w.r.t. θ , with left and right derivatives at every point. Writing $(\lambda_n)'_l(\theta)$ (resp. $(\lambda_n)'_r(\theta)$) for the left (resp. right) derivative, we have the following estimate:

Lemma 4. *For any $n \in \mathbf{Z}_+$ and any $\theta \in \mathbf{R}$,*

$$\max \frac{\|(\lambda_n)'_l(\theta)\|, \|(\lambda_n)'_r(\theta)\|}{[1 + \lambda_n(\theta)]^2} \leq \left\| \frac{d}{d\theta} (I - \Delta_\theta)^{-1} \right\|. \quad (5)$$

Proof. Let $A_\theta = (I - \Delta_\theta)^{-1}$ and $\mu_n = 1/(1 + \lambda_n)$. Pick $n \in \mathbf{Z}_+$ and $\theta > 0$. Then either $\mu_n(\theta)$ belongs to the spectrum $\text{spec}(A_0)$ of A_0 , or it does not. In the former case the operator $I - (\mu_n(\theta) - A_0)^{-1}(A_\theta - A_0)$ is not invertible, which shows that

$$1 \leq \|(\mu_n(\theta) - A_0)^{-1}\| \|A_\theta - A_0\| = \text{dist}(\mu_n(\theta), \text{spec}(A_0))^{-1} \|A_\theta - A_0\|.$$

Since λ_n is known to have a right derivative at $\theta = 0$, the above estimate implies that the right derivative $|(\mu_n)'_r(0)|$ is bounded from above by the r.h.s. of (5). The proof for the left derivative at 0 and for the other values of θ is identical. \square

We now come to the proof of identity (4). For any $n \geq 0$ the function $\theta \mapsto H_h(\lambda_n(\theta) - \frac{1}{4})$ has left and right derivatives at every point $\theta \in \mathbf{R}$. Thanks to assumption (i) the function H_h satisfies $H'_h(x) = h'(\sqrt{x})/2\sqrt{x} = O(x^{-\beta/2-1})$ for some $\beta > 4$, so that using the estimate (5) we find that

$$\sup\|(\lambda_n)'_g(\theta)\|, \|(\lambda_n)'_d(\theta)\| \|H'_h(\lambda_n(\theta) - \frac{1}{4})\| = O(\lambda_n(\theta)^{1-\beta/2}).$$

Now the Weyl asymptotic estimate $\lambda_n(\theta) = O(n)$ is locally uniform w.r.t. θ ; the estimate above, combined with this fact and the dominated convergence theorem vindicates formula (4).

Remark. By the Kato–Rellich theorem quoted above, there exists in fact a re-labeling $\sigma : \mathbf{Z}_+ \rightarrow \mathbf{Z}_+$ of the spectrum such that for any $j \in \mathbf{Z}_+$ the map $\theta \mapsto \lambda_{\sigma(j)}(\theta)$ is real analytic in θ over some j -dependent neighborhood of $\theta = 0$. So for $\theta = 0$ we may in fact write:

$$\sum_{n \geq 0} H'_h(\lambda_n - \frac{1}{4}) \tau_\alpha^\pm(\lambda_n) = \sum_{j \geq 0} h'(\rho_{\sigma(j)}) \tau_\alpha(\rho_{\sigma(j)}),$$

in which $h'(\rho_{\sigma(j)}) \tau_\alpha(\rho_{\sigma(j)})$ is interpreted as $\frac{h'(\rho_{\sigma(j)})}{\rho_{\sigma(j)}} \tau_\alpha(\rho_{\sigma(j)}^2)$. Indeed since h is even and analytic in a neighborhood of the real axis, this is also the case for the function $h'(z)/z$.

4. The geometric side

We turn to the r.h.s. of the variational trace formula (3), with the aim of proving the identity:

$$\tau_\alpha \left(\sum_{\gamma \in \mathcal{P}} \sum_{m=1}^{\infty} \phi_h(\gamma) \right) = \sum_{\gamma \in \mathcal{P}} C(\alpha, \gamma) \left(\sum_{m=1}^{\infty} \phi'_h(m\ell(\gamma)) \right). \quad (6)$$

Given (2) (see Wolpert's lemma in section 1) this amounts to justifying the switching of the sum and the derivative τ_α . To this end we need to bound the intersection term $C(\alpha, \gamma)$ from above. Given the closed geodesics α and γ , let $I(\alpha, \gamma)$ be the number of their intersection points; then clearly $|C(\alpha, \gamma)| \leq I(\alpha, \gamma)$, so that it suffices to bound the last quantity, with the additional assumption that α is simple. In order to do this we need some elementary hyperbolic geometry and first define the quantity $\eta_m = \eta_m(X, \alpha) > 0$ as follows. Given $\eta \geq 0$, let $B_\alpha(\eta)$ be the open strip of width 2η centered around α , in other words the set of points of X whose distance to α is strictly smaller than η . We define η_m as the supremum of those $\eta > 0$ such that $B_\alpha(\eta)$ is isometric to the standard annulus of width 2η . To put it slightly differently, for any $\eta < \eta_m$ and any point $x \in B_\alpha(\eta)$, there is a unique geodesic segment contained in $B_\alpha(\eta)$, originating at x and perpendicular to α ; moreover η_m is maximal with this property.

Then we have the following estimate:

Lemma 5. $I(\alpha, \gamma) \leq \ell(\gamma)/(2\eta_m)$.

Proof. Let $p \in \alpha \cap \gamma$ and $\eta < \eta_m$; then γ traverses the strip $B_\alpha(\eta)$ and defines there a geodesic segment which we denote by I_p . Clearly the length of I_p is at least 2η . Moreover as p runs along the finite set $\alpha \cap \gamma$, the segments I_p are disjoint, except possibly for a finite number of transverse intersections corresponding to self-intersections of γ located in the strip $B_\alpha(\eta)$. The lemma follows from these considerations. \square

Remarks. There is a universal lower bound for η_m , depending only on the length of the simple geodesic loop α . In fact one can show that $\eta_m(\alpha, X) \geq \eta_0(\ell(\alpha))$, where the function η_0 is defined by the equality $\cosh(\eta_0(\ell)) = 1/\tanh(\ell/2)$ (see [5], section I.1.5).

One can also notice that because two geodesics always intersect transversally, the intersection number $I(\alpha, \gamma)$ is in fact constant over T_g . Moreover this constant coincides with the topological intersection number of the isotopy classes α and γ on the model topological surface S . Recall that this number is defined as the minimal number of intersection points of two representatives of the classes α and γ . The fact that $I(\alpha, \gamma)$, which was first defined *via* hyperbolic geometry, coincides with that purely topological quantity is one of the basic results in Thurston's theory of surfaces, leading in particular to his compactification of Teichmüller spaces (see [10], Exposé 3).

We come to the by now easy proof of (6). Wolpert's lemma recalled in section 1 ensures that $|\tau_\alpha(l(\gamma))| = |C(\alpha, \gamma)| \leq I(\alpha, \gamma)$, and this last quantity is bounded from above as in lemma 5. On the other hand, assumptions (i)-(ii) on the test function h show that $\phi'(ml(\gamma)) = O(e^{-c'ml(\gamma)})$ for all $c' \in]1, c + \frac{1}{2}[$. The dominated convergence theorem then justifies term-by-term differentiation of the r.h.s. of (6),

finishing the proof of the validity of (6).

Putting together the results of section 3 and section 4 finishes the proof of the theorem, showing the validity of (3).

5. Amplifications, possible applications, remarks

Let us close with some of the remarks the above might trigger. First we note that we have adopted the point of view of hyperbolic geometry along with the real analytic description of the Teichmüller and moduli spaces. One can at least partially introduce the conformal viewpoint for surfaces along with the complex analytic theory for the classifying spaces by replacing (2) with an analogue in terms of Poincaré series (see in particular [7], theorem 2.9).

The input on the geometric side of the trace formula (1) is the length spectrum of the surface, that is the collection of the numbers $\ell(\gamma)$ for all primitive geodesics $\gamma \in \mathcal{P}$ (counting with multiplicities). By a classical remark originally due to H. Huber and A. Selberg, the trace formula itself implies that the spectrum and the length spectrum determine each other. Now working around a given surface X , the input of the variational trace formula (3) consists of either of the two spectra supplemented by the geometric information conveyed by the numbers $C(\alpha, \gamma)$. Now for any test function h satisfying the assumptions stated in section 1, (3) provides a linear relation between the derivatives $\tau_\alpha^\pm(\lambda_n)$ (see also the remark at the end of section 3). Varying h , one finds that (3) actually determines the collection of these numbers, *i.e.* it determines so to speak the derivative of the spectrum of the surface along any given twist. So either of the length spectrum or the spectrum, supplemented with the intersection data $C(\alpha, \gamma)$ for all pairs of primitive geodesics $\alpha, \gamma \in \mathcal{P}$ determine both the derivatives of the length spectrum (by lemma 1) and of the spectrum (by (3)).

We have discussed only the comparatively elementary case of closed Riemann surfaces. From a more general viewpoint the above is certainly a rank 1 phenomenon, because in higher ranks rigidity results precisely say that the analog simply does not exist. It would however be interesting to investigate the possibility of adapting the above to other base fields (still rank 1 but not over the complex numbers). Coming back to Riemann surfaces (*alias* complex curves) one can add elliptic elements for free, because these are rigid. In other words the first term on the r.h.s. of (1) corresponds to the identity element of the uniformizing group Γ and disappears upon taking the derivative, as it is independent of the curve. In exactly the same way the additional terms which appear when Γ is not assumed to be torsion free depend only on the signature of the group (see *e.g.* [11], theorem 9.1 with $\chi = 1$); so they also disappear in the variation. The extension to the noncompact case however is of course highly non trivial and hopefully interesting, and so is the degenerating process corresponding to – depending on the viewpoint – squeezing a simple closed geodesic, going to the boundary of the moduli space and

the emergence of parabolic elements in the uniformizing Fuchsian group.

It is to be hoped that (3) and possible extensions or variations therof can be useful in several situations. Here are a few tracks, including possible dead ends. One may want to study the variation of the Selberg zeta function or other generating functions. The formula can in principle be iterated or even exponentiated, so as to get information about what happens along a *finite* twist (see [7], theorem 3.4). One can also try to investigate various bifurcation phenomena such as the study of the spectrum near a surface with nontrivial automorphisms, or the splitting of a multiple eigenvalue as the surface varies.

Acknowledgements. We thank Eric Leichtnam and the referee for helpful remarks.

References

- [1] P. Batchelor, Dérivée des petites valeurs propres des surfaces de Riemann, *Comment. Math. Helvetici* **73** (1998), 337–352.
- [2] P. Buser, *Geometry and Spectra of Compact Riemann Surfaces*, Birkhäuser Verlag, 1992.
- [3] P. Cartier and A. Voros, Une nouvelle interprétation de la formule des traces de Selberg, in: P. Cartier et al. eds., *The Grothendieck Festschrift*, Volume II, 1–66, Progress in Mathematics **87**, Birkhäuser, 1990.
- [4] Y. Imayoshi and M. Taniguchi, *An introduction to Teichmüller spaces*, Springer Verlag, 1992.
- [5] X. Buff, J. Fehrenbach and P. Lochak, Éléments de géométrie des espaces de modules de courbes, in: *Espaces de modules de courbes, groupes modulaires et théorie des champs*, Panoramas et Synthèses **7**, Publications SMF, 1999.
- [6] S. Wolpert, An elementary formula for the Fenchel–Nielsen twist, *Comment. Math. Helvetici* **56** (1981), 132–135.
- [7] S. Wolpert, The Fenchel–Nielsen deformation, *Annals of Math.* **115** (1982), 501–528.
- [8] S. Wolpert, Spectral limits for hyperbolic surfaces, I and II, *Invent Math.* **108** (1992), 67–89 and 90–129.
- [9] M. Reed and B. Simon, *Methods of Modern Mathematical Physics*, Vol. 4, Academic Press, 1978.
- [10] *Travaux de Thurston sur les surfaces*, Astérisque **66–67**, A. Fathi, F. Laudenbach and V. Poenaru eds., SMF Publ., 1979 (2nd edition 1991).
- [11] A. B. Venkov, Spectral theory of automorphic functions, *Russian Math. Surveys* **34** (1979), 79–153.

F. Golse
 Institut Universitaire de France
 & Laboratoire Jacques-Louis Lions
 175 rue du Chevaleret
 75013 Paris
 France
 e-mail: golse@math.jussieu.fr

P. Lochak
 Institut Mathématique de Jussieu
 175 rue du Chevaleret
 75013 Paris
 France
 e-mail: lochak@math.jussieu.fr

(Received: May 7, 2001)