

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 82 (2007)

Artikel: The barycenter method on singular spaces
Autor: Storm, Peter
DOI: <https://doi.org/10.5169/seals-98880>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 09.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The barycenter method on singular spaces

Peter A. Storm*

Abstract. Compact convex cores with totally geodesic boundary are proven to uniquely minimize volume over all hyperbolic 3-manifolds in the same homotopy class. This solves a conjecture in Kleinian groups concerning acylindrical 3-manifolds. Closed hyperbolic manifolds are proven to uniquely minimize volume over all compact hyperbolic cone-manifolds in the same homotopy class with cone angles $\leq 2\pi$. Closed hyperbolic manifolds are proven to minimize volume over all compact Alexandrov spaces with curvature bounded below by -1 in the same homotopy class. A version of the Besson–Courtois–Gallot theorem is proven for n -manifolds with boundary. The proofs extend the techniques of Besson–Courtois–Gallot.

1. Introduction

This paper extends the barycenter map machinery of Besson–Courtois–Gallot [BCG1] to a class of singular metric spaces called *convex Riemannian amalgams* (defined in Section 2.8). This class of singular spaces includes cone-manifolds and the metric doubling of hyperbolic convex cores across their boundary. These singular space techniques are used to solve a conjecture in Kleinian groups due to Bonahon. Specifically, we prove that compact convex cores with totally geodesic boundary uniquely minimize volume over all hyperbolic 3-manifolds in the same homotopy class (see Theorem 8.1).

The main tool of this paper is the following extension of Besson–Courtois–Gallot techniques to convex Riemannian amalgams.

Theorem 4.1. *For $n \geq 3$, let Z be a compact n -dimensional convex Riemannian amalgam. Let M_{hyp} be a closed hyperbolic n -manifold. Let $h(\tilde{Z})$ denote the volume growth entropy of the universal cover of Z . If $f: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence then*

$$h(\tilde{Z})^n \text{Vol}(Z) \geq (n-1)^n \text{Vol}(M_{\text{hyp}})$$

with equality if and only if f is homotopic to a homothetic homeomorphism.

By restricting attention to cone-manifolds, we obtain

*This research was partially supported an NSF Postdoctoral Fellowship.

Theorem 8.6. *For $n \geq 3$, let Z be compact n -dimensional cone-manifold built with simplices of constant curvature $K \geq -1$. Assume all its cone angles are $\leq 2\pi$. Let M_{hyp} be a closed hyperbolic n -manifold. If $f: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence then*

$$\text{Vol}(Z) \geq \text{Vol}(M_{\text{hyp}})$$

with equality if and only if f is homotopic to an isometry.

If Z is allowed to be any Alexandrov space with curvature bounded below by -1 (see Section 2.13), we then obtain

Theorem 8.7. *Let Z be a compact n -dimensional ($n \geq 3$) Alexandrov space with curvature bounded below by -1 . Let M_{hyp} be a closed hyperbolic n -manifold. If $f: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence then*

$$\text{Vol}(Z) \geq \text{Vol}(M_{\text{hyp}}).$$

As mentioned above, the paper's main theorems solve a conjecture in Kleinian groups. To state things precisely, let $H(N)$ denote the set of marked oriented isometry classes of hyperbolic 3-manifolds M equipped with a homotopy equivalence $N \rightarrow M$. Define a volume function

$$\text{Vol}: M \in H(N) \mapsto \text{Vol}(C_M).$$

It is a consequence of Thurston's Geometrization Theorem and Mostow Rigidity that N is acylindrical if and only if there exists a convex cocompact $M_g \in H(N)$ such that ∂C_{M_g} is totally geodesic [Th2, p. 14]. Moreover, M_g is unique up to isometry. Let M_g^{opp} denote M_g with the opposite orientation.

Conjecture. M_g and M_g^{opp} are the *only* global minima of Vol over $H(N)$.

Initial progress on this conjecture was made by Bonahon [Bon]. Using different methods, Bonahon proved that M_g is a strict local minimum of Vol (in the quasi-isometric topology on $H(N)$). In [S], the author proved that M_g and M_g^{opp} are global minima of Vol . Here this conjecture is completely solved by

Theorem 8.1. *Let N be a compact acylindrical 3-manifold. Let $M_g \in H(N)$ be a convex cocompact hyperbolic 3-manifold such that the boundary of the convex core $\partial C_{M_g} \subset M_g$ is totally geodesic. Then for all $M \in H(N)$,*

$$\text{Vol}(C_M) \geq \text{Vol}(C_{M_g}),$$

with equality if and only if M and M_g are isometric.

The techniques used to prove Theorem 8.1 immediately generalize to a version of the Besson–Courtois–Gallot theorem for manifolds with boundary. (Perelman's

unpublished Doubling theorem [P, Theorem 5.2] is used in the proof of Theorem 8.9. See Theorem 8.8.)

Theorem 8.9. *Let Z be a compact convex Riemannian n -manifold with boundary ($n \geq 3$). Assume the sectional curvature of $\text{int}(Z)$ is bounded below by -1 . Let Y_{geod} be a compact convex hyperbolic n -manifold with totally geodesic boundary. Let $f: (Z, \partial Z) \rightarrow (Y_{\text{geod}}, \partial Y_{\text{geod}})$ be a homotopy equivalence of pairs. Then*

$$\text{Vol}(Z) \geq \text{Vol}(Y_{\text{geod}}),$$

with equality if and only if f is homotopic to an isometry.

This paper's method of proof may be of independent interest. Following [BCG1], these results are proven by defining a *natural map* from a nice path metric space Z to a hyperbolic manifold M_{hyp} . Instead of obtaining this map as a uniform limit of approximating maps, the natural map is here obtained in a single step. The idea is to emulate the “short” proof of the Besson–Courtois–Gallot theorem found in [BCG2], where it is additionally assumed that Z is nonpositively curved. Here the assumption of nonpositive curvature is removed, but the gist of the “short” proof is retained. As a cost for this generalization, the arguments here require that Z and M_{hyp} be homotopy equivalent. In [BCG1], only a map $Z \rightarrow M_{\text{hyp}}$ of nonzero degree is required.

The results in this paper represent a large portion of the author's Ph.D. thesis, completed at the University of Michigan. The author thanks his advisor, Richard Canary, for his essential assistance at every stage of this project. The author also enjoyed several helpful conversations with Yair Minsky. Thanks to Ralf Spatzier for introducing the author to the work of Besson–Courtois–Gallot.

1.1. Sketch of the proof of the main theorem. Let Z be a compact n -dimensional convex Riemannian amalgam with universal cover X (e.g. a cone-manifold or the double of a convex core), M_{hyp} a closed hyperbolic n -manifold, and $f: Z \rightarrow M_{\text{hyp}}$ a homotopy equivalence. Up to rescaling the metric of Z , we may assume that $h(X) = (n-1) = h(\mathbb{H}^n)$. The first goal is to find a volume decreasing map $F: Z \rightarrow M_{\text{hyp}}$ homotopic to f . The second goal is to show that if the volume decreasing map F is in fact volume preserving, then it is an isometry.

Step 1. Defining “visual measures” on ∂X (Section 3). Following [BM], we define a generalization of Patterson–Sullivan measure $\{\mu_x\}_{x \in X}$ supported on a function theoretic compactification $\mathcal{H}X$ of X by Busemann functions. As Z and M_{hyp} are homotopy equivalent, X must be Gromov hyperbolic with Gromov boundary ∂X . We define an $\text{Isom}(X)$ -equivariant continuous surjection $\pi: \mathcal{H}X \rightarrow \partial X$, and use it to push forward the measures $\{\mu_x\}$ onto ∂X . The resulting family of probability measures $\{\pi_*\mu_x\}$ are the “visual measures” used to define the natural map F .

Step 2. Defining F (Section 4). The homotopy equivalence f lifts to a quasi-isometry between the universal covers $f: X \rightarrow \mathbb{H}^n$, which in turn induces a homeomorphism

of Gromov boundaries $f: \partial X \rightarrow \partial \mathbb{H}^n$. The measure $\pi_*\mu_x$ is pushed forward via the boundary homeomorphism to obtain a measure $f_*\pi_*\mu_x$ on $\partial \mathbb{H}^n$. Finally, $F(x) \in \mathbb{H}^n$ is defined to be the barycenter of the measure $f_*\pi_*\mu_x$. In sum,

$$\begin{aligned} F: X &\rightarrow \mathbb{H}^n, \\ x &\mapsto \text{barycenter of measure } (f \circ \pi)_*\mu_x. \end{aligned}$$

It is relatively easy to show that F descends to a map $F: Z \rightarrow M_{\text{hyp}}$, and is homotopic to $f: Z \rightarrow M_{\text{hyp}}$. In particular, $F: Z \rightarrow M_{\text{hyp}}$ is surjective.

Step 3. F is locally Lipschitz (Section 5). In order to use calculus to study the map F , we must prove it is locally Lipschitz. (For example, a locally Lipschitz map which is infinitesimally volume decreasing almost everywhere must be volume decreasing.) This is done by factoring F as a composition $F = P \circ \Phi$, such that P and Φ can be analyzed directly. $\Phi: X \rightarrow L_+^2(\mathcal{H}X) \subset L^2(\mathcal{H}X)$ equivariantly maps X into the strictly positive functions of the Hilbert space $L^2(\mathcal{H}X)$. $P: L_+^2(\mathcal{H}X) \rightarrow \mathbb{H}^n$ is basically the barycenter map, thinking of $L_+^2(\mathcal{H}X)$ as a space of measures. We show Φ is locally Lipschitz by direct estimates. Applying the implicit function theorem shows P is \mathcal{C}^1 . Together this shows $F = P \circ \Phi$ is locally Lipschitz.

Step 4. F is infinitesimally volume decreasing a.e. (Section 6). With only minor modifications, the arguments of [BCG2] can be applied to show $|\text{Jac } F| \leq 1$ almost everywhere. This accomplishes the first goal of showing F is volume decreasing. The arguments of [BCG2] also show that if $|\text{Jac } F(x)| = 1$ for some x , then dF_x is an infinitesimal isometry. Thus if F is volume preserving, it must be an infinitesimal isometry almost everywhere.

Step 5. F is volume preserving implies it is an isometry (Section 7). Applying the arguments from [BCG1, p. 790–793], we show a volume preserving map F is a local isometry on an open dense set. We show F is injective by using some local properties of convex Riemannian amalgams. Thus F is a homeomorphism. Again using the convex Riemannian structure on Z , we prove F is an isometry. This accomplishes the second goal, and completes the proof of Theorem 4.1.

1.2. Sketch of the applications. With the above machinery established, the theorems concerning cone-manifolds (Section 8.2), Alexandrov spaces (Section 8.3), and manifolds with boundary (Section 8.4) are easy to prove. To apply the machinery to hyperbolic convex cores requires more work (Section 8.1).

Recall the hypotheses of Theorem 8.1. Let M and M_g be homotopy equivalent acylindrical convex cocompact hyperbolic 3-manifolds. Assume the convex core $C_{M_g} \subset M_g$ has totally geodesic boundary. The goal is to prove $\text{Vol}(C_M) \geq \text{Vol}(C_{M_g})$, with equality if and only if M and M_g are isometric. To begin, metrically double the convex cores across their boundaries to obtain the convex Riemannian amalgams DC_M and DC_{M_g} . Notice that DC_{M_g} is in fact a closed hyperbolic 3-manifold. A

short argument proves there exists a π_1 -equivariant quasi-isometric homeomorphism $f: \widehat{DC}_M \rightarrow \widehat{DC}_{M_g}$. Applying the above machinery to f yields Theorem 8.1.

In the case where M is geometrically finite with at least one rank one cusp, geometric (but not hyperbolic) Dehn surgery arguments are used to reduce to the case of closed (non-hyperbolic) manifolds, where Theorem 4.1 can be applied. (These geometric Dehn surgery techniques are based on [Bes], [L].)

2. Preliminaries

The following is a review of the necessary definitions. Throughout this paper, metric spaces are assumed to be complete unless otherwise stated.

2.1. δ -hyperbolicity. This paper will follow the definitions and notation of [GH]. For convenience, we recall a few basic notions. Let (X, d) be a δ -hyperbolic space with basepoint $o \in X$. Then for $x, y \in X$, the *Gromov product* of x and y is

$$(x | y) := \frac{1}{2} \{d(x, o) + d(y, o) - d(x, y)\}.$$

A defining property of δ -hyperbolic spaces is that for any triple $x, y, z \in X$,

$$(x | y) \geq \min\{(x | z), (z | y)\} - \delta. \quad (1)$$

The geometric content of the Gromov product may be difficult to grasp initially. The idea is that geodesic triangles in a δ -hyperbolic space are very close to being tripods. For a tripod the Gromov product has the simple interpretation shown in Figure 1: it is the length of the tripod's “ o ” leg. Since long geodesic triangles in a δ -hyperbolic




Figure 1

space are very close to tripods, on a sufficiently large scale the Gromov product is the length of the “ o ” leg of the geodesic triangle formed by o , x , and y .

The Gromov boundary at infinity of X will be denoted by ∂X . The Gromov product can be extended to all of $X \cup \partial X$ as follows. For $a, b \in X \cup \partial X$, define

$$(a | b) := \sup \liminf_{i,j \rightarrow \infty} (x_i | y_j)$$

where the supremum is taken over all sequences $\{x_i\}, \{y_j\} \subset X$ such that $x_i \rightarrow a$ and $y_j \rightarrow b$. For $a, b \in X$ this reduces to the previous definition. Using inequality (1), one can show that for any given sequences $x_i \rightarrow a, y_j \rightarrow b$,

$$(a | b) - 2\delta \leq \liminf_{i,j \rightarrow \infty} (x_i | y_j) \leq (a | b)$$

(see [GH, p. 122]). Therefore a sequence $\{z_i\} \subset X \cup \partial X$ converges to $a \in \partial X$ if and only if $(z_i | a) \rightarrow \infty$.

2.2. Barycenter. Consider \mathbb{H}^n with basepoint o . For each $\theta \in \partial \mathbb{H}^n$ let B_θ^o be the unique Busemann function of θ such that $B_\theta^o(o) = 0$. Namely, if γ is the geodesic ray based at o asymptotic to θ , then $B_\theta^o(p) = \lim_{t \rightarrow \infty} [d(p, \gamma(t)) - t]$. Let λ be a Radon measure on the compact space $\partial \mathbb{H}^n$. Define the average

$$B_\lambda(p) := \int_{\partial \mathbb{H}^n} B_\theta^o(p) d\lambda(\theta).$$

Proposition 2.1 ([BCG1, Appendix A]). *If λ has no atoms, then B_λ is proper and has a unique critical point in \mathbb{H}^n corresponding to the unique global minimum. Moreover, the Hessian of B_λ is a positive definite bilinear form. Namely, for all $v \in T_p \mathbb{H}^n$,*

$$(\text{Hess } B_\lambda)(v, v) := \langle \nabla_v \nabla B_\lambda, v \rangle > 0.$$

The unique critical point of B_λ is the *barycenter* of λ , denoted $\text{bar } \lambda$. Since the barycenter is the unique critical point of B_λ , it is defined implicitly as the unique point p such that

$$\int_{\partial \mathbb{H}^n} \langle \nabla B_\theta^o, v \rangle_p d\lambda(\theta) = 0$$

for all $v \in T_p \mathbb{H}^n$. Notice that the barycenter map is scale invariant, i.e. the barycenter of the measure λ equals the barycenter of the measure $c\lambda$ for any $c > 0$.

2.3. The brain in a jar lemma. At a crucial stage in the proof of Proposition 4.3, the following linear algebra lemma is needed.

Lemma 2.2. *Let H be an $n \times n$ positive definite symmetric matrix with trace $\text{tr}(H) = 1$. If $n \geq 3$, then*

$$\frac{\det(H)}{[\det(\text{Id} - H)]^2} \leq \left[\frac{n}{(n-1)^2} \right]^n.$$

Moreover, equality holds if and only if $H = \frac{1}{n} \text{Id}$.

This lemma is false for $n = 2$. It is the only part of this paper (and Besson–Courtois–Gallot theory in general) which fails in 2 dimensions.

2.4. Generalized differentiable and Riemannian structures ([OS]). Let X be a topological space, $\Omega \subseteq X$, and $n \in \mathbb{N}$. A family $\{(U_\phi, \phi)\}_{\phi \in \Phi}$ is called a \mathcal{C}^1 -atlas on $\Omega \subseteq X$ if the following hold:

- (1) For each $\phi \in \Phi$, U_ϕ is an open subset of X .
- (2) Each $\phi \in \Phi$ is a homeomorphism from U_ϕ into an open subset of \mathbb{R}^n .
- (3) $\{U_\phi\}_{\phi \in \Phi}$ is a covering of Ω .
- (4) If two maps $\phi, \psi \in \Phi$ satisfy $U_\phi \cap U_\psi \neq \emptyset$, then

$$\psi \circ \phi^{-1}: \phi(U_\phi \cap U_\psi) \rightarrow \psi(U_\phi \cap U_\psi)$$

is \mathcal{C}^1 on $\phi(U_\phi \cap U_\psi \cap \Omega)$.

A family $\{g_\phi\}_{\phi \in \Phi}$ is called a \mathcal{C}^0 -Riemannian metric associated with a \mathcal{C}^1 -atlas $\{(U_\phi, \phi)\}_{\phi \in \Phi}$ on $\Omega \subseteq X$ if the following hold:

- (1) For each $\phi \in \Phi$, g_ϕ is a map from U_ϕ to the set of positive symmetric matrices.
- (2) For each $\phi \in \Phi$, $g_\phi \circ \phi^{-1}$ is continuous on $\phi(U_\phi \cap \Omega)$.
- (3) For any $x \in U_\phi \cap U_\psi$, $\phi, \psi \in \Phi$, we have

$$g_\psi(x) = [d(\phi \circ \psi^{-1})(\psi(x))]^t g_\phi(x) [d(\phi \circ \psi^{-1})(\psi(x))].$$

These two structures induce a distance metric D_g on Ω which we now describe. The length of a piecewise \mathcal{C}^1 path $\gamma: (0, 1) \rightarrow \Omega$ is defined in the usual way by pulling back the metric tensor via γ . A path $\eta: [0, 1] \rightarrow X$ is *admissible* if $\eta^{-1}(X \setminus \Omega)$ is a finite set of points $\{t_1, \dots, t_l\}$, and η is piecewise \mathcal{C}^1 on $(0, 1) \setminus \{t_1, \dots, t_l\}$. For $x, y \in \Omega$, define

$$D_g(x, y) := \inf \{ \text{length}(\gamma) \mid \gamma \text{ joins } x \text{ to } y \text{ and is admissible} \}.$$

If x and y cannot be joined by an admissible path, set $D_g(x, y) = \infty$. In general, the topology of (Ω, D_g) can be quite different from the subspace topology of $\Omega \subseteq X$.

2.5. Almost everywhere Riemannian spaces. Let (X, d) be a geodesic metric space with Hausdorff dimension $n < \infty$. X is an *almost everywhere Riemannian* metric space if there exists $\Omega \subseteq X$, a dense subset of full n -dimensional Hausdorff measure, such that:

- (1) Ω admits a n -dimensional \mathcal{C}^1 -atlas $\{(U_\phi, \phi)\}_{\phi \in \Phi}$.
- (2) Ω admits a \mathcal{C}^0 -Riemannian metric $\{g_\phi\}_{\phi \in \Phi}$.

- (3) Each homeomorphism $\phi \in \Phi$ is in fact locally bilipschitz.
- (4) The identity map $(\Omega, D_g) \rightarrow (\Omega, d)$ is an isometry (see Section 2.4).
- (5) The Riemannian metric induces a volume element $d\text{vol}_\Omega$ on Ω . The measure on X obtained by integrating this element equals n -dimensional Hausdorff measure on X ($X \setminus \Omega$ has zero measure).

Notice X is not assumed to be a topological manifold, and $\Omega \subseteq X$ is not assumed to be open. Conditions (1)–(5) are not as restrictive as they initially appear. For example, Otsu and Shioya proved that a finite dimensional Alexandrov space with curvature bounded below by $k \in \mathbb{R}$ is almost everywhere Riemannian [OS] (see Section 2.13). Condition (3) allows Rademacher's theorem to be applied to almost everywhere Riemannian spaces. This means locally Lipschitz functions on an almost everywhere Riemannian metric space are differentiable almost everywhere. Condition (4) says that the \mathcal{C}^0 -Riemannian metric on Ω reproduces the metric d , in the sense that the metric completion of (Ω, D_g) is isometric to (X, d) .

2.6. Cone-manifolds ([CHK, p. 53]). An n -dimensional cone-manifold M is a manifold which can be triangulated so that the link of each simplex is piecewise linear homeomorphic to a standard sphere and M is equipped with a path metric such that the restriction of the metric to each simplex is isometric to a geodesic simplex of constant curvature K . The singular locus Σ consists of the points with no neighborhood isometric to a ball in a Riemannian manifold.

It follows that

- Σ is a union of totally geodesic closed simplices of dimension $n - 2$.
- At each point of Σ in an open $(n - 2)$ -simplex, there is a *cone angle* which is the sum of dihedral angles of n -simplices containing the point.

In particular, the singular locus of a 3-dimensional cone-manifold forms a graph in the manifold. A cone-manifold is an almost everywhere Riemannian metric space (see Lemma 2.3). Though a definition will not be given here, a cubed-manifold is another example of a cone-manifold (see [AMR]). More abstractly, any manifold admitting a locally finite decomposition into convex geodesic polyhedra is a cone-manifold. This can be seen by adding superfluous faces to the polyhedral decomposition.

The manifold structure of a cone-manifold will not be used in this paper. Theorem 8.6 is equally valid for more general simplicial metric spaces not satisfying the above link condition.

2.7. Convex Riemannian manifolds with boundary. A geodesic metric space C is an n -dimensional *convex Riemannian* (resp. *hyperbolic*) *manifold with boundary* if

- (1) C is topologically an n -manifold with boundary,
- (2) there is an incomplete Riemannian (resp. hyperbolic) metric on the interior of C ,

- (3) the metric on C is the metric completion of the Riemannian (resp. hyperbolic) metric on the interior,
- (4) for any pair of points in the interior of C , the shortest path between them lies in the interior of C , and
- (5) for any compact $K \subseteq C$, the curvature of the Riemannian manifold $K \cap \text{int}(C)$ is bounded from above and below by finite constants.

(Notice no differentiability assumptions have been made on the boundary.)

Property (5) ensures C has “locally bounded geometry”. The lower curvature bound on compact sets guarantees that C is locally an Alexandrov space with curvature bounded below. This implies the boundary ∂C has Hausdorff dimension $n - 1$ [OS]. (In [OS], they prove the singular set of an n -dimensional Alexandrov space with curvature bounded below has Hausdorff dimension $\leq n - 1$. In this case, the boundary *is* the singular set.) The local upper curvature bound gives a *lower* bound on the volume of small metric balls in C . This useful property will be used in Section 7.

Given a convex Riemannian manifold with boundary C , we can metrically double it across its boundary to obtain a metric space DC . Topologically DC is the closed manifold obtained by doubling C across its boundary. The metric on DC is the path metric induced by gluing the two copies of C (one with opposite orientation) along ∂C . The convexity of C (property (4)) insures that the path metric obtained after doubling does not alter the original metric on C . Notice that DC is an almost everywhere Riemannian metric space (see Lemma 2.3).

2.8. Convex Riemannian amalgams. A geodesic metric space Z is an n -dimensional *convex Riemannian amalgam* if Z contains an isometrically embedded locally finite countable collection $\{C_j\} \subseteq Z$ of n -dimensional convex Riemannian manifolds with boundary such that

- (1) $\bigcup_j C_j = Z$,
- (2) $\text{int}(C_j) \cap \text{int}(C_k) = \emptyset$ for $j \neq k$.

(Notice Z is not assumed to be a manifold.) A cone-manifold is a convex Riemannian amalgam (see Section 2.6). Another convex Riemannian amalgam is the metric doubling DC of a convex Riemannian manifold with boundary C (see Section 2.7).

Lemma 2.3. *A convex Riemannian amalgam is an almost everywhere Riemannian metric space.*

Proof. Define $\Omega := \bigcup_j \text{int}(C_j)$. We must check conditions (1)–(5) of Section 2.5. (1)–(3) are trivial. For (4) use the following consequence of convexity: any $x, y \in Z$ can be joined by a path γ such that

- the length of γ is arbitrarily close to $d(x, y)$, and
- $\gamma \cap \partial C_j$ is at most two points for any C_j in the decomposition.

For (5) use that $\bigcup_j \partial C_j$ has measure zero. \square

Convex Riemannian amalgams seem to be the most natural class of metric spaces for which the arguments of Theorem 7.1 are valid.

2.9. Convex cores. Let M be a hyperbolic manifold. Let $S \subseteq M$ be the union of all closed geodesics in M . The *convex core*, C_M , is the smallest closed convex subset of M which contains S , in other words it is the closed convex hull of S in M . The convex core may also be defined as the smallest closed convex subset of M such that the inclusion map is a homotopy equivalence. M is *geometrically finite* if an ε -neighborhood of C_M has finite volume. Otherwise, M is *geometrically infinite*.

For finite volume hyperbolic manifolds, the convex core is the entire manifold. Thus this is a useful object only in the infinite volume case, where C_M is the smallest submanifold which carries all the geometry of M .

2.10. Pared 3-manifolds [M, Def.4.8]. Let N be a compact orientable irreducible 3-manifold with nonempty boundary. Assume N is not a 3-ball. Let $P \subseteq \partial N$. (N, P) is a *pared 3-manifold* if the following three conditions hold.

- (1) Every component of P is an incompressible torus or a compact annulus.
- (2) Every noncyclic abelian subgroup of $\pi_1(N)$ is conjugate into the fundamental group of a component of P .
- (3) Every π_1 -injective cylinder $C: (S^1 \times I, S^1 \times \partial I) \rightarrow (N, P)$ is relatively homotopic to a map ψ such that $\psi(S^1 \times I) \subseteq P$.

By Thurston's Geometrization Theorem [M], (N, P) is a pared 3-manifold if and only if there exists a geometrically finite hyperbolic structure on the interior of N such that $C_M \cong N \setminus P$.

2.11. Pared acylindrical ([Th3, p. 244]). A pared 3-manifold (N, P) is *pared acylindrical* if $\partial N \setminus P$ is incompressible and if every π_1 -injective cylinder

$$C: (S^1 \times I, S^1 \times \partial I) \rightarrow (N, \partial N \setminus P)$$

is homotopic rel boundary to ∂N .

2.12. Deformation theory. Let $\mu > 0$ be the Margulis constant for hyperbolic 3-manifolds. Then for a hyperbolic 3-manifold M , the μ -thin part of M is a disjoint union of bounded Margulis tubes and unbounded cusps [BP]. After possibly making μ smaller, we may also assume that the intersection of ∂C_M and the μ -thin part of M is totally geodesic [M, Lemma 6.9]. Define M° to be M minus the unbounded components of its μ -thin part. In other words, M° is the manifold with boundary obtained by removing the cusps from M .

Let (N, P) be a compact pared 3-manifold. Define the deformation space $H(N, P)$ as follows. For a hyperbolic 3-manifold M and a map $m: N \rightarrow M^o$, $(M, m) \in H(N, P)$ if there exists a union Q_M of components of ∂M^o such that $m: (N, P) \rightarrow (M, Q_M)$ is a relative homotopy equivalence. $(M_1, m_1) = (M_2, m_2)$ in $H(N, P)$ if there exists an orientation preserving isometry $\mathcal{I}: M_1 \rightarrow M_2$ such that $\mathcal{I} \circ m_1 \sim m_2$. ($H(N, P)$ admits several interesting topologies [Th3]. They will not be needed here.) (M, m) has *no additional parabolics* if $Q_M = \partial M^o$. Using the product structure on the complement of C_M [EM] and the thick-thin decomposition, there exists a relative homotopy equivalence $p: (M, Q_M) \rightarrow (C_M \cap M^o, C_M \cap Q_M)$.

Theorem 2.4 (Johannson). *Let (N, P) be a compact pared acylindrical 3-manifold. If $(M, m) \in H(N, P)$ is geometrically finite then $p \circ m$ is homotopic to a homeomorphism $(N, P) \rightarrow (C_M \cap M^o, C_M \cap Q_M)$.*

Proof. Since N is not homotopy equivalent to a surface, C_M is a 3-manifold. Since M is geometrically finite, $(C_M \cap M^o, C_M \cap Q_M)$ is a compact pared 3-manifold [M, Corollary 6.10]. So $p \circ m$ is a relative homotopy equivalence between compact pared 3-manifolds, and the domain is pared acylindrical. By the work of Johannson [J, Lemma X.23, p. 235], $p \circ m$ is homotopic (rel the paring) to a homeomorphism $(N, P) \rightarrow (C_M \cap M^o, C_M \cap Q_M)$. \square

We are interested in pared acylindrical 3-manifolds because of the following corollary of Thurston's Geometrization Theorem and Mostow rigidity.

Corollary 2.5 ([Th2, p. 14]). *Let (N, P) be a pared acylindrical 3-manifold. Then there exist exactly two spaces $(M, m), (M^{\text{opp}}, m^{\text{opp}}) \in H(N, P)$ such that M and M^{opp} are geometrically finite, M and M^{opp} have no additional parabolics, and the convex cores C_M and $C_{M^{\text{opp}}}$ have totally geodesic boundary. Moreover, there exists an orientation reversing isometry $\mathcal{I}: M \rightarrow M^{\text{opp}}$ such that $m^{\text{opp}} \sim \mathcal{I} \circ m$.*

2.13. Alexandrov space with curvature bounded below by -1 . There are many equivalent definitions of Alexandrov spaces. Here we give the most common definition. (See [BBI] for more information.)

Let Y be a path metric space. Y is an *Alexandrov space with curvature bounded below by -1* if about each point in Y there exists a neighborhood U satisfying the following comparison condition. Let $x, y, z \in U$ be distinct points, let w lie on the interior of a geodesic path xy connecting x to y . Let $\tilde{x}, \tilde{y}, \tilde{z}, \tilde{w} \in \mathbb{H}^2$ be such that $d(x, y) = d(\tilde{x}, \tilde{y})$, $d(y, z) = d(\tilde{y}, \tilde{z})$, $d(x, z) = d(\tilde{x}, \tilde{z})$, $d(x, w) = d(\tilde{x}, \tilde{w})$, $d(w, y) = d(\tilde{w}, \tilde{y})$. We then require that $d(z, w) \geq d(\tilde{z}, \tilde{w})$.

This comparison condition guarantees that geodesic triangles in Y are at least as fat as hyperbolic triangles. The *dimension* of an Alexandrov space with curvature bounded below is defined to be its Hausdorff dimension. For finite dimensional




Figure 2

Alexandrov spaces with curvature bounded below, the Hausdorff and topological dimensions agree [BGP, p. 21].

In Section 8.3, we will use

Theorem 2.6 ([OS]). *If Y is a finite dimensional Alexandrov space with curvature bounded below by -1 , then Y is an almost everywhere Riemannian metric space.*

Proof. Following [OS], let S be the singular set of Y . Define $\Omega := Y \setminus S$. Ω is a countable intersection of dense open sets with full measure. It is therefore dense and has full measure. Conditions (1)–(5) of Section 2.5 also follow from results in [OS]. Specifically, conditions (1) and (2) follow from [Theorem B, p. 630], condition (3) follows from [Lemma 5.1.3, p. 651], condition (4) follows from [Theorem B, p. 630] and [Theorem 6.4, p. 654], and condition (5) follows from [Section 7.1]. \square

The main property of these Alexandrov spaces we will use is an upper bound on their volume growth entropy, which we now define.

2.14. Volume growth entropy. Let X be a geodesic metric space of Hausdorff dimension n , \tilde{X} be the universal cover of X , and \mathcal{H}^n be n -dimensional Hausdorff measure. The *volume growth entropy* of \tilde{X} is the number

$$h(\tilde{X}) := \limsup_{R \rightarrow \infty} \frac{1}{R} \log \mathcal{H}^n(B_{\tilde{X}}(x, R)),$$

where x is any point in \tilde{X} , and the ball $B_{\tilde{X}}(x, R)$ is in \tilde{X} .

The volume growth entropy is independent of the choice of $x \in \tilde{X}$.

The following theorem of Burago, Gromov, and Perelman will be important for this paper.

Theorem 2.7 ([BGP, p. 40]). *If X is an Alexandrov space with curvature bounded below by -1 and Hausdorff dimension n , then the volume growth entropy of \tilde{X} is less than or equal to the volume growth entropy of \mathbb{H}^n . In other words*

$$h(\tilde{X}) \leq h(\mathbb{H}^n) = n - 1.$$

3. Horoboundary and densities

In order to define a density in the necessary generality, a second definition of the boundary at infinity must be made. This is a more general definition than of the Gromov boundary at infinity; it makes sense for any proper metric space. In the case of negatively curved Riemannian manifolds, it reduces to the Gromov boundary. (For more information, see [BGS, p. 21], [BM], and [F, Section 2].)

Let Y be a proper metric space. For $p \in Y$, define $d_p(y) := d(p, y)$. Denote the space of continuous (real valued) functions on Y by $C(Y)$, and endow this set with the topology of uniform convergence on compact sets. Define an equivalence relation on $C(Y)$ by $f \sim g$ if and only if $f - g$ is a constant function. Denote the quotient space $C(Y)/\sim$ by $C_*(Y)$. $C_*(Y)$ is Hausdorff. Define a map $\iota: Y \rightarrow C_*(Y)$ by $\iota(p) := [d_p]$. ι is a topological embedding.

Definition 3.1. Let $\text{Cl}(Y)$ denote the closure of $\iota(Y)$ in $C_*(Y)$. The *horoboundary* of Y is

$$\mathcal{H}Y := \text{Cl}(Y) \setminus \iota(Y).$$

A continuous function $h \in C(Y)$ such that $[h] \in \mathcal{H}Y$ is a *horofunction* of Y .

For $\eta \in C_*(Y)$ define a function $b_\eta: Y \times Y \rightarrow \mathbb{R}$ by

$$b_\eta(p, q) := h(p) - h(q) \text{ for any } h \in C(Y) \text{ such that } [h] = \eta.$$

If Y has a fixed basepoint o , then define $b_\eta(p) := b_\eta(p, o)$.

It is a quick check to see that b_η is well-defined, i.e. independent of the choice of h . The functions b_η are 1-Lipschitz. Thus applying the Arzela–Ascoli theorem shows $\text{Cl}(Y)$ is compact, implying $\mathcal{H}Y$ is compact. (If Y is nonpositively curved, then horofunctions and the horoboundary are identical to Busemann functions and the boundary at infinity [BGS, p. 22].)

Isometries of Y extend to homeomorphisms of $\mathcal{H}Y$ in the following simple manner. Consider the $\text{Isom}(Y)$ -action by homeomorphisms on $C(Y)$ given by $\phi.f := f \circ \phi^{-1}$. This action descends to an action on $C_*(Y)$. Since $\phi.[d_p] = [d_{\phi.p}]$, the map $\phi|_{\iota(Y)}: \iota(Y) \rightarrow \iota(Y)$ is a homeomorphism. Since ϕ acts as a homeomorphism on both $\iota(Y)$ and $C_*(Y)$, it also acts as a homeomorphism on $\mathcal{H}Y$. Thus we have defined an $\text{Isom}(Y)$ -action by homeomorphisms on $\mathcal{H}Y$.

Definition 3.2. Let Y be proper metric space. Let G be a closed subgroup of $\text{Isom}(Y)$. A continuous map (under the weak-* topology on measures)

$$\mu: Y \rightarrow \{\text{positive Radon measures on } \mathcal{H}Y\}$$

is an ℓ -dimensional density for G if

- (1) μ is G -equivariant, i.e. $\mu_{g.x} = g_*\mu_x$,
- (2) $\mu_p \ll \mu_q$ for all p, q , and for $\eta \in \mathcal{H}Y$,

$$\frac{d\mu_p}{d\mu_q}(\eta) = e^{-\ell(b_\eta(p,q))}.$$

Example. For \mathbb{H}^n , $\mathcal{H}\mathbb{H}^n = \partial\mathbb{H}^n$. Let $\Phi: S_p\mathbb{H}^n \rightarrow \partial\mathbb{H}^n$ be the standard radial homeomorphism between the unit tangent sphere and the boundary at infinity. Define μ_p to be the push-forward of Lebesgue measure on S_pY by Φ . This is known as the *visual measure at p* , and is an $(n-1)$ -dimensional density for all of $\text{Isom}(\mathbb{H}^n)$. Visual measure is the most natural density in this case.

Our entire reason for defining the horoboundary is the following proposition.

Proposition 3.1 ([BM, Prop. 1.1]). *Let X be a proper metric space of Hausdorff dimension n with basepoint $o \in X$. Let m be n -dimensional Hausdorff measure. Assume $m(X) = \infty$ and X has finite volume growth entropy $h(X)$. Then there exists an $h(X)$ -dimensional density $x \mapsto \mu_x$ for $\text{Isom}(X)$. This density is called Patterson–Sullivan measure.*

By normalizing we may always assume that μ_o is a probability measure. Notice this normalization implies that $\mu_{g.o} = g_*\mu_o$ is also a probability measure for any $g \in \text{Isom}(X)$.

Let X be a δ -hyperbolic proper path metric space. Assume that $h(X) \in (0, \infty)$. We have defined two different compactifications of X ; the Gromov boundary at infinity ∂X and the horoboundary $\mathcal{H}X$. In general, those two compactifications are not homeomorphic. However, they are both necessary for the work of this paper. To connect the two compactifications, we now define a continuous $\text{Isom}(X)$ -equivariant surjection $\pi: \mathcal{H}X \rightarrow \partial X$.

Fix a basepoint $o \in X$. Pick $\xi \in \mathcal{H}X$ and a sequence $\{p_n\}$ such that $[d_{p_n}] \rightarrow \xi$ in $C_*(X)$. Then $\{p_n\}$ leaves every compact set of X . So by the Arzela–Ascoli theorem there exists a subsequence $\{a_l\} \subseteq \{p_n\}$ such that the geodesic segments oa_l converge to some $[\gamma] \in \partial X$ in the compact-open topology, where γ is a geodesic ray based at o .

Define $\pi: \mathcal{H}X \rightarrow \partial X$ by $\pi(\xi) = [\gamma]$. Before proving π is well defined, we need the following lemma.

Lemma 3.2. *In the above notation, the sequence $\{a_l\}$ converges to $[\gamma] \in \partial X$.*

Proof. It is enough to show $([\gamma] | a_l) \rightarrow \infty$. Pick a large $M \gg 0$, and consider the metric ball $B(o, M)$. Find L such that for $l \geq L$, the geodesic segment oa_l is 1-close to γ on $B(o, M)$. Then the picture looks roughly like Figure 3.




Figure 3

As $M \rightarrow \infty$, intuitively $([\gamma] | a_l) \rightarrow \infty$. To prove this carefully, notice that for $k > 1$

$$d(\gamma(kM), a_l) \leq [d(a_l, o) - M] + [d(\gamma(kM), o) - M] + 1.$$

This implies

$$(\gamma(kM) | a_l) \geq M - 1/2.$$

So finally

$$([\gamma] | a_l) \geq \liminf_{k \rightarrow \infty} (\gamma(kM) | a_l) - 2\delta \geq M - 2\delta - 1/2. \quad \square$$

Lemma 3.3. *The map $\pi : \mathcal{H}X \rightarrow \partial X$ is well defined.*

Proof. Suppose π is not well defined. Then there exist sequences $p_i, q_i \rightarrow \xi \in \mathcal{H}X$ such that $\{p_i\}, \{q_i\}$ do not converge to a common point in ∂X . Thus there exists an $M > 0$ such that $(p_i | q_i) < M$ for all i .

Pick a metric closed ball K much larger than $B(o, M)$. $p_i, q_i \rightarrow \xi \in \mathcal{H}X$ implies that for large i and all $x \in K$,

$$d(p_i, x) - d(p_i, o) \approx d(q_i, x) - d(q_i, o). \quad (2)$$

This will lead to a contradiction.

To begin, pick $x \in K \cap oq_i$ such that $d(x, o) \gg M$. Then

$$d(q_i, x) - d(q_i, o) = -d(x, o). \quad (3)$$




Figure 4

Approximate the four point metric space $\{x, o, q_i, p_i\}$ by a tree. Let the corresponding four points in the tree be $\{\tilde{x}, \tilde{o}, \tilde{q}_i, \tilde{p}_i\}$. We may assume this tree is at worst $(1, 2\delta)$ -quasi-isometric to $\{x, o, q_i, p_i\}$ [GH, p. 33] (see Figure 4). We thus obtain the inequalities

$$\begin{aligned}
 d(p_i, x) - d(p_i, o) &\geq d(\tilde{p}_i, \tilde{x}) - d(\tilde{p}_i, \tilde{o}) - 4\delta \\
 &= d(\tilde{x}, \tilde{o}) - 2(\tilde{x} \mid \tilde{p}_i) - 4\delta \\
 &= d(\tilde{x}, \tilde{o}) - 2(\tilde{q}_i \mid \tilde{p}_i) - 4\delta \\
 &\geq d(x, o) - 2\delta - 2(q_i \mid p_i) - 6\delta - 4\delta \\
 &\geq d(x, o) - 2M - 12\delta > \frac{d(x, o)}{2}.
 \end{aligned}$$

Together with equation (3), this contradicts equation (2). \square

Lemma 3.4. π is continuous and surjective.

Proof. To see that π is surjective, pick a geodesic ray γ based at o . By compactness there is a sequence $\{t_i\} \subset (0, \infty)$ such that $\gamma(t_i)$ converges to a point $\xi \in \mathcal{H}X$. Then by definition $\pi(\xi) = [\gamma]$.

Let $\xi^n \rightarrow \xi$ in $\mathcal{H}X$. Pick sequences $\{p_i^n\}$ such that for all n , $\lim_{i \rightarrow \infty} p_i^n = \xi^n$. This implies $\lim_{i \rightarrow \infty} p_i^n = \pi(\xi^n)$ in ∂X for all n . There is an increasing sequence i_1, i_2, i_3, \dots of natural numbers such that

$$\lim_{n \rightarrow \infty} p_{i_n}^n = \xi \text{ in } \mathcal{H}X, \quad \text{and} \quad (\pi(\xi^n) \mid p_{i_n}^n) > n.$$

This implies $\lim_{n \rightarrow \infty} p_{i_n}^n = \pi(\xi)$ in ∂X . By the definition of δ -hyperbolicity

$$(\pi(\xi) | \pi(\xi^n)) \geq [\min\{(\pi(\xi) | p_{i_n}^n), (\pi(\xi^n) | p_{i_n}^n)\} - 3\delta] \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

Therefore $\pi(\xi^n) \rightarrow \pi(\xi)$ in ∂X . \square

The proof of the following lemma is trivial and has been omitted.

Lemma 3.5. *For all $g \in \text{Isom}(X)$, $g \circ \pi = \pi \circ g$.*

Lemma 3.6. *If X admits a cocompact isometric action, then $\pi_*\mu_p$ has no atoms for all $p \in X$.*

Proof. As $\pi_*\mu_p \ll \pi_*\mu_o$, it is sufficient to show $\pi_*\mu_o$ has no atoms. Suppose there exists $\alpha \in \partial X$ such that $\pi_*\mu_o(\alpha) > 0$. Since X admits a cocompact isometric action, there is a constant $D > 0$ such that any $p \in X$ is at most a distance D from an orbit point $g_p.o$ for $g_p \in \text{Isom}(X)$. It follows that the total mass of the measure μ_p is at most $e^{Dh(X)}$.

We first show all the horofunctions in the fiber $\pi^{-1}(\alpha)$ are a bounded distance from each other. Pick a geodesic ray in X based at o asymptotic to α . Pick a sequence p_i of points going to infinity on the geodesic ray. After passing to a subsequence we may assume the points p_i converge to some $\xi \in \pi^{-1}(\alpha) \subset \mathcal{H}X$.

By definition

$$b_\xi(x) = \lim (-2(p_i | x) + d(x, o)).$$

We know

$$(\alpha | x) - 2\delta \leq \liminf (p_i | x) \leq (\alpha | x).$$

Because b_ξ is well defined, the \liminf can be replaced by a limit. Thus

$$-2(\alpha | x) + d(x, o) \leq b_\xi(x) \leq -2(\alpha | x) + d(x, o) + 4\delta.$$

So for any other $\zeta \in \pi^{-1}(\alpha)$ we have

$$|b_\xi(x) - b_\zeta(x)| \leq 4\delta.$$

Using this we obtain the inequality

$$\begin{aligned} \pi_*\mu_{p_i}(\alpha) &= \int_{\pi^{-1}(\alpha)} d\mu_{p_i} = \int_{\pi^{-1}(\alpha)} e^{-h(X)b_\zeta(p_i)} d\mu_o(\zeta) \\ &\geq \int_{\pi^{-1}(\alpha)} e^{-h(X)[b_\xi(p_i)+4\delta]} d\mu_o(\zeta) \\ &= e^{-h(X)[b_\xi(p_i)+4\delta]} \cdot \pi_*\mu_o(\alpha) > 0. \end{aligned}$$

But $\lim_{i \rightarrow \infty} b_\xi(p_i) = -\infty$. So

$$\lim_{i \rightarrow \infty} \pi_* \mu_{p_i}(\alpha) = \infty.$$

Since $\pi_* \mu_{p_i}$ has total mass at most $e^{Dh(X)}$, this is a contradiction. Therefore $\pi_* \mu_o$ has no atoms. \square

4. The Besson–Courtois–Gallot inequality

Theorem 4.1. *Let M_{hyp} be a closed hyperbolic n -manifold for $n \geq 3$. Let Z be a compact n -dimensional almost everywhere Riemannian metric space with universal cover X . (For a definition, see Section 2.5.) Let $f: Z \rightarrow M_{\text{hyp}}$ be a homotopy equivalence. Then*

$$h(X)^n \text{Vol}(Z) \geq (n-1)^n \text{Vol}(M_{\text{hyp}}).$$

Let f also denote the lifted map $f: X \rightarrow \mathbb{H}^n$. Recall that if Z and M_{hyp} are compact, then a homotopy equivalence $f: Z \rightarrow M_{\text{hyp}}$ lifts to a quasi-isometry $f: X \rightarrow \widetilde{M_{\text{hyp}}}$ between the universal covers. This implies that the volume growth entropy of X is a strictly positive and finite (see for example [Gr, Prop.5.10]). Fix a basepoint $o \in X$, and let $f(o) \in \mathbb{H}^n$ be a basepoint of \mathbb{H}^n . Since $f: X \rightarrow \mathbb{H}^n$ is a quasi-isometry, f extends to a homeomorphism between the Gromov boundaries $f: \partial X \rightarrow \partial \mathbb{H}^n$.

We now define the natural map $F: X \rightarrow \mathbb{H}^n$, but postpone proving its regularity properties until later sections. For $x \in X$, let μ_x be the Patterson–Sullivan measure at x . Recall we have defined a continuous map $\pi: \mathcal{H}X \rightarrow \partial X$. Push forward the Patterson–Sullivan measure μ_x on $\mathcal{H}X$ to a probability measure $(f \circ \pi)_* \mu_x$ on $\partial \mathbb{H}^n$. Define $F: X \rightarrow \mathbb{H}^n$ by

$$F(x) := \text{barycenter}((f \circ \pi)_* \mu_x).$$

F is the natural map induced by f . F is a $\Gamma := \pi_1(Z) \cong \pi_1(M_{\text{hyp}})$ -equivariant continuous map. It therefore descends to a continuous map $F: Z \rightarrow M_{\text{hyp}}$.

Remark 4.2. Any additional symmetries of the map $f: Z \rightarrow M_{\text{hyp}}$ also become symmetries of F . Namely, if Z and M_{hyp} possess an isometric involution, and $f: Z \rightarrow M_{\text{hyp}}$ is equivariant with respect to the involutions, then $F: Z \rightarrow M_{\text{hyp}}$ is similarly equivariant. This follows immediately from the definition, because both $\pi: \mathcal{H}X \rightarrow \partial X$ and Patterson–Sullivan measure are $\text{Isom}(X)$ -equivariant. This fact will be used in the proofs of Theorems 8.1 and 8.9.

$F, f: X \rightarrow \mathbb{H}^n$ are Γ -equivariant maps. The straight-line homotopy between them is also Γ -equivariant. Therefore the downstairs maps $F, f: Z \rightarrow M_{\text{hyp}}$ are homotopic. This shows $F: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence. As M_{hyp} is a closed manifold, F must be surjective.

Proving the local regularity properties of the natural map requires some work. These properties are summarized in

Proposition 4.3. *The natural map $F: Z \rightarrow M_{\text{hyp}}$ has the following properties:*

- (1) *It is locally Lipschitz and differentiable almost everywhere.*
- (2) $|\text{Jac } F(p)| \leq \left(\frac{h(X)}{n-1}\right)^n$ *almost everywhere.*
- (3) *If for some p , $|\text{Jac } F(p)| = \left(\frac{h(X)}{n-1}\right)^n$, then the differential dF_p is a homothety of ratio $\left(\frac{h(X)}{n-1}\right)^n$.*

This proposition will be proven in later sections. Specifically, (1) will be proven in Section 5, (2) and (3) will be proven in Section 6. Let us now temporarily assume it, and complete the proof of Theorem 4.1. (In this section, we use only (1) and (2). (3) will not be used until Section 7.)

By assumption, Z is an almost everywhere Riemannian metric space. So by definition Z has a subset Ω of full measure admitting a \mathcal{C}^1 -atlas $\{U_\phi, \phi\}_{\phi \in \Phi}$ and a \mathcal{C}^0 -Riemannian metric $\{g_\phi\}_{\phi \in \Phi}$. This Riemannian metric induces a volume element ω_Z which agrees with n -dimensional Hausdorff measure on Z . Let $\omega_{M_{\text{hyp}}}$ be the volume element on M_{hyp} .

Lemma 4.4.

$$\text{Vol}(M_{\text{hyp}}) = \text{Vol}(F(Z)) = \int_{F(Z)} \omega_{M_{\text{hyp}}} \leq \int_Z |\text{Jac } F| \omega_Z \leq \left(\frac{h(X)}{n-1}\right)^n \text{Vol}(Z).$$

Proof. Assuming Proposition 4.3, the only non-trivial part is to prove

$$\int_{F(Z)} \omega_{M_{\text{hyp}}} \leq \int_Z |\text{Jac } F| \omega_Z.$$

This amounts to justifying the change of variables formula for the singular space Z . To do this, we will unpack the definitions and apply the change of variables formula for Lipschitz maps.

$Z \setminus \Omega$ is measure zero, and F is locally Lipschitz. This implies $F(Z \setminus \Omega) \subset M_{\text{hyp}}$ is also measure zero. The collection of open sets $\{U_\phi\}_{\phi \in \Phi}$ covers Ω . Let $\{E_k\}$ be a countable partition of $\cup_\phi U_\phi \subset Z$ into measurable sets such that each E_k is contained in some open set U_ϕ . Since $F(\cup_k E_k) = F(\cup_\phi U_\phi) \subset M_{\text{hyp}}$ is of full measure, it

suffices to prove the inequality on each measurable set E_k , i.e. it suffices to show that

$$\int_{F(E_k)} \omega_{M_{\text{hyp}}} \leq \int_{E_k} |\text{Jac } F| \omega_Z.$$

Assume that $E_k \subset U_\phi$. Let us also assume without a loss of generality that the image $F(U_\phi) \subset M_{\text{hyp}}$ lies in an open set V_M equipped with a smooth diffeomorphism ψ onto an open subset of \mathbb{R}^n . Let g_M denote the smooth Riemannian metric on $\psi(V_M)$ given by the hyperbolic metric on M_{hyp} .

The volume element $(\phi^{-1})^*(\omega_Z|_{U_\phi})$ is defined on an open subset of \mathbb{R}^n . It is induced by the \mathcal{C}^0 -Riemannian metric g_ϕ . Concretely this means that

$$(\phi^{-1})^*(\omega_Z|_{U_\phi}) = \sqrt{\det g_\phi} dx_1 dx_2 \dots dx_n.$$

Similarly,

$$(\psi^{-1})^*\omega_{M_{\text{hyp}}} = \sqrt{\det g_M} dy_1 dy_2 \dots dy_n.$$

Define the locally Lipschitz map $G := \psi \circ F \circ \phi^{-1}$, which is a map between subsets of Euclidean space. By definition, the Jacobian of $F : Z \rightarrow M_{\text{hyp}}$ at $p \in U_\phi$ is

$$|\text{Jac } F(p)| := \frac{\sqrt{\det g_M(\psi(F(p)))}}{\sqrt{\det g_\phi(\phi(p))}} \cdot |\text{Jac } G(\phi(p))|.$$

Applying the change of variables formula for Lipschitz maps [EG, 3.4.3] to G yields

$$\begin{aligned} & \int_{\phi(U_\phi)} \chi_{\phi(E_k)}(x) \sqrt{\det g_M(G(x))} |\text{Jac } G(x)| dx_1 dx_2 \dots dx_n \\ &= \int_{\psi(V_M)} \#\{G^{-1}(y) \cap \phi(E_k)\} \sqrt{\det g_M(y)} dy_1 dy_2 \dots dy_n. \end{aligned}$$

Since $\det g_\phi$ vanishes on a set of measure zero, we may perform the following step

$$\begin{aligned} & \int_{\phi(U_\phi)} \chi_{\phi(E_k)}(x) \sqrt{\det g_M(G(x))} |\text{Jac } G(x)| dx_1 dx_2 \dots dx_n \\ &= \int_{\phi(U_\phi)} \chi_{\phi(E_k)}(x) \frac{\sqrt{\det g_M(G(x))}}{\sqrt{\det g_\phi(x)}} \sqrt{\det g_\phi(x)} |\text{Jac } G(x)| dx_1 dx_2 \dots dx_n \\ &= \int_{\phi(U_\phi)} \chi_{\phi(E_k)}(x) |\text{Jac } F(\phi^{-1}(x))| \sqrt{\det g_\phi(x)} dx_1 dx_2 \dots dx_n. \end{aligned}$$

Putting this together yields

$$\begin{aligned}
 \int_{E_k} |\text{Jac } F(z)| \omega_Z &= \int_{\phi(U_\phi)} \chi_{\phi(E_k)}(x) |\text{Jac } F(\phi^{-1}(x))| \sqrt{\det g_\phi(x)} dx_1 dx_2 \dots dx_n \\
 &= \int_{\psi(V_M)} \#\{G^{-1}(y) \cap \phi(E_k)\} \sqrt{\det g_M(y)} dy_1 dy_2 \dots dy_n \\
 &= \int_{V_M} \#\{F^{-1}(m) \cap E_k\} \omega_{M_{\text{hyp}}}(m) \geq \int_{F(E_k)} \omega_{M_{\text{hyp}}}. \quad \square
 \end{aligned}$$

The above lemma implies that

$$h(X)^n \text{Vol}(Z) \geq (n-1)^n \text{Vol}(M_{\text{hyp}}).$$

This proves the inequality of Theorem 4.1. \square

5. The Barycenter map is locally Lipschitz

In this section we prove part (1) of Proposition 4.3.

- (1) The natural map $F: Z \rightarrow M_{\text{hyp}}$ is locally Lipschitz and differentiable almost everywhere.

It is only necessary to prove F is locally Lipschitz. Almost everywhere differentiability will then follow by using Rademacher's theorem (see Section 2.5). We will prove the lifted map $F: X \rightarrow \mathbb{H}^n$ is locally Lipschitz by factoring it as a composition of two locally Lipschitz maps. Namely we will define a locally Lipschitz map $\Phi: X \rightarrow L^2(\mathcal{H}X)$ and a \mathcal{C}^1 -map $P: L^2(\mathcal{H}X) \rightarrow \mathbb{H}^n$ such that $F = P \circ \Phi$.

The barycenter map takes a positive atomless measure ν on $\partial\mathbb{H}^n$ to the unique point $x = \text{bar}(\nu)$ defined implicitly by the equation

$$\int_{\partial\mathbb{H}^n} \langle \nabla B_\theta^o, v \rangle_x d\nu(\theta) = 0 \quad \text{for all } v \in T_x \mathbb{H}^n,$$

where B^o is the Busemann function on \mathbb{H}^n (normalized so $B^o(o, \theta) = 0$ for all $\theta \in \partial\mathbb{H}^n$).

Consider the Hilbert space $L^2(\mathcal{H}X)$ of square integrable functions on $\mathcal{H}X$ with respect to the Patterson–Sullivan probability measure μ_o . Define a $\Gamma := \pi_1(Z)$ -action on $L^2(\mathcal{H}X)$ by

$$(\gamma.\phi)(\eta) := \phi(\gamma^{-1}.\eta) \cdot \sqrt{\exp(-h(X)b_\eta(\gamma.o))}.$$

Lemma 5.1. Γ acts by isometries on $L^2(\mathcal{H}X)$.

Proof.

$$\begin{aligned}
 \int_{\mathcal{H}X} (\gamma.\phi)^2(\eta) d\mu_o(\eta) &= \int_{\mathcal{H}X} \phi^2(\gamma^{-1}.\eta) e^{-h(X)b_\eta(\gamma.o)} d\mu_o(\eta) \\
 &= \int_{\mathcal{H}X} \phi^2(\gamma^{-1}.\eta) d\mu_{\gamma.o}(\eta) \\
 &= \int_{\mathcal{H}X} \phi^2(\eta) d(\gamma_*^{-1}\mu_{\gamma.o})(\eta) = \int_{\mathcal{H}X} \phi^2 d\mu_o \quad \square
 \end{aligned}$$

Let $L_+^2(\mathcal{H}X)$ denote the strictly positive functions in $L^2(\mathcal{H}X)$. Notice that Γ acts by isometries on $L_+^2(\mathcal{H}X)$. An element $\phi \in L_+^2(\mathcal{H}X)$ defines a positive atomless measure $\phi^2 d\mu_o$. Push this measure forward via the map $f \circ \pi$ to a measure $(f \circ \pi)_*(\phi^2 d\mu_o)$ on $\partial\mathbb{H}^n$. Define the map $P: L_+^2(\mathcal{H}X) \rightarrow \mathbb{H}^n$ by

$$P: \phi \mapsto \text{bar}((f \circ \pi)_*(\phi^2 d\mu_o)).$$

In other words $P(\phi)$ is the unique point x defined implicitly by the equation

$$\int_{\partial\mathbb{H}^n} \langle \nabla B_\theta^o, v \rangle_x d((f \circ \pi)_*(\phi^2 d\mu_o))(\theta) = \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, v \rangle_x \phi^2(\eta) d\mu_o = 0,$$

for all $v \in T_x\mathbb{H}^n$.

Lemma 5.2. *P is Γ -equivariant.*

Proof. $P(\gamma.\phi)$ is the unique point x such that for all $v \in T_x\mathbb{H}^n$,

$$\begin{aligned}
 0 &= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, v \rangle_x \phi^2(\gamma^{-1}\eta) e^{-h(X)b_\eta(\gamma.o)} d\mu_o(\eta) \\
 &= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, v \rangle_x \phi^2(\gamma^{-1}\eta) d\mu_{\gamma.o}(\eta) \\
 &= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi \circ \gamma(\eta)}^o, v \rangle_x \phi^2(\eta) d\mu_o(\eta) \quad (\text{change of variables}) \\
 &= \int_{\mathcal{H}X} \langle \nabla B_{\gamma \circ f \circ \pi(\eta)}^o, v \rangle_x \phi^2(\eta) d\mu_o(\eta) \quad (f \circ \pi \text{ is } \Gamma\text{-equivariant}) \\
 &= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, d\gamma^{-1}(v) \rangle_{\gamma^{-1}.x} \phi^2(\eta) d\mu_o(\eta) \quad (\nabla B^o \text{ is } \text{Isom}(\mathbb{H}^n)\text{-invariant}).
 \end{aligned}$$

Since $d\gamma^{-1}: T_x\mathbb{H}^n \rightarrow T_{\gamma^{-1}.x}\mathbb{H}^n$ is an isomorphism, this implies $\gamma^{-1}.x = P(\phi)$. Therefore $P(\gamma.\phi) = x = \gamma.P(\phi)$. \square

Pick a \mathcal{C}^∞ frame $\{e_i\}$ on \mathbb{H}^n and define a map $Q: \mathbb{H}^n \times L_+^2(\mathcal{H}X) \rightarrow \mathbb{R}^n$ by

$$Q: (x, \phi) \mapsto$$

$$\left(\int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, e_1 \rangle_x \phi^2(\eta) d\mu_o(\eta), \dots, \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, e_n \rangle_x \phi^2(\eta) d\mu_o(\eta) \right).$$

$\mathcal{H}X$ is compact, $B_\theta^o(x)$ is a \mathcal{C}^∞ function of both θ and x , and $\{e_i\}$ is a \mathcal{C}^∞ frame. Using these facts, applying the Lebesgue dominated convergence theorem proves that Q is \mathcal{C}^∞ . Notice P is defined implicitly by the equation

$$Q(P(\phi), \phi) = (0, \dots, 0).$$

The goal is to show that P is \mathcal{C}^1 . This can be accomplished by employing the implicit function theorem. (The implicit function theorem is true on Banach spaces. See [RS, p. 366].) For each fixed ϕ , we must show the map $Q^\phi: x \mapsto Q(x, \phi)$ has an invertible differential at each point x of the fiber $(Q^\phi)^{-1}(0, \dots, 0)$. Split Q^ϕ into coordinate functions $Q^\phi = (Q_1^\phi, \dots, Q_n^\phi)$. Then

$$\begin{aligned} \frac{\partial}{\partial x_j} Q_i^\phi(x) &= \int_{\mathcal{H}X} (\text{Hess } B_{f \circ \pi(\eta)}^o)_x(e_j, e_i) \phi^2(\eta) d\mu_o(\eta) \\ &\quad + \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \nabla_{e_j} e_i \rangle_x \phi^2(\eta) d\mu_o(\eta). \end{aligned}$$

The second term in this sum satisfies

$$\begin{aligned} &\int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \nabla_{e_j} e_i \rangle_x \phi^2(\eta) d\mu_o(\eta) \\ &= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \sum_{k=1}^n c_{ji}^k e_k \rangle_x \phi^2(\eta) d\mu_o(\eta) = \sum_{k=1}^n c_{ji}^k Q_k^\phi(x) = 0, \end{aligned}$$

for some constants c_{ji}^k depending on the frame $\{e_i\}$. This implies the bilinear form on $T_x \mathbb{H}^n$ determined by the differential of Q^ϕ at x satisfies

$$\begin{aligned} \langle v, dQ_x^\phi(v) \rangle &= \int_{\mathcal{H}X} (\text{Hess } B_{f \circ \pi(\eta)}^o)_x(v, v) \phi^2(\eta) d\mu_o(\eta) \\ &= \int_{\partial \mathbb{H}^n} (\text{Hess } B_\theta^o)_x(v, v) d((f \circ \pi)_*(\phi^2 d\mu_o))(\theta), \end{aligned}$$

for all $v \in T_x \mathbb{H}^n$. The right hand side of this equation is strictly positive by Lemma 2.1. This implies the differential of Q^ϕ at x is positive definite, and thus invertible. Therefore the implicit function theorem may be applied to conclude that P is \mathcal{C}^1 .

Define a map

$$\begin{aligned} \Phi: X &\rightarrow L_+^2(\mathcal{H}X) \subset L^2(\mathcal{H}X) \\ x &\mapsto \sqrt{\exp(-h(X)b_\eta(x))}. \end{aligned}$$

Lemma 5.3. Φ is Lipschitz.

Proof. Let D be the diameter of the downstairs metric space Z covered by X . It follows that for any $p \in X$, $\|\Phi(p)\|$ is at most $e^{Dh(X)}$. Pick points $x, y \in X$. The goal is to control the quantity

$$\int_{\mathcal{H}X} \left| e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)b_\eta(y)} \right|^2 d\mu_o(\eta).$$

For $\eta \in \mathcal{H}X$, b_η is 1-Lipschitz. This implies $b_\eta(y) \leq d(y, x) + b_\eta(x)$. Using this we obtain the inequalities

$$\begin{aligned} e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)b_\eta(y)} &\leq e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)d(y,x)} e^{-\frac{1}{2}h(X)b_\eta(x)} \\ &= e^{-\frac{1}{2}h(X)b_\eta(x)} (1 - e^{-\frac{1}{2}h(X)d(y,x)}), \end{aligned}$$

yielding

$$\left| e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)b_\eta(y)} \right| \leq (e^{-\frac{1}{2}h(X)b_\eta(x)} + e^{-\frac{1}{2}h(X)b_\eta(y)}) (1 - e^{-\frac{1}{2}h(X)d(x,y)}).$$

Therefore,

$$\begin{aligned} &\int_{\mathcal{H}X} \left| e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)b_\eta(y)} \right|^2 d\mu_o(\eta) \\ &\leq (1 - e^{-\frac{1}{2}h(X)d(x,y)})^2 \int_{\mathcal{H}X} \left| e^{-\frac{1}{2}h(X)b_\eta(x)} + e^{-\frac{1}{2}h(X)b_\eta(y)} \right|^2 d\mu_o(\eta) \\ &= (1 - e^{-\frac{1}{2}h(X)d(x,y)})^2 \cdot \|\Phi(x) + \Phi(y)\|^2 \\ &\leq (1 - e^{-\frac{1}{2}h(X)d(x,y)})^2 \cdot [\|\Phi(x)\| + \|\Phi(y)\|]^2 \leq (1 - e^{-\frac{1}{2}h(X)d(x,y)})^2 \cdot 4e^{2Dh(X)} \end{aligned}$$

To complete the proof, notice that for $t \geq 0$, $(1 - e^{-\frac{1}{2}h(X)t}) \leq \frac{1}{2}h(X)t$. Applying this yields

$$\int_{\mathcal{H}X} \left| e^{-\frac{1}{2}h(X)b_\eta(x)} - e^{-\frac{1}{2}h(X)b_\eta(y)} \right|^2 d\mu_o(\eta) \leq \left(\frac{1}{2}h(X)d(x, y) \right)^2 \cdot 4e^{2Dh(X)}.$$

So finally we've obtained

$$\frac{\|\Phi(x) - \Phi(y)\|}{d(x, y)} \leq h(X) \cdot e^{Dh(X)}. \quad \square$$

P is \mathcal{C}^1 , therefore $F = P \circ \Phi$ is locally Lipschitz. Γ -equivariance implies F descends to a locally Lipschitz map $X/\Gamma = Z \rightarrow \mathbb{H}^n/\Gamma = M_{\text{hyp}}$. By Rademacher's theorem, F is differentiable almost everywhere (see Section 2.5).

6. The Jacobian estimate

In this section we prove parts (2) and (3) of Proposition 4.3.

(2) $|\text{Jac } F(p)| \leq \left(\frac{h(X)}{n-1}\right)^n$ almost everywhere.

(3) If for some p , $|\text{Jac } F(p)| = \left(\frac{h(X)}{n-1}\right)^n$, then the differential dF_p is a homothety of ratio $\left(\frac{h(X)}{n-1}\right)^n$.

The proof closely follows Section 5 of [F], which is in turn based on [BCG2, p. 636–639]. Recall that X is the universal cover of Z , $\Gamma := \pi_1(Z) \cong \pi_1(M_{\text{hyp}})$, $\{e_i\}$ is a \mathcal{C}^∞ frame on $T\mathbb{H}^n$, and $\Omega \subset X$ is a subset of full measure possessing a \mathcal{C}^1 -atlas (see Sections 2.4 and 2.5).

In the previous section we have defined a Γ -equivariant locally Lipschitz map $\Phi: X \rightarrow L_+^2(\mathcal{H}X)$, a Γ -equivariant \mathcal{C}^1 barycenter map $P: L_+^2(\mathcal{H}X) \rightarrow \mathbb{H}^n$, and a \mathcal{C}^∞ map $Q = (Q_1, \dots, Q_n): \mathbb{H}^n \times L_+^2(\mathcal{H}X) \rightarrow \mathbb{R}^n$. They satisfied the equations $F = P \circ \Phi$ and $Q(P(\phi), \phi) = 0$ for $\phi \in L_+^2(\mathcal{H}X)$. We thus obtain the implicit equation $Q(F, \Phi) = 0$. Let $\mathcal{O} \subseteq \Omega \subseteq X$ be the set of points where Φ is differentiable. By Rademacher's theorem (see Section 2.5), $\mathcal{O} \subseteq X$ is a subset of full measure. Pick $p \in \mathcal{O}$ and $v \in T_p X$.

Lemma 6.1. *Differentiating the function*

$$x \mapsto Q_i(F(x), \Phi(x)) = \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, e_i \rangle_{F(x)} e^{-h(X)b_\eta(x)} d\mu_o(\eta) = 0$$

at p in the direction of v yields

$$\begin{aligned} 0 &= \int_{\mathcal{H}X} (\text{Hess } B_{f \circ \pi(\eta)}^o)_{F(p)}(dF(v), e_i) e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\ &\quad + \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \nabla_{dF(v)} e_i \rangle_{F(p)} e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\ &\quad + \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, e_i \rangle_{F(p)} [-h(X) \langle \nabla b_\eta, v \rangle_p e^{-h(X)b_\eta(p)}] d\mu_o(\eta). \end{aligned} \quad (\star)$$

Proof. This is an application of the Lebesgue dominated convergence theorem. $\mathcal{H}X$ is compact. So to apply the theorem it is sufficient to find a $c > 0$ such that for all $\eta \in \mathcal{H}X$, the function

$$x \mapsto \langle \nabla B_{f \circ \pi(\eta)}^o, e_i \rangle_{F(x)} e^{-h(X)b_\eta(x)}$$

is locally c -Lipschitz near p . To show the existence of such a constant, use that b_η is 1-Lipschitz, $\mathcal{H}X$ is compact, and $\langle \nabla B_{f \circ \pi(\eta)}^o, e_i \rangle_q: \mathbb{H}^n \times \partial\mathbb{H}^n \rightarrow \mathbb{R}$ is \mathcal{C}^∞ . \square

The second term in equation (★) satisfies

$$\begin{aligned}
& \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \nabla_{dF(v)} e_i \rangle_{F(p)} e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\
&= \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, \sum_{j=1}^n c_{ij} e_j \rangle_{F(p)} e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\
&= \sum_{j=1}^n c_{ij} Q_j(F(p), \Phi(p)) = 0,
\end{aligned}$$

for some constants c_{ij} depending on the frame $\{e_i\}$. Therefore, for $v \in T_p X$, $u \in T_{F(p)} \mathbb{H}^n$ we have

$$\begin{aligned}
& \int_{\mathcal{H}X} (\text{Hess } B_{f \circ \pi(\eta)}^o)_{F(p)}(dF(v), u) e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\
&= h(X) \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, u \rangle_{F(p)} \langle \nabla b_\eta, v \rangle_p e^{-h(X)b_\eta(p)} d\mu_o(\eta).
\end{aligned} \tag{★★}$$

Let $\|\mu_p\|$ denote the total mass $\mu_p(\mathcal{H}X)$ of the measure μ_p . Define a bilinear form K and a quadratic form H on $T_{F(p)} \mathbb{H}^n$ by

$$\begin{aligned}
\langle K(w), u \rangle_{F(p)} &:= \frac{1}{\|\mu_p\|} \int_{\mathcal{H}X} (\text{Hess } B_{f \circ \pi(\eta)}^o)_{F(p)}(w, u) e^{-h(X)b_\eta(p)} d\mu_o(\eta) \\
\langle H(u), u \rangle_{F(p)} &:= \frac{1}{\|\mu_p\|} \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, u \rangle_{F(p)}^2 e^{-h(X)b_\eta(p)} d\mu_o(\eta).
\end{aligned}$$

Note that the symmetric endomorphism K is positive definite by Theorem 2.1. It is therefore invertible. This is used in the proof of Lemma 6.2.

Use equation (★★), the Cauchy–Schwarz inequality, and the definition of K to obtain

$$\begin{aligned}
& |\langle K \circ dF(v), u \rangle_{F(p)}| \\
&= \left| \frac{h(X)}{\|\mu_p\|} \int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, u \rangle_{F(p)} \cdot \langle \nabla b_\eta, v \rangle_p \cdot e^{-h(X)b_\eta(p)} d\mu_o(\eta) \right| \\
&\leq \frac{h(X)}{\|\mu_p\|} \left[\int_{\mathcal{H}X} \langle \nabla B_{f \circ \pi(\eta)}^o, u \rangle_{F(p)}^2 d\mu_p(\eta) \right]^{1/2} \cdot \left[\int_{\mathcal{H}X} \langle \nabla b_\eta, v \rangle_p^2 d\mu_p(\eta) \right]^{1/2} \\
&= \frac{h(X)}{\|\mu_p\|^{1/2}} [\langle H(u), u \rangle_{F(p)}]^{1/2} \cdot \left[\int_{\mathcal{H}X} \langle \nabla b_\eta, v \rangle_p^2 d\mu_p(\eta) \right]^{1/2},
\end{aligned}$$

for all $u \in T_{F(p)} \mathbb{H}^n$ and $v \in T_p \Omega$.

Lemma 6.2 ([BCG2, p. 637]). *For all $p \in \mathcal{O}$,*

$$|\text{Jac } F(p)| \leq \frac{h(X)^n (\det H)^{1/2}}{n^{n/2} \det K}.$$

Proof. The proof is the proof of Lemma 5.4 of [BCG2, p. 637] with two modifications. First replace the Busemann function B_α with the horofunction b_η , and notice that

$$\sum_{i=1}^n \langle \nabla b_\eta, v_i \rangle_p^2 = \|\nabla b_\eta\|_p^2 \leq 1. \quad (\ddagger)$$

Second, the total mass of the Patterson–Sullivan measure $\|\mu_p\|$ must be carried through the estimate, but it cancels itself out in the final step. \square

A key property of hyperbolic space is that Busemann functions on \mathbb{H}^n satisfy the equation

$$(\text{Hess } B_\theta^o)(u, v) = \langle u, v \rangle - \langle \nabla B_\theta^o, u \rangle \cdot \langle \nabla B_\theta^o, v \rangle$$

for all $\theta \in \partial \mathbb{H}^n$ (see [BCG1, p. 750–751]). Integrating this equation over $\mathcal{H}X$ yields

$$K = \text{Id} - H.$$

Lemma 6.3. *The symmetric endomorphism H is positive definite.*

Proof. Suppose there exists $x \in X$ and $u \in T_x X$ such that $\langle H(u), u \rangle_x = 0$. From the definition of H , this implies the support of the measure $(f \circ \pi)_* \mu_0$ on $\partial \mathbb{H}^n$ is contained in a codimension one conformally round sphere in $S \subset \mathbb{H}^n$. By the equivariance of the Patterson–Sullivan measures, the support of $(f \circ \pi)_* \mu_{\gamma.o}$ is contained in $\gamma(S)$ for $\gamma \in \Gamma$. But $\mu_{\gamma.o} \ll \mu_o$ implies the support of $(f \circ \pi)_* \mu_{\gamma.o}$ is contained in S . Therefore Γ preserves $S \subset \partial \mathbb{H}^n$. This contradicts the fact that the limit set of Γ is the entire sphere at infinity. \square

A short computation shows that $\text{tr}(H) = 1$. Therefore we can apply the brain in a jar lemma (Lemma 2.2) to H . This yields

$$\frac{\det(H)}{[\det(\text{Id} - H)]^2} \leq \left[\frac{n}{(n-1)^2} \right]^n,$$

with equality if and only if $H = \frac{1}{n} \text{Id}$. Combining this with Lemma 6.2 proves the desired inequality, namely

$$|\text{Jac } F(p)| \leq \left(\frac{h(X)}{n-1} \right)^n.$$

This completes the proof of part (2) of Proposition 4.3.

The proof of part (3) of Proposition 4.3 is the “equality case” argument on page 639 of [BCG2] with two modifications (as in Lemma 6.2). First replace the Busemann function B_α with the horofunction b_η and use inequality (\ddagger) . Second, the total mass of the Patterson–Sullivan $\|\mu_p\|$ must be carried through the estimate, and again it cancels itself out. This completes the proof of Proposition 4.3.

7. The equality case

Theorem 7.1. *Let M_{hyp} be a closed hyperbolic n -manifold for $n \geq 3$. Let Z be a compact n -dimensional convex Riemannian amalgam (see Section 2.8). Let $f: Z \rightarrow M_{\text{hyp}}$ be a homotopy equivalence. If*

$$h(X)^n \text{Vol}(Z) = (n-1)^n \text{Vol}(M_{\text{hyp}})$$

then the natural map $F: Z \rightarrow M_{\text{hyp}}$ is a homothetic homeomorphism.

Remark 7.2. Recall that if Z is either

- the metric doubling of a convex hyperbolic manifold with boundary (see Section 2.7), or
 - a cone-manifold (see Section 2.6),
- then Z is a convex Riemannian amalgam.

Let X be the universal cover of Z . Lift the convex Riemannian amalgam structure on Z to a convex Riemannian amalgam structure on X with decomposition $\{C_j\}$ into convex Riemannian manifolds with boundary. Define the incomplete disconnected Riemannian manifold $\Omega := \bigcup_j \text{int}(C_j) \subseteq X$.

The equation

$$h(X)^n \text{Vol}(Z) = (n-1)^n \text{Vol}(M_{\text{hyp}})$$

implies the string of inequalities from Lemma 4.4 is in fact a string of equalities. Therefore $|\text{Jac } F(p)| = \left(\frac{h(X)}{n-1}\right)^n$ almost everywhere. By Proposition 4.3, dF is a homothety almost everywhere. The goal is to prove $F: Z \rightarrow M_{\text{hyp}}$ is a homothetic homeomorphism. Without a loss of generality, scale the metric of Z so that dF is an isometry (not necessarily orientation preserving) almost everywhere. This forces $\text{Vol}(Z) = \text{Vol}(M_{\text{hyp}})$. It now suffices to show that $F: Z \rightarrow M_{\text{hyp}}$ is an isometric homeomorphism. This will be done by working on the universal covers and showing $F: X \rightarrow \mathbb{H}^n$ is an equivariant isometric homeomorphism.

Lemma 7.3. *$F: X \rightarrow \mathbb{H}^n$ is a contraction mapping, i.e. for any pair $x, y \in X$, $d_{\mathbb{H}^n}(F(x), F(y)) \leq d_X(x, y)$.*

Proof. Pick a length minimizing geodesic segment α connecting x to y . Define $\mathcal{O} \subseteq \Omega \subseteq X$ to be the set of full measure where dF is an isometry. There exists a small perturbation α_ε of α such that $\alpha_\varepsilon \cap \mathcal{O} \subseteq \alpha_\varepsilon$ is a subset of full measure, and $\text{length}(\alpha_\varepsilon) \leq \text{length}(\alpha) + \varepsilon$. F is locally Lipschitz, so in all Hausdorff dimensions F maps measure zero sets to measure zero sets. This implies F preserves length on α_ε . Therefore

$$d_{\mathbb{H}^n}(F(x), F(y)) \leq \text{length}(F(\alpha_\varepsilon)) = \text{length}(\alpha_\varepsilon) \leq \text{length}(\alpha) + \varepsilon = d_X(x, y) + \varepsilon.$$

Since ε was arbitrary, the result follows. \square

We claim that F is volume preserving. $|\text{Jac } F| = 1$ a.e. and F is locally Lipschitz, so F is volume non-increasing. If it strictly decreased the volume of some measurable set, then we would have $\text{Vol}(Z) > \text{Vol}(M_{\text{hyp}})$. This would be a contradiction. Therefore F is volume preserving.

There exist constants $C_1, \varepsilon_{\text{hyp}} > 0$ such that if $y \in \mathbb{H}^n$ and $\varepsilon < \varepsilon_{\text{hyp}}$ then

$$v_n \varepsilon^n \leq \text{Vol}(B_{\mathbb{H}^n}(y, \varepsilon)) \leq v_n \varepsilon^n (1 + C_1 \varepsilon^2), \quad (4)$$

where v_n is the volume of a unit ball in \mathbb{R}^n . Similarly, $\Omega \subseteq X$ is a Riemannian manifold with sectional curvature bounded from above. (An upper curvature bound follows from the definition of a convex Riemannian amalgam.) For compact $K \subset \Omega$ (possibly a point), define

$$\text{inj}_\Omega(K) := \inf_{z \in K} (\text{injectivity radius of } \Omega \text{ at } z).$$

The upper curvature bound implies there exist constants $C_2, \varepsilon_\Omega > 0$ such that if $\varepsilon < \varepsilon_\Omega$, $\varepsilon < \text{inj}_\Omega(z)$, and $B_X(z, \varepsilon) \subset \Omega$ then

$$v_n \varepsilon^n (1 - C_2 \varepsilon^2) \leq \text{Vol}(B_\Omega(z, \varepsilon)). \quad (5)$$

We now consider the restriction of F to the “smooth” set Ω . Define $F_\Omega := F|_\Omega: \Omega \rightarrow \mathbb{H}^n$.

Lemma 7.4. F_Ω is injective.

Proof. (This proof is an adaptation of [BCG1, Lemma C.4].) Suppose there exist distinct $z_1, z_2 \in \Omega$ such that $F_\Omega(z_1) = F_\Omega(z_2) = y$. Pick

$$\varepsilon_0 < \min\{\varepsilon_\Omega, \text{inj}_\Omega(z_1), \text{inj}_\Omega(z_2), \varepsilon_{\text{hyp}}\}$$

sufficiently small such that

$$B_X(z_1, \varepsilon_0) \cap B_X(z_2, \varepsilon_0) = \emptyset \quad \text{and} \quad B_X(z_1, \varepsilon_0) \cup B_X(z_2, \varepsilon_0) \subset \Omega.$$

In particular, inequality 4 (resp. inequality 5) is valid at y (resp. z_1 and z_2) for $\varepsilon < \varepsilon_0$.

Since F is contracting, $F_\Omega(B_\Omega(z_i, \varepsilon)) \subseteq B_{\mathbb{H}^n}(y, \varepsilon)$ for all $\varepsilon < \varepsilon_0$. Therefore

$$B_\Omega(z_1, \varepsilon) \cup B_\Omega(z_2, \varepsilon) \subseteq F_\Omega^{-1}(B_{\mathbb{H}^n}(y, \varepsilon)).$$

Since F is volume preserving (and $X \setminus \Omega$ is measure zero),

$$\begin{aligned} \text{Vol}(B_{\mathbb{H}^n}(y, \varepsilon)) &= \text{Vol}(F_\Omega^{-1}(B_{\mathbb{H}^n}(y, \varepsilon))) \geq \text{Vol}(B_\Omega(z_1, \varepsilon) \cup B_\Omega(z_2, \varepsilon)) \\ &= \text{Vol}(B_\Omega(z_1, \varepsilon)) + \text{Vol}(B_\Omega(z_2, \varepsilon)). \end{aligned}$$

Now apply inequalities 4 and 5 to obtain

$$v_n \varepsilon^n (1 + C_1 \varepsilon^2) \geq 2v_n \varepsilon^n (1 - C_2 \varepsilon^2).$$

This implies $\varepsilon^2(C_1 + 2C_2) \geq 1$ for all $\varepsilon < \varepsilon_0$, which is a contradiction. \square

Since Ω is locally compact, Lemma 7.4 shows that $F_\Omega: \Omega \rightarrow F_\Omega(\Omega)$ is a homeomorphism.

Lemma 7.5. F_Ω^{-1} is locally Lipschitz.

Proof. (This proof is an adaptation of [BCG1, Lemma C.7].) Pick $z \in \Omega$. Let $y := F_\Omega(z)$. Pick $\varepsilon_0 < \min\{\varepsilon_{\text{hyp}}, \varepsilon_\Omega\}$ sufficiently small such that $B_X(z, 3\varepsilon_0) \subset \Omega$, $\varepsilon_0 < \text{inj}_\Omega(\overline{B_X(z, 2\varepsilon_0)})$, and

$$2\varepsilon_0^2(C_1 + C_2) < \frac{1}{2^n}.$$

Define $V := B_\Omega(z, \varepsilon_0)$. Since F is a contraction mapping, $F(V) \subseteq B_{\mathbb{H}^n}(y, \varepsilon_0)$. We will show that if $z_1, z_2 \in V$ are distinct, then

$$d_X(z_1, z_2) < 2d_{\mathbb{H}^n}(F(z_1), F(z_2)).$$

This will imply $F|_V^{-1}$ is 2-Lipschitz on the open set $F(V)$.

Suppose the above inequality is false, i.e. there exist distinct $z_1, z_2 \in V$ such that $2d_{\mathbb{H}^n}(F(z_1), F(z_2)) \leq d_X(z_1, z_2)$. Set $\varepsilon := d_{\mathbb{H}^n}(F(z_1), F(z_2)) \leq \varepsilon_0$. Since $\varepsilon < \varepsilon_0$, inequality 4 remains valid at $F(z_1)$ and $F(z_2)$ for ε . $B_{\mathbb{H}^n}(F(z_1), \varepsilon)$ and $B_{\mathbb{H}^n}(F(z_2), \varepsilon)$ intersect, and their intersection contains a ball of radius $\varepsilon/2$ centered in the middle of the minimizing geodesic joining $F(z_1)$ to $F(z_2)$ (see Figure 5). Therefore

$$\text{Vol}(B_{\mathbb{H}^n}(F(z_1), \varepsilon) \cup B_{\mathbb{H}^n}(F(z_2), \varepsilon)) \leq v_n \left[2\varepsilon^n (1 + C_1 \varepsilon^2) - \frac{1}{2^n} \varepsilon^n \right].$$

Moreover, $d_X(z_1, z_2) \geq 2\varepsilon$ implies $B_X(z_1, \varepsilon) \cap B_X(z_2, \varepsilon) = \emptyset$. Since $B_X(z_1, \varepsilon) \cup B_X(z_2, \varepsilon) \subset B_X(z, 2\varepsilon_0) \subset \Omega$, we may apply inequality 5 to conclude

$$\begin{aligned} \text{Vol}(F(B_X(z_1, \varepsilon) \cup B_X(z_2, \varepsilon))) &= \text{Vol}(B_X(z_1, \varepsilon) \cup B_X(z_2, \varepsilon)) \\ &\geq v_n [2\varepsilon^n (1 - C_2 \varepsilon^2)]. \end{aligned}$$




Figure 5

F is a contraction mapping. Therefore

$$F(B_X(z_1, \varepsilon)) \subseteq B_{\mathbb{H}^n}(F(z_1), \varepsilon) \quad \text{and} \quad F(B_X(z_2, \varepsilon)) \subseteq B_{\mathbb{H}^n}(F(z_2), \varepsilon).$$

Putting these inequalities together yields

$$v_n \left[2\varepsilon^n (1 + C_1 \varepsilon^2) - \frac{1}{2^n} \varepsilon^n \right] \geq v_n \varepsilon^n [2(1 - C_2 \varepsilon^2)],$$

implying

$$2\varepsilon^2(C_1 + C_2) \geq \frac{1}{2^n}.$$

This contradicts our choice of ε_0 . □

Since $F|_{\Omega}^{-1}$ is locally Lipschitz, it is differentiable almost everywhere. Therefore, for almost every $z \in \Omega$,

$$d(\text{Id}_{\Omega})_z = d(F|_{\Omega}^{-1} \circ F_{\Omega})_z = d(F|_{\Omega}^{-1}) \circ dF_{\Omega}.$$

This implies $d(F|_{\Omega}^{-1})$ is an isometry of tangent spaces almost everywhere. By an argument analogous to Lemma 7.3, and by working on small balls in M_{hyp} , one can see that $F|_{\Omega}^{-1}$ is locally a contraction mapping $F(\Omega) \rightarrow \Omega$. Both F_{Ω} and $F|_{\Omega}^{-1}$ are local contraction maps. Thus they are both local isometries.

Lemma 7.6. *For all j , $F|_{C_j}$ is an isometry onto its image, and $F(\text{int}(C_j))$ is convex.*

Proof. By continuity, it suffices to show $F|_{\text{int}(C_j)}$ is an isometry onto its image. Pick $x, y \in \text{int}(C_j)$. Since $\text{int}(C_j)$ is convex, there exists a geodesic segment $xy \subset \text{int}(C_j)$ joining x to y . F_{Ω} is a local isometry. Thus $\text{length}(F(xy)) = \text{length}(xy)$, and

$F(xy) \subset \mathbb{H}^n$ is locally a hyperbolic geodesic joining $F(x)$ to $F(y)$. In \mathbb{H}^n , a local geodesic is a global geodesic. Therefore

$$d_{\mathbb{H}^n}(F(x), F(y)) = \text{length}(F(xy)) = \text{length}(xy) = d_X(x, y). \quad \square$$

Lemma 7.7. *For each $z \in X$, there exists $\delta_z > 0$ such that $d_X(z, x) < \delta_z$ implies $d_{\mathbb{H}^n}(F(z), F(x)) = d_X(z, x)$.*

(Notice that this lemma proves neither that F is an isometry nor that it is locally injective. To see why, consider the branched double covering of an Euclidean disk by an Euclidean cone with cone angle 4π .)

Proof. Pick $z \in X$. If $z \in \Omega$ then we are done. So assume $z \in X \setminus \Omega$. Up to rearranging the indices we may assume

$$z \in (\partial C_1 \cup \partial C_2 \cup \dots \cup \partial C_l) \setminus (\partial C_{l+1} \cup \dots \cup \partial C_N).$$

By this, there exists $\delta_z > 0$ such that

$$B_X(z, \delta_z) \cap \Omega \subseteq \bigcup_{j=1}^l C_j.$$

For $x \in B_X(z, \delta_z)$, there exists an integer j_x and a geodesic segment $xz \subset C_{j_x}$. $F|_{C_{j_x}}$ is an isometry, so $d_X(z, x) = d_{C_{j_x}}(z, x) = d_{\mathbb{H}^n}(F(z), F(x))$. This proves the lemma. \square

As a corollary of this lemma, we see that $F^{-1}(y)$ is a discrete set for all $y \in \mathbb{H}^n$.

Lemma 7.8. *$F: X \rightarrow \mathbb{H}^n$ is injective.*

Proof. For $y \in \mathbb{H}^n$, $F^{-1}(y)$ is the discrete set $\{z_i\}$. We will show $F^{-1}(y)$ must be a single point. For all i , pick $\delta_i < 1$ so that $d_X(z_i, x) < \delta_i$ implies $d_{\mathbb{H}^n}(F(z_i), F(x)) = d_X(z_i, x)$. Assume the metric balls $B_X(z_i, \delta_i)$ are disjoint.

If for some i , $F|_{B_X(z_i, \delta_i)}: B_X(z_i, \delta_i) \rightarrow B_{\mathbb{H}^n}(F(z_i), \delta_i)$ is not surjective, then F can be properly homotoped to a map taking $B_X(z_i, \delta_i) \rightarrow \partial B_{\mathbb{H}^n}(F(z_i), \delta_i)$. Moreover, this can be done without altering F outside of $B_X(z_i, \delta_i)$.

If for every i , $F|_{B_X(z_i, \delta_i)}: B_X(z_i, \delta_i) \rightarrow B_{\mathbb{H}^n}(F(z_i), \delta_i)$ is not surjective, then F is properly homotopic to a map which does not have y in its image. F is a proper surjective homotopy equivalence. Thus every map properly homotopic to F is surjective. Therefore for some i , $F|_{B_X(z_i, \delta_i)}$ is surjective. We may assume $i = 1$.

F_Ω is injective and open, Ω is open and dense, and $F(B_X(z_1, \delta_1)) = B_{\mathbb{H}^n}(y, \delta_1)$. Thus

$$F(\Omega \setminus B_X(z_1, \delta_1)) \cap B_{\mathbb{H}^n}(y, \delta_1) = \emptyset.$$

By extending to the closure of Ω , this shows that for all $i > 1$,

$$F(z_i) = y \notin B_{\mathbb{H}^n}(y, \delta_1).$$

Therefore $F^{-1}(y)$ must be the single point z_1 . \square

Recall that $F: X \rightarrow \mathbb{H}^n$ is surjective. Therefore $F: X \rightarrow \mathbb{H}^n$ is a continuous bijection. Since X is locally compact, F is a homeomorphism. To prove that F is an isometry, it is sufficient to show F^{-1} is a contraction mapping.

Lemma 7.9. *F^{-1} is a contraction mapping, i.e. it is 1-Lipschitz.*

Proof. By the above lemmas, F imposes a convex Riemannian amalgam structure on \mathbb{H}^n . The collection of convex Riemannian manifolds with boundary is $\{F(C_j)\}$.

Pick $x, y \in \mathbb{H}^n$, $\varepsilon > 0$. Since each $F(C_j)$ is convex, there exists a path $\gamma \subset \mathbb{H}^n$ joining x to y such that:

- $\text{length}(\gamma) \leq d(x, y) + \varepsilon$,
- $\gamma \cap F(\partial C_j)$ is at most two points for any $F(C_j)$ of the decomposition.

As the collection $\{F(C_j)\}$ is locally finite, the set $F^{-1}(\gamma) \setminus \Omega$ is finite. $F|_{C_j}$ is an isometry for all j . Therefore $F^{-1}(\gamma)$ is a path of the same length as γ . This implies F^{-1} is 1-Lipschitz. \square

This completes the proof of Theorem 7.1.

8. Applications

8.1. Kleinian groups. The Kleinian group theory notation used here is defined in Sections 2.9–2.12.

Let N be a compact acylindrical 3-manifold (see Section 2.10). Recall that by Corollary 2.5, there exists a convex cocompact hyperbolic 3-manifold M_g such that C_{M_g} is homeomorphic to N and the boundary of the convex core $\partial C_{M_g} \subset M_g$ is totally geodesic.

As was discussed in Section 1, the following theorem solves a conjecture in Kleinian groups.

Theorem 8.1. *Let N be a compact acylindrical 3-manifold. Let (M_g, m_g) be a convex cocompact hyperbolic 3-manifold such that C_{M_g} is homeomorphic to N and the boundary of the convex core $\partial C_{M_g} \subset M_g$ is totally geodesic. For all $(M, m) \in \mathcal{H}(N)$,*

$$\text{Vol}(C_M) \geq \text{Vol}(C_{M_g}),$$

with equality if and only if M and M_g are isometric.

Fix an $(M, m) \in H(N)$. We may assume without a loss of generality that M is geometrically finite. Since ∂C_{M_g} is totally geodesic, metrically doubling C_{M_g} across its boundary produces a compact hyperbolic manifold DC_{M_g} .

Lemma 8.2. *Metrically doubling the convex core C_M across its boundary yields an Alexandrov space with curvature bounded below by -1 .*

Proof. In [S, Appendix A] it was proven that taking an ε -neighborhood of C_M in M , and metrically doubling that across its boundary to obtain $D\mathcal{N}_\varepsilon C_M$, yields an Alexandrov space with curvature bounded below by -1 . DC_M is the Gromov–Hausdorff limit of these spaces as $\varepsilon \rightarrow 0$. Being an Alexandrov space with curvature bounded below by -1 is a closed property in the Gromov–Hausdorff topology [BBI, Proposition 10.7.1, p. 376]. This proves the lemma. \square

Remark 8.3. Lemma 8.2 also follows from a more general theorem of Perelman (see Theorem 8.8).

In particular, by Lemma 8.2 and Theorem 2.7, the volume growth entropy of $\widetilde{DC_M}$ is not greater than $2 = h(\mathbb{H}^3)$. This will be used later. Let σ denote the boundary preserving isometric involution of DC_M and DC_{M_g} .

Case 1. Assume M is convex cocompact.

Proof of Case 1. Both M nor M_g are convex cocompact. By Theorem 2.4, $m_g \circ m^{-1}$ is homotopic to a homeomorphism $C_M \rightarrow C_{M_g}$. (m^{-1} denotes a relative homotopy inverse of m .) This homeomorphism can be “doubled” to produce a homeomorphism between the doubled manifolds $f: DC_M \rightarrow DC_{M_g}$. Theorem 4.1 may now be applied to $f: DC_M \rightarrow DC_{M_g}$. This proves that

$$h(\widetilde{DC_M})^3 \text{Vol}(DC_M) \geq 2^3 \text{Vol}(DC_{M_g}),$$

with equality if and only if DC_M and DC_{M_g} are isometric. Since $h(\widetilde{DC_M}) \leq 2$ we have

$$\text{Vol}(DC_M) \geq \text{Vol}(DC_{M_g}),$$

with equality if and only if DC_M and DC_{M_g} are isometric. Dividing both sides by 2 yields the desired inequality.

Let us now assume $\text{Vol}(DC_M) = \text{Vol}(DC_{M_g})$. The goal is to show M and M_g are isometric. To do this it is sufficient to show C_M and C_{M_g} are isometric. The map $f: DC_M \rightarrow DC_{M_g}$ is by construction σ -equivariant. Let $F: DC_M \rightarrow DC_{M_g}$ be the natural map induced by f . The σ -equivariance of f implies F is also σ -equivariant (see Remark 4.2). Therefore $F: C_M \rightarrow C_{M_g}$ is an isometry. This completes the proof of Case 1. \square

Case 2. (M, m) is not convex cocompact.

Proof of Case 2. $(M, m) \in H(N)$, so by definition $m: N \rightarrow M^o$ is a homotopy equivalence (see Section 2.12). Moreover, by Theorem 2.4 there exists a homeomorphism $g: N \rightarrow C_M \cap M^o$. Let $P := g^{-1}(C_M \cap \partial M^o) \subset \partial N$. Consider (M, m) as an element of $H(N, P)$ with no additional parabolics. (N, P) is pared acylindrical. Therefore there exists a geometrically finite $M_P \in H(N, P)$ with no additional parabolics such that C_{M_P} has totally geodesic boundary.

Again by Theorem 2.4, there is a homeomorphism $C_{M_P} \rightarrow N \setminus P$, inducing a doubled homeomorphism $DC_{M_P} \rightarrow D(N \setminus P)$ between open manifolds. Let $p \subset P \subset \partial N$ be a finite collection of disjoint simple closed curves such that p is a strong deformation retract of P , i.e. each component of p is a core curve of a component of P . Then there is a homeomorphism $D(N \setminus P) \rightarrow (DN) \setminus p$. Moreover, the manifolds $DC_{M_P} \cong D(N \setminus P)$ are topologically obtained by removing the curves $p \subset \partial N \subset DN$. Conversely, replacing the removed curves of $DN \setminus p$ corresponds to performing a (topological) Dehn surgery on $DN \setminus p$. Therefore the homeomorphism type of $DN \cong DC_{M_g}$ can be obtained by performing a topological Dehn surgery on DC_{M_P} . DC_{M_P} is a finite volume hyperbolic manifold. Therefore Mostow rigidity implies that DC_{M_g} may in fact be obtained by performing a hyperbolic Dehn surgery on DC_{M_P} [Th1].

Hyperbolic Dehn surgery strictly decreases volume [Th1, Theorem 6.5.6], [Bes]. Therefore $\text{Vol}(DC_{M_P}) > \text{Vol}(DC_{M_g})$. Moreover, by [NZ] there exists a closed hyperbolic 3-manifold L obtained from hyperbolic Dehn surgery on DC_{M_P} such that

$$\text{Vol}(DC_{M_P}) > \text{Vol}(L) > \text{Vol}(DC_{M_g}).$$

So to complete the proof it suffices to show that

$$\text{Vol}(DC_M) \geq \text{Vol}(L).$$

This will be accomplished by geometrically filling in the cusps and reducing to the case of closed manifolds.

The geometric (though not hyperbolic) Dehn surgery arguments below are based on techniques from [Bes], [L]. (See also [BCS] for another application of these methods.) The exposition here will roughly follow [Bes].

Outside a compact set, the metric on DC_M is a collection of smooth rank two hyperbolic cusps. For simplicity, let us assume that DC_M has exactly one cusp. The general case follows by performing the following geometric operations on each cusp individually.

Pick a compact exhaustion K_i of DC_M such that each boundary ∂K_i is a smooth horospherical torus. By “opening up” the cusp of DC_M one can construct a family of metric spaces $\{(A, d_\varepsilon)\}_{\varepsilon \in (0,1]}$ such that:

1. A is homeomorphic to K_i .
2. (A, d_ε) is an Alexandrov space with curvature bounded below by $-1 - c_\varepsilon^2$, where $\lim_{\varepsilon \rightarrow 0} c_\varepsilon = 0$.
3. For each i , there is an isometric embedding $K_i \rightarrow (A, d_\varepsilon)$ for all sufficiently small ε .
4. $\lim_{\varepsilon \rightarrow 0} \text{Vol}(A, d_\varepsilon) = \text{Vol}(DC_M)$.
5. Near the boundary of A the metric d_ε is a flat Riemannian metric with totally geodesic boundary ∂A .
6. For any $\varepsilon \leq 1$, $(\partial A, d_\varepsilon) = (\partial A, \varepsilon \cdot d_1)$.

(A careful and clear explanation of this procedure is in [Bes, Section 2.2]. See also [L].)

Let W denote a solid torus. Using the fact that W is a product of a disk and a circle, one can build a product Riemannian metric g on W such that:

1. (W, g) has totally geodesic boundary isometric to an Euclidean torus.
2. (W, g) has sectional curvature bounded below by zero.

A manifold homemorphic to L is obtained from A by appropriately gluing ∂W to ∂A (i.e. by an appropriate Dehn surgery). The goal is to perform this gluing geometrically. To do so we must scale appropriately and interpolate between the flat torus boundaries of (A, d_ε) and (W, g) . Let $T^2 \times [0, 1]$ denote a trivial interval bundle over a torus. Pick diffeomorphisms

$$\phi_0: T^2 \times \{0\} \rightarrow \partial A \text{ and } \phi_1: T^2 \times \{1\} \rightarrow \partial W$$

such that the glued up manifold

$$A \bigcup_{\phi_0} (T^2 \times [0, 1]) \bigcup_{\phi_1} W$$

is homeomorphic to L .

Consider the metrics $\phi_0^* d_\varepsilon$ and $\phi_1^* g$ on the boundary of $T^2 \times [0, 1]$. We now apply a lemma from [Bes].

Lemma 8.4 ([Bes, Appendix A.2]). *For any $n > 0$ there exist $\alpha_n, \varepsilon_n > 0$ and a Riemannian metric σ_n on $T^2 \times [0, 1]$ such that:*

1. *The curvature of σ_n is bounded between $-1/n$ and $1/n$.*
2. *The volume of $(T^2 \times [0, 1], \sigma_n)$ is less than $1/n$.*
3. *$(T^2 \times [0, 1], \sigma_n)$ has totally geodesic boundary given by $(T^2 \times \{0\}, \phi_0^* d_{\varepsilon_n})$ and $(T^2 \times \{1\}, \alpha_n \cdot \phi_1^* g)$.*
4. *α_n and ε_n go to zero as $n \rightarrow \infty$.*

We may now glue geometrically to form

$$(Y, \Delta_n) := (A, d_{\varepsilon_n}) \bigcup_{\phi_0} (T^2 \times [0, 1], \sigma_n) \bigcup_{\phi_1} (W, \alpha_n \cdot g),$$

such that:

1. $\text{Vol}(Y, \Delta_n) \rightarrow \text{Vol}(DC_M)$.
2. (Y, Δ_n) is an Alexandrov space with curvature bounded below by $-1 - c_{\varepsilon_n}^2$.
3. Y is homeomorphic to the closed hyperbolic manifold L .

There is a sequence $\eta_n \rightarrow 1$ such that the homothetically scaled spaces $(Y, \eta_n \cdot \Delta_n)$ are Alexandrov spaces with curvature bounded below by -1 . Theorem 4.1 applied to $(Y, \eta_n \cdot \Delta_n)$ and L yields the inequality

$$\text{Vol}(Y, \eta_n \cdot \Delta_n) \geq \text{Vol}(L).$$

Taking $n \rightarrow \infty$ yields the desired inequality

$$\text{Vol}(DC_M) \geq \text{Vol}(L). \quad \square$$

Using the above geometric surgery arguments, we now prove that the inequality of Theorem 8.1 holds also for pared acylindrical manifolds.

Corollary 8.5. *Let (N, P) be a compact pared acylindrical 3-manifold. Let (M_g, m_g) be a hyperbolic 3-manifold such that C_{M_g} is homeomorphic to $N \setminus P$ and the boundary of the convex core $\partial C_{M_g} \subset M_g$ is totally geodesic. For all $(M, m) \in H(N, P)$,*

$$\text{Vol}(C_M) \geq \text{Vol}(C_{M_g}).$$

Proof. DC_{M_g} is a finite volume hyperbolic manifold. By repeating the arguments from the beginning of Case 2 above, it follows that DC_{M_g} is obtained topologically by performing a (possibly empty) set of Dehn surgeries on the manifold DC_M . (Note that since DC_{M_g} is not compact, Dehn surgery is not performed on every end of DC_M . The ends corresponding to the cusps of DC_{M_g} are not changed.)

By performing an infinite sequence of *hyperbolic* Dehn surgeries on DC_{M_g} we obtain a sequence of *closed* hyperbolic manifolds L_k such that $\text{Vol}(L_k) \nearrow \text{Vol}(DC_{M_g})$ [NZ]. For each k , a manifold homeomorphic to L_k can be obtained from DC_M by performing an appropriate topological Dehn surgery on each end of DC_M . By repeating the geometric surgery arguments of Case 2 above, the closed manifold Y_k obtained by these Dehn surgeries on DC_M can be given a sequence of metrics δ_k^n such that:

1. $\lim_{n \rightarrow \infty} \text{Vol}(Y_k, \delta_k^n) \rightarrow \text{Vol}(DC_M)$.
2. (Y_k, δ_k^n) is an Alexandrov space with curvature bounded below by -1 .

3. Y_k is homeomorphic to the closed hyperbolic manifold L_k .

Applying Theorem 4.1 to the sequence $\{(Y_k, \delta_k^n)\}_n$ and L_k yields

$$\text{Vol}(DC_M) \geq \text{Vol}(L_k).$$

Taking $k \rightarrow \infty$ yields

$$\text{Vol}(DC_M) \geq \text{Vol}(DC_{M_g}). \quad \square$$

8.2. Cone-manifolds

Theorem 8.6. *Let Z be compact n -dimensional ($n \geq 3$) cone-manifold built with simplices of constant curvature $K \geq -1$. Assume all its cone angles are $\leq 2\pi$. Let M_{hyp} be a closed hyperbolic n -manifold. If $f: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence then*

$$\text{Vol}(Z) \geq \text{Vol}(M_{\text{hyp}})$$

with equality if and only if f is homotopic to an isometry.

Proof. Since $K \geq -1$ and all the cone angles of Z are $\leq 2\pi$, this implies Z is an Alexandrov space with curvature bounded below by -1 [BGP, p. 7]. Therefore, by Theorem 2.7, the volume growth entropy of \tilde{Z} is less than or equal to $(n-1)$. Applying Theorems 4.1 and 7.1 proves the theorem. \square

8.3. Alexandrov spaces

Theorem 8.7. *Let Z be a compact n -dimensional ($n \geq 3$) Alexandrov space with curvature bounded below by -1 . Let M_{hyp} be a closed hyperbolic n -manifold. If $f: Z \rightarrow M_{\text{hyp}}$ is a homotopy equivalence then*

$$\text{Vol}(Z) \geq \text{Vol}(M_{\text{hyp}}).$$

Proof. Otsu and Shioya proved that a finite dimensional Alexandrov space with curvature bounded below by -1 is almost everywhere Riemannian [OS]. By Theorem 2.7, the volume growth entropy of \tilde{Z} is $\leq (n-1)$. The theorem now follows from Theorem 4.1. \square

8.4. n -manifolds with boundary. The argument used in Section 8.1 generalizes immediately to prove a version of the Besson–Courtois–Gallot theorem for convex Riemannian n -manifolds with boundary. (For the definition of convex Riemannian manifolds with boundary, see Section 2.7.) A good example of a convex Riemannian manifold with boundary is a convex core with positive volume.

Perelman’s Doubling theorem [P, Theorem 5.2] is used in the proof of Theorem 8.9. Unfortunately, the well known preprint [P] remains unpublished.

Theorem 8.8 ([P, Theorem 5.2]). *Metrically doubling any Alexandrov space with curvature bounded below by k across its boundary produces an Alexandrov space with curvature bounded below by k .*

Theorem 8.9. *Let Z be a compact convex Riemannian n -manifold with boundary ($n \geq 3$). Assume the sectional curvature of $\text{int}(Z)$ is bounded below by -1 . Let Y_{geod} be a compact convex hyperbolic n -manifold with totally geodesic boundary. Let $f: (Z, \partial Z) \rightarrow (Y_{\text{geod}}, \partial Y_{\text{geod}})$ be a homotopy equivalence of pairs. Then*

$$\text{Vol}(Z) \geq \text{Vol}(Y_{\text{geod}}),$$

with equality if and only if f is homotopic to an isometry.

Proof. The homotopy equivalence of a pairs $f: (Z, \partial Z) \rightarrow (Y_{\text{geod}}, \partial Y_{\text{geod}})$ can be extended to a homotopy equivalence between the doubled manifolds $f: DZ \rightarrow DY_{\text{geod}}$. We know the sectional curvature of $\text{int}(Z)$ is bounded below by -1 . So Z is an Alexandrov space with curvature bounded below by -1 . By Perelman's Theorem 8.8, DZ is also an Alexandrov space with curvature bounded below by -1 . Theorem 2.7 then implies the volume growth entropy of \widehat{DZ} is not greater than $h(\mathbb{H}^n) = (n - 1)$. Applying Theorem 4.1 yields the desired inequality.

Assume the inequality is an equality. Theorem 7.1 then implies the natural map $F: DZ \rightarrow DY_{\text{geod}}$ is an isometry. As before, let σ be the boundary preserving isometric involution of DZ and DY_{geod} . Since f is σ -equivariant, F is σ -equivariant (see Remark 4.2). Therefore, by Theorem 7.1, $F: Z \rightarrow Y_{\text{geod}}$ is an isometry. \square

References

- [AMR] I. Aitchison, S. Matsumotoi, J. H. Rubinstein, Immersed surfaces in cubed manifolds. *Asian J. Math.* **1** (1) (1997), 85–95. [Zbl 0935.57033](#) [MR 1480991](#)
- [Bes] L. Bessières, Sur le volume minimal des variétés ouvertes. *Ann. Inst. Fourier* **50** (3) (2000), 965–980. [Zbl 0971.53027](#) [MR 1779901](#)
- [BBI] D. Burago, Y. Burago, and S. Ivanov, *A Course in Metric Geometry*. Grad. Stud. Math. 33, Amer. Math. Soc., Providence, R.I., 2001. [Zbl 0981.51016](#) [MR 1835418](#)
- [BCG1] G. Besson, G. Courtois, and S. Gallot, Entropies et rigidités des espaces localement symétriques de courbure strictement négative. *Geom. Funct. Anal.* **5** (5) (1995), 731–799. [Zbl 0851.53032](#) [MR 1354289](#)
- [BCG2] G. Besson, G. Courtois, and S. Gallot, Minimal entropy and Mostow's rigidity theorems. *Ergodic Theory Dynam. Systems* **16** (1996), 623–649. [Zbl 0887.58030](#) [MR 1406425](#)
- [BCS] J. Boland, C. Connell, and J. Souto, Minimal volume and minimal entropy. In J. Souto Clement *Geometric structures on 3-manifolds and their deformations*, Bonner Math. Schriften 342. Bonn 2001. [Zbl 0998.57040](#) [MR 1934287](#)

- [BGP] Y. Burago, M. Gromov, and G. Perel'man, A. D. Aleksandrov spaces with curvatures bounded below. *Uspekhi Mat. Nauk* **47** (2) (1992), 3–51; English transl. *Russian Math. Surveys* **47** (2) (1992), 1–58. [Zbl 0802.53018](#) [MR 1185284](#)
- [BGS] W. Ballmann, M. Gromov, and V. Schroeder, *Manifolds of nonpositive curvature*. Progr. Math. 61, Birkhäuser, Boston, MA, 1985. [Zbl 0591.53001](#) [MR 0823981](#)
- [BM] M. Burger and S. Mozes, CAT(-1)-spaces, divergence groups and their commensurators. *J. Amer. Math. Soc.* **9** (1) (1996), 57–93. [Zbl 0847.22004](#) [MR 1325797](#)
- [Bon] F. Bonahon, A Schläfli-type formula for convex cores of hyperbolic 3-manifolds. *J. Differential Geom.* **50** (1998), 25–58. [Zbl 0944.53021](#) [MR 1678473](#)
- [BP] R. Benedetti and C. Petronio, *Lectures on Hyperbolic Geometry*. Universitext, Springer-Verlag, Berlin 1992. [Zbl 0768.51018](#) [MR 1219310](#)
- [CHK] D. Cooper, C. Hodgson, and S. Kerckhoff, *Three-dimensional Orbifolds and Cone-Manifolds*. MSJ Memoirs 5, Mathematical Society of Japan, Tokyo 2000. [Zbl 0955.57014](#) [MR 1778789](#)
- [EM] D. Epstein and A. Marden, Convex hulls in hyperbolic space, a theorem of Sullivan, and measured pleated surfaces. In *Analytical and Geometric Aspects of Hyperbolic Spaces*, London Math. Soc. Lecture Note Ser. 111, Cambridge University Press, Cambridge 1987, 3–92. [Zbl 0612.57010](#) [MR 0903852](#)
- [EG] L. Evans and R. Gariepy, *Measure Theory and Fine Properties of Functions*. Stud. Adv. Math., CRC Press, Boca Raton 1992. [Zbl 0804.28001](#) [MR 1158660](#)
- [F] R. Feres, The minimal entropy theorem and Mostow rigidity, after G. Besson, G. Courtois, and S. Gallot. Unpublished.
- [GH] E. Ghys and P. de la Harpe (eds.), *Sur les Groupes Hyperboliques d'après Mikhael Gromov*. Progr. Math. 83, Birkhäuser, Boston, MA, 1990. [Zbl 0731.20025](#) [MR 1086648](#)
- [Gr] M. Gromov, *Metric Structures for Riemannian and Non-Riemannian Spaces*. Progr. Math. 152, Birkhäuser, Boston, MA, 1999. [Zbl 0953.53002](#) [MR 0682063](#)
- [J] W. Jaco, *Lectures on Three-Manifold Topology*. CBMS Reg. Conf. Ser. Math. 43, Amer. Math. Soc., Providence, R.I., 1980. [Zbl 0433.57001](#) [MR 0565450](#)
- [L] B. Leeb, 3-manifolds with(out) metrics of nonpositive curvature. *Invent. Math.* **122** (2) (1995), 277–289. [Zbl 0840.53031](#) [MR 1358977](#)
- [M] J. Morgan, On Thurston's uniformization theorem for three-dimensional manifolds. In *The Smith Conjecture*, ed. by J. Morgan and H. Bass, Pure Appl. Math. 112, Academic Press, Orlando, FL, 1984, 37–125. [Zbl 0599.57002](#) [MR 0758464](#)
- [NZ] W. Neumann and D. Zagier, Volumes of hyperbolic 3-manifolds. *Topology* **24** (3) (1985), 307–332. [Zbl 0815482](#) [MR 0589.57015](#)
- [OS] Y. Otsu and T. Shioya, The Riemannian structure of Alexandrov spaces. *J. Differential Geom.* **39** (1994), 629–658. [Zbl 0808.53061](#) [MR 1274133](#)
- [P] G. Perel'man, Alexandrov spaces with curvatures bounded from below II. Preprint, 1991.
- [RS] M. Reed and B. Simon, *Methods of modern mathematical physics. I: Functional analysis*. Academic Press, New York 1980. [Zbl 0459.46001](#) [MR 0493419](#)

- [S] P. Storm, Minimal volume Alexandrov spaces. *J. Differential Geom.* **61** (2002), 195–225. [Zbl 1070.53023](#) [MR 1972145](#)
- [Th1] W. Thurston, *The topology and geometry of 3-manifolds*. Princeton University, Lecture Notes, 1976–79.
- [Th2] W. Thurston, Hyperbolic geometry and 3-manifolds. In *Low-dimensional topology* (Bangor, 1979), London Math. Soc. Lecture Note Ser. 48, Cambridge University Press, Cambridge, New York 1982, 9–25. [Zbl 0483.57007](#) [MR 0662424](#)
- [Th3] W. Thurston, Hyperbolic structures on 3-manifolds. I. Deformation of acylindrical manifolds. *Ann. of Math.* (2) **124** (2) (1986), 203–246. [Zbl 0668.57015](#) [MR 0855294](#)

Received December 2, 2002; revised May 31, 2005

Peter A. Storm, Stanford University, Department of Mathematics, Bldg. 380, 450 Serra Mall,
Stanford, CA 94305-2125, U.S.A.

E-mail: pastorm@stanford.edu