

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 83 (2008)

Artikel: Cluster mutation via quiver representations
Autor: Buan, Aslak Bakke / Marsh, Robert J. / Reiten, Idun
DOI: <https://doi.org/10.5169/seals-99026>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Cluster mutation via quiver representations

Aslak Bakke Buan*, Robert J. Marsh and Idun Reiten

*Dedicated to Maria Inés Platzeck
on the occasion of her 60th birthday*

Abstract. Matrix mutation appears in the definition of cluster algebras of Fomin and Zelevinsky. We give a representation theoretic interpretation of matrix mutation, using tilting theory in cluster categories of hereditary algebras. Using this, we obtain a representation theoretic interpretation of cluster mutation in case of acyclic cluster algebras.

Mathematics Subject Classification (2000). Primary 16G20, 16G70; Secondary 13A99, 05E99.

Keywords. Cluster algebras, cluster categories, tilting theory.

Introduction

This paper was motivated by the interplay between the recent development of the theory of cluster algebras defined by Fomin and Zelevinsky in [FZ1] (see [Z] for an introduction) and the subsequent theory of cluster categories and cluster-tilted algebras [BMRRT], [BMR]. Our main results can be considered to be interpretations within cluster categories of important concepts in the theory of cluster algebras.

Cluster algebras were introduced in order to explain the connection between the canonical basis of a quantised enveloping algebra as defined by Kashiwara and Lusztig and total positivity for algebraic groups. It was also expected that cluster algebras should model the classical and quantised coordinate rings of varieties associated to algebraic groups – see [BFZ] for an example of this phenomenon (double Bruhat cells). Cluster algebras have been used to define generalisations of the Stasheff polytopes (associahedra) to other Dynkin types [CFZ], [FZ3]; consequently there are likely to be interesting links with operad theory. They have been used to provide the solution [FZ3] of a conjecture of Zamolodchikov concerning Y -systems, a class of

*Aslak Bakke Buan was supported by a grant from the Norwegian Research Council

functional relations important in the theory of the thermodynamic Bethe Ansatz, as well as solution [FZ4] of various recurrence problems involving Laurent polynomials, including a conjecture of Gale and Robinson on the integrality of generalised Somos sequences. Here the remarkable Laurent properties of the distinguished generators of a cluster algebra play an important role. Cluster algebras have also been related to Poisson geometry [GSV1], Teichmüller spaces [GSV2], positive spaces and stacks [FG], dual braid monoids [BES], ad-nilpotent ideals of a Borel subalgebra of a simple Lie algebra [P] as well as representation theory, see amongst others [BMRRT], [BMR], [CC], [CCS1], [CCS2], [MRZ].

A cluster algebra (without coefficients) is defined via a choice of free generating set $\underline{x} = \{x_1, \dots, x_n\}$ in the field \mathcal{F} of rational polynomials over \mathbb{Q} and a skew-symmetrizable integer matrix B indexed by the elements of \underline{x} . The pair (\underline{x}, B) , called a seed, determines the cluster algebra as a subring of \mathcal{F} . More specifically, for each $i = 1, \dots, n$, a new seed $\mu_i(\underline{x}, B) = (\underline{x}', B')$ is obtained by replacing x_i in \underline{x} by $x_i' \in \mathcal{F}$, where x_i' is obtained by a so-called *exchange multiplication rule* and B' is obtained from B by applying so-called *matrix mutation* at row/column i . Mutation in any direction is also defined for the new seed, and by iterating this process one obtains a countable (sometimes finite) number of seeds. For a seed (\underline{x}, B) , the set \underline{x} is called a *cluster*, and the elements in \underline{x} are called *cluster variables*. The desired subring of \mathcal{F} is by definition generated by the cluster variables.

It is an interesting problem to try to find a categorical/ module theoretical setting with a nice interpretation of the concepts of clusters and cluster variables, and of the matrix mutation and multiplication exchange rule for cluster variables. For the case of acyclic cluster variables so-called *cluster categories* were introduced as a candidate for such a model [BMRRT]. Skew-symmetric matrices are in one-one correspondence with finite quivers with no loops or cycles of length two, and the corresponding cluster algebra is called *acyclic* if there is a seed (\underline{x}, B) such that B corresponds to a quiver Q without oriented cycles. There is then, for a field K , an associated finite dimensional path algebra KQ . The corresponding cluster category \mathcal{C} is defined in [BMRRT] as a certain quotient of the bounded derived category of KQ , which is shown to be canonically triangulated by [K]. In [BMRRT] (cluster-)tilting theory is developed in \mathcal{C} , with emphasis on connections to cluster algebras. The analogs of clusters are (cluster-)tilting objects, and the analogs of cluster variables are exceptional objects. In case Q is a Dynkin quiver, it was shown in [BMRRT] that there is a one-one correspondence between cluster variables and exceptional objects in \mathcal{C} (in this case all indecomposables are exceptional) which takes clusters to tilting objects. This was conjectured to hold more generally.

In this paper we show that also the matrix mutation for cluster algebras has a nice interpretation within cluster categories, in terms of the associated cluster-tilted algebras, investigated in [BMR]. Cluster-tilted algebras are endomorphism algebras of tilting objects in cluster categories. It follows from our results that the quivers of

the cluster-tilted algebras arising from a given cluster category are exactly the quivers corresponding to the exchange matrices of the associated cluster algebra. This has further applications to cluster algebras (see [BR]). Another main result of this paper is an interpretation within cluster categories of the exchange multiplication rule of an (acyclic) cluster algebra. So, together with the results from [BMRRT], all the major ingredients involved in the construction of acyclic cluster algebras have now been interpreted in the cluster category.

Tilting theory for hereditary algebras has been a central topic within representation theory since the early eighties. This involves the study of tilted algebras, and various generalisations. An important motivation for this theory was to compare the representation theory of hereditary algebras with the representation theory of other homologically more complex algebras. The main result of [BMR] is also in this spirit, showing a close connection between the representation theory of cluster-tilted algebras and hereditary algebras. It is the hope of the authors that our “dictionary” also can be used to obtain further developments in the representation theory of finite dimensional algebras. Also new links between this field and other fields of mathematics can be expected, having in mind the influence of cluster algebras on other areas.

In [CCS1] an alternative description of the cluster category is given for type A . The cluster category was also the motivation for a Hall-algebra type definition of a cluster algebra of finite type [CC], [CK].

The paper is organised as follows. In Section 1 we give some preliminaries, allowing us to state the main result more precisely. Most of the necessary background on cluster algebras is however postponed until later (Section 6), since most of the paper does not involve cluster algebras. In Section 2 we prove the following: If Γ is cluster-tilted, then so is $\Gamma/\Gamma e\Gamma$ for an idempotent e in Γ . This is an essential ingredient in the proof of the main result, and also an interesting fact in itself. In Section 3 some consequences of this are given. In Section 4 we prepare for the proof of our main result. This involves studying cluster-tilted algebras of rank 3, and a crucial result of Kerner [Ke] on hereditary algebras. The main result is proved in Section 5, while Section 6 deals with the connection to cluster algebras, including necessary background.

The results of this paper have been presented at conferences in Uppsala (June 2004), Mexico (August 2004) and Northeastern University (October 2004).

The first named author spent most of 2004 at the University of Leicester, and would like to thank the Department of Mathematics, and especially Robert J. Marsh, for their kind hospitality. We would like to thank the referee for pointing out an error in an earlier version of this paper, and Bernhard Keller and Otto Kerner for helpful comments and conversations.

1. Preliminaries

1.1. Finite-dimensional algebras. In this section let Λ be a finite dimensional K -algebra, where K is a field. Then $1_\Lambda = e_1 + e_2 + \cdots + e_n$, where all e_i are primitive idempotents. We always assume that Λ is basic, that is, $\Lambda e_i \not\cong \Lambda e_j$ when $i \neq j$. There are then (up to isomorphism) n indecomposable projective Λ -modules, given by Λe_i , and n simple modules, given by $\Lambda e_i / \underline{r} e_i$, where \underline{r} is the Jacobson radical of Λ .

If K is an algebraically closed field, then there is a finite quiver Q , such that Λ is isomorphic to KQ/I , where KQ is the path-algebra, and I is an admissible ideal, that is there is some m , such that $\underline{r}^m \subseteq I \subseteq \underline{r}^2$. We call Q the quiver of Λ . In case Λ is hereditary, the ideal I is 0. For the rest of this paper, except in Section 2, we always assume that K is algebraically closed.

The category $\text{mod } \Lambda$ of finite dimensional left Λ -modules is an abelian category having almost split sequences. In case Λ is hereditary there is a translation functor τ , which is defined on all modules with no projective (non-zero) direct summands.

The bounded derived category of Λ , denoted $D^b(\text{mod } \Lambda)$, is a triangulated category, with suspension given by the shift-functor $[1]$, which is an autoequivalence. We denote its inverse by $[-1]$. In this paper, we only consider derived categories of hereditary algebras H . They have an especially nice structure, since the indecomposable objects are given by shifts of indecomposable modules. In this case we also have a translation functor $\tau : D^b(\text{mod } H) \rightarrow D^b(\text{mod } H)$, extending the functor mentioned above. We have almost split triangles $A \rightarrow B \rightarrow C \rightarrow$ in $D^b(\text{mod } H)$, where $\tau C = A$, for each indecomposable C in $\mathcal{D} = D^b(\text{mod } H)$. We also have the formula $\text{Hom}_{\mathcal{D}}(X, \tau Y) \simeq D \text{Ext}_{\mathcal{D}}^1(Y, X)$, see [H]. Here D denotes the ordinary duality for finite-dimensional algebras. Let H be a hereditary finite-dimensional algebra. Then a module T in $\text{mod } H$ is called a *tilting module* if $\text{Ext}_H^1(T, T) = 0$ and T has, up to isomorphism, n indecomposable direct summands. The endomorphism ring $\text{End}_H(T)^{\text{op}}$ is called a *tilted algebra*.

For an algebra Γ , it will be convenient to call the quiver of Γ , what is usually called the quiver of Γ^{op} .

See [ARS] and [R] for further information on the representation theory of finite dimensional algebras and almost split sequences.

1.2. Approximations. Let \mathcal{E} be an additive category, and \mathcal{X} a full subcategory. Let E be an object in \mathcal{E} . If there is an object X in \mathcal{X} , and a map $f : X \rightarrow E$, such that for every object X' in \mathcal{X} and every map $g : X' \rightarrow E$, there is a map $h : X' \rightarrow X$, such that $g = fh$, then f is called a right \mathcal{X} -approximation [AS]. The approximation map $f : X \rightarrow E$ is called *minimal* if no non-zero direct summand of X is mapped to 0. The concept of (minimal) left \mathcal{X} -approximations is defined dually. If there is a field K , such that $\text{Hom}_{\mathcal{E}}(X, Y)$ is finite dimensional over K , for all $X, Y \in \mathcal{E}$, and

if $\mathcal{X} = \text{add } M$ for an object M in \mathcal{E} , then (minimal) left and right \mathcal{X} -approximations always exist. Here, for an object M in any additive category, we let $\text{add } M$ denote the smallest full additive subcategory closed under direct sums and containing M .

1.3. Cluster categories and cluster-tilted algebras. We remind the reader of the basic definitions and results from [BMRRT]. Let H be a hereditary algebra, and let $\mathcal{D} = D^b(\text{mod } H)$ be the bounded derived category.

The *cluster category* is defined as the orbit category $\mathcal{C}_H = \mathcal{D}/F$, where $F = \tau^{-1}[1]$. The objects of \mathcal{C}_H are the same as the objects of \mathcal{D} , but maps are given by

$$\text{Hom}_{\mathcal{C}}(X, Y) = \coprod_i \text{Hom}_{\mathcal{D}}(X, F^i Y).$$

Let $Q: \mathcal{D} \rightarrow \mathcal{C}$ be the canonical functor. We often denote $Q(X)$ by \hat{X} , and use the same notation for maps. Let \hat{X} be an indecomposable object in the cluster-category. We call $\text{mod } H \vee \text{add } H[1] = \text{mod } H \vee H[1]$ the *standard domain*.

There is (up to isomorphism) a unique object X in $\text{mod } H \vee H[1] \subseteq \mathcal{D}$ such that $Q(X) = \hat{X}$.

Assume X_1, X_2 are indecomposable in the standard domain, then a map $\hat{f}: \hat{X}_1 \rightarrow \hat{X}_2$, can uniquely be written as a sum of maps $\hat{f}_1 + \hat{f}_2 + \cdots + \hat{f}_r$, such that \hat{f}_i is in $\text{Hom}_{\mathcal{D}}(X_1, F^{d_i} X_2)$, for integers d_i . In this case d_i is called the *degree* of \hat{f}_i .

The following summarises properties about cluster categories that will be freely used later.

Theorem 1.1. *Let H be a hereditary algebra, and \mathcal{C}_H the cluster category of H . Then*

- (a) \mathcal{C}_H is a Krull–Schmidt category and Q preserves indecomposable objects;
- (b) \mathcal{C}_H is triangulated and Q is exact;
- (c) \mathcal{C}_H has AR-triangles and Q preserves AR-triangles.

Proof. (b) is due to Keller [K], while (a) and (c) are proved in [BMRRT]. □

Let us now fix a hereditary algebra H , and assume it has, up to isomorphism, n simple modules. A *cluster tilting object* (or for short, tilting object) in the cluster category is an object T with $\text{Ext}_{\mathcal{C}}^1(T, T) = 0$, and with n non-isomorphic indecomposable direct summands. Two tilting objects T and T' are said to be equivalent if and only if $\text{add } T = \text{add } T'$. We only consider tilting objects up to equivalence, and therefore we always assume that if $T = \coprod_i T_i$ is a tilting object, with each T_i indecomposable, then $T_i \not\cong T_j$ for $i \neq j$.

There is a natural embedding of the module category into the bounded derived category, which extends to an embedding of the module category into \mathcal{C} . This embedding is in general not full. It was shown in [BMRRT] that the image of a tilting

module in $\text{mod } H$ is a tilting object in \mathcal{C}_H . It was also shown that if we choose a tilting object T in \mathcal{C}_H , then there is a hereditary algebra H' and an equivalence $D^b(H') \rightarrow D^b(H)$, such that T is the image of a tilting module, under the embedding of $\text{mod } H'$ into $\mathcal{C}_{H'} \simeq \mathcal{C}_H$.

If $\bar{T} \amalg X$ is a tilting object, and X is indecomposable, then \bar{T} is called an almost complete tilting object.

The following was shown in [BMRRT].

Theorem 1.2. *Let \bar{T} be an almost complete tilting object in \mathcal{C}_H . Then there are exactly two complements M and M^* . There are uniquely defined non-split triangles*

$$M^* \rightarrow B \rightarrow M \rightarrow,$$

and

$$M \rightarrow B' \rightarrow M^* \rightarrow.$$

The maps $B \rightarrow M$ and $B' \rightarrow M^*$ are minimal right add \bar{T} -approximations, and the maps $M^* \rightarrow B$ and $M \rightarrow B'$ are minimal left add \bar{T} -approximations.

The endomorphism ring $\text{End}_{\mathcal{C}}(T)^{\text{op}}$ of a tilting object in \mathcal{C} is called a *cluster-tilted algebra*. Using the notation of Theorem 1.2, we want to compare the quivers of the endomorphism rings $\Gamma = \text{End}_{\mathcal{C}}(\bar{T} \amalg M)^{\text{op}}$ and $\Gamma' = \text{End}_{\mathcal{C}}(\bar{T} \amalg M^*)^{\text{op}}$.

1.4. Matrix mutation. Let $X = (x_{ij})$ be an $n \times n$ -skew-symmetric matrix with integer entries. Choose $k \in \{1, 2, \dots, n\}$ and define a new matrix $\mu_k(X) = X' = (x'_{ij})$ by

$$x'_{ij} = \begin{cases} -x_{ij} & \text{if } k = i \text{ or } k = j, \\ x_{ij} + \frac{|x_{ik}|x_{kj} + x_{ik}|x_{kj}|}{2} & \text{otherwise.} \end{cases}$$

The matrix $\mu_k(X) = X'$ is called the mutation of X in direction k , and one can show that

- $\mu_k(X)$ is skew-symmetric, and
- $\mu_k(\mu_k(X)) = X$.

Matrix mutation appears in the definition of cluster algebras by Fomin and Zelevinsky [FZ1].

1.5. Main result. At this point, we have the necessary notation to state the main result of this paper. There are no loops in the quiver of a cluster-tilted algebra [BMRRT], and we also later show that there are no (oriented) cycles of length two. It follows that one can assign to Γ a skew-symmetric integer matrix X_{Γ} . Actually, there is a 1-1 correspondence between the skew-symmetric integer matrices and quivers

with no loops and no cycles of length two. Fixing an ordering of the vertices of the quiver, this 1-1 correspondence determines mutations μ_k also on finite quivers (with no loops and no cycles of length 2). The following will be proved in Section 5. The notation is as earlier in this section, especially the field K is algebraically closed.

Theorem 1.3. *Let \bar{T} be an almost complete tilting object with complements M and M^* and let $\Gamma = \text{End}_{\mathcal{C}}(\bar{T} \amalg M)^{\text{op}}$ and $\Gamma' = \text{End}_{\mathcal{C}}(\bar{T} \amalg M^*)^{\text{op}}$. Let k be the vertex of Γ corresponding to M . Then the quivers Q_{Γ} and $Q_{\Gamma'}$, or equivalently the matrices $X_{\Gamma} = (x_{ij})$ and $X_{\Gamma'} = (x'_{ij})$, are related by the formulas*

$$x'_{ij} = \begin{cases} -x_{ij} & \text{if } k = i \text{ or } k = j, \\ x_{ij} + \frac{|x_{ik}|x_{kj} + x_{ik}|x_{kj}|}{2} & \text{otherwise.} \end{cases}$$

This is the central result from which the connections with cluster algebras mentioned in the introduction follow. An independent proof of Theorem 1.3 in the case of finite representation type is given by Caldero, Chapoton and Schiffler [CCS2].

2. Factors of cluster-tilted algebras

In this section, our main result is that for any cluster-tilted algebra Γ , and any primitive idempotent e , the factor-algebra $\Gamma/\Gamma e\Gamma$ is in a natural way also a cluster-tilted algebra. This will give us a powerful reduction-technique, which is of independent interest, and which we use in the proof of our main result in this paper.

Suppose that Γ is the endomorphism algebra of a tilting object T in the cluster category corresponding to a hereditary algebra H . The main idea of the proof is to show that if we localise $\mathcal{D} = D^b(\text{mod } H)$ at the smallest thick subcategory containing a fixed indecomposable summand M of T , then we obtain a category triangle-equivalent to the derived category of a hereditary algebra H' . The factor-algebra $\Gamma/\Gamma e\Gamma$ (where e is the primitive idempotent of Γ corresponding to M) is then shown to be isomorphic to the endomorphism algebra of a tilting object in the cluster category corresponding to H' .

2.1. Localisation of triangulated categories. We review the basics of localisation in triangulated categories. Let \mathcal{T} be a triangulated category. A subcategory \mathcal{M} of \mathcal{T} is called a *thick subcategory* of \mathcal{T} if it is a full triangulated subcategory of \mathcal{T} closed under taking direct summands.

When \mathcal{M} is a thick subcategory of \mathcal{T} , one can form a new triangulated category $\mathcal{T}_{\mathcal{M}} = \mathcal{T}/\mathcal{M}$, and there is a canonical exact functor $L_{\mathcal{M}}: \mathcal{T} \rightarrow \mathcal{T}_{\mathcal{M}}$. See [Ric] and [V] for details.

For every M' in \mathcal{M} , we have $L_{\mathcal{M}}(M') = 0$, and $L_{\mathcal{M}}$ is universal with respect to this property. We also have the following.

Lemma 2.1. *Assume \mathcal{T} is a triangulated category, and \mathcal{M} is a thick subcategory of \mathcal{T} . Then, for any map f in \mathcal{T} we have $L_{\mathcal{M}}(f) = 0$ if and only if f factors through an object in \mathcal{M} .*

We will need the following result of Verdier [V, Ch. 2, 5-3], [V2]:

Proposition 2.2. *Let \mathcal{T} be a triangulated category with thick subcategory \mathcal{M} , and let $\mathcal{T}_{\mathcal{M}}$ be the quotient category with quotient functor $L_{\mathcal{M}}: \mathcal{T} \rightarrow \mathcal{T}_{\mathcal{M}}$. Fix an object Y of \mathcal{T} . Then every morphism from an object of \mathcal{M} to Y is zero if and only if for every object X of \mathcal{T} the canonical map*

$$\mathrm{Hom}_{\mathcal{T}}(X, Y) \rightarrow \mathrm{Hom}_{\mathcal{T}_{\mathcal{M}}}(L_{\mathcal{M}}(X), L_{\mathcal{M}}(Y))$$

is an isomorphism.

In particular, we note that this implies that $L_{\mathcal{M}}$ is fully faithful on the full subcategory of \mathcal{T} with objects given by those objects of \mathcal{T} which have only zero morphisms from objects of \mathcal{M} .

2.2. Equivalences of module categories. Let H be a hereditary algebra and M an indecomposable H -module with $\mathrm{Ext}_H^1(M, M) = 0$. Then there is (up to isomorphism) a unique module E with the following properties:

- B1) E is a complement of M (that is, $E \amalg M$ is a tilting module).
- B2) For any module X in $\mathrm{mod} H$, we have that $\mathrm{Ext}_H^1(M, X) = 0$ implies also $\mathrm{Ext}_H^1(E, X) = 0$.

This is due to Bongartz [B], and the module E is sometimes called the *Bongartz-complement* of M . For a module X in $\mathrm{mod} H$, we denote by X^\perp the full subcategory of $\mathrm{mod} H$ with objects Y satisfying $\mathrm{Ext}_H^1(X, Y) = 0$. If T is a tilting module, then it is well known that $T^\perp = \mathrm{Fac} T$, where $\mathrm{Fac} T$ is the full subcategory of all modules that are factors of objects in $\mathrm{add} T$. Note that B2) can be reformulated as $M^\perp = (M \amalg E)^\perp$.

The following result can be found in [H] and [HRS].

Proposition 2.3. (a) *Assume M is an indecomposable non-projective H -module with $\mathrm{Ext}_H^1(M, M) = 0$, and let E be the complement as above. Then the endomorphism ring $H' = \mathrm{End}_H(E)^{\mathrm{op}}$ is hereditary, and $\mathrm{Hom}_H(M, E) = 0$.*

(b) *Let \mathcal{U} denote the full subcategory of $\mathrm{mod} H$ with objects X satisfying*

$$\mathrm{Hom}_H(M, X) = 0 = \mathrm{Ext}_H^1(M, X).$$

Then \mathcal{U} is an exact subcategory of $\text{mod } H$ and the functor $\text{Hom}_H(E, -)$ from $\text{mod } H$ to $\text{mod } H'$ restricts to an exact equivalence between \mathcal{U} and $\text{mod } H'$.

We note that the above result does not hold in general in the case when M is projective. For example, consider the quiver of type A_3 with vertices 1, 2 and 3 and arrows from 1 to 2 and 2 to 3. Let $M = P_2$. Then $E = P_1 \oplus P_3$ and $\text{End}_H(E)^{\text{op}}$ has three indecomposable objects while \mathcal{U} has only two. The only other complement of M is $E' = P_1 \oplus (P_2/P_3)$. Then $\text{Hom}_H(E', P_3) = 0$ although P_3 lies in \mathcal{U} and is non-zero. So also in this case the functor $\text{Hom}_H(E', -)$ from \mathcal{U} to $\text{mod } \text{End}_H(E')^{\text{op}}$ is not an equivalence. However, we will need the following result which is along similar lines for the case when M is projective.

Lemma 2.4. *Let M be an indecomposable projective H -module with corresponding idempotent $e_M \in H$. Let $H' = H/He_MH$.*

- (a) *We have $\text{Tor}_1^H(H', U) = 0$, for any object U in \mathcal{U} , where \mathcal{U} is as defined above.*
- (b) *We have that \mathcal{U} is an exact subcategory of $\text{mod } H$ and the functor $H' \otimes_H -$ from $\text{mod } H$ to $\text{mod } H'$ restricts to an exact equivalence between \mathcal{U} and $\text{mod } H'$.*

Proof. We have that \mathcal{U} is an exact subcategory of $\text{mod } H$ as in Proposition 2.3. It is easy to see that the functor $H' \otimes_H -$ is an equivalence between \mathcal{U} and $\text{mod } H'$. To see that it is exact, we consider the following projective resolution of H' as a right H -module:

$$0 \rightarrow He_MH \rightarrow H \rightarrow H' \rightarrow 0.$$

Applying $- \otimes_H U$ to this sequence, where U is an object in \mathcal{U} , we obtain the long exact sequence:

$$\text{Tor}_1^H(H, U) \rightarrow \text{Tor}_1^H(H', U) \rightarrow He_MH \otimes_H U \rightarrow H \otimes_H U \rightarrow H' \otimes_H U \rightarrow 0.$$

Since H is projective, $\text{Tor}_1^H(H, U) = 0$. We also have

$$He_MH \otimes_H U = H \otimes_H He_MU = 0$$

since $e_MU = 0$. It follows that $\text{Tor}_1^H(H', U) = 0$ and hence that $H' \otimes_H -$ is an exact functor on \mathcal{U} . \square

2.3. Localising with respect to an exceptional module. Fix a hereditary algebra H , and an indecomposable module M in $\text{mod } H$, with $\text{Ext}_H^1(M, M) = 0$.

Lemma 2.5. *Let $\mathcal{M} = \text{add}\{M[i]\}_{i \in \mathbb{Z}}$. Then \mathcal{M} is a thick subcategory in $D^b(\text{mod } H)$.*

Proof. Straightforward from the fact that any map between indecomposable objects in \mathcal{M} is either zero or an isomorphism. \square

Let $\mathcal{D} = D^b(\text{mod } H)$, let $\mathcal{D}_{\mathcal{M}}$ be the category obtained from \mathcal{D} by localising with respect to \mathcal{M} , and let $L_{\mathcal{M}}: \mathcal{D} \rightarrow \mathcal{D}_{\mathcal{M}}$ be the localisation functor. Note that \mathcal{U} is the full subcategory of $\text{mod } H$ consisting of modules X with $\text{Hom}_{\mathcal{D}}(M, X[i]) = 0$ for all i .

Theorem 2.6. *Let H be a hereditary algebra with n simple modules up to isomorphism. Let M be an indecomposable H -module with $\text{Ext}_H^1(M, M) = 0$, and let \mathcal{M} denote the thick subcategory generated by M . Then $\mathcal{D}_{\mathcal{M}}$ is equivalent to the derived category of a hereditary algebra with $n - 1$ simple modules (up to isomorphism).*

To prove this we show that $\mathcal{D}_{\mathcal{M}}$ is equivalent to the subcategory $\mathcal{D}_0 = \text{add}\{X[i] \in \mathcal{D} \mid X \in \mathcal{U}, i \in \mathbb{Z}\}$ of \mathcal{D} , and that \mathcal{D}_0 is equivalent to the derived category of a hereditary algebra with $n - 1$ simple modules. This is the content of the following three propositions. We usually denote the object $L_{\mathcal{M}}(X)$ by \tilde{X} .

Proposition 2.7. *In the setting of Theorem 2.6, the localisation functor $L_{\mathcal{M}}$ induces an equivalence $\mathcal{D}_0 \rightarrow \mathcal{D}_{\mathcal{M}}$.*

Proof. First note that by Proposition 2.2 we have that $L_{\mathcal{M}}: \mathcal{D}_0 \rightarrow \mathcal{D}_{\mathcal{M}}$ is fully faithful. Any object in $\mathcal{D}_{\mathcal{M}}$ is of the form $L_{\mathcal{M}}(X)$ for some object X in \mathcal{D} . Let \tilde{X} be an arbitrary object in $\mathcal{D}_{\mathcal{M}}$ (where X is in \mathcal{D}). Then consider the minimal right \mathcal{M} -approximation $M_X \rightarrow X$, and the induced triangle $M_X \rightarrow X \rightarrow X_0 \rightarrow$. It is clear that $\tilde{X} = \tilde{X}_0$. We claim that X_0 is in \mathcal{D}_0 , that is $\text{Hom}_{\mathcal{D}}(M, X_0[i]) = 0$ for all i . To see this, consider the long exact sequence obtained by applying $\text{Hom}_{\mathcal{D}}(M, _)$ to the triangle $M_X \rightarrow X \rightarrow X_0 \rightarrow$. For any i , the map $\text{Hom}_{\mathcal{D}}(M, M_X[i]) \rightarrow \text{Hom}_{\mathcal{D}}(M, X[i])$ is an epimorphism, since $M_X \rightarrow X$ is a right \mathcal{M} -approximation. The map is injective since any element in $\text{Hom}_{\mathcal{D}}(M, M_X[i])$ is either zero or an isomorphism. Thus, X_0 is in \mathcal{D}_0 . This completes the proof that $L_{\mathcal{M}}$ induces an equivalence $\mathcal{D}_0 \rightarrow \mathcal{D}_{\mathcal{M}}$. \square

The next result is an extension of Proposition 2.3 to the setting of derived categories.

Proposition 2.8. *In the setting of Theorem 2.6, assume M is non-projective. Let E be the Bongartz-complement of M , and let $H' = \text{End}_H(E)^{op}$. Then $\mathbb{R}\text{Hom}(E, _)$ induces an equivalence $\mathcal{D}_0 \rightarrow \mathcal{D}' = D^b(\text{mod } H')$.*

Proof. Recall that $\mathcal{U} \subset M^{\perp} = (M \amalg E)^{\perp}$. This implies that for $X \in \mathcal{U}$, we have that $\mathbb{R}\text{Hom}(E, X)$ is concentrated in degree zero with zero-term $\text{Hom}_H(E, X)$. Since $\text{Hom}_H(E, _)$ is a dense functor from \mathcal{U} to $\text{mod } H'$, and $\mathbb{R}\text{Hom}(E, _)$ commutes with $[1]$, it follows that $\mathbb{R}\text{Hom}(E, _)$ restricted to \mathcal{D}_0 is dense.

Assume X, Y are indecomposable objects in the same degree in \mathcal{D}_0 . By the above it now follows directly from Proposition 2.3 that

$$\mathrm{Hom}_{\mathcal{D}}(X, Y) \simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{R}\mathrm{Hom}(E, X), \mathbb{R}\mathrm{Hom}(E, Y)).$$

We also need to show that

$$\mathrm{Hom}_{\mathcal{D}}(X, Y[1]) \simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{R}\mathrm{Hom}(E, X), \mathbb{R}\mathrm{Hom}(E, Y[1])).$$

For this note that by Proposition 2.3, the equivalence $\mathrm{Hom}_H(E, _): \mathcal{U} \rightarrow \mathrm{mod} H'$ is exact, and that the embedding $\mathcal{U} \hookrightarrow \mathrm{mod} H$ is exact. This implies that

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}}(X, Y[1]) &\simeq \mathrm{Hom}_{\mathcal{U}}(X, Y[1]) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(\mathrm{Hom}_H(E, X), \mathrm{Hom}_H(E, Y[1])) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{R}\mathrm{Hom}(E, X), \mathbb{R}\mathrm{Hom}(E, Y[1])) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{R}\mathrm{Hom}(E, X), \mathbb{R}\mathrm{Hom}(E, Y[1])). \end{aligned}$$

Thus the restriction of $\mathbb{R}\mathrm{Hom}(E, _)$ to \mathcal{D}_0 is fully faithful. This completes the proof. \square

Proposition 2.9. *In the setting of Theorem 2.6, assume M is projective. Assume $M \simeq He_M$ for the primitive idempotent e_M in H and let $H' = H/He_M H$. Then $\mathbb{L}(H' \otimes_H -)$ induces an equivalence $\mathcal{D}_0 \rightarrow \mathcal{D}' = D^b(\mathrm{mod} H')$.*

Proof. First recall from Lemma 2.4 that $\mathrm{Tor}_1^H(H', U) = 0$ for any U in \mathcal{U} . This means that the image $\mathbb{L}(H' \otimes_H U)$ is just $H' \otimes_H U$ concentrated in degree 0.

It now follows that $\mathbb{L}(H' \otimes_H -)$ restricted to \mathcal{D}_0 is dense, by using that the functor $H' \otimes_H -: \mathcal{U} \rightarrow \mathrm{mod} H'$ is dense and that $\mathbb{L}(H' \otimes_H -)$ commutes with $[1]$.

Assume X, Y are indecomposable objects in the same degree in \mathcal{D}_0 . It follows from Lemma 2.4 that

$$\mathrm{Hom}_{\mathcal{D}}(X, Y) \simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{L}(H' \otimes_H X), \mathbb{L}(H' \otimes_H Y)).$$

We need also to show that

$$\mathrm{Hom}_{\mathcal{D}}(X, Y[1]) \simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{L}(H' \otimes_H X), \mathbb{L}(H' \otimes_H Y[1])).$$

For this recall that the embedding of \mathcal{U} into $\mathrm{mod} H$ is exact, and that $H' \otimes_H -$ is exact on \mathcal{U} by Lemma 2.4. Thus it follows that:

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}}(X, Y[1]) &\simeq \mathrm{Hom}_{\mathcal{U}}(X, Y[1]) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(H' \otimes_H X, H' \otimes_H Y[1]) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{L}(H' \otimes_H X), \mathbb{L}(H' \otimes_H Y[1])) \\ &\simeq \mathrm{Hom}_{\mathcal{D}'}(\mathbb{L}(H' \otimes_H X), \mathbb{L}(H' \otimes_H Y[1])). \end{aligned}$$

This shows that the functor is fully faithful and finishes the proof. \square

For the remainder of this section, we view the induced equivalence between $\mathcal{D}_{\mathcal{M}}$ and \mathcal{D}' as an identification.

2.4. The factor construction. As before, let M be an indecomposable H -module with $\text{Ext}_H^1(M, M) = 0$, where H is hereditary, and let E be the Bongartz complement of M . We investigate the image of an arbitrary complement \bar{T} of M under the functor $L_{\mathcal{M}}$. For an object X in \mathcal{D} , we use the notation $\tilde{X} = L_{\mathcal{M}}(X)$, as before. Note that $L_{\mathcal{M}}(\bar{T}) = L_{\mathcal{M}}(T) = \tilde{T}$, where $T = \bar{T} \amalg M$.

Lemma 2.10. *Let the notation be as above.*

- (a) $L_{\mathcal{M}}(\bar{T}) = \tilde{T}$ is in $\text{mod } H' \vee H'[1]$.
- (b) $\text{Hom}_{\mathcal{D}'}(\tilde{T}, \tilde{T}[1]) = 0$.

Proof. (a) Let $f: M' \rightarrow \bar{T}$ be a minimal right \mathcal{M} -approximation, and consider the induced triangle:

$$M' \xrightarrow{f} \bar{T} \xrightarrow{g} U_T \rightarrow \quad (1)$$

in \mathcal{D} . Since $\text{Hom}_{\mathcal{D}}(M[-1], \bar{T}) = 0$, we have that M' is in $\text{add } M$. It is clear that $\tilde{T} \simeq \tilde{U}_T$. Now, as in the proof of Proposition 2.7, we get that U_T is in \mathcal{D}_0 . Here it is clear that $U_T = U_1 \amalg U_2[1]$, where $U_1 = \text{Coker } f$ and $U_2 = \text{Ker } f$ are in \mathcal{U} . It is clear that \tilde{U}_1 and \tilde{U}_2 are H' -modules. We only need to show that \tilde{U}_2 is projective. For an arbitrary U in \mathcal{U} , we have that $\text{Ext}_H^1(U_2, U) = 0$, since $\text{Ext}_H^1(M, U) = 0$ and U_2 is a submodule of M' . Using that \mathcal{U} is an exact subcategory of $\text{mod } H$, and that the equivalence $\mathcal{U} \rightarrow \text{mod } H'$ is also exact, it follows that \tilde{U}_2 is projective in $\text{mod } H'$. Hence $\tilde{T} \simeq \tilde{U}_T$ is in $\text{mod } H' \vee H'[1]$.

(b) Using again the triangle (1) we obtain the long exact sequence

$$\text{Hom}_{\mathcal{D}}(\bar{T}, \bar{T}[1]) \rightarrow \text{Hom}_{\mathcal{D}}(\bar{T}, U_T[1]) \rightarrow \text{Hom}_{\mathcal{D}}(\bar{T}, M'[2]).$$

Therefore, $\text{Hom}_{\mathcal{D}}(\bar{T}, U_T[1]) = 0$. Now, by Proposition 2.2, it follows that $\text{Hom}_{\mathcal{D}'}(\tilde{T}, \tilde{U}_T[1]) = 0$ since $U_T[1]$ is in \mathcal{D}_0 , and hence $\text{Hom}_{\mathcal{D}'}(\tilde{T}, \tilde{T}[1]) = 0$. \square

Denote as before by F the functor $\tau^{-1}[1]: \mathcal{D} \rightarrow \mathcal{D}$. When it is not clear which derived category \mathcal{D} we are dealing with, we will denote this functor by $F_{\mathcal{D}}$ and the functor τ^{-1} by $\tau_{\mathcal{D}}^{-1}$.

Lemma 2.11. *Let H be a hereditary algebra, and let X be an object in \mathcal{D} such that X is in $\text{mod } H \vee H[1]$. Then $\text{Hom}_{\mathcal{D}}(X, X[1]) = 0$ if and only if $\text{Ext}_{\mathcal{C}_H}^1(X, X) = 0$.*

Proof. Assume X is in $\text{mod } H \vee H[1]$, and let \hat{X} be the image of X in the cluster category \mathcal{C}_H of H . Then $\text{Ext}_{\mathcal{C}}^1(\hat{X}, \hat{X}) \simeq \text{Hom}_{\mathcal{D}}(X, X[1]) \amalg D \text{Hom}_{\mathcal{D}}(X, X[1])$. This follows from $\text{Hom}_{\mathcal{D}}(X, F^{-1}X[1]) = \text{Hom}_{\mathcal{D}}(X, \tau X) \simeq D \text{Hom}_{\mathcal{D}}(X, X[1])$ and the easily checked fact that $\text{Hom}_{\mathcal{D}}(X, F^i X[1]) = 0$, whenever $i \notin \{-1, 0\}$. \square

Combining these lemmas, and using that a tilting H -module induces a tilting object in the cluster category [BMRRT, 3.3], we obtain the following.

Proposition 2.12. *Let $T = M \amalg \bar{T}$ be a tilting H -module as before. Then the image \hat{T} of \tilde{T} in the cluster category $\mathcal{C}_{H'}$ is a tilting object.*

Proof. By Lemmas 2.10 and 2.11 we have that $\text{Ext}_{\mathcal{C}_{H'}}^1(\hat{T}, \hat{T}) = 0$, and that \tilde{T} is in $\text{mod } H' \vee H'[1]$. It follows from Lemma 2.1 that \tilde{T}_a is non-zero when T_a is a direct summand in \bar{T} . Thus, it suffices to show the following: For two indecomposable non-isomorphic direct summands T_a and T_b of \bar{T} , we have $\tilde{T}_a \not\cong \tilde{T}_b$. We first show that for any map $\alpha: \tilde{T}_a \rightarrow \tilde{T}_b$, there is a map $\beta: T_a \rightarrow T_b$ such that $L_{\mathcal{M}}(\beta) = \tilde{\beta} = \alpha$. Maps in \mathcal{D}' from T_a to T_b can be viewed as certain equivalence classes of pairs (α_1, α_2) of maps in \mathcal{D} , where X is some object in \mathcal{D} , the map $\alpha_1: X \rightarrow T_a$ is a map, which when completed to a triangle $X \rightarrow T_a \rightarrow M' \rightarrow$ in \mathcal{D} , has M' in \mathcal{M} , and $\alpha_2: X \rightarrow T_b$ is a map in \mathcal{D} . Since $\tilde{M} = 0$, we can assume that M does not occur as a direct summand of X . Hence, M' must be in $\text{add } M$, since $\text{Hom}_{\mathcal{D}}(T_a, M[1]) = 0$. Then there is an exact sequence $\text{Hom}_{\mathcal{D}}(T_a, T_b) \rightarrow \text{Hom}_{\mathcal{D}}(X, T_b) \rightarrow \text{Hom}_{\mathcal{D}}(M', T_b[1]) = 0$. Thus, there is some $\beta: T_a \rightarrow T_b$ such that $\alpha_2 = \beta \circ \alpha_1$. It is easily seen that the map in \mathcal{D}' , represented by the pair (id_{T_a}, β) is equivalent to (α_1, α_2) . But the pair (id_{T_a}, β) represents $\tilde{\beta}$, by the definition of the localisation functor, and we have $\tilde{\beta} = \alpha$.

Now assume $\tilde{\beta}$ is an isomorphism. Then the corresponding triangle $T_a \rightarrow T_b \rightarrow M'' \rightarrow$ in \mathcal{D} has M'' in \mathcal{M} . It is clear that M'' actually is in $\text{add } M$, since otherwise some $M[i]$ would be a direct summand of T_a . This means that $M'' = 0$, since $\text{Hom}_{\mathcal{D}}(M, T_a[1]) = 0$. Hence β is an isomorphism. \square

We can now complete the main result of this section. Let e be the idempotent in $\Gamma = \text{End}_{\mathcal{C}}(T)^{\text{op}}$, such that $\Gamma e \simeq \text{Hom}_{\mathcal{C}}(T, M)$.

Theorem 2.13. *With the above notation, there is a natural isomorphism $\Gamma / \Gamma e \Gamma \simeq \text{End}_{\mathcal{C}_{H'}}(\hat{T})^{\text{op}}$.*

The remainder of this section will be devoted to proving this theorem. Since the cluster category is defined using the functor $F = \tau^{-1}[1]$, we need to compare $\widetilde{\tau_{\mathcal{D}}^{-1}(X)}$ and $\tau_{\mathcal{D}'}^{-1}\tilde{X}$ for an indecomposable object X in \mathcal{D} . In general $\widetilde{\tau_{\mathcal{D}}^{-1}X} \not\cong \tau_{\mathcal{D}'}^{-1}\tilde{X}$, but with extra conditions on X , sufficient for our purposes, everything behaves nicely. We

do not include our proof of the next lemma, since it has been generalised by Keller, with a simpler proof [K]. Note that the existence of minimal left almost split maps is equivalent to the existence of a left Serre functor G by [RV], and that $G = \tau^{-1}[-1]$.

Lemma 2.14. *Let X be an indecomposable object in $\mathcal{D}_0 \subset \mathcal{D}$. Then \widetilde{X} is indecomposable and $\tau_{\mathcal{D}}^{-1}X \simeq \tau_{\mathcal{D}'}^{-1}\widetilde{X}$.*

Let T_x be an indecomposable direct summand in T , not isomorphic to M . Let $M_x \rightarrow T_x$ be a minimal right add M -approximation, and consider as before the induced triangle

$$M_x \rightarrow T_x \rightarrow U_x \rightarrow$$

in \mathcal{D} , where we know that U_x is in \mathcal{D}_0 by the proof of Proposition 2.7. Thus, by applying the above lemma to each of the indecomposable direct summands of U_x , we obtain $\tau_{\mathcal{D}}^{-1}U_x \simeq \tau_{\mathcal{D}'}^{-1}\widetilde{U}_x$, and thus $\widetilde{F_{\mathcal{D}}U_x} \simeq F_{\mathcal{D}'}\widetilde{U}_x$. It is also clear that $\widetilde{U}_x \simeq \widetilde{T}_x$.

Now, pick two (not necessarily different) indecomposable direct summands T_a and T_b of \overline{T} . Construct the triangle

$$M_b \rightarrow T_b \rightarrow U_b \rightarrow,$$

as above, and apply F to it, to obtain the triangle

$$FM_b \rightarrow FT_b \rightarrow FU_b \rightarrow .$$

Apply $\text{Hom}_{\mathcal{D}}(T_a, _)$ to this triangle, to obtain the long exact sequence

$$\begin{aligned} \text{Hom}_{\mathcal{D}}(T_a, FM_b) &\rightarrow \text{Hom}_{\mathcal{D}}(T_a, FT_b) \\ &\rightarrow \text{Hom}_{\mathcal{D}}(T_a, FU_b) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, FM_b[1]). \end{aligned} \quad (2)$$

The last term vanishes, since T_a and M_b are modules. Since $M_b \rightarrow T_b$ is a minimal right add M -approximation, it follows that $FM_b \rightarrow FT_b$ is a minimal right add FM -approximation. We have that $\text{Hom}_{\mathcal{D}}(T_a, FU_b) \simeq \text{Hom}_{\mathcal{D}}(T_a, FT_b)/(FM)$, where for an object Z we use the notation $\text{Hom}(X, Y)/(Z)$ to denote the Hom-space modulo maps factoring through an object in add Z .

We claim there is an exact sequence

$$\text{Hom}_{\mathcal{D}}(T_a, FM_b)/(M) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, FT_b)/(M) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, FU_b)/(M) \rightarrow 0$$

induced from the exact sequence (2). For this it is sufficient to show that the kernel of the second map is contained in the image of the first. So let $\alpha \in \text{Hom}_{\mathcal{D}}(T_a, FT_b)/(M)$, and assume there is a commutative diagram

$$\begin{array}{ccccc} T_a & \xrightarrow{\alpha} & FT_b & \longrightarrow & FU_b \\ & \searrow \beta_1 & & \nearrow & \\ & & M' & & \end{array}$$

for some M' in $\text{add } M$. Since $\text{Hom}_{\mathcal{D}}(M, FM_b[1]) = 0$, there is $\beta_2: M' \rightarrow FT_b$, such that $M' \xrightarrow{\beta_2} FT_b \rightarrow FU_b = M' \rightarrow FU_b$. In $\text{Hom}_{\mathcal{D}}(T_a, FT_b)/(M)$ we have $\alpha = \alpha - \beta_2\beta_1$. By using the long exact sequence (2), we obtain that $\alpha = \alpha - \beta_2\beta_1$ factors through $FM_b \rightarrow FT_b$, so the sequence is exact. It follows from this that $\text{Hom}_{\mathcal{D}}(T_a, FT_b)/(M \amalg FM) \simeq \text{Hom}_{\mathcal{D}}(T_a, FU_b)/(M)$.

Let $f: M_1 \rightarrow FU_b$ be a minimal right \mathcal{M} -approximation, and complete to a triangle $M_1 \xrightarrow{f} FU_b \rightarrow (FU_b)'$. Applying $\text{Hom}_{\mathcal{D}}(T_a, _)$, we get an exact sequence

$$\text{Hom}_{\mathcal{D}}(T_a, M_1) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, FU_b) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, (FU_b)') \rightarrow \text{Hom}_{\mathcal{D}}(T_a, M_1[1]).$$

Since U_b is in degree 0 or 1, then FU_b is in degree 1, 2 or 3, so M_1 is in degree 0, 1, 2 or 3. Hence the indecomposable direct summands of $M_1[1]$ are in degree at least 1, so that $\text{Hom}_{\mathcal{D}}(T_a, M_1[1]) = 0$. Since a map $h: T_a \rightarrow FU_b$ factors through an object in \mathcal{M} if and only if it factors through the minimal right \mathcal{M} -approximation of FU_b , we get the isomorphism

$$\text{Hom}_{\mathcal{D}}(T_a, F_{\mathcal{D}}U_b)/(M) \simeq \text{Hom}_{\mathcal{D}}(T_a, (F_{\mathcal{D}}U_b)').$$

We get that this is isomorphic to $\text{Hom}_{\mathcal{D}'}(\widetilde{T}_a, \widetilde{F_{\mathcal{D}}U_b})$, since $(F_{\mathcal{D}}U_b)'$ is in \mathcal{D}_0 . By Lemma 2.14 this is isomorphic to $\text{Hom}_{\mathcal{D}'}(\widetilde{T}_a, F_{\mathcal{D}'}\widetilde{T}_b)$. We thus obtain that

$$\text{Hom}_{\mathcal{D}}(T_a, F_{\mathcal{D}}T_b)/(M \amalg FM) \simeq \text{Hom}_{\mathcal{D}'}(\widetilde{T}_a, F_{\mathcal{D}'}\widetilde{T}_b).$$

We have $\text{Hom}_{\mathcal{D}}(T_a, T_b)/(M \amalg FM) \simeq \text{Hom}_{\mathcal{D}}(T_a, T_b)/(M)$. Consider again the triangle $M_b \xrightarrow{f_b} T_b \rightarrow U_b$ in \mathcal{D} , where $f_b: M_b \rightarrow T_b$ is a minimal right \mathcal{M} -approximation. Applying $\text{Hom}_{\mathcal{D}}(T_a, _)$ gives an exact sequence

$$\text{Hom}_{\mathcal{D}}(T_a, M_b) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, T_b) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, U_b) \rightarrow \text{Hom}_{\mathcal{D}}(T_a, M_b[1]).$$


Since M_b is a module, we have $\text{Hom}_{\mathcal{D}}(T_a, M_b[1]) = 0$, and hence $\text{Hom}_{\mathcal{D}}(T_a, T_b)/(M) \simeq \text{Hom}_{\mathcal{D}}(T_a, U_b)$, which is isomorphic to $\text{Hom}_{\mathcal{D}'}(\widetilde{T}_a, \widetilde{U}_b)$ by Proposition 2.2. We obtain that:

$$\text{Hom}_{\mathcal{D}}(T_a, T_b)/(M \amalg FM) \simeq \text{Hom}_{\mathcal{D}'}(\widetilde{T}_a, \widetilde{T}_b).$$

Hence $\Gamma/\Gamma e\Gamma = \text{Hom}_{\mathcal{D}}(T, T) \amalg \text{Hom}_{\mathcal{D}}(T, FT)/(M \amalg FM) \simeq \text{Hom}_{\mathcal{D}'}(\widetilde{T}, \widetilde{T}) \amalg \text{Hom}_{\mathcal{D}'}(\widetilde{T}, F_{\mathcal{D}'}\widetilde{T})$ as vector spaces. It is straightforward to check that the map is also a ring map. Theorem 2.13 is proved.

2.5. Comparison with tilted algebras. We give an example showing that a result similar to Theorem 2.13 does not hold for tilted algebras. We would like to thank Dieter Happel for providing us with this example. There is a tilting module for

the path algebra of a Dynkin quiver of type D_5 , such that the corresponding tilted algebra Λ has the quiver



with relations $\alpha\beta = \beta\gamma - \delta\epsilon = 0$. If we let e_4 be the primitive idempotent corresponding to vertex 4, then $\Lambda/\Lambda e_4\Lambda$ is not tilted, since it has global dimension three.

It is well known that the endomorphism-ring of a partial tilting module is a tilted algebra. However, a similar result does not hold for cluster-tilted algebras. An example of this is the path algebra of an oriented 4-cycle, modulo the cube of its radical. This is a cluster-tilted algebra of type D_4 .

3. Cluster-tilted algebras of rank at most 2

In this section we apply the main result of the previous section to show that (oriented) cycles in the quiver of a cluster-tilted algebra have length at least three. For the rest of the paper, the field K is assumed to be algebraically closed. Let $T_1 \amalg T_2 \amalg \cdots \amalg T_n$ be a tilting object in the cluster category \mathcal{C} . We denote by $\delta_k(T)$ the tilting object T' obtained by exchanging T_k with the second complement of $T_1 \amalg \cdots \amalg T_{k-1} \amalg T_{k+1} \amalg \cdots \amalg T_n$. Let $\Gamma = \text{End}_{\mathcal{C}}(T)^{\text{op}}$ and $\Gamma' = \text{End}_{\mathcal{C}}(T')^{\text{op}}$ be the corresponding cluster-tilted algebras. Passing from Γ to Γ' depends on the choice of tilting object T . But we still write $\bar{\delta}_k(\Gamma) = \Gamma'$, when either it is clear from the context which tilting object T gives rise to Γ , or when this is irrelevant. We also say that Γ' is obtained from Γ by mutation at k .

From [BMRRT] we know that all tilting objects in \mathcal{C}_H can be obtained from performing a finite number of operations δ_k to H , where H is the hereditary algebra considered as a tilting object in \mathcal{C}_H .

If k is a source or a sink in the quiver of a hereditary algebra, then mutation at k coincides with so-called APR-tilting [APR] (see [BMR]), and the quiver of the mutated algebra $\bar{\delta}_k(H)$ is obtained by reversing all arrows ending or starting in k .

Lemma 3.1. *The cluster-tilted algebras of rank at most 2 are hereditary.*

Proof. This follows from the fact that any cluster-tilted algebra can be obtained by starting with a hereditary algebra, and performing a finite number of mutations. If we

start with a hereditary algebra H of rank at most 2, the algebra obtained by mutating at one of the vertices is isomorphic to H . \square

Proposition 3.2. *The quiver of a cluster-tilted algebra has no loops and no cycles of length 2.*


Proof. This follows directly from combining Lemma 3.1 with Theorem 2.13. \square

Note that the assertion of no loops was proven in [BMRRT, Cor. 6.15], while the second assertion was first proven by Gordana Todorov in case of finite representation type.

4. Cluster-tilted algebras of rank 3

In this section we specialize to connected hereditary algebras of rank 3, and the cluster-tilted algebras obtained from them. We describe the possible quivers, and give some information on the relation-spaces. Later, this will be used to show our main result for algebras of rank 3. In the proof of our main theorem, we use Theorem 2.13 to reduce to the case of rank 3. For hereditary algebras of finite representation type, there is up to derived equivalence only one connected algebra of rank 3, and thus up to equivalence only one cluster category \mathcal{C} . In this case the technically involved results of this section reduce to just checking *one case*: The only cluster-tilted algebra of rank 3 which is not hereditary is given by a quiver which is a cycle of length 3, and with the relations that the composition of any two arrows is zero.

4.1. The quivers. We consider quivers of the form




where $r > 0$, $s > 0$ and $t \geq 0$ denote the number of arrows as indicated in the above figure. For short, we denote such a quiver by Q_{rst} .

Up to derived equivalence, all connected finite dimensional hereditary algebras of rank 3 have a quiver given as above. We first note that factors of path-algebras of such quivers by non-zero admissible ideals are never cluster-tilted. We omit our original proof of this fact since it is a consequence of the more general (recently proven) fact from [KR] that any cluster-tilted algebra is either hereditary or of infinite global dimension. Hence, since Q_{rst} has no oriented cycles, it follows that if it is the quiver of a cluster-tilted algebra, there can be no proper relations. So we have the following.

Lemma 4.1. *If Γ is a cluster-tilted algebra with quiver of type Q_{rst} , then Γ is hereditary.*

This has the following consequence.

Corollary 4.2. *The quiver of a non-hereditary connected cluster-tilted algebra of rank 3 is of the form*



with $r, s, t > 0$.

Proof. Combine Lemma 4.1 with Proposition 3.2. □


In view of this we refer to the cluster-tilted algebras of rank 3 which are non-hereditary as *cyclic cluster-tilted algebras*.

4.2. The relations. We first show that relations are homogeneous.

Proposition 4.3. *Let Γ be a cluster-tilted algebra of rank 3 with Jacobson radical \underline{r} . Then $\underline{r}^6 = 0$, and the relations are homogeneous.*

Proof. Without loss of generality we can assume that there is a tilting module $T = X \amalg Y \amalg Z$ for a hereditary algebra H , such that $\Gamma = \text{End}_{\mathcal{C}_H}(T)^{\text{op}}$.

Using Corollary 4.2 it is clear that we can assume that the quiver of Γ has the form



with $r, s, t > 0$.

Let $\Lambda = \text{End}_H(T)^{\text{op}}$ be the corresponding tilted algebra. There are no cycles in the quiver of a tilted algebra. We can therefore assume that there is a sink in the quiver of Λ , and we assume that this vertex corresponds to Z , that is, $\text{Hom}_H(Z, X \amalg Y) = 0$. We assume that X, Y, Z correspond to the vertices 1, 2, 3, respectively. If \hat{h} is a non-zero map in $\text{Irr}_{\text{add } T}(Z, X)$, it must be of degree 1, that is, the lifting h is in $\text{Hom}_{\mathcal{D}}(Z, FX)$. Since this holds for all maps in $\text{Irr}_{\text{add } T}(Z, X)$, any composition of 6 arrows will correspond to a map of degree ≥ 2 from an indecomposable to itself, and therefore must be the zero-map. This follows from the fact that for any indecomposable module M , we have $\text{Hom}_{\mathcal{D}}(M, F^2 M) = 0$. This gives $\underline{r}^6 = 0$.

We can assume that at least one of the arrows (irreducible maps) $X \rightarrow Y$ and at least one of the arrows $Y \rightarrow Z$ are of degree 0. Otherwise, the tilted algebra Λ would not be connected.

Now let \hat{g} be a map in $\text{Irr}_{\text{add } T}(Y, Z)$. We want to show that it must be of degree 0. Since $X \amalg Y$ is an almost complete tilting object in \mathcal{C}_H , there are exactly two complements. Denote as usual the second one by Z^* . The complement Z^* is either the image of a module or the image of an object of the form $I[-1]$ for an injective indecomposable module I . Furthermore, there is a triangle in \mathcal{C}

$$Z^* \rightarrow Y^s \rightarrow Z \rightarrow, \quad (3)$$

for some $s \geq 0$, which can be lifted to a triangle

$$F^i Z^* \xrightarrow{\begin{pmatrix} \alpha_1 \\ \alpha_2 \end{pmatrix}} Y^{s_1} \amalg (F^{-1}Y)^{s_2} \rightarrow Z \rightarrow$$

in \mathcal{D} for some integer i and with $s = s_1 + s_2$. We need to show that $s_2 = 0$. It is sufficient to show that the map $\alpha_2 = 0$. We have $s_1 \neq 0$, and thus by minimality $\alpha_1 \neq 0$. It is clear that if also $\alpha_2 \neq 0$, then $i = 0$ or $i = -1$.

Assume first $Z^* \simeq I[-1]$, then

$$\text{Hom}_{\mathcal{D}}(I[-1], F^{-1}Y) = \text{Hom}_{\mathcal{D}}(I, \tau Y) = 0,$$

so $i = 0$ gives $\alpha_2 = 0$. On the other hand, it is clear that $i = -1$ gives $\alpha_1 = 0$.

Assume now that Z^* is the image of a module. Then there is an exact sequence of modules

$$0 \rightarrow Z^* \rightarrow Y^v \rightarrow Z \rightarrow 0,$$

and since $\dim_k \text{Hom}_{\mathcal{C}}(Z, Z^*[1]) = 1$ (by [BMRRT]), it follows that the triangle (3) is induced by this sequence, and thus $s_1 = v$ and $s_2 = 0$.

Now we show that all the irreducible maps $X \rightarrow Y$ in \mathcal{C}_H are of degree 0. For this, consider the almost complete tilting object $X \amalg Z$ in \mathcal{C}_H , with complements Y and Y^* . Consider the triangle

$$Y^* \rightarrow X^r \rightarrow Y \rightarrow,$$

and the preimage in \mathcal{D} ,

$$F^i Y^* \xrightarrow{\begin{pmatrix} \beta_1 \\ \beta_2 \end{pmatrix}} X^{r_1} \amalg (F^{-1}X)^{r_2} \rightarrow Y \rightarrow.$$

We need to show that $r_2 = 0$. The case where $Y^* \simeq I[-1]$ is completely similar as for irreducible maps $Y \rightarrow Z$. In case Y^* is the image of a module, it is now more complicated since we have two possibilities. Either there is an exact sequence in $\text{mod } H$ of the form

$$0 \rightarrow Y^* \rightarrow X^a \rightarrow Y \rightarrow 0,$$

or there is an exact sequence of the form

$$0 \rightarrow Y \rightarrow Z^b \rightarrow Y^* \rightarrow 0.$$


If we are in the first case, we can use the same argument as for irreducible maps $Y \rightarrow Z$. If we are in the second case, note that $\text{Hom}_H(Y^*, X) = 0$, since $\text{Hom}_H(Z, X) = 0$. Thus, either $\beta_1 = 0$ or $\beta_2 = 0$ in our triangle. This completes the proof that all irreducible maps $X \rightarrow Y$ are induced by module maps, and thus are of degree 0.

Given that $\underline{r}^6 = 0$, the only possibility for a non-homogeneous relation must involve maps in $\underline{r}^2 \setminus \underline{r}^3$ and maps in \underline{r}^5 . But, by our description of irreducible maps, this is not possible, because it would involve a relation between maps of different degrees. \square


Fix a cyclic cluster-tilted algebra of rank 3, and fix a vertex k . Let α be an arrow ending in k , and β an arrow starting in k . If $\beta\alpha = 0$, as an element of the algebra, for any choice of α and β , then we call k a *zero vertex*.

Proposition 4.4. *Let Γ be a cyclic cluster-tilted algebra, and fix a vertex k . Then k is a zero-vertex if and only if $\bar{\delta}_k(\Gamma)$ is hereditary.*

Proof. We assume the quiver of Γ is



Let $\Gamma = \text{End}_{\mathcal{C}}(T)^{\text{op}}$, and let T_i be the direct summand of T corresponding to the vertex i . Assume that 2 is a zero-vertex. Then it is clear that $\text{Hom}_{\mathcal{C}}(T_1, T_3) = 0$, so the quiver of $\bar{\delta}_2(\Gamma)$ must be



with $t' \geq 0$. Now $\bar{\delta}_2(\Gamma)$ is hereditary, by Lemma 4.1.

Conversely, assume $\bar{\delta}_k(\Gamma)$ is hereditary. The quiver of $\bar{\delta}_k(\Gamma)$ must be as above, with $t' \geq 0$. This means $\text{Hom}_{\mathcal{C}}(T_1, T_3) = 0$, so 2 is a zero-vertex. \square

4.3. Kerner's Theorem. The following result by Kerner [Ke] turns out to be crucial for the proof of the main theorem of this section. There is a more general version of this theorem in [Ke]. We include a proof, for the convenience of the reader. This proof is also due to Kerner, and we thank him for providing us with it.

Theorem 4.5. *Let X, Y be regular indecomposable modules over a wild hereditary algebra H of rank 3. If $\text{Hom}_H(X, \tau Y) = 0$, then also $\text{Hom}_H(X, \tau^{-1}Y) = 0$.*

Proof. We first prove the following.

Lemma 4.6. *Let U be an indecomposable regular module over a wild hereditary algebra of rank 3. Then $\text{Hom}_H(U, \tau^2 U) \neq 0$.*

Proof. Assume first that $\text{Ext}_H^1(U, U) \neq 0$. Then also $\text{Hom}_H(U, \tau U) \neq 0$ by the AR-formula. Assume now $\text{Hom}_H(U, \tau^2 U) = 0$. Then also $\text{Ext}_H^1(\tau U, U) = 0$ and, by the Happel–Ringel lemma [HR], a non-zero map $f: U \rightarrow \tau U$ is either surjective or injective. In either case, $g = \tau(f) \circ f: U \rightarrow \tau^2 U$ is non-zero. This contradicts $\text{Hom}_H(U, \tau^2 U) = 0$.

Now assume $\text{Ext}_H^1(U, U) = 0$. Then by [Ho], U is quasi-simple. Thus, there is an almost split sequence $0 \rightarrow \tau U \rightarrow V \rightarrow U \rightarrow 0$, where V is indecomposable, and by [Ke2] we have $\text{End}_H(V) \simeq K$, while $\text{Ext}_H^1(V, V) \neq 0$. Applying $\text{Hom}_H(U, _)$ to the almost split sequence, we obtain the exact sequence

$$\text{Hom}_H(U, \tau U) \rightarrow \text{Hom}_H(U, V) \rightarrow \text{Hom}_H(U, U) \rightarrow \text{Ext}_H^1(U, \tau U).$$

Since $\text{Hom}_H(U, U) \rightarrow \text{Ext}_H^1(U, \tau U)$ is an isomorphism and $\text{Hom}_H(U, \tau U) = 0$, we have that also $\text{Hom}_H(U, V) = 0$. The long exact sequence obtained by applying $\text{Hom}_H(_, \tau U)$ to the almost split sequence, gives $\text{Hom}_H(V, \tau U) = 0$. Now, this gives $\text{Hom}_H(V, \tau^2 U) \neq 0$, since there is an exact sequence

$$0 \rightarrow \text{Hom}_H(V, \tau^2 U) \rightarrow \text{Hom}_H(V, \tau V) \rightarrow \text{Hom}_H(V, \tau U)$$


and the last term is zero. There is also the long exact sequence

$$0 \rightarrow \text{Hom}_H(U, \tau^2 U) \rightarrow \text{Hom}_H(V, \tau^2 U) \rightarrow \text{Hom}_H(\tau U, \tau^2 U)$$


where the last term is zero. This proves $\text{Hom}_H(U, \tau^2 U) \neq 0$. \square

Let us now complete the proof of the theorem. Let X, Y be regular indecomposable modules. It suffices to show that $\text{Hom}_H(X, Y) \neq 0$ implies $\text{Hom}_H(X, \tau^2 Y) \neq 0$. Let $z: X \rightarrow Y$ be a non-zero map. Then we can assume there is an indecomposable regular module U , such that z factors as $X \xrightarrow{p} U \xrightarrow{i} Y$, where p is surjective and i is injective. Also $\tau^2 i: \tau^2 U \rightarrow \tau^2 Y$ is injective. By Lemma 4.6, there is a non-zero map $f: U \rightarrow \tau^2 U$. The composition $\tau^2 i \circ f \circ p$ is non-zero. This completes the proof of the theorem. \square

4.4. The dimensions of relation-spaces. Let H be a connected hereditary algebra of rank 3. The following notation is used for the rest of this section. Let \bar{T} be an almost complete tilting object with complements M and M^* , and assume there are triangles as in Theorem 1.2. Let $T = \bar{T} \amalg M$ and $T' = \bar{T} \amalg M^*$ and let $\Gamma = \text{End}_{\mathcal{C}}(T)^{\text{op}}$ and $\Gamma' = \text{End}_{\mathcal{C}}(T')^{\text{op}}$. By now, we know that the quiver of Γ is either



with $r, s, t > 0$ or the quiver Q_{rst}



with $r, s > 0$ and $t \geq 0$. We let M correspond to vertex 2. Then $\bar{T} = T_B \amalg T_{B'}$ where T_B corresponds to the vertex 1 and $T_{B'}$ to 3. It is then clear that $B = (T_B)^r$ and $B' = (T_{B'})^s$. We label the vertices with the corresponding modules, then the arrows represent irreducible maps in $\text{add } T$.


We let I denote the ideal such that $\Gamma \simeq KQ/I$. In case Γ is cyclic, we say that Γ is *balanced* at the vertex 2 if

$$\dim((\text{Irr}(T_B, M) \otimes_K \text{Irr}(M, T_{B'}) \cap I) = t.$$

We will show that any vertex of a cyclic cluster-tilted algebra is either balanced or a zero-vertex. We first discuss the algebras obtained by mutating hereditary algebras.

Lemma 4.7. *Let H be a hereditary algebra with quiver Q_{rst} where $r, s > 0$ and $t \geq 0$. Then the following hold.*

- (a) *The cluster-tilted algebra $\Gamma' = \bar{\delta}_2(H)$ is balanced at the vertices 1 and 3.*
- (b) *The new vertex 2^* is a zero-vertex.*
- (c) *The quiver of Γ' is*



Proof. Part (b) and (c) follow directly from Lemma 4.1 and Proposition 4.4. Let P_i be the indecomposable projective H -module corresponding to vertex i , and S_i the simple H -module $P_i / \mathfrak{r} P_i$. Then $P_1 = S_1$ is simple. Consider $P_1 \amalg P_3$ as an almost complete tilting object. There is an exact sequence

$$0 \rightarrow P_2 \rightarrow (P_1)^s \rightarrow P_2^* \rightarrow 0,$$

such that the induced triangle in \mathcal{C} is the exchange-triangle of Theorem 1.2. Let $T' = P_1 \amalg P_2^* \amalg P_3$. Using the definition of τ , one can show that $S_2 = \tau P_2^*$, and thus $\text{Hom}_{\mathcal{D}}(P_2^*, P_3) = 0$. Since 2^* is a zero-vertex, $\text{Irr}_{\text{add } T'}(P_1, P_3) = \text{Hom}_H(P_1, P_3)$, with dimension $rs + t$. We want to compute $\text{Irr}_{\text{add } T'}(P_2^*, P_1) \otimes_{\Gamma'} \text{Irr}_{\text{add } T'}(P_1, P_3) = \text{Hom}_{\mathcal{C}}(P_2^*, P_3) \simeq \text{Hom}_{\mathcal{D}}(F^{-1}P_2^*, P_3)$. We have

$$\text{Hom}_{\mathcal{D}}(F^{-1}P_2^*, P_3) = \text{Hom}_{\mathcal{D}}(\tau P_2^*[-1], P_3) = \text{Ext}_H^1(S_2, P_3).$$

There is an exact sequence

$$0 \rightarrow (P_1)^r \rightarrow P_2 \rightarrow S_2 \rightarrow 0.$$

Apply $\text{Hom}_H(\ , P_3)$ to it, to obtain the long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Hom}_H(S_2, P_3) \rightarrow \text{Hom}_H(P_2, P_3) \\ \rightarrow \text{Hom}_H((P_1)^r, P_3) \rightarrow \text{Ext}_H^1(S_2, P_3) \rightarrow 0. \end{aligned}$$

Since $\dim \text{Hom}_H(P_2, P_3) = s$, and $\dim \text{Hom}_H((P_1)^r, P_3) = (rs + t)r$, it follows that $\dim(\text{Irr}_{\text{add } T'}(P_2^*, P_1) \otimes_{\Gamma'} \text{Irr}_{\text{add } T'}(P_1, P_3)) = r(rs + t) - s$, and therefore $\dim(\text{Irr}_{\text{add } T'}(P_2^*, P_1) \otimes_K \text{Irr}_{\text{add } T'}(P_1, P_3) \cap I) = s$, and Γ' is balanced at 3.

Now apply $\text{Hom}_H(P_1, \)$ to the exact sequence $0 \rightarrow P_2 \rightarrow (P_3)^s \rightarrow P_2^* \rightarrow 0$ to obtain the exact sequence


$$0 \rightarrow \text{Hom}_H(P_1, P_2) \rightarrow \text{Hom}_H(P_1, P_3^s) \rightarrow \text{Hom}_H(P_1, P_2^*) \rightarrow 0.$$

Since $\dim \text{Hom}_H(P_1, P_2) = r$ and $\dim \text{Hom}_H(P_1, P_3^s) = (rs + t)s$, we have


$$\dim(\text{Irr}_{\text{add } T'}(P_1, P_3) \otimes_{\Gamma'} \text{Irr}_{\text{add } T'}(P_3, P_2^*)) = \dim \text{Hom}_{\mathcal{C}}(P_1, P_2^*) = (rs + t)s - r.$$

This means $\dim(\text{Irr}_{\text{add } T'}(P_1, P_3) \otimes_K \text{Irr}_{\text{add } T'}(P_3, P_2^*) \cap I) = r$, and Γ' is balanced also at 1. \square

Proposition 4.8. *Let Γ be a non-hereditary cluster-tilted algebra with quiver*




- (a) If Γ is balanced at the vertex 2, then $\Gamma' = \bar{\delta}_2(\Gamma)$ is non-hereditary, and thus cyclic, with quiver




It is balanced at the new vertex 2^* . Each of the other vertices of Γ' is either balanced or a zero-vertex.

- (b) If Γ has a zero-vertex at 2, then $\bar{\delta}_2(\Gamma)$ is hereditary with quiver




Proof. Part (b) follows from Propositions 4.3 and 4.4.

To prove part (a), we adopt our earlier notation and conventions. Especially, $\Gamma = \text{End}_{\mathcal{C}}(T_B \amalg T_{B'} \amalg M)^{\text{op}}$, and we have the quiver



The quiver of the mutated algebra $\bar{\delta}_2(\Gamma) = \text{End}_{\mathcal{C}}(T_B \amalg T_{B'} \amalg M^*)^{\text{op}}$ is



Using that Γ is balanced at 2, and Proposition 4.3, it follows that $t' = rs - t$. Also by assumption, M does not correspond to a zero-vertex, so there is at least one non-zero composition $T_B \rightarrow M \rightarrow T_{B'}$. Therefore $rs - t > 0$.

We have $\dim(\text{Irr}_{\text{add } T'}(T_{B'}, M^*) \otimes_{\Gamma'} \text{Irr}_{\text{add } T'}(M^*, T_B)) = \dim \text{Irr}_{\text{add } T}(T_{B'}, T_B) + \dim(\text{Irr}_{\text{add } T}(T_{B'}, M) \otimes_{\Gamma'} \text{Irr}_{\text{add } T}(M, T_B)) = t + 0$. Hence $\dim(\text{Irr}_{\text{add } T'}(T_{B'}, M^*) \otimes_K \text{Irr}_{\text{add } T'}(M^*, T_B) \cap I) = rs - t$, so Γ' is balanced at 2^* .

We now proceed to show that for each of the vertices 1 and 3, Γ' is either balanced, or a zero-vertex. We assume $T_{B'}$ is not a zero-vertex in Γ' .

The tilted algebra $\Lambda = \text{End}_H(T_B \amalg T_{B'} \amalg M)^{\text{op}}$ has a unique sink. There is an induced total ordering on the triple $T_B, T_{B'}, M$, where the last element in the ordering corresponds to the sink. Also, by considering the preimage of M^* in the standard domain of \mathcal{D} , the ordering can be extended to the quadruple B', M^*, B, M . Note that we get the following four possible orderings

- $(M, T_{B'}, M^*, T_B)$,
- $(T_{B'}, M^*, T_B, M)$,
- $(M^*, T_B, M, T_{B'})$,
- $(T_B, M, T_{B'}, M^*)$.

First we show the claim for the vertex corresponding to $T_{B'}$.

Lemma 4.9. *Assume that $T_{B'}$ does not correspond to a zero-vertex in Γ' and that M^* is before T_B in the above ordering. Then $\text{Hom}_{\mathcal{D}}(T_B, \tau^{-1}M^*) = 0$.*

Proof. Since $\bar{\delta}_2(\Gamma)$ is not hereditary, we have $\text{Hom}_H(M^*, T_B) \neq 0$. Assume now that $\text{Hom}_H(T_B, \tau^{-1}M^*) \neq 0$. Assume first that T_B is regular, then M^* is also regular. In case H is tame, then there are at most two exceptional modules which are regular. This follows from the fact that H has three simples. But in case there are two exceptional modules which are regular, there is an extension between them. This gives a contradiction. In case H is wild we can apply Kerner's Theorem, which says that $\text{Hom}_H(T_B, \tau M^*) \neq 0$. We have a contradiction, since $\text{Hom}_H(T_B, \tau M^*) \simeq D \text{Ext}_H^1(M^*, T_B) = 0$.

If B is a preprojective or a preinjective module, then $\text{Hom}_H(M^*, T_B) \neq 0$ and $\text{Hom}_H(T_B, \tau^{-1}M^*) \neq 0$ implies that the map $M^* \rightarrow T_B$ is irreducible in the module-category. Thus $M^* \amalg T_B$ can be complemented to a tilting module with hereditary endomorphism ring. We have seen that the mutated algebra $\bar{\delta}_2(\Gamma)$ is by assumption not hereditary, i.e., $T_{B'}$ must correspond to a zero-vertex, so we have a contradiction to $\text{Hom}_H(B, \tau^{-1}M^*) \neq 0$ also for T_B being preprojective or preinjective. \square

Now, let $M \rightarrow B' \rightarrow M^* \rightarrow$ be the usual triangle. We recall that $\bar{T} = T_B \amalg T_{B'}$. Let $\widetilde{\text{Hom}}(T_B, M) = \text{Irr}_{\text{add } T}(T_B, M)$, let $\widetilde{\text{Hom}}(T_B, B') = \text{Irr}_{\text{add } \bar{T}}(T_B, B')$ and let $\widetilde{\text{Hom}}(T_B, M^*) = \text{Irr}_{\text{add } T'}(T_B, M^*)$. Then we claim that there is an exact sequence

$$0 \rightarrow \widetilde{\text{Hom}}(T_B, M) \xrightarrow{\alpha} \widetilde{\text{Hom}}(T_B, B') \rightarrow \widetilde{\text{Hom}}(T_B, M^*) \rightarrow 0. \quad (4)$$

It is clear from Proposition 4.3, and the fact that

$$\text{Hom}_{\mathcal{D}}(T_B, M) \rightarrow \text{Hom}_{\mathcal{D}}(T_B, B') \rightarrow \text{Hom}_{\mathcal{D}}(T_B, M^*) \rightarrow 0$$

is exact, that we only need to show that the map α is a monomorphism. We first assume M^* is a module. To prove the claim for this case, we consider the four

orderings on the quadruple $\{M, B', M^*, B\}$. For each case we show that a map in $\text{Irr}_{\text{add } T}(T_B, M)$ cannot factor via $M^*[-1]$ in \mathcal{C} .

$(M, T_{B'}, M^*, T_B)$: In this case, a map in $\text{Irr}_{\text{add } T}(T_B, M)$ is of degree 1. Assume the lifting is $f: \tau T_B[-1] \rightarrow M$. There is a non-split exact sequence $0 \rightarrow M \rightarrow B' \rightarrow M^* \rightarrow 0$. We have $\dim \text{Hom}_{\mathcal{C}}(M^*, M[1]) = 1$, by [BMRRT], and therefore $\text{Hom}_{\mathcal{D}}(FM^*, M[1]) = \text{Hom}_{\mathcal{D}}(\tau^{-1}M^*, M) = 0$. Therefore, if $f: T_B \rightarrow M$ factors through $M^*[-1]$ in \mathcal{C} , there must be a map $g: \tau T_B[-1] \rightarrow M$ in \mathcal{D} , such that there is a commutative diagram

$$\begin{array}{ccc} & \tau T_B[-1] & \\ g \swarrow & \downarrow f & \\ M^*[-1] & \longrightarrow & M. \end{array}$$

By Lemma 4.9, we have that $\text{Hom}_{\mathcal{D}}(\tau T_B, M^*) = \text{Hom}_{\mathcal{D}}(T_B, \tau^{-1}M^*) = 0$, and thus we obtain $f = 0$.

$(T_{B'}, M^*, T_B, M)$ or $(M^*, T_B, M, T_{B'})$: In these cases, a map in $\text{Irr}_{\text{add } T}(T_B, M)$ is of degree 0. Assume the lifting of it is $f: T_B \rightarrow M$. The preimage of M^* in \mathcal{D} is a module in these cases, so a factorisation of f must be of the form

$$\begin{array}{ccc} & T_B & \\ g \swarrow & \downarrow f & \\ \tau^{-1}M^* & \longrightarrow & M. \end{array}$$

Lemma 4.9 gives $f = 0$.

$(T_B, M, T_{B'}, M^*)$: In this case the preimage of M^* in \mathcal{D} is either a module or $P[1]$, for an indecomposable projective H -module P . In both cases, a map in $\text{Irr}_{\text{add } T}(T_B, M)$ is a map of degree 0. Assume the lifting is $f: T_B \rightarrow M$. In the first case there is a non-split exact sequence $0 \rightarrow M \rightarrow B' \rightarrow M^* \rightarrow 0$. Therefore, since $\dim \text{Hom}_{\mathcal{C}}(M^*, M[1]) = 1$, we have $\text{Hom}_{\mathcal{D}}(\tau^{-1}M^*, M) = 0$. Since $\text{Hom}_{\mathcal{D}}(T_B, M^*[-1]) = 0$, we must have $f = 0$, if f factors as below

$$\begin{array}{ccc} & T_B & \\ g \swarrow & \downarrow f & \\ M^*[-1] & \longrightarrow & M. \end{array}$$

Assume $M^* \simeq P[1]$, with P projective. If $\text{Hom}_{\mathcal{D}}(T_B, P) \neq 0$, then T_B is also projective. Therefore $\text{Hom}_{\mathcal{D}}(T_B, P[1]) = 0$, and thus $\text{Hom}_{\mathcal{C}}(T_B, M^*) = 0$, which means that $T_{B'}$ is a zero-vertex in Γ' , a contradiction. Then $\text{Hom}_{\mathcal{D}}(T_B, P) = 0$, but

this means that $f: T_B \rightarrow M$ factors through $M^*[-1] \simeq P$ only for $f = 0$. Thus, the map α is a monomorphism, and the sequence (4) is exact.

Thus, $\dim \widetilde{\text{Hom}}(T_B, M^*) = \dim \widetilde{\text{Hom}}(T_B, B') - \dim \widetilde{\text{Hom}}(T_B, M) = t's - r$. This means that $\dim(\text{Irr}_{\text{add } T'}(T_B, T_{B'}) \otimes_K \text{Irr}_{\text{add } T'}(T_{B'}, M^*) \cap I) = r$, so Γ' is balanced at the vertex 3, corresponding to $T_{B'}$.

We now show that Γ' is balanced at the vertex 1, corresponding to T_B , or 1 is a zero-vertex. Assume it is not a zero-vertex. We have the dual version of Lemma 4.9.

Lemma 4.10. *Assume that T_B does not correspond to a zero-vertex and that M^* is before T_B in the above ordering. Then $\text{Hom}_{\mathcal{D}}(M^*, \tau T_{B'}) = 0$.*

Proof. Similar to the proof of Lemma 4.9. □

Now, consider the triangle

$$M^* \rightarrow B \rightarrow M \rightarrow .$$

We need to show that there is an exact sequence

$$0 \rightarrow \widetilde{\text{Hom}}(M, T_{B'}) \rightarrow \widetilde{\text{Hom}}(B, T_{B'}) \rightarrow \widetilde{\text{Hom}}(M^*, T_{B'}) \rightarrow 0, \quad (5)$$

where $\widetilde{\text{Hom}}(M, T_{B'}) = \text{Irr}_{\text{add } T}(M, T_{B'})$, while $\widetilde{\text{Hom}}(B, T_{B'}) = \text{Irr}_{\text{add } \bar{T}}(B, T_{B'})$ and $\widetilde{\text{Hom}}(M^*, T_{B'}) = \text{Irr}_{\text{add } T'}(M^*, T_{B'})$. The proof of this is parallel to the proof for the sequence (4), and therefore omitted. Using the exact sequence (5), one obtains that Γ' is balanced at the vertex 1. □

We summarise the results of this section.

Theorem 4.11. *Let Γ be a cluster-tilted algebra of rank 3.*

- (a) Γ is either hereditary, or it is cyclic.
- (b) If Γ is cyclic, then each vertex of Γ is either balanced or a zero-vertex.
- (c) Let k be a vertex of Γ , let $\bar{\delta}_k(\Gamma)$ be the mutation in direction k , and let k^* be the new vertex of $\bar{\delta}_k(\Gamma)$. Then there are the following possible cases:
 - I. Both Γ and $\bar{\delta}_k(\Gamma)$ are hereditary.
 - II. Γ is hereditary, while $\bar{\delta}_k(\Gamma)$ is cyclic with a zero-vertex at k^* ,
 - III. Γ is cyclic with a zero-vertex at k , and $\bar{\delta}_k(\Gamma)$ is hereditary, or
 - IV. Γ is cyclic and balanced at k , and $\bar{\delta}_k(\Gamma)$ is cyclic and balanced at k^* .

Proof. This follows directly from the previous results in this section, and the fact that all cluster-tilted algebras can be obtained by starting with a hereditary algebra, and then performing a finite number of mutations [BMRRT], [BMR]. □

The above theorem is very easily verified for algebras of finite type, as indicated in the introduction of this section.

5. Mutation

As mentioned in the introduction, in view of Proposition 3.2 it is possible to assign to a cluster-tilted algebra Γ a skew-symmetric matrix $X_\Gamma = (x_{ij})$. More precisely, if there is at least one arrow from i to j in the quiver of the endomorphism-algebra Γ , let x_{ij} be the number of arrows from i to j . If there are no arrows between i and j , let $x_{ij} = 0$. Let $x_{ij} = -x_{ji}$ otherwise.

Now let \bar{T} be an almost complete tilting object in the cluster category \mathcal{C} with complements M and M^* . Let $T = \bar{T} \amalg M$, let $T' = \bar{T} \amalg M^*$, let $\Gamma = \text{End}_{\mathcal{C}}(T)^{\text{op}}$ and let $\Gamma' = \text{End}_{\mathcal{C}}(T')^{\text{op}}$. Then we want to show that the quivers of Γ and Γ' are related by the cluster-mutation formula. We use the results of Section 4 to show this for cluster-tilted algebras of rank 3, and Theorem 2.13 to extend to the general case. There is an independent proof of this for finite representation type in [CCS2].


Theorem 5.1. *Let H be a hereditary algebra, and let \bar{T}, M, M^*, Γ and Γ' be as above. Then the quivers of Γ and Γ' , or equivalently the matrices X_Γ and $X_{\Gamma'}$, are related by the cluster mutation formula.*

Proof. First, assume H has rank 3. In case H is not connected, the claim is easily checked. Assume H is connected. Fix k , the vertex where we mutate. By Theorem 1.2, it is clear that $x'_{ik} = -x_{ik}$ for $i = 1, 2, 3$, and that $x'_{kj} = -x_{kj}$ for $j = 1, 2, 3$.


Now assume $i \neq k$ and $j \neq k$. By Theorem 4.11, there are four possible cases.

Case I: This happens if and only if k is a source or a sink. In this case it is clear that either $x_{ik} = 0$ or $x_{kj} = 0$. For $i \neq k$ and $j \neq k$, it is clear that $x_{ij} = x'_{ij}$, since in this case mutation at k is the same as so-called APR-tilting at k . Thus the formula holds.

Case II: Since k is now not a source or a sink, we can assume Γ is the path algebra of




where $r > 0$ and $s > 0$ and $t \geq 0$ and with $k = 2$. Then, by Lemma 4.7, the quiver of $\bar{\delta}_2(\Gamma)$ is




with $t' = rs + t$. So $x'_{13} = t' = rs + t$, and

$$x_{13} + \frac{|x_{12}|x_{23} + x_{12}|x_{23}|}{2} = t + \frac{rs + rs}{2} = t + rs.$$

Case III: We assume that the quiver of Γ is



By Proposition 4.8, the quiver of Γ' is




with $t' = t - rs$. That is $x'_{13} = t - rs$, and

$$x_{13} + \frac{|x_{12}|x_{23} + x_{12}|x_{23}|}{2} = t + \frac{|-r|(-s) + (-r)|-s|}{2} = t - rs,$$

and the formula holds.

Case IV: We assume the quiver of Γ is the same as in case III. Now the quiver of Γ' is



where $t' = rs - t$. That is $x'_{13} = -t' = t - rs$, while

$$x_{13} + \frac{|x_{12}|x_{23} + x_{12}|x_{23}|}{2} = t + \frac{|-r|(-s) + (-r)|-s|}{2} = t - rs,$$

thus the formula holds true also in this case.

Now, assume that H has arbitrary rank. Fix k , the vertex where we mutate. By Theorem 1.2, it is clear that $x'_{ik} = -x_{ik}$ for any value of i , and that $x'_{kj} = -x_{kj}$ for any value of j . Assume now that $k \neq i$ and $k \neq j$. Let e_i, e_j, e_k be the primitive idempotents in Γ corresponding to the vertices i, j, k of the quiver of Γ . Assume $1_\Gamma = f + e_i + e_j + e_k$. and let $\Gamma_{\text{red}} = \Gamma / \Gamma f \Gamma$. Let e_i, e_j, e_k^* be the primitive

idempotents corresponding to the vertices i, j, k^* of the quiver of Γ' . Assume $1_{\Gamma'} = f' + e_i + e_j + e_{k^*}$. and let $\Gamma'_{\text{red}} = \Gamma' / \Gamma' f' \Gamma'$. It is clear that the number of arrows from i to j in the quiver of Γ'_{red} is x_{ij} and the number of arrows from i to j in the quiver of Γ'_{red} is x'_{ij} . So, by the first part of the proof, x_{ij} and x'_{ij} are related by the matrix mutation formula. \square

There are some nice direct consequences of this. For the first one, one uses in addition to Theorem 5.1 that the tilting graph (see [BMRRT, Section 3]) is connected, [BMRRT, Prop. 3.5].

Corollary 5.2. *Let Q be a finite quiver with no oriented cycles. Then a quiver Q' can be obtained from Q by repeated mutation if and only if Q' is the quiver of a cluster-tilted algebra $\text{End}_{\mathcal{C}}(T)^{\text{op}}$ for some tilting object T in $\mathcal{C} = \mathcal{C}_{KQ}$.*

We thank A. Seven for pointing out to us the following direct consequence of Theorem 2.13, and Corollary 5.2.

Corollary 5.3. *Let Q be a quiver which can be obtained by repeated mutation from an acyclic quiver. Then all full subquivers of Q have the same property.*

6. Connections to cluster algebras

Our main motivation for studying matrix mutation for quivers/matrices associated with tilting objects in cluster categories is the connection to cluster algebras. In this section we explain how Theorem 5.1 gives such a connection. In order to formulate our result we first need to give a short introduction to a special type of cluster algebras [FZ1], relevant to our setting [BFZ]. See also [FZ2] for an overview of the theory of cluster algebras.

Let $\mathcal{F} = \mathbb{Q}(u_1, \dots, u_n)$ be the field of rational functions in indeterminates u_1, \dots, u_n , let $\underline{x} = \{x_1, \dots, x_n\} \subset \mathcal{F}$ be a transcendence basis over \mathbb{Q} , and $B = (b_{ij})$ an $n \times n$ skew-symmetric integer matrix. A pair (\underline{x}, B) is called a *seed*. The *cluster algebra* associated to the seed (\underline{x}, B) is by definition a certain subring $\mathcal{A}(\underline{x}, B)$ of \mathcal{F} , as we shall describe. Given such a seed (\underline{x}, B) and some i , with $1 \leq i \leq n$, define a new element of \mathcal{F} by

$$x_i x'_i = \prod_{j; b_{ji} > 0} x^{b_{ji}} + \prod_{j; b_{ji} < 0} x^{-b_{ji}}.$$

We say that x_i, x'_i form an *exchange pair*. We obtain a new transcendence basis $\underline{x}' = \{x_1, \dots, x_n\} \cup \{x'_i\} \setminus \{x_i\}$ of \mathcal{F} . Then define a new matrix $B' = (b'_{ij})$ associated

with B by

$$b'_{ij} = \begin{cases} -b_{ij} & \text{if } k = i \text{ or } k = j, \\ b_{ij} + \frac{|b_{ik}|b_{kj} + b_{ik}|b_{kj}|}{2} & \text{otherwise.} \end{cases}$$

The pair (\underline{x}', B') is called the *mutation* of the seed (\underline{x}, B) in direction i , written $\mu_i(\underline{x}, B) = (\underline{x}', B')$. Let \mathcal{S} be the set of seeds obtained by iterated mutations of (\underline{x}, B) (in all possible directions). The set of *cluster variables* is by definition the union of all transcendence bases appearing in all the seeds in \mathcal{S} , and the cluster algebra $\mathcal{A}(\underline{x}, B)$ is the subring of \mathcal{F} generated by the cluster variables. The transcendence bases appearing in the seeds are called *clusters*.

As mentioned earlier, there is a 1-1 correspondence between finite quivers with no loops and no oriented cycle of length two and skew-symmetric integer matrices (up to reordering the columns). The vertices of the quiver of a matrix $B = (b_{ij})$ are $1, \dots, n$, and there are b_{ij} arrows from i to j if $b_{ij} > 0$. The cluster algebra is said to be *acyclic* if there is some seed where the quiver associated with the matrix has no oriented cycles [BFZ]. We take the corresponding seed as an initial seed. In this case, let $H = KQ$ be the hereditary path algebra associated with an initial seed (\underline{x}, B) . Let $\mathcal{C} = \mathcal{C}_H$ be the corresponding cluster category, and let T be a tilting object in \mathcal{C} . Similar to the above we can associate with T a *tilting seed* (T, Q_T) , where Q_T is the quiver of the endomorphism algebra $\text{End}_{\mathcal{C}}(T)^{\text{op}}$. Let T_1, \dots, T_n be the non-isomorphic indecomposable direct summands of T . Fix k , and let as before $\delta_k(T) = T'$ be the tilting object of \mathcal{C} obtained by exchanging T_k with T_k^* (using our earlier notation from Theorem 1.2). Define mutation of (T, Q_T) in direction k to be given by $\delta_k(T, Q_T) = (T', Q_{T'})$.

We now want to associate tilting seeds with seeds for acyclic cluster algebras. We first associate $(H[1], Q_H)$ with a fixed initial seed (\underline{x}, B) , where Q is the quiver for B and $H = KQ$. Let (\underline{x}', B') be some seed. We then have $(\underline{x}', B') = \mu_{k_t} \dots \mu_{k_1}(\underline{x}, B)$ for some ordered sequence (k_1, \dots, k_t) . There are in general several such sequences, and we choose one of minimal length. Associated with (\underline{x}, B) is the sequence of length 0, that is the empty set \emptyset . We define $\alpha((\underline{x}, B), \emptyset) = (H[1], Q_H)$, and $\alpha((\underline{x}', B'), (k_1, \dots, k_t)) = \delta_{k_t} \dots \delta_{k_1}(H[1], Q_H) = (T', Q_{T'})$. Fix an ordering on the cluster variables in the cluster $\underline{x} = \{x_1, \dots, x_n\}$ of the chosen initial seed and choose a corresponding indexing for the H_i in $H = H_1 \amalg \dots \amalg H_n$, so that we have a correspondence between x_i and H_i . This induces a correspondence between the cluster variables x'_i in the cluster \underline{x}' and the indecomposable direct summands T'_i in T' , which we also denote by α . We do not know in general if the definition of α only depends on the seed (\underline{x}', B') .

We can now formulate the connection between cluster algebras and tilting in cluster categories implied by our main result.

Theorem 6.1. *Let the notation be as above, with (\underline{x}, B) an initial seed for an acyclic cluster algebra, and $(T', Q_{T'})$ a tilting seed corresponding to a seed (\underline{x}', B') , via*

the correspondence α , inducing a correspondence $x'_i \leftrightarrow T'_i$ for $x'_i \in \underline{x}'$ and T'_i an indecomposable direct summand of T' .

(a) For any $k \in \{1, \dots, n\}$ we have a commutative diagram

$$\begin{array}{ccc} ((\underline{x}', B')(k_1, \dots, k_t)) & \xrightarrow{\alpha} & (T', Q_{T'}) \\ \mu_k \downarrow & & \delta_k \downarrow \\ ((\underline{x}'', B'')(k_1, \dots, k_t, k)) & \xrightarrow{\alpha} & (T'', Q_{T''}) \end{array}$$

where \underline{x}'' is the cluster obtained from \underline{x}' by replacing $x'_k \in \underline{x}'$ by x''_k , and T'' is the tilting object in \mathcal{C} obtained by exchanging the indecomposable summand T'_k by T''_k where $T' = \bar{T} \amalg T'_k$ and $T'' = \bar{T} \amalg T''_k$ are non-isomorphic tilting objects.

(b) Identifying x'_k with T'_k and x''_k with T''_k , the multiplication rule for $x'_k x''_k$ is given by

$$T'_k T''_k = \prod (T'_j)^{a_j} + \prod (T'_l)^{c_l}$$

where a_j and c_l are determined by the minimal respectively right and left add \bar{T} -approximations $\amalg (T'_j)^{a_j} \rightarrow T'_k$ and $T'_k \rightarrow \amalg (T'_l)^{c_l}$.

Proof. (a): This follows by induction, using Theorem 5.1, where δ_k is interpreted as given by a mutation rule like μ_k .

(b) Let T'_k be the direct summand of T' corresponding to x'_k . By (a), $Q_{T'}$ is the quiver of B' , and the monomials M_1 and M_2 are given by the entries of the matrix B' , hence by the arrows in the quiver $Q_{T'}$. In particular, the arrows entering and leaving k , are given by the minimal right and minimal left add \bar{T} -approximations of T'_k . \square

Note that with the appropriate formulation, this solves Conjecture 9.3 in [BMRRT].

For algebras of finite type we know from [BMRRT] that the map α gives a one-one correspondence between the seeds and tilting seeds, in particular it does not depend on the t -tuple (i_1, \dots, i_t) . In fact, we have in this case a 1-1 correspondence between cluster variables and indecomposable objects of \mathcal{C} , inducing a 1-1 correspondence between clusters and tilting objects.

Two cluster variables x_k and x_k^* are said to form an exchange pair if there are $n-1$ cluster variables $\{y_1, \dots, y_{n-1}\}$ such that $\{x_k, y_1, \dots, y_{n-1}\}$ and $\{x_k^*, y_1, \dots, y_{n-1}\}$ are clusters. Similarly we have exchange pairs with respect to tilting objects. If α identifies x_k and x_k^* with T_k and T_k^* , respectively, we then have the following.

Theorem 6.2. *For a cluster algebra of finite type, let α be the above correspondence between seeds and tilting seeds, and between cluster variables and indecomposable objects in the cluster category.*

(a) For any $k \in \{1, \dots, n\}$ we have a commutative diagram

$$\begin{array}{ccc} (\underline{x}', B') & \xrightarrow{\alpha} & (T', Q_{T'}) \\ \mu_k \downarrow & & \delta_k \downarrow \\ (\underline{x}'', B'') & \xrightarrow{\alpha} & (T'', Q_{T''}). \end{array}$$

(b) Identify the cluster variables with the indecomposable objects in \mathcal{C} via α . We have

$$T_k T_k^* = \prod (T_j)^{a_j} + \prod (T_l^*)^{c_l}$$

for an exchange pair T_k and T_k^* where the a_j and c_l appear in the unique non-split triangles

$$T_k^* \rightarrow \coprod T_j^{a_j} \rightarrow T_k \rightarrow,$$

and

$$T_i \rightarrow \coprod T_l^{c_l} \rightarrow T_i^* \rightarrow$$

in \mathcal{C} .

References

- [APR] M. Auslander, M. I. Platzeck, I. Reiten, Coxeter functors without diagrams. *Trans. Amer. Math. Soc.* **250** (1979), 1–46. [Zbl 0421.16016](#) [MR 0530043](#)
- [AR] M. Auslander, I. Reiten, Representation theory of Artin algebras. III. Almost split sequences. *Comm. Algebra* **3** (1975), 239–294. [Zbl 0331.16027](#) [MR 0379599](#)
- [ARS] M. Auslander, I. Reiten, S. O. Smalø, *Representation Theory of Artin Algebras*. Cambridge Stud. Adv. Math. 36, Cambridge University Press, Cambridge 1995. [Zbl 0834.16001](#) [MR 1314422](#)
- [AS] M. Auslander, S. O. Smalø, Preprojective modules over Artin algebras. *J. Algebra* **66** (1980), 6–122. [Zbl 0477.16013](#) [MR 0591246](#)
- [BFZ] A. Berenstein, S. Fomin, A. Zelevinsky, Cluster Algebras III: Upper bounds and double Bruhat cells. *Duke Math. J.* **126** (2005), 1–52. [Zbl 02147024](#) [MR 2110627](#)
- [BES] D. Bessis, The dual braid monoid. *Ann. Sci. Ecole Norm. Sup. (4)* **36** (2003), 647–683. [Zbl 1064.20039](#) [MR 2032983](#)
- [B] K. Bongartz, Tilted algebras. In *Representations of algebras* (Puebla, 1980), Lecture Notes in Math. 903, Springer-Verlag, Berlin 1981, 26–38. [Zbl 0478.16025](#) [MR 0654701](#)
- [BMRRT] A. Buan, R. Marsh, Reineke M., I. Reiten, G. Todorov, Tilting theory and cluster combinatorics. *Adv. Math.* **204** (2006), 572–618. [Zbl 05047155](#) [MR 2249625](#)
- [BMR] A. Buan, R. Marsh, I. Reiten, Cluster-tilted algebras. *Trans. Amer. Math. Soc.* **359** (2007), 323–332. [Zbl 05120550](#) [MR 2247893](#)

- [BR] A. Buan, I. Reiten, Acyclic quivers of finite mutation type. *Internat. Math. Res. Notices* **13** (2006), article ID 12804, 1–10. [Zbl 1116.16013](#) [MR 2249997](#)
- [CC] P. Caldero, F. Chapoton, Cluster algebras as Hall algebras of quiver representations. *Comment. Math. Helv.* **81** (2006), 595–616. [Zbl 1119.16013](#) [MR 2250855](#)
- [CCS1] P. Caldero, F. Chapoton, R. Schiffler, Quivers with relations arising from clusters (A_n case). *Trans. Amer. Math. Soc.* **358** (2006), 1347–1364. [Zbl 02237892](#) [MR 2187656](#)
- [CCS2] P. Caldero, F. Chapoton, R. Schiffler, Quivers with relations and cluster-tilted algebras. *Algebr. Represent. Theory* **9** (2006), 359–376. [Zbl 05125411](#) [MR 2250652](#)
- [CK] P. Caldero, B. Keller, From triangulated categories to cluster algebras. *Invent. Math.*, to appear. [arXiv:math.RT/0506018](#)
- [CFZ] F. Chapoton, S. Fomin, A. Zelevinsky, *Polytopal realizations of generalized associahedra*, *Canad. Math. Bull.* **45** (2002), 537–566. [Zbl 1018.52007](#) [MR 1941227](#)
- [FG] V. V. Fock, A. B. Goncharov, Cluster ensembles, quantization and the dilogarithm. Preprint 2003/2007. [arXiv:math.AG/0311245](#)
- [FZ1] S. Fomin, A. Zelevinsky, Cluster Algebras I: Foundations. *J. Amer. Math. Soc.* **15** (2002), 497–529. [Zbl 1021.16017](#) [MR 1887642](#)
- [FZ2] S. Fomin, A. Zelevinsky, Cluster Algebras: notes for the CDM-conference. In *Current developments in mathematics, 2003*, International Press, Somerville, MA, 2003. [Zbl 1119.05108](#) [MR 2132323](#)
- [FZ3] S. Fomin, A. Zelevinsky, Y -systems and generalized associahedra. *Ann. of Math.* (2) **158** (2003), 977–1018. [Zbl 1057.52003](#) [MR 2031858](#)
- [FZ4] S. Fomin, A. Zelevinsky, The Laurent phenomenon. *Adv. in Appl. Math.* **28** (2002), 119–144. [Zbl 1012.05012](#) [MR 1888840](#)
- [GSV1] M. Gekhtman, M. Shapiro, A. Vainshtein, Cluster algebras and Poisson geometry. *Moscow Math. J.* **3** (2003), 899–934. [Zbl 1057.53064](#) [MR 2078567](#)
- [GSV2] M. Gekhtman, M. Shapiro, A. Vainshtein, Cluster algebras and Weil-Petersson forms. *Duke Math. J.* **127** (2005), 291–311. [Zbl 1079.53124](#) [MR 2130414](#)
- [H] D. Happel, *Triangulated categories in the representation theory of finite-dimensional algebras*. London Math. Soc. Lecture Note Ser. 119, Cambridge University Press, Cambridge 1988. [Zbl 0635.16017](#) [MR 0935124](#)
- [H2] D. Happel, A characterization of hereditary categories with tilting object. *Invent. Math.* **144** (2001), 381–398. [Zbl 1015.18006](#) [MR 1827736](#)
- [HR] D. Happel, C. Ringel, Tilted algebras. *Trans. Amer. Math. Soc.* **274** (1982), 399–443. [Zbl 0503.16024](#) [MR 0675063](#)
- [HRS] D. Happel, J. Rickard, A. Schofield, Piecewise hereditary algebras. *Bull. London Math. Soc.* **20** (1988), 23–28. [Zbl 0638.16018](#) [MR 0916069](#)
- [HU] D. Happel, L. Unger, On the set of tilting objects in hereditary categories. In *Representations of algebras and related topics*, Fields Inst. Commun. 45, Amer. Math. Soc., Providence, RI, 2005, 141–159. [Zbl 1082.16019](#) [MR 2146246](#)
- [Ho] M. Hoshino, Modules without self-extensions and Nakayama’s conjecture. *Arch. Math. (Basel)* **43** (1984), 493–500. [Zbl 0547.16014](#) [MR 0775735](#)

- [K] B. Keller, On triangulated orbit categories. *Doc. Math.* **10** (2005), 551–581. [Zbl 02238184](#) [MR 2184464](#)
- [KR] B. Keller, I. Reiten, Cluster-tilted algebras are Gorenstein and stably Calabi-Yau. *Adv. Math.* **211** (2007), 123–151. [Zbl 05144368](#) [MR 2313531](#)
- [Ke] O. Kerner, Exceptional components of wild hereditary algebras. *J. Algebra* **152** (1997), 184–206. [Zbl 0779.16006](#) [MR 1190411](#)
- [Ke2] O. Kerner, Representations of wild quivers. In *Representation theory of algebras and related topics* (Mexico City, 1994), CMS Conf. Proc. 19, Amer. Math. Soc., Providence, RI, 1996, 65–107. [Zbl 0863.16010](#) [MR 1388560](#)
- [MRZ] R. Marsh, M. Reineke, A. Zelevinsky, Generalized associahedra via quiver representations. *Trans. Amer. Math. Soc.* **355** (2003), 4171–4186. [Zbl 1042.52007](#) [MR 1990581](#)
- [P] D. I. Panyushev, ad-nilpotent ideals of a Borel subalgebra: generators and duality. *J. Algebra* **274** (2004), 822–846. [Zbl 1067.17005](#) [MR 2043377](#)
- [RV] I. Reiten, M. van den Bergh, Noetherian hereditary categories satisfying Serre duality. *J. Amer. Math. Soc.* **15** (2002), 295–366. [Zbl 0991.18009](#) [MR 1887637](#)
- [Ric] J. L. Rickard, Derived categories and stable equivalence. *J. Pure. Appl. Algebra* **61** (1989), 303–317. [Zbl 0685.16016](#) [MR 1027750](#)
- [R] C. M. Ringel, *Tame algebras and integral quadratic forms*. Lecture Notes in Math. 1099, Springer-Verlag, Berlin 1984. [Zbl 0546.16013](#) [MR 0774589](#)
- [V] J. L. Verdier, Catégories dérivées, état 0. In *Semin. Geom. Algebr. Bois-Marie, SGA 4 1/2*, Lecture Notes in Math. 569, Springer-Verlag, Berlin 1977, 262–311. [Zbl 0407.18008](#)
- [V2] J. L. Verdier, Des catégories dérivées des catégories abéliennes. *Astérisque* **239** (1996). [Zbl 0882.18010](#) [MR 1453167](#)
- [Z] A. Zelevinsky, Cluster algebras. Notes for 2004 IMCC (Chonju, Korea, August 2004), Preprint 2004. [arXiv:math.RT/0407414](#)

Received March 22, 2005; revised September 9, 2005

Aslak Bakke Buan, Institutt for matematiske fag, Norges teknisk-naturvitenskapelige universitet, 7491 Trondheim, Norway

E-mail: aslakb@math.ntnu.no

Robert J. Marsh, Department of Pure Mathematics, University of Leeds, Leeds, LS2 9JT, UK

E-mail: marsh@maths.leeds.ac.uk

Idun Reiten, Institutt for matematiske fag, Norges teknisk-naturvitenskapelige universitet, 7491 Trondheim, Norway

E-mail: idunr@math.ntnu.no