

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 83 (2008)

Artikel: A volume maximizing canonical surface in 3-space
Autor: Bauer, Ingrid C. / Catanese, Fabrizio
DOI: <https://doi.org/10.5169/seals-99034>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 15.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A volume maximizing canonical surface in 3-space

Ingrid C. Bauer and Fabrizio Catanese

Abstract. Answering a question posed by Enriques, we construct a minimal smooth algebraic surface S of general type over the complex numbers with $K^2 = 45$ and $p_g = 4$, and with birational canonical map. The canonical system $|K_S|$ has a fixed part and the degree of the canonical image is 19. The surface we construct is rigid, S is indeed a ball quotient. It is obtained as an Abelian covering of the plane branched over an arrangement of lines already considered by Hirzebruch, and it is the first such example which is regular ($q = 0$).

Mathematics Subject Classification (2000). 14J25, 14J29, 14J80.

Keywords. Regular ball quotients, configurations of lines.

Introduction

At the onset of surface theory surfaces in 3-space, and especially canonical surfaces in 3-space, occupied a central role.

In particular, this study led to the famous Noether inequality $K^2 \geq 2p_g - 4$, while Castelnuovo observed that if the canonical map of a minimal smooth surface S is birational (obviously then $p_g \geq 4$) the inequality $K^2 \geq 3p_g - 7$ must hold true.

These are the lower bounds for surface geography, but upper bounds played a decisive role in the investigations of the last 30 years, leading to the so called Bogomolov–Miyaoka–Yau inequality

$$K^2 \leq 9\chi := 9(p_g - q + 1)$$

(cf. [BPV], Chapter VII, Section 4).

For instance, the BMY inequality gives a hypothetical upper bound for a question raised by F. Enriques (cf. [Enr], Chapter VIII, p. 284).

Problem. Which are the possible values of K^2 , in particular which is the highest possible value of K^2 for minimal surfaces with geometric genus $p_g = 4$ having a birational canonical map (so-called *simple canonical* surfaces)?

In fact, Enriques even conjectured that the highest possible value for K^2 should be 24, based on the conjecture that the expected number of moduli should be strictly positive. The second author showed in [Cat2] that this bound does not hold true, constructing simple canonical surfaces with geometric genus $p_g = 4$ and $11 \leq K^2 \leq 28$ (in these examples K^2 equals the 'canonical degree', i.e., the degree of the canonical image). This bound was improved by C. Liedtke (cf. [Lie]) who showed the existence of a simple canonical surface with $p_g = 4$ and $K^2 = 31$ (and canonical degree 12).

Simple canonical surfaces have $K^2 \geq 5$, and for $5 \leq K^2 \leq 7$ they were constructed by Enriques, Franchetta, Kodaira, Maxwell, and for $6 \leq K^2 \leq 16$ by Burniat, while Ciliberto was able to show for $5 \leq K^2 \leq 10$ the existence of simple canonical surfaces with ordinary singularities (cf. [Enr], [Fran], [Max], [Kod], [Bur], [Cil]).

If we try to go up with K^2 , the BMY inequality tells us that $K^2 \leq 45$, and that, if equality holds, then necessarily S is regular ($q(S) = 0$).

The main result of this paper is the following

Main Theorem. *There exists a minimal smooth algebraic surface S of general type over the complex numbers with $K^2 = 45$ and $p_g = 4$, and with birational canonical map. S is rigid, the canonical system $|K_S|$ has a fixed part and the degree of the canonical image is 19.*

More precisely, consider the smooth $(\mathbb{Z}/5)^2$ -Galois covers of $\mathbb{P}_{\mathbb{C}}^2$ branched on a complete quadrangle. These are 4 ball quotients S_i , $i = 1, 2, 3, 4$, with $\chi(S_i) = 5$, $K_{S_i}^2 = 45$ and $q(S_3) = 0$, $q(S_i) = 2$ for $i = 1, 2, 4$.

For S_3 the canonical system $|K_{S_3}|$ has a fixed part Ψ , the movable part $|M| := |K_{S_3} - \Psi|$ has 5 base points, and the canonical map $\phi_K : S_3 \dashrightarrow \Sigma$ is birational onto a surface Σ of degree 19.

Finally, the first homology groups are as follows:

$$H_1(S_3, \mathbb{Z}) \cong (\mathbb{Z}/5)^4, \quad H_1(S_i, \mathbb{Z}) \cong (\mathbb{Z}/5)^2 \oplus \mathbb{Z}^4, \quad i \neq 3.$$

The rigidity of S is due to the fact that, by Yau's proof of the inequality $K^2 \leq 9\chi$, it follows (cf. also [Miy]) that $K^2 = 9\chi$ if and only if the universal covering of S is the 2-dimensional complex ball \mathfrak{B}_2 .

It was for long time extremely hard to give direct algebro geometric constructions of such ball quotients, until a breakthrough came via the explicit constructions by Hirzebruch as Kummer coverings of the complex projective plane branched in a configuration of lines ([Hir]). These examples were extended and generalized in the book [BHH], which amply describes three examples of such (compact) ball quotients. The configurations occurring are quite classical: a complete quadrangle, the Hesse configuration and the dual Hesse configuration. Even if it is possible to determine the numerical data which a configuration has to fulfill in order to give rise to a ball quotient, it is less easy to compute the holomorphic invariants.

In fact, already the determination of the irregularities q of the Hirzebruch examples and of some étale quotients of them required further work by M.-N. Ishida (cf. [Ish1], [Ish2]), but no regular examples were indeed found (except Mumford's fake projective plane, whose construction however was not so explicit as Hirzebruch's one, see [Mum]).

The example of [BHH] we are interested in here is the $(\mathbb{Z}/5\mathbb{Z})^5$ -covering \hat{S} of \mathbb{P}^2 branched exactly in a complete quadrangle Δ . This surface has the invariants $K^2 = 45 \cdot 125$ and $\chi = 5 \cdot 125$. It is clear that an étale $(\mathbb{Z}/5\mathbb{Z})^3$ quotient or, equivalently, a smooth $(\mathbb{Z}/5\mathbb{Z})^2$ covering of \mathbb{P}^2 branched exactly in a complete quadrangle has the invariants $K^2 = 45$ and $\chi = 5$. Since, as we observed, $\chi = p_g - q + 1$, we have to produce an example of a surface S which is regular (i.e., $q = 0$) in order to get the desired example of a surface with $K^2 = 45$ and $p_g = 4$. In fact, we will show that up to isomorphisms there are exactly four smooth surfaces with $K^2 = 45$, $\chi = 5$, obtained as $(\mathbb{Z}/5\mathbb{Z})^2$ coverings of \mathbb{P}^2 branched exactly in a complete quadrangle: but only one of them is regular (has $q = 0$).

The main ingredient of our investigation is the theory of Abelian Galois coverings, developed by Pardini (cf. [Par]), but apparently not sufficiently known. Since the treatment by Pardini is very algebraic, and at some points not so explicit, we devote Section 1 to explain the structure theorem for such Abelian coverings, and especially the relation occurring between the topological data (which allow to construct the examples) and the explicit determination of the character sheaves (or eigensheaves) of the covering (these determine not only the topological but also the holomorphic invariants of the constructed surface).

Sections 2 and 3 are devoted to the construction of our surfaces, and to the investigation of the symmetries of our construction. This study allows us to classify all the examples up to isomorphisms.

Section 4 is devoted to the study of the canonical map of S_3 , while in Section 5 we give a description of the fundamental group $\pi_1(S)$ of our surfaces, from which one can calculate the first homology group $H_1(S, \mathbb{Z})$.

1. Abelian covers

In this section we shall recall the structure theorem for normal Abelian Galois ramified coverings. We shall give a more direct presentation than the one in the original paper by R. Pardini (cf. [Par]). This will turn out to be more suitable for our purposes.

Let X, Y be normal projective varieties, assume Y to be smooth and let $\pi : X \rightarrow Y$ be a finite Galois cover with Abelian Galois group G . By the theorem on the purity of the branch locus the critical set of π is a divisor R , the ramification divisor, whose image $D := \pi(R)$ is called the branch divisor. In the case where also X is smooth we have the following result (cf. [Cat1], Proposition 1.1).

Proposition 1.1. *If X is smooth, then R is a normal crossing divisor with smooth components. Moreover, if $x \in X$, then the stabilizer of x is the direct sum of the stabilizers of the components of R passing through x and these last groups are cyclic.*

We may assume without loss of generality, and will assume in the following, that Y is smooth and D is a normal crossing divisor.

We remark that π factors canonically as

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ & \searrow p & \nearrow p' \\ & X' & \end{array}$$

where X' is maximal such that $p': X' \rightarrow Y$ is unramified. In fact, one takes $X' := X/G'$, where G' is the subgroup of G generated by the stabilizers G_x of points $x \in X$.

Definition 1.2. π is called *totally ramified* iff p' is an isomorphism (i.e., $G = G'$).

Observe that π is necessarily totally ramified if Y has a trivial algebraic fundamental group.

Now, π is determined by the surjective homomorphism $\phi: \pi_1(Y - D) \rightarrow G$, which factors through $\varphi: H_1(Y - D, \mathbb{Z}) \rightarrow G$, since G is assumed to be Abelian.

We denote by G^* the group of characters of G , and we shall use the additive notation for the group operation in G^* . Recall that π is flat (for this it suffices that Y is smooth and X is normal) and that the action of G induces a splitting of the direct image of \mathcal{O}_X into eigensheaves

$$\pi_*\mathcal{O}_X = \bigoplus_{\chi \in G^*} \mathcal{L}_\chi^{-1},$$

where G acts on the invertible sheaf \mathcal{L}_χ^{-1} via the character χ .

Note that $\mathcal{L}_1 \cong \mathcal{O}_Y$ and denote by L_χ a divisor associated to the eigensheaf \mathcal{L}_χ (thus $\mathcal{L}_\chi \cong \mathcal{O}(L_\chi)$).

We shall show how

- (1) one calculates $H_1(Y - D, \mathbb{Z})$;
- (2) one calculates the character sheaves $\mathcal{L}_\chi = \mathcal{O}(L_\chi)$ in terms of the surjective homomorphism $\varphi: H_1(Y - D, \mathbb{Z}) \rightarrow G$.

Consider the exact sequence

$$0 \rightarrow K \rightarrow H_1(Y - D, \mathbb{Z}) \rightarrow H_1(Y, \mathbb{Z}) \rightarrow 0. \tag{1}$$

Remark 1.3. If π is totally ramified, $\varphi|_K: K \rightarrow G$ is surjective.

Following the arguments in [Cat1] we obtain

Proposition 1.4.

$$K = \ker(H_1(Y - D) \rightarrow H_1(Y)) = \text{coker}(r : H^{2n-2}(Y) \rightarrow H^{2n-2}(D)).$$

In particular, if $H_1(Y, \mathbb{Z}) = 0$, then

$$H_1(Y - D, \mathbb{Z}) = \text{coker}(r : H^{2n-2}(Y) \rightarrow H^{2n-2}(D)).$$

Proof. Let V be an open tubular neighbourhood of D and denote by ∂V its boundary. Then we have the exact sequence

$$\dots \rightarrow H^{2n-2}(Y) \rightarrow H^{2n-2}(D) \rightarrow H^{2n-1}(Y, \bar{V}) \rightarrow H^{2n-1}(Y) \rightarrow \dots$$

Observing that $H^{2n-1}(Y, \bar{V}) \cong H_1(Y - D, \mathbb{Z})$, we see that

$$\begin{aligned} K &= \ker(H_1(Y - D, \mathbb{Z}) \rightarrow H_1(Y, \mathbb{Z})) \\ &\cong \text{coker}\left(r : H^{2n-2}(Y) \rightarrow H^{2n-2}(D) \cong \bigoplus_{i=1}^k [D_i] \mathbb{Z}\right). \quad \square \end{aligned}$$

Remark 1.5. Applying $\text{Hom}_{\mathbb{Z}}(\cdot, G)$ to the short exact sequence (1) above we get

$$\begin{aligned} 0 &\rightarrow \text{Hom}(H_1(Y, \mathbb{Z}), G) \rightarrow \text{Hom}(H_1(Y - D, \mathbb{Z}), G) \rightarrow \text{Hom}(K, G) \\ &\rightarrow \text{Ext}^1(H_1(Y, \mathbb{Z}), G) \rightarrow \text{Ext}^1(H_1(Y - D, \mathbb{Z}), G) \rightarrow \text{Ext}^1(K, G) \rightarrow 0. \end{aligned}$$

Hence an Abelian covering of Y ramified in D is uniquely determined by a surjective morphism $\varphi : K \rightarrow G$ if and only if $\text{Hom}(H_1(Y), G) = 0$ and $\text{Ext}^1(H_1(Y), G) \rightarrow \text{Ext}^1(H_1(Y - D), G)$ is injective. This happens, for instance, if $H_1(Y, \mathbb{Z}) = 0$, or more generally if $H_1(Y, \mathbb{Z})$ is a finite group whose exponent is relatively prime to the exponent of G .

Let us determine the character sheaves of the Abelian covering determined by $\varphi : H_1(Y - D, \mathbb{Z}) \rightarrow G$.

Let $\chi \in G^*$ be a character of G , i.e., $\chi : G \rightarrow C \subset \mathbb{C}^*$, where C is cyclic. Then χ induces a surjective morphism $\chi \circ \varphi : H_1(Y - D, \mathbb{Z}) \rightarrow C$, whence a factorization of π as

$$\begin{array}{ccc} X & \xrightarrow{\pi} & Y \\ & \searrow & \nearrow \pi_\chi \\ & Z := X/(\ker(\chi \circ \varphi)) & \end{array}$$

where $\pi_\chi : Z \rightarrow Y$ is a cyclic covering with group C .

Remark 1.6. $\mathcal{L}_\chi(Z) = \mathcal{L}_\chi(X)$, and we are reduced to calculate the character sheaves for cyclic coverings.

Write $D = \bigcup_{i=1}^k D_i$ as a union of smooth irreducible components and denote by δ_i the image of a small loop around D_i in $H_1(Y - D, \mathbb{Z})$.

Let d be the order of C and let us identify C with \mathbb{Z}/d ; then we have the well known formula (cf. the proof of Proposition 4.5 of [Ba-Ca])

$$\mathcal{O}_Y(dL_\chi) \cong \mathcal{O}_Y\left(\sum_{i=1}^k (\chi \circ \varphi)(\delta_i) D_i\right).$$

Remark 1.7. We remark that the above linear equivalence

$$dL_\chi \equiv \sum_{i=1}^k (\chi \circ \varphi)(\delta_i) D_i,$$

depends only on $\chi \circ (\varphi|_K)$ and does not uniquely determine the character sheaf \mathcal{L}_χ . In fact, if $\mathcal{L}_\chi \in \text{Pic}(Y)$ satisfies the above equation, then also $\mathcal{L}_\chi \otimes \eta$ does, for each d -torsion sheaf $\eta \in \text{Pic}(Y)$. If η corresponds to an element $\alpha \in \text{Hom}(H_1(Y, \mathbb{Z}), \mathbb{Z}/d\mathbb{Z})$, then $\mathcal{L}_\chi \otimes \eta$ is the character sheaf of the cyclic covering corresponding to $\chi \circ \varphi + \alpha \circ p$, where $p: H_1(Y - D, \mathbb{Z}) \rightarrow H_1(Y, \mathbb{Z})$. Clearly $(\chi \circ \varphi + \alpha \circ p)|_K = (\chi \circ \varphi)|_K$.

The observation that p is surjective shows however once more that \mathcal{L}_χ is uniquely determined by ϕ .

Choosing a fixed system of representatives of $\mathbb{Z}/d\mathbb{Z}$, e.g., $\mathbb{Z}/d\mathbb{Z} = \{0, \dots, d - 1\}$, we get then a unique representation

$$dL_\chi \equiv \sum_{i=1}^k (\chi \circ \varphi)(\delta_i) D_i = \sum_{i=1}^k \Delta_i D_i, \quad 0 \leq \Delta_i \leq d - 1.$$

We will now use the above approach in order to write explicit equations for X as a subvariety in the geometric vector bundle corresponding to the locally free sheaf $\bigoplus_{\chi \in G^* \setminus \{1\}} \mathcal{L}_\chi$.

Remark 1.8. Let $\chi: G \rightarrow C \cong \mathbb{Z}/d$, $\chi': G \rightarrow C \cong \mathbb{Z}/d'$ be two characters of G . Then $\text{ord}(\chi + \chi') = \text{l.c.m.}(d, d') =: M$. Write M as $M = \lambda \cdot d = \lambda' \cdot d'$. Consider the linear equivalences

$$\begin{aligned} dL_\chi &\equiv \sum_{i=1}^k (\chi \circ \varphi)(\delta_i) D_i = \sum_{i=1}^k \Delta_i D_i, \\ d'L_{\chi'} &\equiv \sum_{i=1}^k (\chi' \circ \varphi)(\delta_i) D_i = \sum_{i=1}^k \Delta'_i D_i, \end{aligned}$$

and

$$M(L_{\chi+\chi'}) \equiv \sum_{i=1}^k ((\chi + \chi') \circ \varphi)(\delta_i) D_i \equiv \sum_{i=1}^k ((\lambda \Delta_i + \lambda' \Delta'_i) \bmod (M)) \cdot D_i.$$

Since moreover $0 < \lambda \Delta_i + \lambda' \Delta'_i < 2M$, we may write (identifying the divisor L_χ with the divisor $(\sum_{i=1}^k \frac{\Delta_i}{d} D_i) \in \oplus_{i=1}^k \mathbb{Q} D_i$)

$$L_\chi + L_{\chi'} - L_{\chi+\chi'} = \sum_{i=1}^k \epsilon_{D_i}^{\chi, \chi'} D_i,$$

where $\epsilon_{D_i}^{\chi, \chi'} := 1$ if $\lambda \Delta_i + \lambda' \Delta'_i \geq M$ and $\epsilon_{D_i}^{\chi, \chi'} := 0$ otherwise.

The above equality is equivalent (as shown in [Par]) to the existence of the multiplication maps

$$\mu_{\chi, \chi'}: \mathcal{L}_\chi^{-1} \otimes \mathcal{L}_{\chi'}^{-1} \rightarrow \mathcal{L}_{\chi+\chi'}^{-1}$$

which correspond to global sections of $\mathcal{L}_\chi \otimes \mathcal{L}_{\chi'} \otimes \mathcal{L}_{\chi+\chi'}^{-1}$ whose divisor is exactly equal to $\sum_{i=1}^k \epsilon_{D_i}^{\chi, \chi'} D_i$.

Let in fact $\sigma_i \in \Gamma(X, \mathcal{O}(D_i))$ be a section with $\text{div}(\sigma_i) = D_i$ and set

$$\epsilon_{\chi, \chi'}^i := \epsilon_{D_i}^{\chi, \chi'}.$$

Then $\prod_i \sigma_i^{\epsilon_{\chi, \chi'}^i}$ is a global section of $\mathcal{L}_\chi \otimes \mathcal{L}_{\chi'} \otimes \mathcal{L}_{\chi+\chi'}^{-1}$ yielding the multiplication maps.

These sections define equations for the natural embedding

$$i: X \hookrightarrow W := \bigoplus_{\chi \in G^+ \setminus \{1\}} \mathbb{V}(\mathcal{L}_\chi^{-1}).$$

In fact, let w_χ be a fibre coordinate of $\mathbb{V}(\mathcal{L}_\chi^{-1})$: then $i(X)$ is defined by the equations

$$w_\chi w_{\chi'} = \prod_v \sigma_v^{\epsilon_{\chi, \chi'}^v} w_{\chi+\chi'}.$$

We infer the following

Corollary 1.9. *If Y and the D_i 's are defined over a field K , then also X is defined over K .*

2. The construction

We consider in $\mathbb{P}^2 = \mathbb{P}^2_{\mathbb{C}}$ a complete quadrangle Δ , i.e., the union of the six lines through four points P_0, \dots, P_3 in general position.

Let $\pi : Y := \hat{\mathbb{P}}^2(P_0, \dots, P_3) \rightarrow \mathbb{P}^2$ be the Del Pezzo surface of degree 5 which is the blow up of the plane in the points P_0, \dots, P_3 . Denote by E_0, \dots, E_3 the exceptional curves. Moreover, for $j = 1, 2, 3$, let $L'_j := H - E_0 - E_j$, where H is the total transform in Y of a line on \mathbb{P}^2 , and let $L_j := H - \sum_{i=1}^3 E_i + E_j$. I.e., L'_j is the strict transform of the line in \mathbb{P}^2 through P_0 and P_j , whereas L_j is the strict transform of the line in \mathbb{P}^2 through E_i and E_k , where $\{i, j, k\} = \{1, 2, 3\}$.

The divisor $L_1 + L_2 + L_3 + L'_1 + L'_2 + L'_3 + E_0 + E_1 + E_2 + E_3$ on Y has simple normal crossings and we shall denote it by D .

Remark 2.1. It is well known that $H^2(Y, \mathbb{Z})$ is freely generated by H, E_0, \dots, E_3 . Since

$$H_1(Y - D, \mathbb{Z}) \cong \text{coker}(r : H^{2n-2}(Y) \rightarrow H^{2n-2}(D)) \cong \bigoplus_{i=1}^k [D_i] \mathbb{Z}, \tag{2}$$

where r is given by the intersection matrix

	H	E_0	E_1	E_2	E_3
L'_1	1	1	1	0	0
L'_2	1	1	0	1	0
L'_3	1	1	0	0	1
L_1	1	0	0	1	1
L_2	1	0	1	0	1
L_3	1	0	1	1	0
E_0	0	-1	0	0	0
E_1	0	0	-1	0	0
E_2	0	0	0	-1	0
E_3	0	0	0	0	-1

we obtain

$$H_1(Y - D, \mathbb{Z}) \cong \left(\bigoplus_{i=0}^3 \mathbb{Z}e_i \oplus \bigoplus_{i=1}^3 \mathbb{Z}l_i \oplus \bigoplus_{i=0}^3 \mathbb{Z}l'_i \right) / H^2(Y, \mathbb{Z}),$$

where e_j (resp. l_i, l'_i) is a (small) simple loop oriented counterclockwise around E_j (resp. L_i, L'_i). I.e., $H_1(Y - D, \mathbb{Z})$ has generators $e_0, \dots, e_3, l_1, l_2, l_3, l'_1, l'_2, l'_3$ and the relations are $e_0 = l'_1 + l'_2 + l'_3, e_i = l'_i + l_j + l_k$ (here and in the following, $\{i, j, k\} = \{1, 2, 3\}$), $\sum l'_i + \sum l_i = 0$. In particular, $H_1(Y - D, \mathbb{Z})$ is free of rank 5.

We want to construct a smooth Galois cover $p: S \rightarrow Y$ with group $(\mathbb{Z}/5\mathbb{Z})^2$ branched exactly in D .

Such a Galois cover is determined by a surjective homomorphism $\varphi: \mathbb{Z}^5 \cong H_1(Y - D, \mathbb{Z}) \rightarrow (\mathbb{Z}/5\mathbb{Z})^2$ with certain conditions ensuring that S is smooth and that the covering branches exactly in D .

We write

$$\varphi(l_i) =: \begin{pmatrix} x_i \\ y_i \end{pmatrix} =: u_i, \quad \varphi(l'_j) =: \begin{pmatrix} z_j \\ w_j \end{pmatrix} =: v_j, \quad \varphi(e_h) =: \epsilon_h,$$

where $x_i, y_i, z_j, w_j \in \{0, \dots, 4\} \cong \mathbb{Z}/5\mathbb{Z}$.

Remark 2.2. It obviously follows from Remark 2.1 that each ϵ_h is determined by the u_i 's, v_j 's and that $\sum_i u_i + \sum_j v_j = 0$.

In order to calculate the invariants (i.e., p_g, K_S^2, q) of the Galois covering given by the homomorphism φ , we have to calculate for each character $\chi \in ((\mathbb{Z}/5\mathbb{Z})^2)^*$ the eigensheaf \mathcal{L}_χ .

Before doing this let us work out first the two sets of conditions ensuring that our covering is

- 1) branched exactly in $D = L_1 + L_2 + L_3 + L'_1 + L'_2 + L'_3 + E_0 + E_1 + E_2 + E_3$; and that
- 2) S is smooth.

Lemma 2.3. 1) If for all $i, u_i, v_i, \sum u_h, \epsilon_i = u_i + v_j + v_k$ are different from zero in $(\mathbb{Z}/5\mathbb{Z})^2$, then the covering $p: S \rightarrow Y$ is branched exactly in $L_1, L_2, L_3, L_1, L'_2, L'_3, E_0, \dots, E_3$.

2) If the following pairs of vectors in $(\mathbb{Z}/5\mathbb{Z})^2$

$$\left. \begin{aligned} &(u_i, v_i) \text{ for } i \in \{1, 2, 3\}, \\ &(u_1, u_1 + u_2 + u_3), (u_2, u_1 + u_2 + u_3), (u_3, u_1 + u_2 + u_3), \\ &(u_1, u_1 + v_2 + v_3), (u_2, u_2 + v_1 + v_3), (u_3, u_3 + v_1 + v_2), \\ &(u_1 + v_2 + v_3, v_i), \\ &(u_2 + v_1 + v_3, v_i) \text{ for } i = 1, 3, \\ &(u_3 + v_1 + v_2, v_i) \text{ for } i = 1, 2 \end{aligned} \right\} \text{ for } i = 2, 3,$$

are linearly independent, then S is smooth.

Proof. 1) is obvious.

2) follows from the fact that S given by the homomorphism φ is smooth if and only if the following condition holds: let D_1, D_2 be two non trivial irreducible subdivisors of the branch divisor of $p: S \rightarrow Y$ and let d_i a small loop around D_i , then $\varphi(d_1)$ and $\varphi(d_2)$ are not in the same cyclic subgroup of $(\mathbb{Z}/5\mathbb{Z})^2$. □

Remark 2.4. Let $p: S \rightarrow Y$ be a $(\mathbb{Z}/5\mathbb{Z})^2$ -Galois cover with u_i and v_j satisfying the two conditions of the above lemma. Then S is a smooth minimal surface with $K_S^2 = 45$ and $\chi = 5$. In fact, since the canonical divisor K_S is the pull back of the rational divisor

$$-3H + \sum E_i + 4/5 \sum E_i + 4/5 \sum (L_j + L'_j) \equiv 9/5H - 3/5 \sum E_i$$

we get $K_S^2 = (9^2 - 4 \cdot 3^2) = 45$. Moreover, by the Noether's formula it suffices to compute that the Euler number $e(S) = 15$. In fact, we get 25 times the Euler number of \mathbb{P}^2 blown up in 4 points, minus 20 times the Euler number of 10 \mathbb{P}^1 's plus 16 times the Euler number of 15 points: i.e., $25 \cdot 7 - 400 + 240 = 15$.

We are interested to find such surfaces with $q = 0$, because then they will have geometric genus equal to 4.

Given a character $\chi = (a, b) \in (\mathbb{Z}/5\mathbb{Z})^2$, let us determine $\mathcal{L}_\chi = \mathcal{L}_{(a,b)}$.

By the results of Section 1, we get

Proposition 2.5. Denoting by $[z]$ the residue class of z modulo 5, we have the following formulae:

$$\begin{aligned} 5\mathcal{L}_\chi &\equiv \sum_{i=1}^3 \chi(l_i)L_i + \sum_{i=1}^3 \chi(l'_i)L'_i + \sum_{i=0}^3 \chi(e_i)E_i. \\ 5\mathcal{L}_{(a,b)} &\equiv \sum_{i=1}^3 [ax_i + by_i]L_i + \sum_{i=1}^3 [az_i + bw_i]L'_i + [a(x_1 + x_2 + x_3) \\ &\quad + b(y_1 + y_2 + y_3)]E_0 + \sum_{i=1}^3 [a(x_i + z_j + z_k) + b(y_i + w_j + w_k)]E_i. \end{aligned}$$

3. The symmetries of the construction

Definition 3.1. A six-tuple $\mathcal{U} := (u_1, u_2, u_3, v_1, v_2, v_3) \in ((\mathbb{Z}/5\mathbb{Z})^2 \setminus \{0\})^6$ is said to be *admissible* if and only if

$$u_1 + u_2 + u_3 + v_1 + v_2 + v_3 = 0,$$

and moreover the two conditions of Lemma 2.3 are satisfied, i.e.:

- a) for all i , $u_i, v_i, \sum u_h, \epsilon_i = u_i + v_j + v_k$ are different from zero in $(\mathbb{Z}/5\mathbb{Z})^2$;
- b) the pairs of vectors listed in Lemma 2.3, 2) are linearly independent in $(\mathbb{Z}/5\mathbb{Z})^2$.

Remark 3.2. We have seen in the previous section that an admissible six-tuple \mathfrak{U} induces a smooth Galois cover $p: S \rightarrow Y$ with Galois group $(\mathbb{Z}/5\mathbb{Z})^2$. Moreover, S is a minimal surface of general type with $K_S^2 = 45$ and $\chi = 5$. We recall that S is a ball quotient, hence rigid.

Using MAGMA one sees that there are exactly 201600 admissible six-tuples. But of course a lot of them will lead to isomorphic surfaces. In order to understand how many non isomorphic surfaces (with $p_g = 4$) we will get by this construction, we have to understand the symmetries.

Two admissible six-tuples $\mathfrak{U}, \mathfrak{U}'$ obviously give isomorphic surfaces if there is an automorphism $\phi \in \text{Gl}(2, \mathbb{Z}/5\mathbb{Z})$ such that $\phi(\mathfrak{U}) = \mathfrak{U}'$. On the other hand the group of biholomorphic automorphisms of $\mathbb{P}^2 \setminus \{L_1, L_2, L_3, L'_1, L'_2, L'_3\}$ equals \mathfrak{S}_5 (cf. [Ter]). The action of \mathfrak{S}_5 on the set of admissible six tuples is generated by the following transformations:

$$(01): (u_1, u_2, u_3, v_1, v_2, v_3) \rightarrow (u_1, u_3 + v_1 + v_2, u_2 + v_1 + v_3, u_1 + u_2 + u_3, v_2, v_3);$$

$$(02): (u_1, u_2, u_3, v_1, v_2, v_3) \rightarrow (u_3 + v_1 + v_2, u_2, u_1 + v_2 + v_3, v_1, u_1 + u_2 + u_3, v_3);$$

$$(03): (u_1, u_2, u_3, v_1, v_2, v_3) \rightarrow (u_2 + v_1 + v_3, u_3 + v_1 + v_2, u_3, v_1, v_2, u_1 + u_2 + u_3);$$

$$(04): (u_1, u_2, u_3, v_1, v_2, v_3) \rightarrow (u_1, u_2, u_3, u_1 + v_2 + v_3, u_2 + v_1 + v_3, u_3 + v_1 + v_2).$$

It is easy to see that these four transpositions generate the action of a group isomorphic to \mathfrak{S}_5 .

We consider now the group \mathfrak{G} acting on the set of admissible six-tuples \mathfrak{S} , which is generated by \mathfrak{S}_5 and $\text{Gl}(2, \mathbb{Z}/5\mathbb{Z})$. Then \mathfrak{G} is a quotient of $\text{Gl}(2, \mathbb{Z}/5\mathbb{Z}) \times \mathfrak{S}_5$ (the actions commute, being given by multiplication on the right, respectively on the left). A MAGMA computation, reproduced in the appendix, reveals that \mathfrak{G} has four orbits on \mathfrak{S} (indeed, the simple minded idea is to calculate the cardinality of these 4 orbits, and see that their sum equals the cardinality of the set of admissible six-tuples).

Representatives for these orbits are

$$\mathfrak{U}_1 = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix} \right);$$

$$\mathfrak{U}_2 = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 2 \end{pmatrix}, \begin{pmatrix} 2 \\ 1 \end{pmatrix} \right);$$

$$\mathfrak{U}_3 = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 4 \\ 1 \end{pmatrix}, \begin{pmatrix} 3 \\ 2 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right);$$

$$\mathfrak{U}_4 = \left(\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 1 \end{pmatrix}, \begin{pmatrix} 0 \\ 3 \end{pmatrix}, \begin{pmatrix} 2 \\ 0 \end{pmatrix} \right).$$

The orbit of \mathfrak{U}_1 has length 28800, whereas the orbits of $\mathfrak{U}_2, \mathfrak{U}_3, \mathfrak{U}_4$ have respective length 57600.

In particular we see that $\mathcal{G} \cong \text{Gl}(2, \mathbb{Z}/5\mathbb{Z}) \times \mathfrak{S}_5$.

We have moreover:

Theorem 3.3. *Let S_i be the minimal smooth surface of general type with $K^2 = 45$ and $\chi = 5$ obtained from the covering induced by the admissible six-tuple \mathfrak{U}_i , where $i \in \{1, 2, 3, 4\}$. Then we have that S_3 is regular (i.e., $q(S_3) = 0$), whereas $q(S_i) = 2$ for $i \neq 3$.*

In particular, S_3 is the unique minimal surface with $K_S^2 = 45$ and $p_g = 4$ obtained as a $(\mathbb{Z}/5\mathbb{Z})^2$ -cover of \mathbb{P}^2 branched exactly in a complete quadrangle of the complex projective plane.

Proof. We will calculate the geometric genus of $S = S_3$, using the formula

$$H^0(S, \mathcal{O}_S(K_S)) = \bigoplus_{(a,b) \in G^*} H^0(Y, \mathcal{O}_Y(K_Y) \otimes \mathcal{L}_{(a,b)}).$$

Applying Proposition 2.5 we obtain the following table for the divisors $L_{a,b}$ yielding the character sheaves $\mathcal{L}_{(a,b)} = \mathcal{O}_Y(L_{a,b})$:

$\mathcal{L}_{(a,b)}$	$a = 0$	$a = 1$	$a = 2$	$a = 3$	$a = 4$
$b = 0$	0	$2H - E_1 - E_2 - E_3$	$2H - E_1 - E_2$	$3H - E_0 - 2E_1 - E_2 - E_3$	$3H - E_0 - 2E_1 - E_2$
$b = 1$	H	H	$3H - E_0 - E_1 - E_2 - E_3$	$3H - E_0 - E_1 - 2E_2 - E_3$	$3H - E_0 - E_1 - E_2 - E_3$
$b = 2$	$2H - E_1 - E_3$	$2H - E_1 - E_2 - E_3$	$2H - E_0 - E_1 - E_2$	$3H - E_0 - E_1 - E_2 - E_3$	$3H - 2E_0 - E_1 - E_2 - E_3$
$b = 3$	$2H - E_2 - E_3$	$3H - E_0 - E_1 - E_2 - E_3$	$2H - E_0 - E_3$	$2H - E_0$	$4H - 2E_0 - E_1 - 2E_2 - 2E_3$
$b = 4$	$3H - E_1 - E_2 - 2E_3$	$2H - E_0 - E_3$	$3H - E_0 - E_1 - E_2 - 2E_3$	$3H - 2E_0 - E_1 - E_2 - E_3$	$3H - 2E_0 - E_1 - E_2$

We see immediately that $H^0(Y, \mathcal{O}_Y(K_Y) \otimes \mathcal{L}_{(a,b)}) = 0$ for all $(a, b) \notin \{(2, 1), (3, 2), (1, 3), (4, 1)\}$ and $H^0(Y, \mathcal{O}_Y(K_Y) \otimes \mathcal{L}_{(a,b)}) \cong \mathbb{C}$ for $(a, b) \in \{(2, 1), (3, 2), (1, 3), (4, 1)\}$, i.e., $p_g(S_3) = 4$. This proves the claim for S_3 .

The geometric genus of the remaining surfaces is calculated in exactly the same way. □

4. The canonical map

In the previous section we have constructed a minimal surface S of general type with $K_S^2 = 45$, $p_g = 4$ and $q(S) = 0$. We want now to understand the behaviour of the canonical map of S .

For $(a, b) \in (\mathbb{Z}/5)^2$ we write

$$\begin{aligned} \delta_i(a, b) &:= [ax_i + by_i], \\ \lambda_j(a, b) &:= [az_j + bw_j], \\ \mu_0(a, b) &:= [a(x_1 + x_2 + x_3) + b(y_1 + y_2 + y_3)], \\ \mu_h(a, b) &:= [a(x_h + z_j + z_k) + b(y_h + w_j + w_k)]. \end{aligned}$$

Then we know that

$$5\mathcal{L}_{(a,b)} = \sum_{i=1}^3 \delta_i(a, b)L'_i + \sum_{j=1}^3 \lambda_j(a, b)L_j + \sum_{h=0}^3 \mu_h(a, b)E_h.$$

Denote by R_1, \dots, R_{10} the ramification divisors of $p: S \rightarrow Y$ lying over $L'_1, L'_2, L'_3, L_1, L_2, L_3, E_0, \dots, E_3$: it is easy to see that they are all irreducible genus 2 curves. Further, let x_i be a local equation of R_i . We already saw that $H^0(S, \mathcal{O}_S(K_S))$ is the direct sum of 4 one dimensional eigenspaces $H^0(S, \mathcal{O}_S(K_S))_{(a,b)} \cong H^0((Y, \mathcal{O}_Y(K_Y) \otimes \mathcal{L}_{(a,b)}) \cong H^0(\mathbb{P}^2, \mathcal{O}_{\mathbb{P}^2})$. Then a basis of $H^0(S, \mathcal{O}_S(K_S))$ is given by

$$\{x_1^{4-\delta_1(a,b)} \cdot x_2^{4-\delta_2(a,b)} \cdot x_3^{4-\delta_3(a,b)} \cdot x_4^{4-\lambda_1(a,b)} \cdot x_5^{4-\lambda_2(a,b)} \cdot x_6^{4-\lambda_3(a,b)} \cdot x_7^{4-\mu_0(a,b)} \dots x_{10}^{4-\mu_3(a,b)} \mid H^0(Y, \mathcal{O}_Y(K_Y) \otimes \mathcal{L}_{(a,b)}) \neq 0\}.$$

It is easy to compute the table giving the numbers λ_i, δ_j and μ_h for $(a, b) \in \{(2, 1), (3, 2), (1, 3), (4, 1)\}$:

(a, b)	δ_1	δ_2	δ_3	λ_1	λ_2	λ_3	μ_0	μ_1	μ_2	μ_3
(1, 3)	1	1	3	2	4	4	0	4	2	4
(2, 1)	2	2	1	4	3	3	0	3	4	3
(3, 2)	3	3	2	4	3	0	3	1	2	4
(4, 1)	4	4	1	2	4	0	4	3	1	2

Therefore we have the following result.

Lemma 4.1. *A basis for $H^0(S, \mathcal{O}_S(K_S))$ is given by*

$$\{x_1^3 x_2^3 x_3 x_4^2 x_7^4 x_9^2, x_1^2 x_2^2 x_3^3 x_5 x_6 x_7^4 x_8 x_{10}, x_1 x_2 x_3^2 x_5 x_6^4 x_7 x_8^3 x_9^2, x_3^3 x_4^2 x_6^4 x_8 x_9^3 x_{10}^2\}.$$

We can now prove the following

Theorem 4.2. 1) *The canonical map ϕ_K of S has R_3 as fixed part and its movable part has five base points. We have $R_3^2 = -1$ and $K_S \cdot R_3 = 3$. The base points of $K_S - R_3$ are $x_1 \cap x_4$ (of type $(1, 1)$), $x_1 \cap x_8$ (of type $(1, 1, 1)$), $x_2 \cap x_9$ (of type $(2, 1, 1)$), $x_3 \cap x_7$ (of type $(2, 1, 1)$), $x_6 \cap x_9$ (of type $(1, 1)$).*

2) *The canonical map is birational and its image in \mathbb{P}^3 has degree 19.*

In order to keep the formulation of the above theorem as simple as possible we adopted the notation: a base point p of $K - R_3$ on S is of type (n_1, n_2, \dots, n_k) iff p is a n_1 -tuple base point of $K - R_3$, after one blow up the strict transform of $K - R_3$ has a n_2 -tuple base point and so on.

Proof. It is immediate from the description of the basis of $H^0(S, \mathcal{O}_S(K_S))$ given in Lemma 4.1 that R_3 is exactly the fixed part of $|K_S|$. It is easy to see that the base points of $|K - R_3|$ are exactly $x_1 \cap x_4, x_1 \cap x_8, x_2 \cap x_9, x_3 \cap x_7, x_6 \cap x_9$. Next we will see which kind of base points we have and whether there are still infinitely near base points.

1) $x_1 \cap x_4$: locally around this point $|K - R_3|$ is given by $x_1^3 x_4^2, x_1^2, x_1, x_4^2$. The ideal generated is the ideal (x_1, x_4^2) , thus $K - R_3$ has a base point of type $(1, 1)$ in $x_1 \cap x_4$.

2) $x_1 \cap x_8$: locally around this point $|K - R_3|$ is given by $x_1^3, x_1^2 x_8, x_1 x_8^3, x_8$. The ideal generated is the ideal (x_1^3, x_8) , thus $K - R_3$ has a base point of type $(1, 1, 1)$ in $x_1 \cap x_8$.

Similarly in the three remaining cases we see that:

3) $|K - R_3|$ has a base point of type $(2, 1, 1)$ in $x_2 \cap x_9$ (ideal $(x_2^2, x_2 x_9^2)$).

4) $|K - R_3|$ has a base point of type $(2, 1, 1)$ in $x_3 \cap x_7$ (ideal $(x_3^2, x_7^4, x_3 x_7)$).

5) $|K - R_3|$ has a base point of type $(1, 1)$ in $x_6 \cap x_9$ (ideal (x_6, x_9^2)).

Therefore we get $\deg \phi_K \deg \phi_K(S) = (K - R_3)^2 - 2 \cdot 4 - 11 = 19$. Here we use that R_3 has self intersection -1 and genus 2. It follows immediately that $\deg \phi_K = 1$ and that the canonical image has degree 19. \square

The following is an answer to a question by Junho Lee.

Corollary 4.3. *There exist surfaces of general type S with birational canonical map such that the canonical system of each deformation of S has base points.*

There remains finally the statement of the main theorem concerning the first homology group of the four ball quotients which are obtained as $(\mathbb{Z}/5)^2$ -Galois covers of \mathbb{P}^2 branched on a complete quadrangle Δ . This will be done in the next section.

5. Fundamental group and first homology group of S

Let S be a surface as above with $K^2 = 45$ and $\chi = 5$ constructed as $(\mathbb{Z}/5\mathbb{Z})^2$ cover of the plane ramified exactly over a complete quadrangle. The first aim of this section is to find a description of the fundamental group of S .

Recall that the covering $p: S \rightarrow Y = \hat{\mathbb{P}}^2(P_0, \dots, P_3)$ is given by the homomorphism $\varphi: \pi_1(\mathbb{P}^2 - \Delta) \rightarrow H^1(\mathbb{P}^2 - \Delta, \mathbb{Z}/5\mathbb{Z}) \rightarrow (\mathbb{Z}/5\mathbb{Z})^2$. I.e., we have the exact sequence

$$1 \rightarrow \pi_1(S - R) \cong \ker(\varphi) \rightarrow \pi_1(\mathbb{P}^2 - \Delta) \rightarrow (\mathbb{Z}/5\mathbb{Z})^2 \rightarrow 0,$$

where $R \subset S$ is the ramification divisor of $p: S \rightarrow Y$, i.e., $R = \sum_{i=1}^{10} R_i$. Let α_i be a small geometric loop around R_i , for each $i \in \{1, \dots, 10\}$, and denote by $\langle \alpha_1, \dots, \alpha_{10} \rangle$ the normal subgroup of $\pi_1(S - R)$ generated by $\alpha_1, \dots, \alpha_{10}$.

Then $\pi_1(S) = \pi_1(S - R) / \langle \alpha_1, \dots, \alpha_{10} \rangle$.

We first have to give a suitable presentation of $\pi_1(\mathbb{P}^2 - \Delta)$. For this we use a result of T. Terada ([Ter]). We change the notation and use Terada's notation (cf. Figure 1).




Figure 1. A complete quadrangle in \mathbb{P}^2 .

Let $a \in \mathbb{P}^2 - \Delta$ and denote by $\rho(ij)$ resp. $\rho(ijk)$, $0 \leq i < j < k \leq 3$ a small loop starting from a around (ij) resp. around the exceptional divisor of the blow up in (ijk) defined as in [Ter]. Then Terada shows the following result.

Theorem 5.1 (Terada). $\pi_1(\mathbb{P}^2 - \Delta, a)$ is generated by $\rho(ij)$, $i < j$ and the relations are

$$\begin{aligned} [\rho(ij)\rho(ik)\rho(jk), \rho(ij)] &= 1, \\ [\rho(ij)\rho(ik)\rho(jk), \rho(ik)] &= 1, \\ [\rho(ij)\rho(ik)\rho(jk), \rho(jk)] &= 1, \quad \text{for all } i < j < k; \\ \rho(01)\rho(02)\rho(12)\rho(03)\rho(13)\rho(23) &= 1. \end{aligned}$$

On the other hand, it is well known that $\pi_1(\mathbb{P}^2 - \Delta)$ is isomorphic to the quotient of the pure braid group of four points by its center \mathcal{P}_4/Z_4 . Let σ_i be the standard generators of Artin’s braid group. Then we have

$$\rho(ij) = \sigma_i^{-1}\sigma_{i+1}^{-1}\dots\sigma_{j-2}^{-1}\sigma_{j-1}^2\sigma_{j-2}\dots\sigma_i.$$

We will show that $\rho(ij)\rho(ik)\rho(jk)$ generates the center of $\mathcal{P}_3 \cong \pi_1(\mathbb{C}^2 - Z)$, where Z is the union of three lines through one point. This implies, in view of the geometric exact sequence

$$1 \rightarrow Z_3 \rightarrow \mathcal{P}_3 \rightarrow \pi_1(\mathbb{P}^1 \setminus \{0, 1, \infty\}) \rightarrow 1,$$

that $\rho(ij)\rho(ik)\rho(jk)$ represents a simple loop around the exceptional divisor over (ijk) in $\pi_1(\mathbb{P}^2 - \Delta)$.

We have in fact that $\rho(01)\rho(02)\rho(12) = \sigma_0^2\sigma_0^{-1}\sigma_1^2\sigma_0\sigma_1^2$, is the standard generator of the centre of \mathcal{P}_3 according to Chow’s theorem (see Corollary 1.8.4 , p. 28 of [Bir]).

The remaining three cases are treated in the same way.

We summarize our result in the following

Proposition 5.2. Let $\mathfrak{U} := (u_1, u_2, u_3, v_1, v_2, v_3)$ be an admissible six-tuple and let S be the smooth surface constructed from \mathfrak{U} .

Let $\Phi : \pi_1(\mathbb{P}^2 - \Delta) \rightarrow (\mathbb{Z}/5)^2$ be the morphism given by $\rho(01) \mapsto v_2$, $\rho(02) \mapsto v_3$, $\rho(12) \mapsto u_1$, $\rho(03) \mapsto v_1$, $\rho(13) \mapsto u_3$, $\rho(23) \mapsto u_2$.

Then $\pi_1(S) \cong \ker \Phi/N$, where N is the subgroup of $\ker \Phi$ normally generated by

$$\begin{aligned} &\rho(01)^5, \quad \rho(02)^5, \quad \rho(12)^5, \quad \rho(02)^5, \quad \rho(13)^5, \quad \rho(23)^5, \quad (\rho(01)\rho(02)\rho(12))^5, \\ &(\rho(02)\rho(03)\rho(23))^5, \quad (\rho(01)\rho(03)\rho(13))^5, \quad (\rho(12)\rho(13)\rho(23))^5. \end{aligned}$$

From the above description of the fundamental group of S , using the method of Reidemeister–Schreier, its abelianization $H_1(S, \mathbb{Z})$ can be calculated via a MAGMA program which is reproduced in the appendix.

6. Appendix

The following program calculates the admissible six-tuples.

```

R:=Integers;
f:=FiniteField(5);
V:=VectorSpace(f,2);
aal:=0;LVU:={@ @};
for x1 in f do for y1 in f do for x2 in f do for y2 in f do
for x3 in f do for y3 in f do for z1 in f do
for z2 in f do for w1 in f do for w2 in f do
M:={@ @};
u1:=V![x1,y1];u2:=V![x2,y2];
u3:=V![x3,y3];u4:=V![z1,w1];u5:=V![z2,w2];
u6:=- (u1+u2+u3+u4+u5);e0:=u1+u2+u3;e1:=u1+u5+u6;
e2:=u2+u4+u6;e3:=u3+u4+u5;
R2:=[u1,u2,u3,u4,u5,u6];
U1:=sub<V|e0,u1>;U2:=sub<V|e0,u2>;U3:=sub<V|e0,u3>;
U4:=sub<V|u1,u4>;U5:=sub<V|u2,u5>;U6:=sub<V|u3,u6>;
U7:=sub<V|u1,e1>;U8:=sub<V|u2,e2>;U9:=sub<V|u3,e3>;
U11:=sub<V|u4,e2>;U12:=sub<V|u4,e3>;
U13:=sub<V|u5,e1>;U15:=sub<V|u5,e3>;
U16:=sub<V|u6,e1>;U17:=sub<V|u6,e2>;
Include(~M,Dimension(U1));Include(~M,Dimension(U2));
Include(~M,Dimension(U3));Include(~M,Dimension(U4));
Include(~M,Dimension(U5));Include(~M,Dimension(U6));
Include(~M,Dimension(U7));Include(~M,Dimension(U8));
Include(~M,Dimension(U9));
Include(~M,Dimension(U11));Include(~M,Dimension(U12));
Include(~M,Dimension(U13));Include(~M,Dimension(U15));
Include(~M,Dimension(U16));Include(~M,Dimension(U17));
if M eq {@ 2 @} then aal:=aal+1;Include(~LVU,R2);R2;end if;
aal;
end for;end for;end for;end for;end for;end for;
end for;end for;end for;end for;

```

The following script calculates the length of the orbits of four admissible six-tuples. We skip the verification that these are four distinct orbits.

```

G:=GeneralLinearGroup(2,f);
l:=LVU[1];
L1:={@ l @};
l1:=l[1];l2:=l[2];l3:=l[3];l4:=l[4];l5:=l[5];l6:=l[6];
e0:=l1+l2+l3;e1:=l1+l5+l6;e2:=l2+l4+l6;e3:=l3+l4+l5;
for g in G do
Rg:=[l1*g,l2*g,l3*g,l4*g,l5*g,l6*g];
Include(~L1,Rg);
end for;
for j in [1..10] do

```

```

for l in L1 do
l1:=l[1];l2:=l[2];l3:=l[3];l4:=l[4];l5:=l[5];l6:=l[6];
e0:=l1+l2+l3;e1:=l1+l5+l6;e2:=l2+l4+l6;e3:=l3+l4+l5;
Rn1:=[l1,l2,l3,e1,e2,e3];Rn2:=[e1,e2,l3,l5,l4,e0];
Rn3:=[l1,e3,e2,e0,l5,l6];Rn4:=[e3,l2,e1,l4,e0,l6];
Include(~L1,Rn2);Include(~L1,Rn3);Include(~L1,Rn4);
Include(~L1,Rn1);
end for;
print #L1;
end for;

```

```

LVU[1]:
[(0 1), (0 1), (1 0), (1 2), (1 2), (2 4)]
has an orbit of length 28800.

```

```

LVU[3]:
[(0 1), (0 1), (1 0), (1 2), (2 4), (1 2)]
has an orbit of length 57600.

```

```

LVU[4]:
[(0 1), (0 1), (1,0), (1 4), (2 3), (1 1)]
has an orbit of length 57600.

```

```

LVU[7]:
[(0 1), (0 1), (1 0), (1 1), (3 0), (0 2)]
has an orbit of length 57600.

```

The following script calculates the abelianization of the fundamental group of S_3 .

```

G:=CyclicGroup(5);
GG:=DirectProduct(G,G);
P:=Group<x01,x02,x12,x03,x13,x23 | x01*x02*x12*x03*x13*x23,
x01*x02*x12*x01*(x01*x02*x12)^-1*x01^-1,
x01*x02*x12*x02*(x01*x02*x12)^-1*x02^-1,
x01*x02*x12*x12*(x01*x02*x12)^-1*x12^-1,
x01*x03*x13*x01*(x01*x03*x13)^-1*x01^-1,
x01*x03*x13*x03*(x01*x03*x13)^-1*x03^-1,
x01*x03*x13*x13*(x01*x03*x13)^-1*x13^-1,
x02*x03*x23*x02*(x02*x03*x23)^-1*x02^-1,
x02*x03*x23*x03*(x02*x03*x23)^-1*x03^-1,
x02*x03*x23*x23*(x02*x03*x23)^-1*x23^-1,
x12*x13*x23*x12*(x12*x13*x23)^-1*x12^-1,
x12*x13*x23*x13*(x12*x13*x23)^-1*x13^-1,
x12*x13*x23*x23*(x12*x13*x23)^-1*x23^-1>;
S:=Sym(10);
ims := [ S!(1,4,2,5,3)(6,8,10,7,9), S!(1,2,3,4,5)(6,7,8,9,10),
S!(1,2,3,4,5), S!(1,5,4,3,2)(6,7,8,9,10), S!(6,7,8,9,10),
S!(1,2,3,4,5)];
rels := Relations(P);

```

```

rels;
IsSatisfied(rels, ims);
f := hom< P->S | ims >;
K:=Kernel(f);
KK:=Rewrite(P,K);
R1:=P.1^5;
R2:=P.2^5;
R3:=P.3^5;
R4:=P.4^5;
R5:=P.5^5;
R6:=P.6^5;
RR1:=(P.1*P.2*P.3)^5;
RR2:=(P.2*P.4*P.6)^5;
RR3:=(P.1*P.4*P.5)^5;
RR4:=(P.3*P.5*P.6)^5;
PP1:=quo<KK|R1,R2,R3,R4,R5,R6,RR1,RR2,RR3,RR4>;
AbelianQuotient(PP1);

```

Acknowledgement. We are grateful to Fritz Grunewald for his invaluable help in the explicit calculations leading to Theorem 3.3.

This research was performed in the realm of the DFG Schwerpunkt “Globale methoden in der komplexen Geometrie”.

References

- [BPV] W. Barth, C. Peters, A. Van de Ven, *Compact complex surfaces*. Ergeb. Math. Grenzgeb. (3) 4, Springer-Verlag, Berlin 1984. [Zbl 0718.14023](#) [MR 0749574](#)
- [BHH] G. Barthel, F. Hirzebruch, T. Höfer, *Geradenkonfigurationen und Algebraische Flächen*. Aspekte der Math. D4, Vieweg, Braunschweig 1987. [Zbl 0645.14016](#) [MR 0912097](#)
- [Ba-Ca] I. Bauer, F. Catanese, Symmetry and Variation of Hodge Structures. *Asian J. Math.* **8** (2) (2004), 363–390. [Zbl 1070.14013](#) [MR 2129542](#)
- [Bea] A. Beauville, Surfaces algébriques complexes. *Astérisque* **54** (1978). [Zbl 0394.14014](#) [MR 0485887](#)
- [Bir] J. Birman, *Braids, links and mapping class groups*. Ann. of Math. Stud. 82, Princeton University Press, Princeton, N. J., 1975. [Zbl 0305.57013](#) [MR 0375281](#)
- [Bur] P. Burniat, Modèles de surfaces canoniques normales de S_3 et de genre linéaire $11 \leq p^{(1)} \leq 17$. In *Deuxième Colloque de Géométrie Algébrique* (Liège, 1952), Georges Thone, Liège; Masson & Cie, Paris 1952, 185–210. [Zbl 0047.39901](#) [MR 0054285](#)
- [Cat1] F. Catanese, On the moduli space of surfaces of general type. *J. Differential Geom.* **19** (1984), 483–515. [Zbl 0549.14012](#) [MR 0755236](#)
- [Cat2] F. Catanese, Singular bidouble covers and the construction of interesting algebraic surfaces. In *Algebraic geometry: Hirzebruch 70*, Contemp. Math. 241, Amer. Math. Soc., Providence, RI, 1999, 97–120. [Zbl 0964.14012](#) [MR 1718139](#)

- [Cil] C. Ciliberto, Canonical surfaces with $p_g = p_a = 4$ and $K^2 = 5, \dots, 10$. *Duke Math. J.* **48** (1) (1981), 121–157. [Zbl 0468.14011](#) [MR 0610180](#)
- [Enr] F. Enriques, *Le Superficie Algebriche*. Zanichelli, Bologna 1949. [Zbl 0036.37102](#) [MR 0031770](#)
- [Fran] A. Franchetta, Su alcuni esempi di superficie canoniche. *Rend. Sem. Mat. Roma* **3** (1939), 23–28. [JFM 65.0722.04](#) [Zbl 0021.15202](#)
- [Hir] F. Hirzebruch, Arrangements of lines and algebraic surfaces. In *Arithmetic and geometry*, Vol. II, Progr. Math 36, Birkhäuser, Boston, Mass., 1983, 113–140. [Zbl 0527.14033](#) [MR 0717609](#)
- [Ish1] M.-N. Ishida, The irregularities of Hirzebruch's examples of surfaces of general type with $c_1^2 = 3c_2$. *Math. Ann.* **262** (1983), 407–420. [Zbl 0488.14009](#) [MR 0692865](#)
- [Ish2] M.-N. Ishida, Hirzebruch's examples of surfaces of general type with $c_1^2 = 3c_2$. In *Algebraic Geometry*, Lecture Notes in Math. 1016, Springer-Verlag, Berlin 1983, 412–431. [Zbl 0558.14026](#) [MR 0726436](#)
- [Kod] K. Kodaira, On characteristic systems of families of surfaces with ordinary singularities in a projective space. *Amer. J. Math.* **87** (1965), 227–256. [Zbl 0145.43802](#) [MR 0176380](#)
- [Lie] C. Liedtke, Singular abelian covers of algebraic surfaces. *Manuscripta Math.* **112** (2003), 375–390. [Zbl 1044.14016](#) [MR 2067044](#)
- [Max] E. A. Maxwell, Regular canonical surfaces of genus three and four. *Proc. Cambridge Philos. Soc.* **23** (1937), 306–310. [Zbl 0017.03103](#)
- [Miy] Y. Miyaoka, *Algebraic surfaces with positive indices*. In *Classification of algebraic and analytic manifolds*. (Katata, 1982), Progr. Math. 39, Birkhäuser, Boston, Mass., 281–301. [Zbl 0584.14022](#) [MR 0728611](#)
- [Mum] D. Mumford, An algebraic surface with K ample, $K^2 = 9$, $p_g = q = 0$. *Amer. J. Math.* **101** (1) (1979), 233–244. [Zbl 0433.14021](#) [MR 0527834](#)
- [Par] R. Pardini, Abelian covers of algebraic varieties. *J. Reine Angew. Math.* **417** (1991), 191–213. [Zbl 0721.14009](#) [MR 1103912](#)
- [Ter] T. Terada, Quelques propriétés géométriques du domaine de F_1 et le groupe de tresses colorées. *Publ. Res. Inst. Math. Sci.* **17** (1981), 95–111. [Zbl 0466.33010](#) [MR 0613935](#)

Received August 1, 2006

Ingrid C. Bauer, Mathematisches Institut der Universität Bayreuth, Universität Bayreuth, NWII, 95440 Bayreuth, Germany

E-mail: Ingrid.Bauer@uni-bayreuth.de

Fabrizio Catanese, Lehrstuhl Mathematik VIII, Universität Bayreuth, NWII, 95440 Bayreuth, Germany

E-mail: Fabrizio.Catanese@uni-bayreuth.de