

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 83 (2008)

Artikel: Volume growth, curvature decay, and critical metrics
Autor: Tian, Gang / Viaclovsky, Jeff
DOI: <https://doi.org/10.5169/seals-99052>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Volume growth, curvature decay, and critical metrics

Gang Tian* and Jeff Viaclovsky**

Abstract. We make some improvements to our previous results in [TV05a] and [TV05b]. First, we prove a version of our volume growth theorem which does not require any assumption on the first Betti number. Second, we show that our local regularity theorem only requires a lower volume growth assumption, not a full Sobolev constant bound. As an application of these results, we can weaken the assumptions of several of our theorems in [TV05a] and [TV05b].

Mathematics Subject Classification (2000). Anti-self-dual metrics, ALE spaces, curvature decay, ε -regularity, orbifold compactness, volume growth.

Keywords. Primary 53C21; Secondary 58D27, 58E11.

1. Introduction

Riemannian spaces with quadratic curvature decay have been widely studied in the literature, see for example [Abr85], [Abr87], [AG90], [BKN89], [Kas88], [Kas89], [GPZ94], [Gro81], [Zhu94]. All of these works assume that the curvature decay is strictly better than quadratic in the sense that

$$|\mathrm{Rm}| = O(r^{-(2+\delta)}), \quad \text{as } r \rightarrow \infty, \quad (1.1)$$

for some $\delta > 0$, where $r(x) = d(p, x)$ is the distance to a basepoint p , or the weaker assumption that

$$\mathrm{Rm} \geq -\frac{k(r)}{r^2}, \quad (1.2)$$

(meaning all of the sectional curvatures are bounded below accordingly), and the function $k(r)$ satisfies

$$\int_1^\infty \frac{k(r)}{r} dr < \infty. \quad (1.3)$$

*The research of the first author was partially supported by NSF Grant DMS-0302744.

**The research of the second author was partially supported by NSF Grant DMS-0503506.

Spaces satisfying such a curvature decay condition are said to have *asymptotically nonnegative curvature*. We remark that by standard comparison theory, (1.2) and (1.3) imply an upper volume growth estimate,

$$\text{Vol}(B(p, r)) \leq V_0 r^n,$$

for some constant V_0 . Moreover, only a lower bound on the Ricci curvature is needed for this upper volume growth estimate [Zhu94].

It has been understood that such a volume estimate is the core of extending orbifold-compactness theorems for Einstein metrics to metrics satisfying more general equations, such as anti-self-dual or harmonic curvature. To this end, in our investigation of critical metrics in [TV05a], [TV05b], and in the work of [And05], spaces arise with curvature decay as in (1.2), but the function $k(r)$ only satisfies $k(r) \rightarrow 0$ as $r \rightarrow \infty$, and standard comparison arguments do not apply. In [TV05a] we proved an upper volume growth estimate in this case, but our proof required finiteness of the first Betti number to rule out the presence of so-called “bad” annuli. In this paper, we show that adding the condition (1.4) below, eliminates this pathology. For M non-compact, C_S is defined to be the best constant so that

$$\|f\|_{L^{\frac{2n}{n-2}}} \leq C_S \|\nabla f\|_{L^2},$$

for all $f \in C^{0,1}(M)$ with compact support. Let Ric_- denote the negative part of the Ricci tensor.

Theorem 1.1. *Let (M, g) be a complete, noncompact, n -dimensional Riemannian manifold with base point p . Assume that*

$$C_S < \infty,$$

and that

$$\sup_{S(r)} |\text{Rm}_g| = o(r^{-2}),$$

as $r \rightarrow \infty$, where $S(r)$ denotes the sphere of radius r centered at p . If

$$\int_M |\text{Ric}_-|^{\frac{n}{2}} dV_g < \Lambda, \tag{1.4}$$

for some constant $\Lambda \in \mathbb{R}$, then (M, g) has finitely many ends, and there exists a constant C_2 (depending on g) so that

$$\text{Vol}(B(p, r)) \leq C_2 r^n.$$

Furthermore, each end is ALE of order 0.

We have another generalization of our previous results. Consider any system of the type

$$\Delta \text{Ric} = \text{Rm} * \text{Ric}, \quad (1.5)$$

where the right hand side is shorthand notation for a linear combination of terms of the form $A_{ijkl} B_{jl}$, where A_{ijkl} depends on the full curvature tensor, and B_{jl} depends only on the Ricci tensor. We call any metric satisfying a system of the form (1.5) a *critical metric*. In [TV05a, Theorem 3.1], we proved an ε -regularity theorem in dimension four which depended on the Sobolev constant. Here we relax this condition and require only a lower volume growth assumption:

Theorem 1.2. *Let (M, g) be a four-dimensional Riemannian manifold. Assume that (1.5) is satisfied, let $r < \text{diam}(M)/2$, $B(p, r)$ be a geodesic ball around the point p , and $k \geq 0$. If there exists a constant $V_0 > 0$ so that*

$$\text{Vol}(B(q, s)) \geq V_0 s^4$$

for all $q \in B(p, r/2)$ and $s \leq r/2$, then there exist constants ε_0, C_k (depending upon V_0) so that if

$$\|\text{Rm}\|_{L^2(B(p, r))} = \left\{ \int_{B(p, r)} |\text{Rm}|^2 dV_g \right\}^{1/2} \leq \varepsilon_0,$$

then

$$\sup_{B(p, r/2)} |\nabla^k \text{Rm}| \leq \frac{C_k}{r^{2+k}} \left\{ \int_{B(p, r)} |\text{Rm}|^2 dV_g \right\}^{1/2} \leq \frac{C_k \varepsilon_0}{r^{2+k}}.$$

Theorems 1.1 and 1.2 are the main results in this paper. A consequence is that we can (i) remove the Betti number assumption from our volume growth theorem from [TV05b, Theorem 1.2], or (ii) we can relax the Sobolev constant assumption to only an assumption on lower volume growth.

Recall that a metric g is called anti-self-dual if $W_g^+ \equiv 0$, and self-dual if $W_g^- \equiv 0$. As in [TV05b], we specialize to the class of

- (a) self-dual or anti-self-dual metrics with constant scalar curvature,
- (b) metrics with harmonic curvature ($\delta \text{Rm} \equiv 0$),
- (c) Kähler metrics with constant scalar curvature.

We have the following notion of local Sobolev constant. For $p \in M$, and $r > 0$, we define $C_S(p, r)$ to be the best constant such that

$$\|f\|_{L^4} \leq C_S(p, r) \|\nabla f\|_{L^2},$$

for all $f \in C^{0,1}$ with compact support in $B(p, r)$, and define

$$C_S(r) = \sup_{p \in M} C_S(p, r).$$

Let $b_1(M)$ denote the first Betti number of M .

Theorem 1.3. *Let (M, g) be a metric of class (a), (b), or (c) on a smooth, complete four-dimensional manifold M satisfying*

$$\int_M |\text{Rm}_g|^2 dV_g \leq \Lambda, \quad (1.6)$$

for some constant Λ .

Assume that

$$\begin{aligned} \text{Vol}(B(q, s)) &\geq V_0 s^4, \quad \text{for all } q \in M, \text{ and } s \leq \text{diam}(M)/2, \\ b_1(M) &< B_1, \end{aligned}$$

where V_0, B_1 are constants. Then there exists a constant V_1 , depending only upon V_0, Λ, B_1 , such that $\text{Vol}(B(p, r)) \leq V_1 \cdot r^4$, for all $p \in M$ and $r > 0$.

Assume instead that

$$C_S(r) < C_1, \quad \text{for } r < \text{diam}(M)/2,$$

where C_1 is a constant. Then there exists a constant V_2 , depending only upon C_1, Λ , such that $\text{Vol}(B(p, r)) \leq V_2 \cdot r^4$, for all $p \in M$ and $r > 0$.

We have the following improvement of our compactness theorem [TV05b, Theorem 1.3] (we refer to that work for the definition of a multi-fold):

Theorem 1.4. *Let (M_i, g_i) be a sequence of unit-volume metrics of class (a), (b) or (c) on smooth, closed four-dimensional manifolds M_i satisfying*

$$\int_{M_i} |\text{Rm}_{g_i}|^2 dV_{g_i} \leq \Lambda, \quad (1.7)$$

where Λ is a constant.

Assume that

$$\begin{aligned} \text{Vol}(B(q, s)) &\geq V_0 s^4, \quad \text{for all } q \in M, \text{ and } s \leq \text{diam}(M)/2, \\ b_1(M_i) &< B_1, \end{aligned} \quad (1.8)$$

where C_1, Λ, B_1 are constants. Then a subsequence converges to a limit metric space (M_∞, g_∞) which is a compact, connected, critical Riemannian multi-fold. The convergence is smooth away from a finite singular set.

If we assume instead that

$$C_{S_{g_i}}(r) < C_1, \quad \text{for } r < \text{diam}(M)/2,$$

then the same conclusion is true.

Remark. The first Betti number assumption from [TV05a, Theorem 1.1] (which allows non-compact limits) can similarly be removed, and the Sobolev constant assumption relaxed to lower volume growth.

This type of uniform volume estimate, compactness theorem, and related bubbling phenomena follows in a standard fashion from an ε -regularity theorem and a volume growth estimate of the type in Theorem 1.1 and [TV05a, Theorem 1.1]. This passage was first carried out for Einstein metrics [And89], [BKN89], [Tia90], [AC91], [Ban90a], [Ban90b], [Nak92] and extends to more general cases with careful attention to the number of components of geodesic spheres [TV05b], [And05], [CQY07]. In Section 4, we discuss the main technical details of this step under the weaker assumptions in Theorem 1.3 and 1.4.

Let R_g denote the scalar curvature of the metric g . Using the above in case (a), we note the following special corollary.

Corollary 1.5. *Let (M, g_i) be a sequence of unit volume constant scalar curvature anti-self-dual metrics on a fixed closed 4-manifold M . Assume that*

$$|R_{g_i}| < C, \\ \text{Vol}(B(q, s)) \geq V_0 s^4, \quad \text{for all } q \in M, \text{ and } s \leq \text{diam}(M)/2,$$

where C, V_0 are constants. Then a subsequence converges to a limit space (M_∞, g_∞) which is a compact, connected, anti-self-dual Riemannian multi-fold.

Acknowledgements. The authors would like to thank Gilles Carron, John Lott and Joao Santos for enlightening discussions. We also thank Xiuxiong Chen, Edward Fan and Brian Weber for insightful questions and comments on our previous work.

2. Proof of Theorem 1.1

The space of L^2 -harmonic k -forms $\mathcal{H}^k(M)$ is defined to be those $\omega \in \Lambda^k(T^*M)$ satisfying $\Delta\omega = 0$, and $\omega \in L^2(M)$. It is well-known that $\mathcal{H}^k(M) \simeq \bar{H}_{(2)}^k(M)$, the *reduced* L^2 -cohomology, see [Car99]. We next quote the following finiteness theorem.

Theorem 2.1 (Carron, [Car98], [Car99]). *Let (M, g) be a complete Riemannian manifold satisfying for $p > 2$ the Sobolev inequality*

$$\mu_p(M) \left(\int_M |u|^{\frac{2p}{p-2}}(x) dV_g \right)^{1-\frac{2}{p}} \leq \int_M |du|^2(x) dV_g, \quad \text{for all } u \in C_{cpt}^\infty(M). \quad (2.1)$$

If the negative part of the Ricci satisfies

$$\int_M |\text{Ric}_-|^{p/2} dV_g < \infty, \quad (2.2)$$

then $\mathcal{H}^1(M)$ is finite dimensional. If the full Riemannian tensor curvature satisfies

$$\int_M |\text{Rm}|^{p/2} dV_g < \infty, \quad (2.3)$$

then $\mathcal{H}^k(M)$ is finite dimensional for each $1 \leq k \leq n$.

Remark. For convenience of the reader, we give an indication of the proof in [Car99]. A crucial estimate is that the Sobolev constant bound yields an estimate on the heat kernel: there exists a constant C such that for all $x \in M$, and any $t > 0$,

$$k(t, x, x) \leq Ct^{-n/2}.$$

It is then shown that $\mathcal{H}^k(M)$ is finite dimensional, using the Cwickel–Lieb–Rosenbljum estimate adapted to the Riemannian setting. Similar results were also obtained in [BB90]. The Weitzenböck formula for a 1-form is

$$\nabla^* \nabla = \Delta_H + \text{Ric},$$

so for the case of harmonic 1-forms, only the Ricci assumption (2.2) is necessary. For $k > 1$, the Weitzenböck formula depends upon the full curvature tensor, which is why the full curvature assumption (2.3) is required. We note that this method also gives an explicit estimate on the dimension of $\mathcal{H}^k(M)$ in terms of the Sobolev constant $\mu_p(M)$, and the $L^{p/2}$ curvature integral.

We recall a definition.

Definition 2.2. We say a component $A_0(r_1, r_2)$ of an annulus $A(r_1, r_2) = \{q \in M \mid r_1 < d(p, q) < r_2\}$ is *bad* if $S(r_1) \cap \overline{A_0(r_1, r_2)}$ has more than one component, where $S(r_1)$ is the sphere of radius r_1 centered at p .

For an annulus $A_0(r_1, r_2)$, we call a component of $S(r_1) \cap \overline{A_0(r_1, r_2)}$ an *inner sphere*. Note that this may have several components – indeed, this is one of the main subtleties in proving the volume growth estimate in Theorem 1.1.

Theorem 2.3. *Under the assumption in Theorem 1.1, there exists a constant N_0 (depending upon C_S and Λ) such that if A_1, A_2, \dots, A_N is a collection of disjoint connected bad annuli, then $N < N_0$.*

Proof. From the assumptions in Theorem 1.1, the estimate (2.1) is satisfied for $p = n$, therefore Theorem 2.1 says that \mathcal{H}^1 is finite dimensional. Letting $H^1(M)$ and $H_c^1(M)$ denote the first cohomology and first cohomology with compact support, respectively, this implies that

$$\text{Image}(H_c^1(M) \rightarrow H^1(M)) \quad (2.4)$$

has finite dimension, since the space in (2.4) injects into \mathcal{H}^1 (see [And88]). The rest of the argument just uses the finiteness of (2.4). Let $A_i = A(r_i, s_i)$. Without loss of generality, let us assume that the sequence of radii r_i is non-decreasing. Since A_i is bad, the inner sphere $S(r_i) \cap A_i$ has $N_i > 1$ components, call them $\mathcal{C}_{i,j}$, $j = 1, \dots, N_i$. Take any two components, say $\mathcal{C}_{i,1}$ and $\mathcal{C}_{i,2}$, and let $p_{i,j}$ be any point of $\mathcal{C}_{i,j}$. For fixed i , let $\gamma_{i,2,1}(t)$ be a curve in A_i connecting $p_{i,2}$ with $p_{i,1}$. We can always find such a curve since, by assumption, A_i is connected. We can also find a curve $\alpha_{i,1,2}(t)$ connecting $p_{i,1}$ with $p_{i,2}$, with image in $B(p, r_i)$. Joining these curves, we find a closed loop $\beta_i = \gamma_{i,2,1} \# \alpha_{i,1,2}$ based at $p_{i,1}$.

Next, we find a function f_i defined on \bar{A}_i such that f_i is supported in a neighborhood of $\mathcal{C}_{i,1}$, with $f_i = 1$ on smaller neighborhood of $\mathcal{C}_{i,1}$, and $f_i = 0$ in a neighborhood of all other components $\mathcal{C}_{i,j}$. Then the 1-form $\alpha_i = df_i$ clearly has an extension to a smooth 1-form on M , which is closed (but not exact), and is supported on A_i . We claim that

$$\int_{\beta_i} \alpha_i = 1. \quad (2.5)$$

To see this, since α_i is supported on A_i ,

$$\int_{\beta_i} \alpha_i = \int_{\gamma_{i,2,1}(t)} \alpha_i = \int_{\gamma_{i,2,1}(t)} df = f(p_{i,1}) - f(p_{i,2}) = 1.$$

Furthermore,

$$\int_{\beta_i} \alpha_j = 0, \quad i < j.$$

This is true since we have assumed the annuli are indexed by increasing radius, the α_j forms are supported either on a different component, or on an annulus which is further out. We have shown that the α_i define non-zero independent cohomology classes in $\text{Image}(H_c^1(M) \rightarrow H^1(M))$, and we therefore have

$$N_0 \leq \dim \{ \text{Image}(H_c^1(M) \rightarrow H^1(M)) \}. \quad \square$$

Remark. We thank Gilles Carron for providing the above argument, which was much simpler than our original proof. Note also that the finiteness of the dimension of (2.4) will be automatically satisfied if either $H^1(M)$, $H_c^1(M)$, or $\mathcal{H}^1(M)$ is finite dimensional.

We quote the following theorem from our previous work.

Theorem 2.4 ([TV05b, Theorem 5.2]). *Let (M, g) be a complete, non-compact, n -dimensional Riemannian orbifold (with finitely many singular points) with base point p . Assume that there exists a constant $C_1 > 0$ so that*

$$\text{Vol}(B(q, s)) \geq C_1 s^n, \quad (2.6)$$

for any $q \in M$, and all $s \geq 0$. Assume furthermore that as $r \rightarrow \infty$,

$$\sup_{S(r)} |\text{Rm}_g| = o(r^{-2}), \quad (2.7)$$

where $S(r)$ denotes the sphere of radius r centered at p . If (M, g) contains only finitely many disjoint bad annuli, then (M, g) has finitely many ends, and there exists a constant C_2 so that

$$\text{Vol}(B(p, r)) \leq C_2 r^n,$$

Furthermore, each end is ALE of order 0.

Theorem 1.1 follows from the above, since we have shown there are only finitely many bad annuli, and noting that the lower volume growth estimate is implied by the Sobolev constant bound (see [TV05a, Lemma 6.1]). Note also that there is an explicit bound on the number of ends in terms of C_S and Λ , see [Car98, Theorem 3.3].

Theorem 2.5. *Theorem 1.1 is valid if (M, g) is assumed to be a smooth orbifold with finitely many singular points.*

Proof. It can be verified that Carron's arguments are valid for smooth orbifolds, and since Theorem 2.4 is also valid for orbifolds, the proof is identical to the smooth case. Alternatively, instead of using Carron's result for orbifolds, one can argue, albeit non-effectively, as follows. Take a smoothing of (M, g) by cutting out a small ball about each orbifold singularity and gluing in any smooth metric. This can be done because each boundary S^3/Γ certainly bounds some smooth manifold. We now have a smooth manifold (\tilde{M}, \tilde{g}) which is isometric to (M, g) outside of a large ball $B(p, R)$. The manifold (\tilde{M}, \tilde{g}) has the same asymptotic behaviour as the original (M, g) , so all of the assumptions of Theorem 1.1 are satisfied by (\tilde{M}, \tilde{g}) . Applying Theorem 1.1, we obtain an upper volume estimate for balls in (\tilde{M}, \tilde{g}) , which clearly implies an upper volume estimate for the original (M, g) , since the asymptotics are the same. \square

3. Sobolev constant and local regularity

Recall from the introduction that we have the following notion of local Sobolev constant. For $p \in M$, and $r > 0$, we define $C_S(p, r)$ to be the best constant such that

$$\|f\|_{L^4} \leq C_S(p, r) \|\nabla f\|_{L^2},$$

for all $f \in C^{0,1}$ with compact support in $B(p, r)$. Clearly

$$C_S(p, r) \text{ is an increasing function of } r,$$

and

$$\lim_{r \rightarrow 0} C_S(p, r) = C_E, \quad (3.1)$$

where C_E is the best constant for the Euclidean Sobolev inequality.

Proposition 3.1. *Assume that (1.5) is satisfied, and let $B(p_0, 2r)$ be a geodesic ball around the point p_0 . Assume there exists a constant $V_0 > 0$ so that*

$$\text{Vol}(B(q, s)) \geq V_0 s^4 \quad (3.2)$$

for all $q \in B(p_0, r)$, and $s \leq r$.

Then there exists a constant ε_0 (depending upon V_0) so that if

$$\|\text{Rm}\|_{L^2(B(p_0, 2r))} = \left\{ \int_{B(p_0, 2r)} |\text{Rm}|^2 dV_g \right\}^{1/2} \leq \varepsilon_0,$$

then

$$C_S(p_0, r/2) \leq C V_0^{-1/4},$$

and C does not depend on V_0 .

Remark. We conjecture that for (1.5), ε_0 can moreover be taken independent of V_0 and assumption (3.2) is not necessary. This was recently proved for Einstein metrics by Cheeger–Tian [CT06]. In fact, in [CT06, Section 11] this was already conjectured to hold for anti-self-dual metrics and Kähler metrics with constant scalar curvature.

Proof. Without loss of generality, rescale so that $r = 1$. The proof goes by contradiction. Assume we have a sequence of critical metrics g_i , $i = 1, \dots, \infty$, and a sequence $\varepsilon_i \rightarrow 0$ as $i \rightarrow \infty$, with

$$\|\text{Rm}\|_{L^2(B(p_0, 2))} \leq \varepsilon_i,$$

and that

$$C_S(p_0, 1/2) > CV_0^{-1/4}$$

(we will choose C later).

We first choose a “nice” ball, one for which the Sobolev constant is controlled in the ball, and also for nearby points. The following lemma is for a fixed metric.

Lemma 3.2. *There exist a point $p_\infty \in B(p, 1)$ and a radius $0 < r_\infty \leq 1/2$ such that $C_S(p_\infty, r_\infty) > CV_0^{-1/4}$, and $C_S(p, r_\infty/2) \leq CV_0^{-1/4}$ for all $p \in B(p_\infty, r_\infty)$.*

Proof. From assumption, we have $C_S(p_0, 1/2) > CV_0^{-1/4}$. If $C_S(p, 1/4) \leq CV_0^{-1/4}$ for all $p \in B(p_0, 1/2)$, then let $B = B(p_0, 1/2)$. Otherwise, there exists a point $p_1 \in B(p_0, 1/2)$ with $C_S(p_1, 1/4) > CV_0^{-1/4}$. If $C_S(p, 1/8) \leq CV_0^{-1/4}$ for all $p \in B(p_1, 1/4)$, then we let $B = B(p_1, 1/4)$. We continue inductively, assume we have chosen p_{i-1} with $p_{i-1} \in B(p_{i-2}, 2^{-i+1})$ and $C_S(p_{i-1}, 2^{-i}) > CV_0^{-1/4}$. If $C_S(p, 2^{-i}) \leq CV_0^{-1/4}$ for all $p \in B(p_{i-1}, 2^{-i})$ then we stop, and let $B = B(p_{i-1}, 2^{-i})$. Otherwise, there exists a point $p_i \in B(p_{i-1}, 2^{-i})$ and $C_S(p_i, 2^{-i-1}) > CV_0^{-1/4}$.

We claim this procedure must stop in finitely many steps. We have

$$\begin{aligned} d(p_0, p_i) &\leq d(p_0, p_1) + \sum_{j=1}^{i-1} d(p_j, p_{j+1}) \\ &< 1/2 + \sum_{j=1}^{i-1} 2^{-j-1} \\ &< 1/2 + 1/2 = 1. \end{aligned}$$

The sequence of points $\{p_i\}$ are therefore all contained in the unit ball $B(p, 1)$. If the process did not stop, then there would exist a limit point q . Our first restriction on C is that $CV_0^{-1/4} > C_E$. But from the choices, we would clearly have $\lim_{r \rightarrow 0} C_S(q, r) > C_E$, which contradicts (3.1). \square

Now we apply the lemma to each metric g_i , to find points $p_i \in B(p, 1)$ (ball in the g_i metric) and $r_i < 1/2$, with $C_S(p_i, r_i) > CV_0^{-1/4}$, and $C_S(p, r_i/2) \leq CV_0^{-1/4}$ for all $p \in B(p_i, r_i)$ (Sobolev constant with respect to the g_i metric).

Now we rescale to make r_i unit size, that is, define $\tilde{g}_i = r_i^{-2}g_i$. and consider the sequence of balls $\tilde{B}(p_i, 2)$. By scale invariance, we have

$$\|\text{Rm}\|_{L^2(\tilde{B}(p_i, 2))} \leq \varepsilon_i.$$

By our previous ε -regularity theorem [TV05a, Theorem 3.1], the curvature and all of its derivatives are uniformly bounded in $B(p_i, 3/2)$

$$|\nabla^k \text{Rm}| \leq C_k \varepsilon_i,$$

where C_k depends upon k and $C V_0^{-1/4}$. Since $\varepsilon_i \rightarrow 0$, we use the theorem of Cheeger–Gromov to extract a flat limit space, $B(p_\infty, 3/2)$, with smooth convergence to the limit. From the assumption $\text{Vol}(B(p, r)) \geq V_0 r^n$, and since the limit space is flat, we therefore get a bound on the Sobolev constant of the limit space, $C_S \leq C_1 V_0^{-1/4}$. Choosing $C > C_1$, we arrive at a contradiction. \square

Theorem 1.2 then follows from Proposition 3.1 and [TV05a, Theorem 3.1].

4. Main volume estimate

In this section, we discuss the proof of Theorems 1.3 and 1.4, and Corollary 1.5. We first prove Theorem 1.3.

Proof of Theorem 1.3. The proof is based on the argument from [TV05b], with some modifications. First, let us assume that the volume growth estimate from Theorem 1.3 holds for $r \leq r_0$, where r_0 is some fixed scale. That is, let us assume that

$$\text{Vol}(B_{g_i}(p, r)) \leq V r^4 \quad (4.1)$$

for all $p \in M_i$, and all $r \leq r_0$.

From the ε -regularity Theorem 1.2, (M_j, g_j, p_j) converges to a limiting multi-fold $(M_\infty, g_\infty, p_\infty)$ for some subsequence $\{j\} \subset \{i\}$, with finitely many C^0 -multi-fold singular points (recall that a C^0 -multi-fold point means that for each cone at a singular point, the metric has a continuous extension to the universal cover of the punctured cone). The argument for this is the same as in [TV05b].

Proposition 4.1. *The singular points are smooth orbifold points. That is, if x is a singular point, then for some $\delta > 0$, the universal cover of $B(x, \delta) \setminus \{x\}$ is diffeomorphic to a punctured ball $B^4 \setminus \{0\}$ in \mathbb{R}^4 , and the lift of g , after diffeomorphism, extends to a smooth critical metric \tilde{g} on B^4 .*

Proof. In the case we assume a bound on the Sobolev constant (but no Betti number bound), this follows directly from [TV05b, Theorem 6.4]. In the case where we only assume a lower volume growth bound, we claim that for any singular point p , there exists a constant C_s so that

$$\|u\|_{L^4(B(p, \varepsilon))} \leq C_s \|\nabla u\|_{L^2(B(p, \varepsilon))}, \quad u \in C_{cpt}^{0,1}(B(p, \varepsilon)).$$

Indeed, in a neighborhood of a singular point, a C^0 -orbifold is just a C^0 -perturbation of a flat cone, so clearly it satisfies a Sobolev inequality. The result then follows again directly from [TV05b, Theorem 6.4]. \square

Proposition 4.2. *If (4.1) is satisfied, and (M_j, g_j, p_j) converges to a limiting multi-fold $(M_\infty, g_\infty, p_\infty)$, as $j \rightarrow \infty$, then there is a bound on the number of cones at a singular point of convergence, depending only upon Λ , V_0 , and B_1 in the first case, and depending only upon Λ and $C_S(r)$ in the second case. The bound does not depend upon the constant V in (4.1).*

Proof. We consider the first case. At any limiting multi-fold point p'_∞ , for $\delta > 0$ small, we look at the balls $U_j = B(p'_j, \delta) \subset M_j$ where $p'_j \rightarrow p'_\infty$ as $j \rightarrow \infty$. These are manifolds with boundary components spherical space forms, with the metrics arbitrarily close to the limiting multi-fold cone metric. We apply the Gauss–Bonnet Theorem to conclude

$$\chi(U_j) \sim \int_{U_j} \text{Rm} * \text{Rm} + \sum_k \frac{1}{|\Gamma_k|}.$$

where $\text{Rm} * \text{Rm}$ is a quadratic curvature expression, the second sum is over the boundary components, and Γ_k denotes the orbifold group for each cone. Note the final term is an approximation of the boundary integral, with vanishing error term as $\delta \rightarrow 0$. In terms of Betti numbers,

$$1 - b_1(U_j) + b_2(U_j) - b_3(U_j) \sim \int_{U_j} \text{Rm} * \text{Rm} + \sum_k \frac{1}{|\Gamma_k|}. \quad (4.2)$$

We write $M_j = U_j \cup V_j$, where

$$U_j \cap V_j \sim \coprod_k S^3 / \Gamma_k.$$

The Mayer–Vietoris sequence for U_j and V_j is

$$0 = H_2(U_j \cap V_j) \rightarrow H_2(U_j) \oplus H_2(V_j) \rightarrow H_2(M_j) \rightarrow H_1(U_j \cap V_j) = 0,$$

which gives $b_2(M_j) = b_2(U_j) + b_2(V_j)$.

Using (4.2), we can estimate

$$\sum_k \frac{1}{|\Gamma_k|} \leq 1 + b_2(U_j) + C \cdot \Lambda \leq 1 + b_2(M_j) + C \cdot \Lambda.$$

Applying the Gauss–Bonnet formula to the oriented manifold M_j ,

$$2 - 2b_1(M_j) + b_2(M_j) = \int_{M_j} \text{Rm} * \text{Rm},$$

yields the estimate

$$b_2(M_j) \leq C \cdot \Lambda + 2b_1(M_j) - 2,$$

so we have

$$\sum_k \frac{1}{|\Gamma_k|} \leq 2C \cdot \Lambda + 2B_1 - 1.$$

The lower volume growth estimate (1.8) clearly implies that the orders of the orbifold groups are bounded strictly from below, so the proposition follows in this case.

In the second case, we use Carron's bound on the number of ends N of a complete space X [Car98, Theorem 0.4]

$$N \leq 1 + C \cdot C_S \int_X |\text{Ric}_-|^2 dx \leq 1 + C \cdot C_S \cdot \Lambda, \quad (4.3)$$

which, as mentioned above, is also valid for smooth orbifolds. Since we have a volume growth bound (4.1) we can perform a standard bubbling analysis. This analysis was carried out for Einstein metrics in [Ban90a], [Ban90b], see also [Nak92] for a nice description of this bubbling process. The same method works in our case, with a few modifications.

We recall the main steps of the bubbling analysis. Let S denote the singular set of convergence. Note that in contrast to the Einstein case, a point $p \in S$ may actually be a smooth point of the limit. For $0 < r_1 < r_2$, we let $D(r_1, r_2)$ denote $B(p, r_2) \setminus B(p, r_1)$. Given a singular point $x \in S$, take a sequence $x_j \in (M_j, g_j)$ such that $\lim_{j \rightarrow \infty} x_j = x$ and $B(x_j, \delta)$ converges to $B(x, \delta)$ for all $\delta > 0$. We choose a radius r_∞ sufficiently small and the sequence x_j to satisfy

$$\sup_{B(x_j, r_\infty)} |\text{Rm}_{g_j}|^2 = |R_{g_j}|^2(x_j) \rightarrow \infty \quad \text{as } j \rightarrow \infty,$$

and

$$\int_{B(x, r_\infty)} |R_{g_\infty}|^2 dV_{g_\infty} \leq \varepsilon_0/2,$$

where ε_0 is the constant in the ε -regularity Theorem 1.2.

We next choose $r_0(j)$ so that

$$\int_{D(r_0, r_\infty)} |R_{g_j}|^2 dV_{g_j} = \varepsilon_0,$$

and again $D_j(r_0, r_\infty) = B(x_j, r_\infty) \setminus B(x_j, r_0)$. An important note, which differs from the Einstein case, the annulus $D(r_0, r_\infty)$ may have several components.

Since the curvature is concentrating at x , $r_0(j) \rightarrow 0$ as $j \rightarrow \infty$, the rescaled sequence $(M_j, r_0(j)^{-2}g_j, x_j)$ has a subsequence which converges to a complete, non-compact multi-fold with finitely many singular points, which we denote by

$$M_{\infty, i_1}, \quad 1 \leq i_1 \leq \#\{S\}.$$

Note that by assumption, M_{∞, i_1} has bounded Sobolev constant. Since

$$\int_{D(1, \infty)} |\text{Rm}|^2 dV_g \leq \varepsilon_0,$$

there are no singular points outside of $B(x, 1)$.

On the noncompact ends, since we are assuming an upper volume growth estimate, the proof of [TV05a, Theorem 1.3] allows us to conclude that the metric is ALE of order τ for any $\tau < 2$. As in [Ban90a, Proposition 4], we conclude that the neck regions (for large j) will be arbitrarily close to a portion of a flat cones \mathbb{R}^4/Γ , possibly several cones if M_{i_1} has several ends (again in contrast with the Einstein case). The convergence at a singular point x_{i_1} is that the ALE multi-fold M_{∞, i_1} is bubbling off, or scaled down to a point in the limit, with each end of M_{∞, i_1} corresponding exactly to a cone at a multi-fold point of the limit (M_{∞}, g_{∞}) . An important fact is that the fundamental group of an end and the group of the orbifold cone onto which the end is glued must be isomorphic (together with their actions on \mathbb{R}^4), this also follows from the proof in [Ban90a], see also [Nak92, Theorem 2.5]. In particular, the number of ends of M_{∞, i_1} is the same as the number of components of $\partial B(x_{i_1}, r_{\infty})$.

To further analyze the degeneration at the singular points, we look at the multi-fold M_{∞, i_1} with singular set S_{i_1} . If S_{i_1} is empty, then we can stop, as the number of cones must be bounded. If not, we do the same process as above for each singular point of M_{∞, i_1} , and obtain ALE multi-folds

$$M_{\infty, i_1, i_2}, \quad 1 \leq i_2 \leq \#\{S_{i_1}\},$$

each of which are just limits of re-scalings of the original sequence around the appropriate basepoints. If M_{∞, i_1, i_2} has singularities, then we repeat the procedure. This process must terminate in finite steps, since in this construction, each singularity takes at least ε_0 of curvature: the L^2 -curvature bound (1.6) clearly implies the number of steps in the procedure is bounded by the number $b = \Lambda/\varepsilon_0$. As pointed out in [Ban90b], there could be some overlap if any singular point lies on the boundary of $B(1)$ at some stage in the above construction. But there can only be finitely many, and then there must also be a singular point in the interior of $B(1)$, so we still take away at least ε_0 of curvature at each step.

Note that in each step of the bubbling process, each end of the multi-fold obtained in the k th step will be glued to one of the cones at a multi-fold singular point of the $(k-1)$ st step, along a neck region which is close to a portion of a flat cone. Again,

we use the fact that the fundamental group of an end and the group of the orbifold cone onto which the end is glued must be isomorphic (with isomorphic actions on \mathbb{R}^4). Consequently, the bound N on the number of ends (4.3) and the bound b on the number of steps in the process, together imply that the number of cones at any singular point must be bounded by N^b . \square

Remark. As was discussed in [TV05b, page 369], in the Kähler case only irreducible singular points can occur in limit. We remark that this still holds under the weaker assumptions in this paper. That is, in the Kähler case there are never multiple cones at a singular point of convergence.

Remark. The above proposition was proved using an alternative method in [TV05b, Proposition 7.2]. However, that argument required both a Sobolev constant and a first Betti number bound.

Next, we have

Proposition 4.3. *Let (M, g) be a smooth multi-fold with finitely many singular points and g a critical metric. Assume that the number of singular points is uniformly bounded by the number N_1 , that the number of cones at any singular point is uniformly bounded by the number N_2 , and that there exists a constant $V_0 > 0$ so that*

$$\text{Vol}(B(q, s)) \geq V_0 s^4$$

for all $q \in B(p, 2)$, and $s \leq 2$. If

$$\|\text{Rm}\|_{L^2(B(p, 2))} = \left\{ \int_{B(p, 2)} |\text{Rm}|^2 dV_g \right\}^{1/2} \leq \varepsilon_0,$$

then there exists a constant A_0 such that

$$\text{Vol}(B(p, 1)) \leq A_0,$$

where A_0 depends only upon N_1 , N_2 and V_0 .

Proof. If (M, g) is smooth, then by Theorem 1.2,

$$\sup_{B(p, 1)} |\text{Rm}| \leq \frac{1}{4} C \varepsilon_0. \quad (4.4)$$

By Bishop's volume comparison theorem, we must have $\text{Vol}(B(p, 1)) \leq A'$, where A' depends only lower volume growth constant V_0 (since ε_0 only depends on V_0).

In the case that (M, g) is a smooth orbifold, we claim that the ε -regularity theorem still holds in this setting. This is because the argument in [TV05a, Theorem 3.1] uses

integration by parts. One then performs a similar argument, by cutting out small balls of radius δ around the singular points and verifying that the resulting boundary terms vanish as $\delta \rightarrow 0$ (note that it is crucial that the orbifold points be *smooth* for this to be valid). Furthermore, Proposition 3.1 remains valid for a smooth orbifold with a bounded number of singular points. The details, which we omit, are straightforward. Consequently, the estimate (4.4) holds. Next, Bishop's volume comparison theorem remains valid for smooth orbifolds (see for example [Bor93]), with the same constant (or better) as in the smooth case. So for an orbifold (no multiple cone points), we still obtain

$$\text{Vol}(B(p, 1)) \leq A'.$$

In the more general situation of a multi-fold, since there are at most N_2 cones at each of the N_1 singular points, clearly we obtain the estimate

$$\text{Vol}(B(p, 1)) \leq N_1 N_2 A' \equiv A_0. \quad \square$$

We also note the following fact, for any metric,

$$\lim_{r \rightarrow 0} \text{Vol}(B(p, r)) r^{-4} = \omega_4,$$

where ω_4 is the volume ratio of the Euclidean metric on \mathbb{R}^4 . Clearly, $A_0 \geq \omega_4$.

For any metric (M, g) , define the maximal volume ratio as

$$MV(g) = \max_{x \in M, r \in \mathbb{R}^+} \frac{\text{Vol}(B(x, r))}{r^4}.$$

If the theorem is not true, then there exists a sequence of critical manifolds (M_i, g_i) , with $MV(g_i) \rightarrow \infty$, that is, there exist points $x_i \in M_i$, and $t_i \in \mathbb{R}^+$ so that

$$\text{Vol}(B(x_i, t_i)) \cdot t_i^{-4} \rightarrow \infty, \quad (4.5)$$

as $i \rightarrow \infty$. We choose a subsequence (which for simplicity we continue to denote by the index i) and radii r_i so that

$$2 \cdot A_0 = \text{Vol}(B(x_i, r_i)) \cdot r_i^{-4} = \max_{r \leq r_i} \text{Vol}(B(x_i, r)) \cdot r^{-4},$$

We furthermore assume that x_i is chosen so that r_i is minimal, that is, the smallest radius for which there exists some $p \in M_i$ such that $\text{Vol}(B_{g_i}(p, r)) \leq 2A_0 r^4$, for all $r \leq r_i$.

First let us assume that r_i has a subsequence converging to zero. For this subsequence (which we continue to index by i), we consider the rescaled metric $\tilde{g}_i = r_i^{-2} g_i$, so that $B_{g_i}(x_i, r_i) = B_{\tilde{g}_i}(x_i, 1)$. From the choice of x_i and r_i , the metrics \tilde{g}_i have bounded volume ratio, in all balls of unit size.

From the argument above, some subsequence converges on compact subsets to a complete length space $(M_\infty, g_\infty, p_\infty)$ with finitely many point singularities. The limit could either be compact or non-compact. In either case, the arguments above imply that the limit is a Riemannian multi-fold.

Claim 4.4. *The conclusions of Theorem 1.1 hold for the limit $(M_\infty, g_\infty, p_\infty)$*

Proof. In the case that M_∞ is compact the claim is trivial. For M_∞ non-compact, the remarks at the end of [TV05a, Section 3] shows that assumption (2.7) is satisfied. Also, from [TV05a, Lemma 6.1], the Sobolev constant bound implies a lower volume growth bound (this is also valid for orbifolds), so (2.6) is satisfied in both cases.

If we make the assumption on $b_1(M_i)$ (but no Sobolev constant assumption), then the proof is the same as in our previous work [TV05b, Claim 7.1]: since $b_1(M_i) < \infty$, the limit must have finitely many bad annuli. The finiteness of ends and upper volume growth estimate then follow from Theorem 2.4. In the case where we assume the Sobolev constant bound (but no b_1 bound), the result is contained in Theorem 2.5. \square

From Claim 4.4, we have that M_∞ has only finitely many ends, and that there exists a constant $A_1 \geq 2A_0$ so that

$$\text{Vol}(B_{g_\infty}(p_\infty, r)) \leq A_1 r^4, \quad \text{for all } r > 0. \quad (4.6)$$

If M_∞ is compact, then clearly the estimate (4.6) is valid for some constant $A_1 \geq 2A_0$, since the limit has finite diameter and volume, and the estimate holds for $r \leq 1$.

The inequality

$$\int_{B_{g_i}(x_i, 2r_i)} |\text{Rm}_i|^2 dV_i > \varepsilon_0, \quad (4.7)$$

must hold; otherwise, as remarked above, we would have $\text{Vol}(B_{g_i}(x_i, r_i)) \leq A_0 r_i^4$, which violates Proposition 4.3.

If the r_i are bounded away from zero then there exists a radius t so that

$$\text{Vol}(B_{g_i}(p, r)) \leq 2A_0 r^4, \quad \text{for all } r \leq t, p \in M_i.$$

We repeat the argument from the first case, but without any rescaling. Since the maximal volume ratio is bounded on small scales, we can extract an multi-fold limit. The limit can either be compact or non-compact, but the inequality (4.6) will also be satisfied for some A_1 . Following the same argument, we find a sequence of balls satisfying (4.7).

We next return to the (sub)sequence (M_i, g_i) and extract another subsequence so that

$$2600 \cdot A_1 = \text{Vol}(B(x'_i, r'_i)) \cdot (r'_i)^{-4} = \max_{r \leq r'_i} \text{Vol}(B(x'_i, r)) \cdot r^{-4}. \quad (4.8)$$

Again, we assume that x'_i is chosen so that r'_i is minimal, that is, the smallest radius for which there exists some $p \in M_i$ such that $\text{Vol}(B_{g_i}(p, r)) \leq 2600A_1r^4$, for all $r \leq r_i$. Clearly, $r_i < r'_i$.

Arguing as above, if $r'_i \rightarrow 0$ as $i \rightarrow \infty$, then we repeat the rescaled limit construction, but now with scaled metric $\tilde{g}_i = (r'_i)^{-2}g_i$, and basepoint x'_i . We find a limiting multi-fold $(M'_\infty, g'_\infty, p'_\infty)$, and a constant $A_2 \geq 2600A_1$ so that

$$\text{Vol}(B_{g'_\infty}(p'_\infty, r)) \leq A_2r^4 \quad \text{for all } r > 0.$$

For the same reason as above, we must have

$$\int_{B_{g_j}(x'_j, 2r'_j)} |\text{Rm}_j|^2 dV_j > \varepsilon_0.$$

If r'_i is bounded below, we argue similarly, but without any rescaling.

We next consider the ratio r'_i/r_i . There are 2 possible cases.

Case (i): there exists a subsequence (which we continue to index with i) satisfying $r'_i < Cr_i$ for some constant C .

Case (ii):

$$\lim_{i \rightarrow \infty} \frac{r'_i}{r_i} = \infty.$$

In Case (i) we proceed as follows: We claim that for i sufficiently large, the balls $B(x_i, 2r_i)$ (from the first subsequence) and $B(x'_i, 2r'_i)$ (from the second) must be disjoint because of the choice in (4.8). To see this, if $B(x_i, 2r_i) \cap B(x'_i, 2r'_i) \neq \emptyset$, then $B(x'_i, 2r'_i) \subset B(x_i, 6r'_i)$. Then (4.6) and (4.8) yield

$$\begin{aligned} 2600A_1(r'_i)^4 &= \text{Vol}(B(x'_i, r'_i)) < \text{Vol}(B(x'_i, 2r'_i)) \\ &< \text{Vol}(B(x_i, 6r'_i)) \leq 2A_1(6r'_i)^4 = 2592A_1(r'_i)^4, \end{aligned}$$

which is a contradiction (note the last inequality is true for i sufficiently large since (4.6) holds for the limit, which is valid only in Case (i)).

In Case (ii) we argue as follows. If the balls $B(x_i, 2r_i)$ (from the first subsequence) and $B(x'_i, 2r'_i)$ (from the second) are disjoint for all i sufficiently large, then we proceed to the next step. Otherwise, we look again at the scaling so that $r'_i = 1$: $\tilde{g}_i = (r'_i)^{-2}g_i$, and basepoint x'_i . Then in this rescaled metric,

$$\text{Vol}(B(x'_i, 1)) = 2600A_1.$$

As above, we have a limiting multi-fold $(M'_\infty, g'_\infty, p'_\infty)$, satisfying

$$\text{Vol}(B(p'_\infty, 1)) = 2600A_1.$$

Proposition 4.2 implies the number of cones at a multi-fold point is a priori bounded, so from Proposition 4.3, we conclude that

$$\int_{B_{g'_\infty}(p'_\infty, 2)} |\text{Rm}|^2 dV > \varepsilon_0,$$

There is now a singular point of convergence corresponding to the balls $B(x_i, r_i)$ in the first subsequence. But since we are in Case (ii), in the g'_i metric, these balls must limit to a point in M'_∞ . The only possibility is that the original sequence satisfied

$$\int_{B_{g_i}(x'_i, 2r'_i)} |\text{Rm}_{g_i}|^2 dV_i > 2\varepsilon_0,$$

for all i sufficiently large.

We repeat the above procedure, considering possible Cases (i) and (ii) at each step. At the k th step, we can always account for at least $k \cdot \varepsilon_0$ of L^2 -curvature. The process must terminate in finitely many steps from the bound $\|\text{Rm}_i\|_{L^2} < \Lambda$. This contradicts (4.5), which finishes the proof. \square

Remark. In the proof of [TV05b, Theorem 1.2], we neglected to consider the possibility of Case (ii), the above fixes this minor oversight. Another point is that Propositions 4.2 and 4.3 depend crucially on the limiting multi-folds being *smooth*, which requires the removable singularity result [TV05b, Theorem 6.4]. It will be interesting to find a valid proof not using the removable singularity theorem. Furthermore, it has been proposed long ago that the upper volume growth estimate in Theorem 1.4 should hold without requiring either a Betti number assumption or a Sobolev constant bound (only assuming a lower volume growth bound). However, there is no *complete* proof of this yet. An extension of the existing techniques will require a highly nontrivial analysis of the connectedness properties of geodesic spheres and annuli. This paper overcomes the difficulty in several important cases, but we emphasize that the result is still not known in full generality.

Theorem 1.4 is proved in a similar manner as the corresponding theorem in [TV05b], using Theorems 1.1, 1.2, and the volume growth estimate in Theorem 1.3.

We next prove Corollary 1.5, which is a simple consequence of the Gauss–Bonnet Theorem and Hirzebruch Signature Theorem in dimension four (see [Bes87]):

$$\begin{aligned} 8\pi^2 \chi(M) &= \frac{1}{6} \int_M R^2 - \frac{1}{2} \int_M |\text{Ric}|^2 + \int_M |W|^2, \\ 12\pi^2 \tau(M) &= \int_M |W^+|^2 - \int_M |W^-|^2. \end{aligned}$$

In the anti-self-dual case, $W^+ \equiv 0$, so we have

$$\begin{aligned} 8\pi^2 \chi(M) &= \frac{1}{6} \int_M R^2 - \frac{1}{2} \int_M |\text{Ric}|^2 + \int_M |W^-|^2, \\ 12\pi^2 \tau(M) &= - \int_M |W^-|^2. \end{aligned} \quad (4.9)$$

Add these equations together to obtain

$$\begin{aligned} 8\pi^2 \chi(M) + 12\pi^2 \tau(M) &= \frac{R^2}{6} \text{Vol}(M) - \frac{1}{2} \int_M |\text{Ric}|^2 \\ &= \frac{R^2}{6} - \frac{1}{2} \int_M |\text{Ric}|^2, \end{aligned}$$

since the scalar curvature is constant, and $\text{Vol} = 1$. The topology of the manifold M is fixed, and the scalar curvature is uniformly bounded, so we find that $\frac{1}{2} \int_M |\text{Ric}|^2$ is bounded. Also (4.9) yields a bound on $\int_M |W^-|^2$, so we have the estimate

$$\int_M |\text{Rm}|^2 dV_g < \Lambda,$$

for some constant Λ . Therefore the assumptions of Theorem 1.4 are satisfied, which finishes the proof.

5. ALE metrics and removable singularities

A related problem is to find geometric conditions so that each end of a complete space will be ALE of order $\tau > 0$, and to determine the optimal order of decay. In [TV05a] we examined this problem for the following cases:

- (a) Self-dual or anti-self-dual metrics with zero scalar curvature.
- (b) Scalar-flat metrics with harmonic curvature.

By using Theorems 1.1, 1.2, and the volume growth estimate in Theorem 1.3, we have the following improvement of [TV05a, Theorem 1.3]:

Theorem 5.1. *Let (M, g) be a complete, noncompact four-dimensional Riemannian manifold with g of class (a) or (b) satisfying,*

$$\int_M |\text{Rm}_g|^2 dV_g < \infty.$$

Assume that

$$\begin{aligned}\mathrm{Vol}(B(q, s)) &\geq V_0 s^4, \quad \text{for all } q \in M \text{ and } s > 0, \\ b_1(M) &< \infty,\end{aligned}$$

then (M, g) has finitely many ends, and each end is ALE of order τ for any $\tau < 2$. If we assume instead that

$$C_S < \infty,$$

then the same conclusion holds.

To conclude, we mention that the following removable singularity theorem for critical metrics is expected:

Let (M, g) be a C^0 -orbifold with singular point at x , and g be a critical metric satisfying (1.5). Suppose that

$$\int_{B(x, 1)} |\mathrm{Rm}_g|^2 dV_g < \infty.$$

Then the metric g extends to $B(x, 1)$ as a smooth orbifold metric. That is, for some small $\delta > 0$, the universal cover of $B(x, \delta) \setminus \{x\}$ is diffeomorphic to a punctured ball $B^4 \setminus \{0\}$ in \mathbb{R}^4 , and the lift of g , after diffeomorphism, extends to a smooth critical metric \tilde{g} on B^4 .

This will be addressed in a forthcoming paper¹. Such a removable singularity theorem was proved for the special cases of (a), (b), and (c) in [TV05b, Theorem 6.4]. The above generalization is crucial in extending Theorems 1.3, 1.4, and 5.1 to the more general class of critical metrics satisfying (1.5), in particular, Bach-flat metrics.

References

- [Abr85] Uwe Abresch, Lower curvature bounds, Toponogov's theorem, and bounded topology. *Ann. Sci. École Norm. Sup. (4)* **18** (4) (1985), 651–670. [Zbl 0595.53043](#) [MR 0839689](#)
- [Abr87] U. Abresch, Lower curvature bounds, Toponogov's theorem, and bounded topology. II. *Ann. Sci. École Norm. Sup. (4)* **20** (3) (1987), 475–502. [Zbl 0651.53031](#) [MR 0925724](#)
- [AC91] Michael T. Anderson and Jeff Cheeger, Diffeomorphism finiteness for manifolds with Ricci curvature and $L^{n/2}$ -norm of curvature bounded. *Geom. Funct. Anal.* **1** (3) (1991), 231–252. [Zbl 0764.53026](#) [MR 1118730](#)

¹After completion of this paper, this result was proved by Jeffrey Streets [Str07].

- [AG90] Uwe Abresch and Detlef Gromoll, On complete manifolds with nonnegative Ricci curvature. *J. Amer. Math. Soc.* **3** (2) (1990), 355–374. [Zbl 0704.53032](#) [MR 1030656](#)
- [And88] Michael T. Anderson, L^2 harmonic forms on complete Riemannian manifolds. In *Geometry and analysis on manifolds* (Katata/Kyoto, 1987), Lecture Notes in Math. 1339, Springer-Verlag, Berlin 1988, 1–19. [Zbl 0652.53030](#) [MR 0961469](#)
- [And89] Michael T. Anderson, Ricci curvature bounds and Einstein metrics on compact manifolds. *J. Amer. Math. Soc.* **2** (3) (1989), 455–490. [Zbl 0999661](#) [MR 0999661](#)
- [And05] Michael T. Anderson, Orbifold compactness for spaces of Riemannian metrics and applications. *Math. Ann.* **331** (4) (2005), 739–778. [Zbl 1071.53025](#) [MR 2148795](#)
- [Ban90a] Shigetoshi Bando, Bubbling out of Einstein manifolds. *Tohoku Math. J.* (2) **42** (2) (1990), 205–216. [Zbl 0719.53025](#) [MR 1053949](#)
- [Ban90b] Shigetoshi Bando, Correction and addition: “Bubbling out of Einstein manifolds”. *Tohoku Math. J.* (2) **42** (4) (1990), 587–588. [Zbl 0762.53030](#) [MR 1053949](#)
- [BB90] P. Bérard and G. Besson, Number of bound states and estimates on some geometric invariants. *J. Funct. Anal.* **94** (2) (1990), 375–396. [Zbl 0723.53026](#) [MR 1081650](#)
- [Bes87] Arthur L. Besse, *Einstein manifolds*. Springer-Verlag, Berlin 1987. [Zbl 0613.53001](#) [MR 0867684](#)
- [BKN89] Shigetoshi Bando, Atsushi Kasue, and Hiraku Nakajima, On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth. *Invent. Math.* **97** (2) (1989), 313–349. [Zbl 0682.53045](#) [MR 1001844](#)
- [Bor93] Joseph E. Borzellino, Orbifolds of maximal diameter. *Indiana Univ. Math. J.* **42** (1) (1993), 37–53. [Zbl 0801.53031](#) [MR 1218706](#)
- [Car98] Gilles Carron, Une suite exacte en L^2 -cohomologie. *Duke Math. J.* **95** (2) (1998), 343–372. [Zbl 0951.58024](#) [MR 1652017](#)
- [Car99] Gilles Carron, L^2 -cohomologie et inégalités de Sobolev. *Math. Ann.* **314** (4) (1999), 613–639. [Zbl 0933.35054](#) [MR 1709104](#)
- [CQY07] Alice Chang, Jie Qing, and Paul Yang, On a conformal gap and finiteness theorem for a class of four manifolds. *Geom. Funct. Anal.* **17** (2) (2007), 404–434. [Zbl 1124.53020](#) [MR 2322490](#)
- [CT06] Jeff Cheeger and Gang Tian, Curvature and injectivity radius estimates for Einstein 4-manifolds. *J. Amer. Math. Soc.* **19** (2) (2006), 487–525. [Zbl 1092.53034](#) [MR 2188134](#)
- [GPZ94] Robert E. Greene, Peter Petersen, and Shun-Hui Zhu, Riemannian manifolds of faster-than-quadratic curvature decay. *Internat. Math. Res. Notices* **1994** (9) (1994), 363–377. [Zbl 0833.53037](#) [MR 1301436](#)
- [Gro81] Michael Gromov, Curvature, diameter and Betti numbers. *Comment. Math. Helv.* **56** (2) (1981), 179–195. [Zbl 0467.53021](#) [MR 0630949](#)
- [Kas88] Atsushi Kasue, A compactification of a manifold with asymptotically nonnegative curvature. *Ann. Sci. École Norm. Sup.* (4) **21** (4) (1988), 593–622. [Zbl 0662.53032](#) [MR 0982335](#)
- [Kas89] Atsushi Kasue, A convergence theorem for Riemannian manifolds and some applications. *Nagoya Math. J.* **114** (1989), 21–51. [Zbl 0682.53042](#) [MR 1001487](#)

- [Nak92] Hiraku Nakajima, Convergence theorem of Einstein metrics and ALE spaces. *Sūgaku* **44** (2) (1992), 133–146. [Zbl 0789.53029](#) [MR 1193019](#)
- [Str07] Jeffrey Streets, Asymptotic curvature decay and removal of singularities of Bach-flat metrics. Preprint, 2007. [arXiv:0708.0869](#)
- [Tia90] G. Tian, On Calabi's conjecture for complex surfaces with positive first Chern class. *Invent. Math.* **101** (1) (1990), 101–172. [Zbl 0716.32019](#) [MR 1055713](#)
- [TV05a] Gang Tian and Jeff Viaclovsky, Bach-flat asymptotically locally Euclidean metrics. *Invent. Math.* **160** (2) (2005), 357–415. [Zbl 1085.53030](#) [MR 2138071](#)
- [TV05b] Gang Tian and Jeff Viaclovsky, Moduli spaces of critical Riemannian metrics in dimension four. *Adv. Math.* **196** (2) (2005), 346–372. [Zbl 02213018](#) [MR 2166311](#)
- [Zhu94] Shun-Hui Zhu, A volume comparison theorem for manifolds with asymptotically nonnegative curvature and its applications. *Amer. J. Math.* **116** (3) (1994), 669–682. [Zbl 0953.53027](#) [MR 1277451](#)

Received January 10, 2007

Gang Tian, 702 Fine Hall, Princeton University, Princeton, NJ 08544, U.S.A.

E-mail: tian@math.princeton.edu

Jeff Viaclovsky, Department of Mathematics, University of Wisconsin, Madison, WI 53706, U.S.A.

E-mail: jeffv@math.wisc.edu