

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 84 (2009)

Artikel: Transcendental submanifolds of projective space
Autor: Kucharz, Wojciech
DOI: <https://doi.org/10.5169/seals-99112>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 02.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Transcendental submanifolds of projective space

Wojciech Kucharz*

Abstract. Given integers m and c satisfying $m-2 \geq c \geq 2$, we explicitly construct a nonsingular m -dimensional algebraic subset of $\mathbb{P}^{m+c}(\mathbb{R})$ that is not isotopic to the set of real points of any nonsingular complex algebraic subset of $\mathbb{P}^{m+c}(\mathbb{C})$ defined over \mathbb{R} . The first examples of this type were obtained by Akbulut and King in a more complicated and nonconstructive way, and only for certain large integers m and c .

Mathematics Subject Classification (2000). 57R55, 14P25.

Keywords. Smooth manifold, algebraic set, isotopy.

1. Introduction

Denote by $\mathbb{P}^n(\mathbb{R})$ and $\mathbb{P}^n(\mathbb{C})$ real and complex projective n -spaces. We regard $\mathbb{P}^n(\mathbb{R})$ as a subset of $\mathbb{P}^n(\mathbb{C})$. A smooth (of class \mathcal{C}^∞) submanifold M of $\mathbb{P}^n(\mathbb{R})$ is said to be of *algebraic type* if it is isotopic in $\mathbb{P}^n(\mathbb{R})$ to the set of real points of a nonsingular complex algebraic subset of $\mathbb{P}^n(\mathbb{C})$ defined over \mathbb{R} ; otherwise M is said to be *transcendental*. It is not at all obvious that transcendental submanifolds exist. However, Akbulut and King [2] proved the existence of transcendental submanifolds M of $\mathbb{P}^n(\mathbb{R})$ which can even be realized as nonsingular algebraic subsets of $\mathbb{P}^n(\mathbb{R})$. Their examples are obtained in a nonconstructive way, by a method which requires both $m = \dim M$ and $n - m$ to be large integers satisfying $2m - n \geq 2$. In the present paper we explicitly construct such examples, assuming only $n - m \geq 2$ and $2m - n \geq 2$. Moreover, we verify that M is a transcendental submanifold of $\mathbb{P}^n(\mathbb{R})$ using only the Barth–Larsen theorem [6, Corollary 6.5] and completely avoiding all results of [1], [2]. More precisely, denote by S^k the unit k -sphere,

$$S^k = \{(y_1, \dots, y_{k+1}) \in \mathbb{R}^{k+1} \mid y_1^2 + \dots + y_{k+1}^2 = 1\}.$$

In Section 3 we prove the following:

*The paper was completed at the Max-Planck-Institut für Mathematik in Bonn, whose support and hospitality are gratefully acknowledged.

Theorem 1.1. *Let m and n be positive integers satisfying $n - m \geq 2$ and $2m - n \geq 2$. Let*

$$\varphi: \mathbb{P}^2(\mathbb{R}) \times S^{m-2} \longrightarrow \mathbb{P}^n(\mathbb{R})$$

be defined by

$$\begin{aligned} \varphi((x_1 : x_2 : x_3), (y_1, \dots, y_{m-1})) \\ = (x_1^2 + x_2^2 + x_3^2 : x_1 x_2 : x_1 x_3 : x_2 x_3 : \sigma y_1 : \dots : \sigma y_{m-1} : 0 : \dots : 0), \end{aligned}$$

where 0 is repeated $n - m - 2$ times and $\sigma = x_1^2 + 2x_2^2 + 3x_3^2$. Then:

- (i) *The image $M = \varphi(\mathbb{P}^2(\mathbb{R}) \times S^{m-2})$ is an m -dimensional nonsingular algebraic subset of $\mathbb{P}^n(\mathbb{R})$.*
- (ii) *$\varphi: \mathbb{P}^2(\mathbb{R}) \times S^{m-2} \rightarrow M$ is a biregular isomorphism.*
- (iii) *M is a transcendental submanifold of $\mathbb{P}^n(\mathbb{R})$.*

It follows directly from Theorem 1.1 that for any integers m and c satisfying $m - 2 \geq c \geq 2$, there is a nonsingular algebraic set M in $\mathbb{P}^{m+c}(\mathbb{R})$ such that $\dim M = m$ and M is a transcendental submanifold. In particular, there are transcendental submanifolds of arbitrary dimension $m \geq 4$. The existence of transcendental submanifolds of dimension 2 or 3 remains unsettled at this time. There are no transcendental submanifolds of dimension 1 or of codimension 1. The last assertion is a special case of the following well known fact.

Remark 1.2. Let M be a smooth m -dimensional submanifold of $\mathbb{P}^n(\mathbb{R})$. If either $n - m = 1$ or $2m + 1 \leq n$, then there exists a smooth embedding $e: M \rightarrow \mathbb{P}^n(\mathbb{R})$, arbitrarily close in the \mathcal{C}^∞ topology to the inclusion map $M \hookrightarrow \mathbb{P}^n(\mathbb{R})$, such that $e(M)$ is the set of real points of a nonsingular complex algebraic subset of $\mathbb{P}^n(\mathbb{C})$ defined over \mathbb{R} .

If $n - m = 1$, the claim is explicitly established for example in [3, Theorem 7.1].

For the second case, consider $\mathbb{P}^n(\mathbb{R})$ as a subset of $\mathbb{P}^k(\mathbb{R})$, where k is a large integer. By [8], there exists a smooth embedding $j: M \rightarrow \mathbb{P}^k(\mathbb{R})$, arbitrarily close in the \mathcal{C}^∞ topology to the inclusion map $M \hookrightarrow \mathbb{P}^k(\mathbb{R})$, such that $j(M)$ is a nonsingular algebraic subset of $\mathbb{P}^k(\mathbb{R})$. Increasing k if necessary and making use of Hironaka's resolution of singularities theorem [7], we may assume that the Zariski complex closure of $j(M)$ in $\mathbb{P}^k(\mathbb{C})$ is nonsingular. If $2m + 1 \leq n$, we obtain an embedding $e: M \rightarrow \mathbb{P}^n(\mathbb{R})$ with the required properties by composing j with an appropriate generic projection onto $\mathbb{P}^n(\mathbb{R})$.

2. A criterion for transcendence

First we need some results related to the Picard group. Following the current custom, we state them in the language of schemes.

Let V be a smooth projective scheme over \mathbb{R} . Assume that the set $V(\mathbb{R})$ of \mathbb{R} -rational points of V is nonempty. We regard $V(\mathbb{R})$ as a compact smooth manifold. Every invertible sheaf \mathcal{L} on V determines a real line bundle on $V(\mathbb{R})$, denoted $\mathcal{L}(\mathbb{R})$. The correspondence which assigns to each invertible sheaf \mathcal{L} on V the first Stiefel–Whitney class $w_1(\mathcal{L}(\mathbb{R}))$ of $\mathcal{L}(\mathbb{R})$ gives rise to a canonical homomorphism

$$w_1: \mathrm{Pic}(V) \longrightarrow H^1(V(\mathbb{R}), \mathbb{Z}/2),$$

defined on the Picard group $\mathrm{Pic}(V)$ of isomorphism classes of invertible sheaves on V . We set

$$H_{\mathrm{alg}}^1(V(\mathbb{R}), \mathbb{Z}/2) = w_1(\mathrm{Pic}(V)).$$

It will be convenient to recall another description of $\mathrm{Pic}(V)$. Consider the scheme $V_{\mathbb{C}} = V \times_{\mathbb{R}} \mathbb{C}$ over \mathbb{C} and its Picard group $\mathrm{Pic}(V_{\mathbb{C}})$. The Galois group $G = \mathrm{Gal}(\mathbb{C}/\mathbb{R})$ of \mathbb{C} over \mathbb{R} acts on $\mathrm{Pic}(V_{\mathbb{C}})$. We denote by $\mathrm{Pic}(V_{\mathbb{C}})^G$ the subgroup of $\mathrm{Pic}(V_{\mathbb{C}})$ consisting of the elements fixed by G . Given an invertible sheaf \mathcal{L} on V , we write $\mathcal{L}_{\mathbb{C}}$ for the corresponding sheaf on $V_{\mathbb{C}}$. The correspondence $\mathcal{L} \rightarrow \mathcal{L}_{\mathbb{C}}$ defines a canonical group homomorphism

$$\alpha: \mathrm{Pic}(V) \longrightarrow \mathrm{Pic}(V_{\mathbb{C}})^G.$$

It follows from the general theory of descent [4] that α is an isomorphism (a simple treatment of the case under consideration can also be found in [5]).

As usual, we set $\mathbb{P}_{\mathbb{R}}^n = \mathrm{Proj}(\mathbb{R}[T_0, \dots, T_n])$ and identify $\mathbb{P}_{\mathbb{R}}^n(\mathbb{R})$ with $\mathbb{P}^n(\mathbb{R})$. Thus if V is a subscheme of $\mathbb{P}_{\mathbb{R}}^n$, then $V(\mathbb{R})$ is a subset of $\mathbb{P}^n(\mathbb{R})$.

Proposition 2.1. *Let V be a closed smooth m -dimensional subscheme of $\mathbb{P}_{\mathbb{R}}^n$. If $2m - n \geq 2$, then*

$$H_{\mathrm{alg}}^1(V(\mathbb{R}), \mathbb{Z}/2) = i^*(H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2)),$$

where $i: V(\mathbb{R}) \hookrightarrow \mathbb{P}^n(\mathbb{R})$ is the inclusion map.

Proof. Let $j: V \hookrightarrow \mathbb{P}_{\mathbb{R}}^n$ and $j_{\mathbb{C}}: V_{\mathbb{C}} \hookrightarrow \mathbb{P}_{\mathbb{C}}^n = \mathbb{P}_{\mathbb{R}}^n \times_{\mathbb{R}} \mathbb{C}$ be the inclusion morphisms. By the Barth–Larsen theorem [6, Corollary 6.5], the induced homomorphism

$$j_{\mathbb{C}}^*: \mathrm{Pic}(\mathbb{P}_{\mathbb{C}}^n) \longrightarrow \mathrm{Pic}(V_{\mathbb{C}})$$

is an isomorphism. Since $j_{\mathbb{C}}^*$ is G -equivariant, the restriction

$$j_{\mathbb{C}}^*: \mathrm{Pic}(\mathbb{P}_{\mathbb{C}}^n)^G \longrightarrow \mathrm{Pic}(V_{\mathbb{C}})^G$$

is an isomorphism. We have the following commutative diagram:

$$\begin{array}{ccc}
 \mathrm{Pic}(\mathbb{P}_{\mathbb{C}}^n)^G & \xrightarrow{j_{\mathbb{C}}^*} & \mathrm{Pic}(V_{\mathbb{C}})^G \\
 \alpha \uparrow & & \uparrow \alpha \\
 \mathrm{Pic}(\mathbb{P}_{\mathbb{R}}^n) & \xrightarrow{j^*} & \mathrm{Pic}(V) \\
 w_1 \downarrow & & \downarrow w_1 \\
 H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2) & \xrightarrow{i^*} & H^1(V(\mathbb{R}), \mathbb{Z}/2).
 \end{array}$$

Since the homomorphisms α are isomorphisms and

$$H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2) = H_{\mathrm{alg}}^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2),$$

it follows that

$$H_{\mathrm{alg}}^1(V(\mathbb{R}), \mathbb{Z}/2) = i^*(H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2)),$$

as required. \square

Note that a smooth submanifold of $\mathbb{P}^n(\mathbb{R})$ is of algebraic type if and only if it is isotopic in $\mathbb{P}^n(\mathbb{R})$ to $V(\mathbb{R})$ for some closed smooth subscheme V of $\mathbb{P}_{\mathbb{R}}^n$. Hence Proposition 2.1 yields the following criterion for transcendence.

Proposition 2.2. *Let M be a compact smooth m -dimensional submanifold of $\mathbb{P}^n(\mathbb{R})$. Assume that the inclusion map $e: M \hookrightarrow \mathbb{P}^n(\mathbb{R})$ induces a trivial homomorphism*

$$e^*: H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2) \longrightarrow H^1(M, \mathbb{Z}/2),$$

that is, $e^ = 0$. If M is nonorientable and $2m - n \geq 2$, then M is a transcendental submanifold of $\mathbb{P}^n(\mathbb{R})$.*

Proof. Suppose to the contrary that M is of algebraic type. Let V be a closed smooth subscheme of $\mathbb{P}_{\mathbb{R}}^n$ with $V(\mathbb{R})$ isotopic to M in $\mathbb{P}^n(\mathbb{R})$. Then the homomorphism

$$i^*: H^1(\mathbb{P}^n(\mathbb{R}), \mathbb{Z}/2) \longrightarrow H^1(V(\mathbb{R}), \mathbb{Z}/2),$$

induced by the inclusion map $i: V(\mathbb{R}) \hookrightarrow \mathbb{P}^n(\mathbb{R})$, is trivial. Since $\dim V = m$ and $2m - n \geq 2$, Proposition 2.1 implies

$$H_{\mathrm{alg}}^1(V(\mathbb{R}), \mathbb{Z}/2) = 0.$$

On the other hand, the first Stiefel–Whitney class $w_1(V(\mathbb{R}))$ of $V(\mathbb{R})$ is nonzero, $V(\mathbb{R})$ being a nonorientable manifold. Moreover, $w_1(V(\mathbb{R})) = w_1(\mathcal{K}(\mathbb{R}))$, where \mathcal{K} is the canonical invertible sheaf of V , and hence, $w_1(V(\mathbb{R}))$ is in $H_{\mathrm{alg}}^1(V(\mathbb{R}), \mathbb{Z}/2)$. In view of this contradiction, the proof is complete. \square

3. Transcendental submanifolds

We begin with some preliminary observations. Identify \mathbb{R}^n with its image under the map

$$\mathbb{R}^n \longrightarrow \mathbb{P}^n(\mathbb{R}), \quad (x_1, \dots, x_n) \longmapsto (1 : x_1 : \dots : x_n);$$

thus $\mathbb{R}^n \subset \mathbb{P}^n(\mathbb{R})$. An algebraic subset X of \mathbb{R}^n is said to be *projectively closed* if X is also an algebraic subset of $\mathbb{P}^n(\mathbb{R})$. One readily checks that X is projectively closed if and only if it can be defined by a real polynomial equation

$$f(x_1, \dots, x_n) = 0,$$

where the homogeneous form of top degree in f vanishes only at 0 in \mathbb{R}^n .

Lemma 3.1. *Let X be an algebraic subset of \mathbb{R}^k contained in the open half-space*

$$H = \{(x_1, \dots, x_k) \in \mathbb{R}^k \mid x_k > 0\}.$$

Then the map $\psi: X \times S^\ell \rightarrow \mathbb{R}^{k+\ell}$ defined by

$$\psi((x_1, \dots, x_k), (y_1, \dots, y_{\ell+1})) = (x_1, \dots, x_{k-1}, x_k y_1, \dots, x_k y_{\ell+1})$$

is an algebraic embedding, that is, the image $Y = \psi(X \times S^\ell)$ is an algebraic subset of $\mathbb{R}^{k+\ell}$ and $\psi: X \times S^\ell \rightarrow Y$ is a biregular isomorphism. Moreover, if X is projectively closed in \mathbb{R}^k , then Y is projectively closed in $\mathbb{R}^{k+\ell}$.

Proof. Let

$$f(u, v) = 0$$

be a real polynomial equation defining X , where $u = (x_1, \dots, x_{k-1})$ and $v = x_k$. Since

$$X \subset H, \tag{1}$$

the subset Y of $\mathbb{R}^{k+\ell}$ is defined by the equation

$$f(u, \rho) = 0, \tag{2}$$

where

$$\rho = (x_k^2 + x_{k+1}^2 + \dots + x_{k+\ell}^2)^{\frac{1}{2}}.$$

We will now show that (2) can be replaced by a polynomial equation in x_1, \dots, x_{k-1} , $x_k, \dots, x_{k+\ell}$. To this end we write

$$f(u, v) = g(u, v^2) + vh(u, v^2), \tag{3}$$

where g and h are real polynomials in (u, v) . Then (2) is equivalent to

$$g(u, \rho^2) + \rho h(u, \rho^2) = 0, \quad (4)$$

and in view of (1) also to

$$(g(u, \rho^2))^2 - \rho^2(h(u, \rho^2))^2 = 0, \quad (5)$$

which is a polynomial equation, as required. Consequently, Y is an algebraic subset of $\mathbb{R}^{k+\ell}$.

It is clear that ψ is injective and $\theta: Y \rightarrow X$,

$$\theta(x_1, \dots, x_{k-1}, x_k, \dots, x_{k+\ell}) = \left(x_1, \dots, x_{k-1}, \frac{x_k}{\rho}, \dots, \frac{x_{k+\ell}}{\rho} \right),$$

is the inverse of $\psi: X \rightarrow Y$. By (4),

$$\rho = -\frac{g(x_1, \dots, x_{k-1}, x_k^2 + \dots + x_{k+\ell}^2)}{h(x_1, \dots, x_{k-1}, x_k^2 + \dots + x_{k+\ell}^2)}$$

for $(x_1, \dots, x_{k-1}, x_k, \dots, x_{k+\ell})$ in Y , and hence θ is a regular map. Thus $\psi: X \rightarrow Y$ is a biregular isomorphism.

Assume now that X is projectively closed in \mathbb{R}^k . We may also assume that the homogeneous form of top degree in f , denoted F , vanishes only at 0 in \mathbb{R}^k . Note that $F(u, \rho^2)F(u, -\rho^2)$ is the homogeneous form of top degree in equation (5). This form vanishes only at 0 in $\mathbb{R}^{k+\ell}$, and hence Y is projectively closed in $\mathbb{R}^{k+\ell}$. \square

Lemma 3.2. *The map $g: \mathbb{P}^2(\mathbb{C}) \rightarrow \mathbb{P}^4(\mathbb{C})$,*

$$g((x_1 : x_2 : x_3)) = (x_1^2 + x_2^2 + x_3^2 : x_1 x_2 : x_1 x_3 : x_2 x_3 : x_1^2 + 2x_2^2 + 3x_3^2),$$

is an algebraic embedding. In particular, the restriction $f: \mathbb{P}^2(\mathbb{R}) \rightarrow \mathbb{P}^4(\mathbb{R})$ of g is an algebraic embedding.

Proof. One readily checks that g is injective. Moreover, the (complex) differential of g at each point of $\mathbb{P}^2(\mathbb{C})$ is of rank 2. It follows that g is an algebraic embedding, and hence f is an algebraic embedding. \square

Proof of Theorem 1.1. Let $f: \mathbb{P}^2(\mathbb{R}) \rightarrow \mathbb{P}^4(\mathbb{R})$ be the algebraic embedding of Lemma 3.2. Note that the image $X = f(\mathbb{P}^2(\mathbb{R}))$ is a projectively closed algebraic subset of $\mathbb{R}^4 \subset \mathbb{P}^4(\mathbb{R})$, contained in the open half-space

$$\{(u_1, u_2, u_3, u_4) \in \mathbb{R}^4 \mid u_4 > 0\}.$$

Let

$$\psi: X \times S^{m-2} \longrightarrow \mathbb{R}^{4+(m-2)} = \mathbb{R}^{m+2} \subset \mathbb{P}^{m+2}(\mathbb{R})$$

be the algebraic embedding of Lemma 3.1 (with $k = 4$ and $\ell = m - 2$). Note that $\psi(X \times S^{m-2})$ is projectively closed in \mathbb{R}^{m+2} , and hence is an algebraic subset of $\mathbb{P}^{m+2}(\mathbb{R})$.

Clearly, if $i : S^{m-2} \rightarrow S^{m-2}$ is the identity map, then

$$f \times i : \mathbb{P}^2(\mathbb{R}) \times S^{m-2} \longrightarrow X \times S^{m-2}$$

is a biregular isomorphism. Denoting by $j : \mathbb{P}^{m+2}(\mathbb{R}) \rightarrow \mathbb{P}^n(\mathbb{R})$ the standard embedding,

$$j((v_0 : \dots : v_{m+2})) = (v_0 : \dots : v_{m+2} : 0 : \dots : 0),$$

we obtain

$$\varphi = j \circ \psi \circ (f \times i),$$

which implies that φ is an algebraic embedding. In other words, conditions (i) and (ii) are satisfied. Moreover, $M \subset \mathbb{R}^n \subseteq \mathbb{P}^n(\mathbb{R})$. Since M is nonorientable and $2m - n \geq 2$, condition (iii) follows from Proposition 2.2. \square

References

- [1] S. Akbulut and H. King, Transcendental submanifolds of \mathbb{R}^n . *Comment. Math. Helv.* **68** (1993), 308–318. [Zbl 0806.57017](#) [MR 1214234](#)
- [2] S. Akbulut and H. King, Transcendental submanifolds of $\mathbb{R}\mathbb{P}^n$. *Comment. Math. Helv.* **80** (2005), 427–432. [Zbl 1071.57026](#) [MR 2142249](#)
- [3] J. Bochnak, M. Buchner and W. Kucharz, Vector bundles over real algebraic varieties. *K-Theory* **3** (1989), 271–298; Erratum, *K-Theory* **4** (1990), 103. [Zbl 0761.14020](#) [MR 1076527](#)
- [4] A. Grothendieck, *Technique de descente et théorèmes d'existence en géométrie algébrique*, I–VI. Séminaire Bourbaki, Exposés 190, 195, 212, 221, 232, 236, 1959–62. [Zbl 0234.14007](#) [MR 1603475](#)
- [5] J. van Hamel, Algebraic cycles and topology of real algebraic varieties. Dissertation, Vrije Universiteit Amsterdam. CWI Tract. 129, Stichting Mathematisch Centrum, Centrum voor Wiskunde en informatica, Amsterdam 2000. [Zbl 0986.14042](#) [MR 1760347](#)
- [6] R. Hartshorne, Equivalence relations on algebraic cycles and subvarieties of small codimension. *Proc. Sympos. Pure Math.* **29** (1975), 129–164. [Zbl 0314.14001](#) [MR 0369359](#)
- [7] H. Hironaka, Resolution of singularities of an algebraic variety over a field of characteristic zero. *Ann. of Math.* **79** (1964), 109–203. [Zbl 0122.38603](#) [MR 0199184](#)
- [8] H. King, Approximating submanifolds of real projective space by varieties. *Topology* **15** (1976), 81–85. [Zbl 0316.57015](#) [MR 0396572](#)

Received February 28, 2007

Wojciech Kucharz, Department of Mathematics and Statistics, University of New Mexico, Albuquerque, New Mexico 87131-0001, U.S.A.

E-mail: kucharz@math.unm.edu