

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 84 (2009)

Artikel: Diameter pinching in almost positive Ricci curvature
Autor: Aubry, Erwann
DOI: <https://doi.org/10.5169/seals-99118>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Diameter pinching in almost positive Ricci curvature

Erwann Aubry

Abstract. In this paper we prove a diameter sphere theorem and its corresponding λ_1 sphere theorem under L^p control of the curvature. They are generalizations of some results due to S. Ilias [8].

Mathematics Subject Classification (2000). 53C20.

Keywords. Ricci curvature, comparison theorems, integral bounds on the curvature, sphere theorems.

1. Introduction

Let (M^n, g) be a complete manifold with Ricci curvature $\text{Ric} \geq n - 1$. Then (M^n, g) satisfies the following classical results (the proofs can be found in [13] for instance):

- $\text{Diam}(M^n, g) \leq \pi$ (S. Myers) with equality iff $(M^n, g) = (\mathbb{S}^n, \text{can})$ (S. Cheng),
- $\lambda_1(M^n, g) \geq n$ (A. Lichnerowicz) with equality iff $(M^n, g) = (\mathbb{S}^n, \text{can})$ (M. Obata),

where Diam is the diameter and λ_1 is the first positive eigenvalue.

Studying the properties of the sphere kept by manifolds with $\text{Ric} \geq n - 1$ and almost extremal diameter or λ_1 , S. Ilias proved in [8] the following results:

Theorem 1.1 (S. Ilias). *For any $A > 0$, there exists $\epsilon(A, n) > 0$ such that any n -manifold with $\text{Ric} \geq n - 1$, sectional curvature $\sigma \leq A$ and $\lambda_1 \leq n + \epsilon$ is homeomorphic to \mathbb{S}^n .*

Theorem 1.2 (S. Ilias). *For any $A > 0$, there exists $\epsilon(A, n) > 0$ such that any n -manifold with $\text{Ric} \geq n - 1$, $\sigma \leq A$ and $\text{Diam}(M) \geq \pi - \epsilon$ is homeomorphic to \mathbb{S}^n .*

Remark 1.3. C. Croke proves in [7] that for n -manifolds with $\text{Ric} \geq n - 1$, $\lambda_1(M)$ close to n implies $\text{Diam}(M)$ close to π . The converse is proved in [8] (using a spectral inequality due to S. Cheng [6]).

Remark 1.4. For $n \geq 4$, M. Anderson [1] and Y. Otsu [10] construct sequences of complete metrics g_i with $\text{Ric}(g_i) \geq n - 1$, $\lambda_1(g_i) \rightarrow n$ and $\text{Diam}(g_i) \rightarrow \pi$ on manifolds that are not homotopic to \mathbb{S}^n (more precisely, Otsu shows that if $n \geq 5$, these manifolds can have infinitely many different fundamental groups).

Remark 1.5. The two results of S. Ilias have been improved by G. Perelman in [11], where the assumption $\sigma \leq A$ is replaced by $\sigma \geq -A$ (note that under the Ilias's assumptions $\sigma \leq A$ and $\text{Ric} \geq n - 1$ we have $|\sigma| \leq (n - 2)A$).

Subsequently, we denote $\underline{\text{Ric}}(x)$ the lowest eigenvalue of the Ricci tensor and $\bar{\sigma}(x)$ the maximal sectional curvature at x . In [4], we prove the following generalization of the theorems of Myers and Lichnerowicz:

Theorem 1.6. For any $p > n/2$, there exists $C(p, n)$ such that if (M^n, g) is a complete manifold with $\int_M (\underline{\text{Ric}} - (n - 1))_-^p < \frac{\text{Vol } M}{C(p, n)}$, then M is compact, has finite fundamental group and satisfies

$$\begin{aligned}\text{Diam}(M) &\leq \pi \left[1 + C(p, n) \left(\frac{\rho_p}{\text{Vol } M} \right)^{\frac{1}{10}} \right], \\ \lambda_1(M) &\geq n \left[1 - C(p, n) \left(\frac{\rho_p}{\text{Vol } M} \right)^{\frac{1}{p}} \right],\end{aligned}$$

where $\rho_p = \int_M (\underline{\text{Ric}} - (n - 1))_-^p$ and $x_- = \max(0, -x)$.

Remark 1.7. It follows from [4] that the constant $C(p, n)$ is computable, that if $\int_M (\underline{\text{Ric}} - (n - 1))_-^p$ is finite (for $p > n/2$), then $\text{Vol } M$ is finite, and that we cannot bound the diameter or the first non zero eigenvalue under the assumption $\rho_p \leq \frac{1}{C(p, n)}$ or $\rho_{\frac{n}{2}}$ small (see [4]).

In this paper we prove the following extensions of Ilias's stability results.

Theorem 1.8. Let $n \geq 2$ be an integer, $A > 0$ and $p > n$ be some reals. There exists a positive constant $C(p, n, A)$ such that any complete n -manifold which satisfies

$$\int_M (\underline{\text{Ric}} - (n - 1))_-^p < C(p, n, A) \text{Vol } M, \quad \int_M \bar{\sigma}_+^p < A \text{Vol } M$$

and

$$\text{Diam}(M) \geq \pi (1 - C(p, n, A))$$

is homeomorphic to \mathbb{S}^n (where $x_+ = \max(0, x)$).

Theorem 1.9. Let $n \geq 2$ be an integer, $A > 0$ and $p > n$ be some reals. There exists a positive constant $C(p, n, A)$ such that any complete n -manifold which satisfies

$$\int_M (\underline{\text{Ric}} - (n - 1))_-^p < C(p, n, A) \text{Vol } M, \quad \int_M \bar{\sigma}_+^p < A \text{Vol } M$$

and

$$\lambda_1(M) \leq n(1 + C(p, n, A))$$

is homeomorphic to \mathbb{S}^n .

Remark 1.10. By the Hölder inequality, the two curvature assumptions of Theorem 1.9 can be replaced by

$$\int_M (\underline{\text{Ric}} - (n-1))_- < C(p, n, A) \text{Vol } M \quad \text{and} \quad \int_M \sigma^p < A \text{Vol } M,$$

where $\sigma(x)$ is an upper bound for the absolute value of the sectional curvatures at x .

2. Comparison results in almost positive Ricci curvature

Subsequently we denote $B(x, r)$ (resp. $S(x, r)$) the geodesic ball (resp. sphere) with center x and radius r and $L_k(r)$ (resp. $A_k(r)$) the volume of a geodesic sphere (resp. ball) of radius r in $(\mathbb{S}^n, \frac{1}{k}g)$. Besides Theorem 1.6, we will need the following comparison results for manifolds of almost positive Ricci curvature (see [4] for a proof).

Proposition 2.1. *For any $n \geq 2$ and $p > n/2$ ($p \geq 1$ if $n = 2$) there exists a constant $C(p, n)$ such that for any complete Riemannian n -manifold (M^n, g) with $\eta^{10} = \frac{\rho_p}{\text{Vol } M} \leq \frac{1}{C(p, n)}$, we have*

$$\begin{aligned} \left(\frac{\text{Vol}_{n-1} S(x, R)}{L_{1-\eta}(R)} \right)^{\frac{1}{2p-1}} - \left(\frac{\text{Vol}_{n-1} S(x, r)}{L_{1-\eta}(r)} \right)^{\frac{1}{2p-1}} &\leq C(p, n)\eta^2(R-r)^{\frac{2p-n}{2p-1}}, \\ \frac{\text{Vol } B(x, r)}{\text{Vol } B(x, R)} &\geq (1 - C(p, n)\eta) \frac{A_1(r)}{A_1(R)}, \\ \text{Vol}_{n-1} S(x, R) &\leq (1 + \eta^2)L_{1-\eta}(R), \\ \text{Vol } B(x, R) &\leq (1 + \eta)A_1(R) \end{aligned}$$

for all $x \in M$ and all radii $0 \leq r \leq R$.

For any $n \geq 2$ and $p > n/2$ there exists a constant $C(p, n)$ such that if (M^n, g) is a complete n -manifold with $\bar{\rho}_p \leq \frac{1}{C(p, n)}$, then $\|u\|_{\frac{2n}{n-2}} \leq \text{Diam}(M)C(p, n)\|du\|_2 + \|u\|_2$, for any $u \in H^{1,2}(M)$. In the case $n = 2$, we have $\|u\|_4 \leq \text{Diam}(M)C\|du\|_2 + \|u\|_2$ if $\bar{\rho}_1 \leq \frac{1}{C}$.

Similar estimates are proved in [12] under the assumption that M is compact and $\text{Diam}(M)^{2p} \frac{\rho_p}{\text{Vol } M} \leq \frac{1}{C(p, n)}$.

3. Theorem 1.9 implies Theorem 1.8

Proposition 3.1. *Let $n \geq 2$ and $p > n/2$. There exists $C(p, n) > 0$ such that if (M^n, g) is a complete n -manifold with $\eta^{10} = \bar{\rho}_p \leq \frac{1}{C(p, n)}$ and $\text{Diam}(M) \geq \pi - \frac{1}{C(p, n)}$, then we have*

$$\lambda_1(M) \leq n + C(p, n)[\eta + (\text{Diam}(M) - \pi)_-].$$

The main tool to prove this proposition is the following lemma:

Lemma 3.2. *Let $n \geq 2$ and $p > n/2$ ($p \geq 1$ if $n = 2$) and $\bar{x}_0 \in \mathbb{S}^n$. There exists a constant $C(p, n)$ such that if (M^n, g) is a complete n -manifold with $\eta^{10} = \bar{\rho}_p \leq \frac{1}{C(p, n)}$, then there exists $x_0 \in M$ such that for any C^1 -function $u: [0, 2\pi] \rightarrow \mathbb{R}$ we have*

$$\left| \frac{1}{\text{Vol } M} \int_M u \circ d_M(x_0, \cdot) dv_g - \frac{1}{\text{Vol } \mathbb{S}^n} \int_{\mathbb{S}^n} u \circ d_{\mathbb{S}^n}(\bar{x}_0, \cdot) dv_{\mathbb{S}^n} \right| \\ \leq \|u'\|_\infty C(p, n)[\eta + (\text{Diam}(M) - \pi)_-].$$

Proof. Let $(x_0, y_0) \in M^2$ such that $d = \text{Diam}(M) = d(x_0, y_0)$. The functions A , L , A_1 and L_1 are defined in Proposition 2.1 and prolonged by 0 to \mathbb{R} (note that the diameter of M can be greater than π). The function $r \rightarrow u(r)A(r)$ is continuous and has right differential on \mathbb{R} equal to $u'A + uL$. We infer the equalities

$$u(d) \text{Vol } M = \int_0^d u(r)L(r) dr + \int_0^d u'(r)A(r) dr, \\ u(\pi) \text{Vol } \mathbb{S}^n = \int_0^\pi u(r)L_1(r) dr + \int_0^\pi u'(r)A_1(r) dr$$

which imply

$$\left| \frac{1}{\text{Vol } M} \int_M u \circ d_M(x_0, x) dv_g - \frac{1}{\text{Vol } \mathbb{S}^n} \int_{\mathbb{S}^n} u \circ d_{\mathbb{S}^n}(\bar{x}_0, x) dv_{\mathbb{S}^n} \right| \\ = \left| \int_0^d \frac{u(r)L(r)}{\text{Vol } M} dr - \int_0^\pi \frac{u(r)L_1(r)}{\text{Vol } \mathbb{S}^n} dr \right| \\ = \left| u(d) - u(\pi) + \int_0^\pi \frac{u' A_1}{\text{Vol } \mathbb{S}^n} - \int_0^d \frac{u' A}{\text{Vol } M} \right| \\ = \left| \int_0^d u' \left(\frac{A_1}{\text{Vol } \mathbb{S}^n} - \frac{A}{\text{Vol } M} \right) + \int_d^\pi u' \left(\frac{A_1}{\text{Vol } \mathbb{S}^n} - 1 \right) \right| \\ \leq \|u'\|_\infty \left(\int_0^d \left| \frac{A_1}{\text{Vol } \mathbb{S}^n} - \frac{A}{\text{Vol } M} \right| dr + |\pi - d| \right).$$

By Proposition 2.1 we have, for all $r \leq d$:

$$\begin{aligned} (1 - C(p, n)\eta) \frac{A_1(r)}{\text{Vol } \mathbb{S}^n} &\leq \frac{A(r)}{\text{Vol } M} \leq 1 - \frac{\text{Vol } B(y_0, d - r)}{\text{Vol } M} \\ &\leq 1 - (1 - C(p, n)\eta) \frac{A_1(d - r)}{\text{Vol } \mathbb{S}^n} \\ &\leq \frac{A_1(r + \pi - d)}{\text{Vol } \mathbb{S}^n} + C(p, n)\eta. \end{aligned}$$

Hence $\left| \frac{A(r)}{\text{Vol } M} - \frac{A_1(r)}{\text{Vol } \mathbb{S}^n} \right| \leq C(p, n)\eta + \frac{(A_1(r) - A_1(r + \pi - d))_-}{\text{Vol } \mathbb{S}^n}$. An easy computation gives $\left\| \frac{(A_1(\cdot) - A_1(\cdot + h))_-}{\text{Vol } \mathbb{S}^n} \right\|_\infty \leq C(n)(-h)_-$, and by Proposition 2.1 we get:

$$\begin{aligned} &\left| \frac{1}{\text{Vol } M} \int_M u \circ d_M(x_0, x) dv_g - \frac{1}{\text{Vol } \mathbb{S}^n} \int_{\mathbb{S}^n} u \circ d_{\mathbb{S}^n}(\bar{x}_0, x) dv_{\mathbb{S}^n} \right| \\ &\leq \|u'\|_\infty C(p, n)[\eta + (d - \pi)_-]. \end{aligned} \quad \square$$

We now finish the proof of Proposition 3.1.

Proof. Lemma 3.2 applied to $u = \sin^2$, $u = \cos^2$ and $u = \cos$ gives

$$\begin{aligned} \left| \int_M \frac{\sin^2 d_M(x_0, \cdot)}{\text{Vol } M} - \int_{\mathbb{S}^n} \frac{\sin^2 d_{\mathbb{S}^n}(\bar{x}_0, \cdot)}{\text{Vol } \mathbb{S}^n} \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1, \\ \left| \int_M \frac{\cos^2 d_M(x_0, \cdot)}{\text{Vol } M} - \int_{\mathbb{S}^n} \frac{\cos^2 d_{\mathbb{S}^n}(\bar{x}_0, \cdot)}{\text{Vol } \mathbb{S}^n} \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1, \\ \left| \int_M \frac{\cos d_M(x_0, \cdot)}{\text{Vol } M} - \int_{\mathbb{S}^n} \frac{\cos d_{\mathbb{S}^n}(\bar{x}_0, \cdot)}{\text{Vol } \mathbb{S}^n} \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1. \end{aligned}$$

Hence, if we set $f = \cos d_M(x_0, \cdot)$, we get

$$\begin{aligned} \left| \|\nabla f\|_2^2 - \frac{n}{n+1} \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1, \\ \left| \|f\|_2^2 - \frac{1}{n+1} \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1, \\ \left| \frac{1}{\text{Vol } M} \int_M f \right| &\leq C(p, n)(\eta + (d - \pi)_-) \leq 1, \end{aligned}$$

which readily implies that

$$\lambda_1(M) \leq \frac{\|\nabla(f - \bar{f})\|_2^2}{\|f - \bar{f}\|_2} \leq n(1 + C(p, n)(\eta + (d - \pi)_-)),$$

where we have set $\bar{f} = \frac{1}{\text{Vol } M} \int_M f$.

\square

Remark 3.3. The same technique as in [5] can be used to prove that manifolds with almost positive Ricci curvature and λ_1 is close to n have a diameter close to π (see [12]).

4. Proof of Theorem 1.9

4.1. Fiber bundle E . Let E be the fiber bundle $TM \oplus \mathbb{R} e \rightarrow M$ endowed with the following scalar product and linear connection:

$$\begin{aligned}\langle X + fe, Y + he \rangle_E &= g(X, Y) + fh, \\ D_Z^E(X + fe) &= D_Z^M X + fZ + (df(Z) - g(Z, X)) \cdot e,\end{aligned}$$

where D^M is the Levi-Civita connection of the metric g on M . We denote by p the orthogonal projection of E on TM , $\text{Ric}'(S) = \text{Ric}_M(p(S)) - (n-1)p(S)$ and $\Delta_{\text{sph}} = \bar{\Delta}^E + \text{Ric}'$.

The following lemma is proved in [3]:

Lemma 4.1. If $f: M \rightarrow \mathbb{R}$ satisfies $\Delta f = \lambda f$, then $S_f = \nabla f + fe$ satisfies $\Delta_{\text{sph}}(S_f) = (\lambda - n)(\nabla f - fe)$ and $\langle D_X^E S_f, X \rangle = Ddf(X, X) + fg(X, X)$.

Note also that we have

$$R_{(Z,Y)}^E(X + fe) = R^M(Z, Y)X - (g(Y, X)Z - g(Z, X)Y).$$

4.2. Bound on the Hessian of the first eigenfunction. To prove Theorem 1.9 we need an L^∞ bound on the Hessian of the first eigenfunction. For that purpose, we will modify the proof of Theorem 2.4 in [2] (whose proof would give us only a bound on $\|DS_f\|_{n+\epsilon}/\|S_f\|_\infty$ for a given $\epsilon = \epsilon(p, n)$). In our case we really need to perform a Moser iteration.

Proposition 4.2. Let $n \geq 2$ and $\infty \geq p > n/2$. There exists a constant $C(p, n)$ such that if (M^n, g) is any manifold with $\bar{\rho}_p \leq \frac{1}{C(p,n)}$ and $\lambda_1 \leq n + \frac{1}{C(p,n)}$, then for $f: M \rightarrow \mathbb{R}$ such that $\Delta f = \lambda_1 f$ we have

$$\frac{\|D^E S_f\|_\infty}{\|S_f\|_\infty} \leq C(p, n)(\lambda_1 + \|R\|_{2p})^\gamma(|\lambda_1 - n| + \bar{\rho}_p)^{\frac{1}{2(1+\gamma)}},$$

where $S_f = \nabla f + f \cdot e$ and $\gamma = \frac{pn}{2p-n}$.

To prove Proposition 4.2 we need a commutation lemma (see [2]):

Lemma 4.3. *For any section $S \in \Gamma(E)$ we have*

$$\begin{aligned} & \frac{1}{2}\Delta(|DS|^2) + |D^2S|^2 \\ & \leq \langle D^*R^E S, DS \rangle + \underline{\text{Ric}}^-|DS|^2 + \langle D\bar{\Delta}S, DS \rangle + \|R^E\| \cdot |DS|^2, \end{aligned}$$

where $\|R^E\|$ is the norm of the linear map $R^E: \bigwedge^2 T_m M \rightarrow \bigwedge^2 E_m^*$ defined by $R^E(u \wedge v)(T, S) = \langle R^E(u, v)T, S \rangle$.

Remark 4.4. This lemma is valid for any Riemannian fiber bundle $(E, D, \langle \cdot, \cdot \rangle)$.

We now give the proof of Proposition 4.2.

Proof of Proposition 4.2. We set $u = \sqrt{|DS|^2 + \epsilon^2}$. We have

$$\begin{aligned} u\Delta u &= \frac{1}{2}\Delta(u^2) + |du|^2 = \frac{1}{2}\Delta(u^2) + \frac{|\langle D^2S, DS \rangle|^2}{|DS|^2 + \epsilon^2} \\ &\leq \frac{1}{2}\Delta(|DS|^2) + |D^2S|^2. \end{aligned}$$

Hence, by Lemma 4.3

$$\begin{aligned} \int_M |d(u^k)|^2 &\leq \frac{k^2}{2k-1} \int_M \left(\frac{1}{2}\Delta|DS|^2 + |D^2S|^2 \right) u^{2(k-1)} \\ &\leq \frac{k^2}{2k-1} \left(\int_M \underline{\text{Ric}}^- u^{2k} + \int_M \langle D\bar{\Delta}S, DS \rangle u^{2(k-1)} \right. \\ &\quad \left. + \int_M \langle D^*R^E S, DS \rangle u^{2(k-1)} + \int_M \|R^E\| u^{2k} \right). \end{aligned}$$

We now apply the divergence theorem to the form $u^{2(k-1)} \langle \bar{\Delta}S, D_{\bullet}S \rangle$, and get for any $k \geq 1$:

$$\begin{aligned} & \int_M \langle D\bar{\Delta}S, DS \rangle u^{2(k-1)} \\ &= \int_M |\bar{\Delta}S|^2 u^{2(k-1)} - 2(k-1) \sum_i \int_M \langle \bar{\Delta}S, DS(i) \rangle du(i) \cdot u^{2k-3} \\ &\leq \int_M |\bar{\Delta}S|^2 u^{2(k-1)} + 2(k-1) \int_M |\bar{\Delta}S| |du| u^{2(k-1)} \\ &\leq \frac{k-1}{2} \int_M |du|^2 u^{2(k-1)} + (2k-1) \int_M |\bar{\Delta}S|^2 u^{2(k-1)}. \end{aligned}$$

We do the same with the form $u^{2(k-1)}(tr_{1,3}(\langle R_{(\bullet,\bullet)}^E S, D_\bullet S \rangle))$ and get

$$\begin{aligned} & \int_M \langle D^* R^E S, D S \rangle u^{2(k-1)} \\ &= \int_M \frac{1}{2} |R^E S|^2 u^{2(k-1)} + 2(k-1) \sum_{i,j} \int_M \langle R^E(i, j) S, D_j S \rangle du(i) u^{2k-3} \\ &\leq \frac{k-1}{2} \int_M |du|^2 u^{2(k-1)} + (2k-1) \int_M |R^E S|^2 u^{2(k-1)}, \end{aligned}$$

where we have used $\sum_{i,j} \langle R^E S(i, j), D^2 S(i, j) \rangle = \frac{1}{2} |R^E S|^2$.

Since $\int_M |du|^2 u^{2(k-1)} = \frac{1}{k^2} \int_M |d(u^k)|^2$, the three last inequalities give, for any $k \geq 1$,

$$\begin{aligned} \|d(u^k)\|_2^2 &\leq k \left(\int_M \underline{\text{Ric}}^- u^{2k} + \int_M \|R^E\| u^{2k} \right) \\ &\quad + k(2k-1) \left(\int_M \|R^E S\|^2 u^{2k-2} + \int_M \|\bar{\Delta} S\|^2 u^{2k-2} \right) \\ &\leq 4k^2 \left(B_1 \|u\|_{\frac{2kp}{p-1}}^{2k} + B_2 \|S\|_\infty^2 \|u\|_{\frac{2(k-1)p}{p-1}}^{2(k-1)} \right), \end{aligned}$$

where we have set

$$\begin{aligned} B_1 &= \|\underline{\text{Ric}}^-\|_p + \|R^E\|_p \leq C(n) (\|R^M\|_{2p}^2 + \lambda_1^2) = B^2, \\ B_2 &= \frac{\|\bar{\Delta} S\|_{2p}^2}{\|S\|_\infty^2} + \frac{\|R^E S\|_{2p}^2}{\|S\|_\infty^2} \leq \frac{\|\Delta_{\text{sph}} S\|_{2p}^2}{\|S\|_\infty^2} + \|\text{Ric}'\|_{2p}^2 + \|R^E\|_{2p}^2 \\ &\leq C(n) (\lambda_1^2 + \|R^M\|_{2p}^2) = B^2. \end{aligned}$$

By the Sobolev inequality given by Proposition 2.1 we get

$$\|DS\|_{\frac{2kn}{n-2}}^k \leq \|DS\|_{2k}^k + C(p, n) B k \sqrt{\|DS\|_{\frac{2kp}{p-1}}^{2k} + \|S\|_\infty^2 \|DS\|_{\frac{2(k-1)p}{p-1}}^{2(k-1)}},$$

and by $\|DS\|_{2k} \leq \|DS\|_{\frac{2kp}{p-1}} \leq \|DS\|_\infty^{1/k} \|DS\|_{\frac{2(k-1)p}{p-1}}^{(1-1/k)}$ we have

$$\begin{aligned} & \left(\frac{\|DS\|_{\frac{2kn}{n-2}}}{\|DS\|_\infty} \right)^{\frac{2kn}{n-2}} \\ &\leq \left[1 + B k C(p, n) \left(1 + \frac{\|S\|_\infty^2}{\|DS\|_\infty^2} \right) \right]^{\frac{2n}{n-2}} \left(\frac{\|DS\|_{\frac{2(k-1)p}{p-1}}}{\|DS\|_\infty} \right)^{v \frac{2(k-1)p}{p-1}}, \end{aligned}$$

where $\nu = \frac{n(p-1)}{2p(n-2)} > 1$. We set $k = \frac{a_n(p-1)}{2p} + 1$, where $(a_n)_n$ is the sequence defined by $a_0 = \frac{2p}{p-1}$ and $a_{n+1} = \nu a_n + \frac{2n}{n-2}$. Then we get

$$\begin{aligned} & \left(\frac{\|DS\|_{a_{n+1}}}{\|DS\|_\infty} \right)^{\frac{a_{n+1}}{\nu^{n+1}}} \\ & \leq \left[1 + a_n C(p, n) B \left(1 + \frac{\|S\|_\infty^2}{\|DS\|_\infty^2} \right) \right]^{\frac{2n}{(n-2)\nu^{n+1}}} \left(\frac{\|DS\|_{a_n}}{\|DS\|_\infty} \right)^{\frac{a_n}{\nu^n}}. \end{aligned}$$

Hence

$$\begin{aligned} 1 &= \lim_{n \rightarrow +\infty} \left(\frac{\|DS\|_{a_n}}{\|DS\|_\infty} \right)^{\frac{a_n}{\nu^n}} \\ &\leq \prod_{i=1}^{\infty} \left(1 + C(p, n) a_i B \left(1 + \frac{\|S\|_\infty^2}{\|DS\|_\infty^2} \right) \right)^{\frac{2n}{(n-2)\nu^i}} \left(\frac{\|DS\|_{a_0}}{\|DS\|_\infty} \right)^{a_0}. \end{aligned}$$

The Hölder inequality $\|DS\|_{a_0} \leq \|DS\|_2^{1-\frac{1}{p}} \|DS\|_\infty^{\frac{1}{p}}$, gives

$$\|DS\|_\infty \leq \prod_{i=1}^{\infty} \left(1 + C(p, n) a_i B \left(1 + \frac{\|S\|_\infty^2}{\|DS\|_\infty^2} \right) \right)^{\frac{n}{(n-2)\nu^i}} \|DS\|_2. \quad (*)$$

If $\|DS\|_\infty \geq \|S\|_\infty$, then inequality $(*)$ gives

$$\begin{aligned} \|DS\|_\infty &\leq \prod_{i=1}^{\infty} \left(1 + C(p, n) a_i B \right)^{\frac{n}{(n-2)\nu^i}} \|DS\|_2 \\ &\leq C(p, n) (\lambda_1 + \|R\|_{2p})^{\frac{pn}{2p-n}} \|DS\|_2. \end{aligned}$$

If $\|DS\|_\infty \leq \|S\|_\infty$, then inequality $(*)$ gives

$$\frac{\|DS\|_\infty}{\|DS\|_2} \leq \left(\frac{\|S\|_\infty}{\|DS\|_\infty} \right)^{\frac{2pn}{2p-n}} \prod_{i=1}^{\infty} \left(1 + C(p, n) a_i B \right)^{\frac{n}{(n-2)\nu^i}},$$

hence

$$\frac{\|DS\|_\infty}{\|S\|_\infty} \leq C(p, n) (\lambda_1 + \|R\|_{2p})^{\frac{pn}{2p-n}} \left(\frac{\|DS\|_2}{\|S\|_\infty} \right)^{\frac{2p-n}{2p-n+2pn}}.$$

At this stage note that, by Lemma 4.1 we have

$$\begin{aligned} \|DS\|_2^2 &= \langle \bar{\Delta}_{\text{sph}}(S), S \rangle_{L^2} - \langle \text{Ric}'(S), S \rangle_{L^2} \\ &\leq |\lambda_1 - n| \|S\|_2^2 + \int_M \frac{(\underline{\text{Ric}} - (n-1))^+}{\text{Vol } M} |S|^2 \leq (|\lambda_1 - n| + \bar{\rho}_p) \|S\|_\infty^2. \end{aligned}$$

Since we have $\frac{2p-n}{2p-n+2pn} \leq 1$, we get the result. \square

4.3. Critical points of the first eigenfunction. By Proposition 4.2, the section $S_f = \nabla f + fe$ of E satisfies $\|D^E S_f\|_\infty \leq C(p, n, A)(|\lambda_1 - n| + \bar{\rho}_p)^{\frac{1}{1+\gamma}} \|S_f\|_\infty$. Since we can suppose the pinching on $|\lambda_1 - n|$ and $\bar{\rho}_p$ small enough to have

$$C(p, n, A)(|\lambda_1 - n| + \bar{\rho}_p)^{\frac{1}{1+\gamma}} \leq 1/4,$$

the previous inequality and Theorem 1.6 give

$$\begin{aligned} \inf |S_f| &\geq [1 - C(p, n, A)(|\lambda_1 - n| + \bar{\rho}_p)^{\frac{1}{1+\gamma}}] \|S_f\|_\infty \\ &> C(p, n, A)(|\lambda_1 - n| + \bar{\rho}_p)^{\frac{1}{1+\gamma}} \|S_f\|_\infty \geq \|D^E S_f\|_\infty. \end{aligned}$$

We infer that if x_0 is a critical point of f , then by Lemma 4.1 we have

$$|Ddf_{x_0}(X, X) + f(x_0)| = |\langle D_X^E S_f, X \rangle_E| \leq \|D^E S_f\|_\infty < |S_f(x_0)| = |f(x_0)|$$

for any unit vector X of $T_{x_0} M$. Hence we have $-|f(x_0)| - f(x_0) < Ddf_{x_0}(X, X) < |f(x_0)| - f(x_0)$ for any critical point x_0 of f . So the only critical points of f are non degenerate global extrema, which implies that M is homeomorphic to \mathbb{S}^n by Reeb's theorem.

References

- [1] M. Anderson, Metrics of positive Ricci curvature with large diameter. *Manuscripta Math.* **68** (1990), 405–415. [Zbl 0711.53036](#) [MR 1068264](#)
- [2] E. Aubry, B. Colbois, P. Ghanaat, E. Ruh, Curvature, Harnack's inequality, and a spectral characterization of nilmanifolds. *Ann. Glob. Anal. Geom.* **23** (2003), 227–246. [Zbl 1026.58021](#) [MR 1966846](#)
- [3] E. Aubry, Pincement sur le spectre et le volume en courbure de Ricci positive. *Ann. Sci. École Norm. Sup.* **38** (2005) 387–405. [Zbl 1085.53024](#) [MR 2166339](#)
- [4] E. Aubry, Finiteness of π_1 and geometric inequalities in almost positive Ricci curvature. *Ann. Sci. École Norm. Sup.* **40** (2007), 675–695. [Zbl 1141.53034](#)
- [5] P. Bérard, G. Besson, S. Gallot, Sur une inégalité isopérimétrique qui généralise celle de Paul Levy-Gromov. *Invent. Math.* **80** (1985), 295–308. [Zbl 0571.53027](#) [MR 0788412](#)
- [6] S. Cheng, Eigenvalue comparison theorems and its geometric applications. *Math. Z.* **143** (1975), 289–297. [Zbl 0329.53035](#) [MR 0378001](#)
- [7] C. Croke, An eigenvalue pinching theorem. *Invent. Math.* **68** (1982), 253–256. [Zbl 0505.53018](#) [MR 0666162](#)
- [8] S. Ilias, Un nouveau résultat de pincement de la première valeur propre du laplacien et preuve de la conjecture du diamètre pincé. *Ann. Inst. Fourier* **43** (1993), 843–863. [Zbl 0783.53024](#) [MR 1242618](#)

- [9] M. Le Couturier, G. Robert, L^p -pinching and the geometry of compact Riemannian manifolds. *Comment. Math. Helv.* **69** (1994), 249–271. [Zbl 0821.53033](#) [MR 1282370](#)
- [10] Y. Otsu, On manifolds of positive Ricci curvature with large diameter. *Math. Z.* **206** (1991), 255–264. [Zbl 0697.53042](#) [MR 1091941](#)
- [11] G. Perelman, A diameter sphere theorem for manifolds of positive Ricci curvature. *Math. Z.* **218** (1998), 595–596. [Zbl 0831.53033](#) [MR 1326988](#)
- [12] P. Petersen, C. Sprouse, Integral curvature bounds, distance estimates and applications. *J. Differential Geom.* **50** (1998), 269–298. [Zbl 0969.53017](#) [MR 1684981](#)
- [13] T. Sakai, *Riemannian Geometry*. Transl. Math. Monogr. 149, Amer. Math. Soc., Providence, RI, 1996. [Zbl 0886.53002](#) [MR 1390760](#)

Received March 26, 2007

Erwann Aubry, Laboratoire Dieudonné, Université Nice Sophia-Antipolis, Parc Valrose,
06108 Nice, France
E-mail: eaubry@math.unice.fr