

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 84 (2009)

Artikel: On the Weinstein conjecture in higher dimensions
Autor: Albers, Peter / Hofer, Helmut
DOI: <https://doi.org/10.5169/seals-99125>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 05.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

On the Weinstein conjecture in higher dimensions

Peter Albers and Helmut Hofer*

Abstract. The existence of a “Plastikstufe” for a contact structure implies the Weinstein conjecture for all supporting contact forms.

Mathematics Subject Classification (2000). 53D10, 53D35, 37J45.

Keywords. Contact structure, Reeb vector field, over-twisted, Plastikstufe, Weinstein conjecture.

1. Introduction and main result

A one-form λ on an odd-dimensional manifold M^{2n-1} is called a contact form, provided $\lambda \wedge d\lambda^{n-1}$ is a volume-form. Associated to a contact form λ we have the Reeb vector field X defined by

$$i_X \lambda = 1 \quad \text{and} \quad i_X d\lambda = 0$$

and the contact structure $\xi = \ker(\lambda)$. In 1978, A. Weinstein, [21], motivated by a result of P. Rabinowitz, [16], and one of his own results, [20], made the following conjecture:

A Reeb vector field on a closed manifold M^{2n-1} admits a periodic orbit.

The first break-through on this conjecture was obtained by C. Viterbo, [19], showing that compact energy surfaces in \mathbb{R}^{2n} of contact-type have periodic orbits. Extending Gromov’s theory of pseudoholomorphic curves, [3], to symplectized contact manifolds, H. Hofer, [4], related the Weinstein conjecture to the existence of certain pseudoholomorphic curves. He showed that in dimension three the Weinstein conjectures holds in many cases. In particular, he showed that Reeb vector fields associated to over-twisted contact structures admit periodic orbits. Recently the Weinstein conjecture in dimension three was completely settled by C. Taubes, [17, 18], who exploited relationships between Seiberg–Witten–Floer homology, [12], and embedded contact

*Research partially supported by NSF grant DMS-0603957 and DFG grant AL 904/1-1.

homology, [11], in order to construct holomorphic curves in the symplectized contact manifold out of nontrivial Seiberg–Witten–Floer homology classes. For more references on the Weinstein conjecture see [6].

In this note we show that many Reeb vector fields on higher dimensional closed manifolds have periodic orbits generalizing the main result from [4]. Our existence result is closely connected to the interesting attempt by K. Niederkrüger [13] to generalize the three-dimensional notion of an overtwisted contact structure. He introduced the concept of a *Plastikstufe* which currently seems to be the most compelling generalisation given recent further developments by F. Presas, [15], and K. Niederkrüger and O. van Koert, [14].

Let us denote by (M, ξ) a pair consisting of a closed manifold M of dimension $2n - 1$ and a co-oriented contact structure ξ . We denote by \mathbb{D}^2 the closed unit disk in \mathbb{C} with coordinates $x + iy$.

Definition 1.1. We say that (M, ξ) contains a *Plastikstufe* with singular set S provided M admits a closed submanifold S of dimension $n - 2$ and an embedding $\iota: \mathbb{D}^2 \times S \rightarrow M$ with $\iota(\{0\} \times S) = S$ having the following properties:

- (1) There exists a contact form λ_{PS} inducing ξ so that the one-form $\beta := \iota^* \lambda_{\text{PS}}$ satisfies $\beta \wedge d\beta = 0$ and moreover $\beta \neq 0$ on $(\mathbb{D}^2 \setminus \{0\}) \times S$. Near $\{0\} \times S$ the form β is given by $\beta = xdy - ydx$ and the pull-back of β to $\partial\mathbb{D}^2 \times S$ vanishes.
- (2) The complement of $\{0\} \times S$ in $(\mathbb{D}^2 \setminus \partial\mathbb{D}^2) \times S$ is smoothly foliated by β via an S^1 -family of leaves diffeomorphic to $(0, 1) \times S$, where one of the ends converges to the singular set $\{0\} \times S$ and the other is asymptotic to the leaf $\partial\mathbb{D}^2 \times S$.

The set $\mathcal{PS}(S) = \iota(\mathbb{D}^2 \times S)$ is called the *Plastikstufe*.

Let us observe that the existence of a *Plastikstufe* for a given contact structure involves the existence of a certain inducing contact form. This is different from the three-dimensional case where an over-twisted disk is defined only in terms of the contact structure and does not require the existence of a particular contact form. In the following we shall call a closed co-oriented contact manifold (M, ξ) PS-overtwisted provided there exists a contact form λ_{PS} inducing ξ containing a *Plastikstufe*. Recently Niederkrüger and van Koert showed that every odd-dimensional sphere S^{2n-1} with $n \geq 3$ has a contact structure admitting a *Plastikstufe*. If now (M^{2n-1}, ξ) is a co-oriented contact manifold then a connected sum with a PS-overtwisted sphere admits by standard arguments a contact structure which is PS-overtwisted. In particular, any closed manifold of dimension $2n - 1$ admitting a co-oriented contact structure also admits a PS-overtwisted contact structure. Our main result is the following theorem.

Theorem. *Let (M, ξ) be a closed PS-overtwisted contact manifold. Then every Reeb vector field associated to a contact form λ inducing ξ has a contractible periodic orbit.*

Remark 1.2. In [13] Niederkrüger shows that a PS-overtwisted contact structure does not have a semi-positive symplectic filling. We noticed that some of his idea combined with ideas from [4] lead to the above theorem. We also observed that the limitation to semi-positive fillings is not necessary and can be removed using polyfolds [5]. This will be discussed in a forthcoming paper.

2. Background

All material in this section is taken from [13].

2.1. Local normal form. Let (M, λ) contain a Plastikstufe $\mathcal{PS}(S)$. In [13, Section 3.1] it is proved that there exist constants $\varepsilon, C > 0$ and an open set V in the symplectic manifold $((-\varepsilon, 0] \times M, d(e^s \lambda))$ such that $\{0\} \times S \subset V$ and V is symplectomorphic to the subset

$$U := \left\{ ((z_1, z_2), (q, p)) \in \mathbb{C}^2 \times T^*S \mid \begin{array}{l} -C < \operatorname{Re}(z_1) \leq 0, \\ -C < \operatorname{Im}(z_1) < C, \\ \operatorname{Re}(z_1) + \frac{1}{4}|z_2|^2 + \frac{1}{2}\|p\|^2 \leq 0 \end{array} \right\} \quad (2.1)$$

of $\mathbb{C}^2 \times T^*S$ which carries its natural symplectic structure. Moreover, $M \cap V$ corresponds to equality in the last equation and $\mathcal{PS}(S) \cap V$ to equality and $\operatorname{Im}(z_1) = 0, p = 0$.

2.2. Bishop family. The local model U contains a natural $(n-1)$ -dimensional *Bishop family* given by

$$\begin{aligned} u_{t_0, q_0} : \mathbb{D}^2 &\longrightarrow \mathbb{C}^2 \times T^*S, \\ z &\longmapsto ((-t_0, 2\sqrt{t_0}z), (q_0, 0)), \end{aligned} \quad (2.2)$$

where $0 \leq t_0 < C$ is a real parameter and $q_0 \in S$. The maps $u_{(t_0, q_0)}$ are $(i \times j)$ -holomorphic, where j denotes the natural almost complex structure on T^*S induced by the Levi-Civita connection of a Riemannian metric on S . Moreover, they have boundary on the set corresponding to $\mathcal{PS}(S)$.

We denote by J the almost complex structure on V obtained by pulling back the almost complex structure $i \times j$ from $\mathbb{C}^2 \times T^*S$. Then we can pull back the Bishop family to holomorphic maps (denoted by the same symbols)

$$u_{t_0, q_0} : \mathbb{D}^2 \longrightarrow V \subset (-\varepsilon, 0] \times M. \quad (2.3)$$

2.3. Uniqueness results for holomorphic disks. We extend the almost complex structure J from the set V to a compatible almost complex structure on $(W := (-\infty, 0] \times M, d(e^s \lambda))$. We introduce the notation

$$\widehat{\mathcal{PS}}(S) = \mathcal{PS}(S) \setminus (\partial \mathcal{PS}(S) \cup S) \quad (2.4)$$

and remark that $\widehat{\mathcal{PS}}(S)$ is totally real with respect to J . The following proposition is taken from [13, Proposition 7].

Proposition 2.1. *Let $u: (\mathbb{D}^2, \partial \mathbb{D}^2) \rightarrow (W, \widehat{\mathcal{PS}}(S))$ be a J -holomorphic disk which is simple. Moreover, we assume that $u(S^1) \subset \mathcal{PS}(S)$ bounds a disk in $\mathcal{PS}(S)$ and*

$$\text{image}(u) \cap V \neq \emptyset. \quad (2.5)$$

Then, up to an element in $\text{Aut}(\mathbb{D}^2)$, we have

$$u = u_{t_0, q_0}, \quad (2.6)$$

that is, after reparametrization, the holomorphic disk u is a member of the Bishop family.

3. Proof of the theorem

By assumption there exists a contact form λ_{PS} on M containing a Plastikstufe. Let λ be another contact form inducing the same contact structure.

We assume by contradiction that there exists no contractible closed Reeb orbit for λ .

3.1. The set-up. We choose a function $f: M \rightarrow \mathbb{R}$ such that $\lambda = f \lambda_{\text{PS}}$. Since multiplying λ with a non-zero constant does not change its Reeb orbits (up to reparametrization) we may assume without loss of generality that the function f takes only values in $(0, 1)$. Then we can choose a smooth family of functions $f_s: M \rightarrow (0, \infty)$ for $s \in [-1, -\varepsilon]$ satisfying

$$f_s = \begin{cases} 1 & \text{near } s = -\varepsilon, \\ f & \text{near } s = -1, \end{cases} \quad \text{and moreover} \quad \frac{\partial f_s}{\partial s} \geq 0. \quad (3.1)$$

This gives rise to a smooth family $\lambda_s = f_s \lambda_{\text{PS}}$ of contact forms which we extend by λ_{PS} for $s \geq -\varepsilon$ and by λ for $s \leq -1$. On $W = (-\infty, 0] \times M$ we choose an exact symplectic form Ω which satisfies

$$\Omega = \begin{cases} d(e^s \lambda_{\text{PS}}) & \text{on } [-\varepsilon, 0] \times M, \\ d(e^s \lambda) & \text{on } (-\infty, -1] \times M. \end{cases} \quad (3.2)$$

This is possible due to our choice of the family f_s . This has been used in the literature many times, see for instance [7]. We modify the almost complex structure J from above to a compatible almost complex structure J on (W, Ω) by requiring that on $(-\infty, -2] \times M$ the almost complex structure is adapted to the negative part of the symplectization of λ , in the sense of [1]. On V it remains as defined in the previous section. In particular, (W, Ω) still contains the Bishop family u_{t_0, q_0} . We denote the relative homotopy class given by the Bishop disks by $a \in \pi_2(W, \widehat{\mathcal{PS}}(S))$ and set

$$\mathcal{M}(J) := \{u : (\mathbb{D}^2, \partial\mathbb{D}^2) \longrightarrow (W, \widehat{\mathcal{PS}}(S)) \mid \bar{\partial}_J u = 0, [u] = a, \text{lk}(u, S) = 1\}, \quad (3.3)$$

$$\widehat{\mathcal{M}}(J) := \mathcal{M}(J)/\text{Aut}(\mathbb{D}^2), \quad (3.4)$$

where $\text{lk}(u, S)$ is the linking number of $u(S^1)$ in $\mathcal{PS}(S)$ with the set S . This is defined as follows. By definition $\widehat{\mathcal{PS}}(S)$ is foliated by an S^1 -family of Legendrian submanifolds, thus there exists a natural map $\theta : \widehat{\mathcal{PS}}(S) \rightarrow S^1$. We set $\text{lk}(u, S) := \deg(\theta \circ u|_{S^1})$.

3.2. The proof. We need the following three facts established in Propositions 8–10 in [13].

- (1) The Maslov index of a equals $\mu_{\text{Maslov}}(a) = 2$,
- (2) the almost complex structure J is regular at members of the Bishop family,
- (3) the energy of all elements in $\mathcal{M}(J)$ is uniformly bounded.

The totally real submanifold $\widehat{\mathcal{PS}}(S)$ is non-compact. Since $\partial\mathcal{PS}(S)$ is a closed leaf of the characteristic foliation the maximum principle implies that no holomorphic maps intersect $\partial\mathcal{PS}(S)$ at an interior point. According to Proposition 2.1 near S the only holomorphic disks are members of the Bishop family. Therefore, the non-compactness of $\widehat{\mathcal{PS}}(S)$ poses no problem. Moreover, due to the energy bounds and the specific structure of the almost complex structure J on the end of W we can apply the ideas of the SFT-compactness theorem [1]. Since we assumed that there exists no contractible closed Reeb orbits bubbling-off cannot occur in the interior. Therefore, the only non-compactness of the moduli space $\widehat{\mathcal{M}}(J)$ comes from bubbling-off of holomorphic disks having boundary on $\widehat{\mathcal{PS}}(S)$. The next proposition is taken from [13, Proposition 11] and shows that there exists no bubbling-off of holomorphic disks.

Proposition 3.1. *Given a sequence $(u_n) \subset \widehat{\mathcal{M}}(J)$ there exists a subsequence either converging to an element in $\widehat{\mathcal{M}}(J)$ or to a point in S .*

The latter case occurs if a family of Bishop disks shrinks to a point in S . We remark that in the former case the limit is simple.

Proposition 3.2. *For a compatible almost complex structure J , which is generic on the subset $((-2, 0] \times M) \setminus V$ of (W, Ω) , the moduli space $\mathcal{M}(J)$ is a smooth manifold of dimension*

$$\dim \mathcal{M}(J) = n + 2. \quad (3.5)$$

Proof. We pick $u \in \mathcal{M}(J)$. In case that $\text{image}(u) \cap V \neq \emptyset$ we conclude from Proposition 2.1 that u is a member of the Bishop family. In particular, $\text{image}(u) \subset V$. Moreover, J is already regular for members in the Bishop family.

If $\text{image}(u) \cap V = \emptyset$ then u has to pass through the region $((-2, 0] \times M) \setminus V$. Since all the disks are simple a generic J will be regular, see for example [2]. The dimension formula follows from the fact that $\mu_{\text{Maslov}}(a) = 2$ and $\dim \widehat{\mathcal{PS}}(S) = n$. \square

We consider the evaluation map

$$\begin{aligned} \text{ev}: \widehat{\mathcal{M}}(J)_{S^1} &:= \mathcal{M}(J) \times_{\text{Aut}(\mathbb{D}^2)} S^1 \longrightarrow \widehat{\mathcal{PS}}(S) \subset M, \\ [u, t] &\longmapsto u(e^{2\pi i t}), \end{aligned} \quad (3.6)$$

defined on the smooth manifolds $\widehat{\mathcal{M}}(J)_{S^1}$ of dimension $\dim \widehat{\mathcal{M}}(J)_{S^1} = n$. The following proposition is proved analogously to the transversality result above.

Proposition 3.3. *For a generic J as in the previous proposition the evaluation map is smooth.*

To derive the contradiction to the assumption that λ has no closed Reeb orbits we make the following

Definition 3.4. For a point $p = \iota(z, s) \in \iota(\mathbb{D}^2 \times S) = \mathcal{PS}(S)$ we define the distance of p to S by $d(p, S) = |z|$ and set for $\delta > 0$ sufficiently small

$$\widehat{\mathcal{M}}(J)_{S^1}^\delta := \{[u, t] \in \widehat{\mathcal{M}}(J)_{S^1} \mid d(\text{ev}([u, t]), S) \geq \delta\}. \quad (3.7)$$

Then we have by construction of the Bishop family

$$\text{ev}(\partial \widehat{\mathcal{M}}(J)_{S^1}^\delta) = \iota(S_\delta^1 \times S), \quad (3.8)$$

where $S_\delta^1 = \{z \in \mathbb{D}^2 \mid |z| = \delta\}$. We conclude that

$$[\text{ev}(\partial \widehat{\mathcal{M}}(J)_{S^1}^\delta)] \in H_{n-1}(\widehat{\mathcal{PS}}(S), \mathbb{Z}/2)$$

is the generator. On the other hand the set $\text{ev}(\partial \widehat{\mathcal{M}}(J)_{S^1}^\delta)$ is clearly the boundary of the compact, see Proposition 3.1, manifold $\text{ev}(\widehat{\mathcal{M}}(J)_{S^1}^\delta)$. This implies, that

$$[\text{ev}(\partial \widehat{\mathcal{M}}(J)_{S^1}^\delta)] = 0 \in H_{n-1}(\widehat{\mathcal{PS}}(S), \mathbb{Z}/2).$$

This contradiction concludes the proof of the theorem.

References

- [1] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, and E. Zehnder, Compactness results in symplectic field theory. *Geom. Topol.* **7** (2003), 799–888. [Zbl 1131.53312](#) [MR 2026549](#)
- [2] D. L. Dragnev, Fredholm theory and transversality for noncompact pseudoholomorphic maps in symplectizations. *Comm. Pure Appl. Math.* **57** (2004), 726–763. [Zbl 1063.53086](#) [MR 2038115](#)
- [3] M. Gromov, Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.* **82** (1985), 307–347. [Zbl 0592.53025](#) [MR 0809718](#)
- [4] H. Hofer, Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three. *Invent. Math.* **114** (1993), 515–563. [Zbl 0797.58023](#) [MR 1244912](#)
- [5] —, A general Fredholm theory and Applications. In *Proceedings of the CDM* (Harvard), 2004.
- [6] H. Hofer, K. Wysocki, and E. Zehnder, Pseudoholomorphic curves and dynamics in three dimensions. In *Handbook of dynamical systems*, Vol. 1A, North-Holland, Amsterdam 2002, 1129–1188. [Zbl 1052.53063](#) [MR 1928532](#)
- [7] —, Finite energy foliations of tight three-spheres and Hamiltonian dynamics. *Ann. of Math.* (2) **157** (2003), 125–255. [Zbl 01933157](#) [MR 1954266](#)
- [8] —, A General Fredholm Theory I: A Splicing-Based Differential Geometry. *J. Eur. Math. Soc.* (JEMS) **9** (2007), 841–876. [Zbl 05255318](#) [MR 2341834](#)
- [9] —, A General Fredholm Theory II: Implicit Function Theorems. *Geom. Funct. Anal.*, to appear; [arXiv:0705.1310v2](#).
- [10] —, Applications of Polyfold Theory I: Gromov-Witten Theory. Article in preparation.
- [11] M. Hutchings, An index inequality for embedded pseudoholomorphic curves in symplectizations. *J. Eur. Math. Soc.* (JEMS) **4** (2002), 313–361. [Zbl 1017.58005](#) [MR 1941088](#)
- [12] P. B. Kronheimer and T. S. Mrowka, *Floer homology for Seiberg-Witten monopoles*. Cambridge University Press, to appear.
- [13] K. Niederkrüger, The plastikstufe – a generalization of the overtwisted disk to higher dimensions. *Algebr. Geom. Topol.* **6** (2006), 2473–2508. [Zbl 1129.53056](#) [MR 2286033](#)
- [14] K. Niederkrüger and O. van Koert, Every contact manifold can be given a non-fillable contact structure. *Internat. Math. Res. Not.* (2007), article ID rnm115. [Zbl 1133.53053](#) [MR 2380008](#)
- [15] F. Presas, A class of non-fillable contact structures. *Geom. Topol.* **11** (2007), 2203–2225. [Zbl 1132.57023](#) [MR 2372846](#)
- [16] P. H. Rabinowitz, Periodic solutions of a Hamiltonian system on a prescribed energy surface. *J. Differential Equations* **33** (1979), 336–352. [Zbl 0424.34043](#) [MR 0543703](#)
- [17] C. H. Taubes, The Seiberg-Witten equations and the Weinstein conjecture. *Geom. Topol.* **11** (2007), 2117–2202. [Zbl 1135.57015](#) [MR 2350473](#)
- [18] —, The Seiberg-Witten equations and the Weinstein conjecture II: More closed integral curves of the Reeb vector field. Preprint 2007; [arXiv:math/0702366v2](#).

- [19] C. Viterbo, A proof of Weinstein's conjecture in \mathbf{R}^{2n} . *Ann. Inst. H. Poincaré Anal. Non Linéaire* **4** (1987), 337–356. [Zbl 0631.58013](#) [MR 0917741](#)
- [20] A. Weinstein, Periodic orbits for convex Hamiltonian systems. *Ann. of Math. (2)* **108** (1978), 507–518. [Zbl 0403.58001](#) [MR 0512430](#)
- [21] —, On the hypotheses of Rabinowitz' periodic orbit theorems. *J. Differential Equations* **33** (1979), 353–358. [Zbl 0388.58020](#) [MR 0543704](#)

Received June 9, 2007

Peter Albers, Courant Institute, New York University, 251 Mercer Street, NY 10012, U.S.A.

E-mail: albers@cims.nyu.edu

Helmut Hofer, Courant Institute, New York University, 251 Mercer Street, NY 10012, U.S.A.

E-mail: hofer@cims.nyu.edu