

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 85 (2010)

Artikel: Reconstructing p -divisible groups from their truncations of small level
Autor: Vasiu, Adrian
DOI: <https://doi.org/10.5169/seals-130660>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Reconstructing p -divisible groups from their truncations of small level

Adrian Vasiu

Dedicated to the memory of Angela Vasiu

Abstract. Let k be an algebraically closed field of characteristic $p > 0$. Let D be a p -divisible group over k . Let n_D be the smallest non-negative integer for which the following statement holds: if C is a p -divisible group over k of the same codimension and dimension as D and such that $C[p^{n_D}]$ is isomorphic to $D[p^{n_D}]$, then C is isomorphic to D . To the Dieudonné module of D we associate a non-negative integer ℓ_D which is a computable upper bound of n_D . If D is a product $\prod_{i \in I} D_i$ of isoclinic p -divisible groups, we show that $n_D = \ell_D$; if the set I has at least two elements we also show that $n_D \leq \max\{1, n_{D_i}, n_{D_i} + n_{D_j} - 1 \mid i, j \in I, j \neq i\}$. We show that we have $n_D \leq 1$ if and only if $\ell_D \leq 1$; this recovers the classification of minimal p -divisible groups obtained by Oort. If D is quasi-special, we prove the Traverso truncation conjecture for D . If D is F -cyclic, we explicitly compute n_D . Many results are proved in the general context of latticed F -isocrystals with a (certain) group over k .

Mathematics Subject Classification (2000). 11E57, 11G10, 11G18, 11G25, 14F30, 14G35, 14L05, 14L30, 20G25.

Keywords. p -divisible groups, F -crystals, algebras, affine group schemes.

Contents

1	Introduction	165
2	Preliminaries	172
3	The proof of Main Theorem A	175
4	Direct applications to p -divisible groups	189
5	On Main Theorem B	198
	References	201

1. Introduction

Let $p \in \mathbb{N}$ be a prime. Let k be an algebraically closed field of characteristic p . Let $c, d \in \mathbb{N} \cup \{0\}$ be such that $r := c + d > 0$. Let D be a p -divisible group over k of codimension c and dimension d . The height of D is r . Let $n_D \in \mathbb{N} \cup \{0\}$ be the

smallest number for which the following statement holds: if C is a p -divisible group of codimension c and dimension d over k such that $C[p^{n_D}]$ is isomorphic to $D[p^{n_D}]$, then C is isomorphic to D . We have $n_D = 0$ if and only if $cd = 0$. For the existence of n_D we refer to [Ma, Chapter III, Section 3], [Tr1, Theorem 3], [Tr2, Theorem 1], [Va1, Corollary 1.3], or [Oo2, Corollary 1.7]. For instance, one has the following gross estimate $n_D \leq cd + 1$ (cf. [Tr1, Theorem 3]). The classical Dieudonné theory says that the category of p -divisible groups over k is antiequivalent to the category of Dieudonné modules over k . Thus the existence of n_D gets translated into a suitable problem pertaining to Dieudonné modules and thus to a particular type of latticed F -isocrystals over k (see Section 1.1 below for precise definitions).

Traverso's truncation conjecture predicts that $n_D \leq \min\{c, d\}$, cf. [Tr3, Section 40, Conjecture 4]. This surprising and old conjecture is known to hold only in few cases (like for supersingular p -divisible groups over k ; see [NV, Theorem 1.2]). To prove different refinements of this conjecture, one needs to have easy ways to compute and estimate n_D . Each estimate of n_D represents progress towards the classification of p -divisible groups over k ; implicitly, it represents progress towards the understanding of the ultimate stratifications defined in [Va1, Section 5.3] and (thus also) of the special fibres of all integral canonical models of Shimura varieties of Hodge type. The goal of the paper is to put forward *basic principles* that compute either n_D or some very sharp upper bounds of n_D .

For the sake of generality, a great part of this paper will be worked out in the context of latticed F -isocrystals with a (certain) group over k .

1.1. Latticed F -isocrystals. Let $W(k)$ be the ring of Witt vectors with coefficients in k . Let $B(k)$ be the field of fractions of $W(k)$. Let σ be the Frobenius automorphism of $W(k)$ and $B(k)$ induced from k .

By a *latticed F -isocrystal* over k we mean a pair (M, ϕ) , where M is a free $W(k)$ -module of finite rank and $\phi: M[\frac{1}{p}] \xrightarrow{\sim} M[\frac{1}{p}]$ is a σ -linear automorphism. We recall that if $\phi(M) \subseteq M$, then the pair (M, ϕ) is called an *F -crystal* over k . We also recall that if $pM \subseteq \phi(M) \subseteq M$, then the pair (M, ϕ) is called a *Dieudonné module* over k and $\vartheta := p\phi^{-1}: M \rightarrow M$ is called the *Verschiebung map* of (M, ϕ) .

The composite of $W(k)$ -linear maps endows $\text{End}(M)$ with a natural structure of a $W(k)$ -algebra (and thus also of a Lie algebra over $W(k)$). We denote also by ϕ the σ -linear automorphism of $\text{End}(M[\frac{1}{p}])$ that takes $e \in \text{End}(M[\frac{1}{p}])$ to $\phi(e) := \phi \circ e \circ \phi^{-1}$. Let $G_{B(k)}$ be a connected subgroup of $\text{GL}_{M[\frac{1}{p}]}$ such that its Lie algebra $\text{Lie}(G_{B(k)})$ is left invariant by ϕ i.e., we have $\phi(\text{Lie}(G_{B(k)})) = \text{Lie}(G_{B(k)})$. Let G be the schematic closure of $G_{B(k)}$ in GL_M . The triple (M, ϕ, G) is called a *latticed F -isocrystal with a group* over k , cf. [Va1, Definition 1.1 (a)]. Let $\mathfrak{g} := \text{Lie}(G_{B(k)}) \cap \text{End}(M)$; it is a Lie subalgebra of $\text{End}(M)$ which as a $W(k)$ -submodule is a direct summand. If G is smooth over $\text{Spec}(W(k))$, then $\mathfrak{g} = \text{Lie}(G)$. Let $n_G \in \mathbb{N} \cup \{0\}$ be the *i -number* of (M, ϕ, G) introduced in [Va1, Definition 3.1.4]. Thus n_G is the smallest non-negative

integer for which the following statement holds:

- If $g \in G(W(k))$ is congruent to 1_M modulo p^{n_G} , then there exists $h \in G(W(k))$ which is an isomorphism between $(M, g\phi, G)$ and (M, ϕ, G) (equivalently, between $(M, g\phi)$ and (M, ϕ)). In other words, we have $hg\phi h^{-1} = \phi$ (equivalently, $hg\phi(h)^{-1} = 1_M$).

In [Va1] we developed methods that provide good upper bounds of n_G (see [Va1, Section 3.1.3 and Example 3.1.5]). The methods used exponential maps and applied to all possible types of affine, integral group schemes G over $\text{Spec}(W(k))$. But when the type of G is simple (like when G is GL_M), then one can obtain significantly better bounds. This idea was exploited to some extent in [Va1, Section 3.3] and it is brought to full fruition in this paper. Accordingly, in the whole paper we will work under the following assumption:

1.1.1. Assumption. We have $M \neq 0$, the $W(k)$ -submodule \mathfrak{g} of $\text{End}(M)$ is a $W(k)$ -subalgebra of $\text{End}(M)$ (and not only a Lie subalgebra of $\text{End}(M)$), and (thus) G is the group scheme over $\text{Spec}(W(k))$ of invertible elements of \mathfrak{g} .

Typical cases we have in mind: (i) G is either GL_M or a parabolic subgroup scheme of GL_M ; (ii) G is the centralizer in GL_M of a semisimple $W(k)$ -subalgebra of $\text{End}(M)$; and (iii) \mathfrak{g} is $W(k)1_M \oplus \mathfrak{n}$, with \mathfrak{n} a nilpotent subalgebra (without unit) of $\text{End}(M)$.

1.1.2. Newton polygon slopes. Dieudonné's classification of F -isocrystals over k (see [Di, Theorems 1 and 2], [Ma, Chapter 2, Section 4], [De], etc.) implies that we have a direct sum decomposition $M[\frac{1}{p}] = \bigoplus_{\alpha \in \mathbb{Q}} W(\alpha)$ that is left invariant by ϕ and that has the property that all Newton polygon slopes of $(W(\alpha), \phi)$ are α . We recall that if $m \in \mathbb{N}$ is the smallest number such that $m\alpha \in \mathbb{Z}$, then there exists a $B(k)$ -basis for $W(\alpha)$ which is formed by elements fixed by $p^{-m\alpha}\phi^m$. One says that (M, ϕ) is *isoclinic* if there exists a rational number α such that we have $M[\frac{1}{p}] = W(\alpha)$. We consider the direct sum decomposition into $B(k)$ -vector spaces

$$\text{End}(M[\frac{1}{p}]) = L_+ \oplus L_0 \oplus L_-$$

that is left invariant by ϕ and such that all Newton polygon slopes of (L_+, ϕ) are positive, all Newton polygon slopes of (L_-, ϕ) are negative, and finally all Newton polygon slopes of (L_0, ϕ) are 0. We have direct sum decompositions $L_+ = \bigoplus_{\alpha, \beta \in \mathbb{Q}, \alpha < \beta} \text{Hom}(W(\alpha), W(\beta))$, $L_- = \bigoplus_{\alpha, \beta \in \mathbb{Q}, \alpha < \beta} \text{Hom}(W(\beta), W(\alpha))$, and $L_0 = \bigoplus_{\alpha \in \mathbb{Q}} \text{End}(W(\alpha))$. Thus both L_+ and L_- are nilpotent subalgebras (without unit) of $\text{End}(M)$.

We have $L_0 = \text{End}(M[\frac{1}{p}])$ if and only if (M, ϕ) is isoclinic.

1.2. Level modules and torsions. We define

$$\begin{aligned} O_+ &:= \{x \in \text{End}(M) \cap L_+ \mid \phi^q(x) \in \text{End}(M) \cap L_+ \text{ for all } q \in \mathbb{N}\} \\ &= \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^{-q}(\text{End}(M) \cap L_+). \end{aligned}$$

Let $A_0 := \{e \in \text{End}(M) \mid \phi(e) = e\}$ be the \mathbb{Z}_p -algebra of endomorphisms of (M, ϕ) . Let O_0 be the $W(k)$ -span of A_0 ; it is a $W(k)$ -subalgebra of $\text{End}(M) \cap L_0$. We have identities

$$O_0 = A_0 \otimes_{\mathbb{Z}_p} W(k) = \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^q(\text{End}(M) \cap L_0) = \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^{-q}(\text{End}(M) \cap L_0).$$

We also define

$$\begin{aligned} O_- &:= \{x \in \text{End}(M) \cap L_- \mid \phi^{-q}(x) \in \text{End}(M) \cap L_- \text{ for all } q \in \mathbb{N}\} \\ &= \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^q(\text{End}(M) \cap L_-). \end{aligned}$$

As all Newton polygon slopes of (L_+, ϕ) are positive, for each $x \in L_+$ the sequence $(\phi^q(x))_{q \in \mathbb{N}}$ of elements of L_+ converges to 0 in the p -adic topology. This implies that there exists $s \in \mathbb{N}$ such that $p^s x \in O_+$. Thus we have $O_+[\frac{1}{p}] = L_+$. As O_+ is a $W(k)$ -submodule of the finitely generated $W(k)$ -module $\text{End}(M)$, we conclude that O_+ is a lattice of L_+ . A similar argument shows that O_0 and O_- are lattices of L_0 and L_- (respectively). We have the following relations: $\phi(O_+) \subseteq O_+$, $\phi(O_0) = O_0 = \phi^{-1}(O_0)$, $\phi^{-1}(O_-) \subseteq O_-$, $L_+L_0 + L_0L_+ \subseteq L_+$, $L_0L_0 \subseteq L_0$, and $L_0L_- + L_-L_0 \subseteq L_-$. These relations imply the following:

- (i) Both O_+ and O_- are left and right O_0 -modules.
- (ii) The direct sum $O_+ \oplus O_0$ (resp. $O_0 \oplus O_-$) is a $W(k)$ -subalgebra of $\text{End}(M)$ that has O_+ (resp. O_-) as a nilpotent, two-sided ideal.

Let $O := O_+ \oplus O_0 \oplus O_-$; it is a lattice of $\text{End}(M)[\frac{1}{p}]$ contained in $\text{End}(M)$. In general, O is not a $W(k)$ -subalgebra of $\text{End}(M)$ (see Example 2.2). Thus we call O the *level module* of (M, ϕ) .

Let $O_G := (\mathfrak{g} \cap O_+) \oplus (\mathfrak{g} \cap O_0) \oplus (\mathfrak{g} \cap O_-)$; it is a lattice of $\mathfrak{g}[\frac{1}{p}]$ contained in \mathfrak{g} . We refer to O_G as the *level module* of (M, ϕ, G) . We note down that $O = O_{\text{GL}_M}$.

By the *level torsion* of (M, ϕ, G) we mean the unique number $\ell_G \in \mathbb{N} \cup \{0\}$ for which the following inclusions hold

$$p^{\ell_G} \mathfrak{g} \subseteq O_G \subseteq \mathfrak{g} \tag{1}$$

and which obeys the following two disjoint rules:

- (a) if $\mathfrak{g} = O_G$ and if the two-sided ideal of the $W(k)$ -algebra \mathfrak{g} generated by $(\mathfrak{g} \cap O_+) \oplus (\mathfrak{g} \cap O_-)$ is not topologically nilpotent, then $\ell_G := 1$;
- (b) in all other cases, ℓ_G is the smallest non-negative integer for which (1) holds.

1.2.1. A connection to [Va1]. Let $m_G := T(\mathfrak{g}, \phi)$ be the Fontaine–Dieudonné torsion of (\mathfrak{g}, ϕ) introduced in [Va1, Definitions 2.2.2 (a) and (b)]. We recall that m_G is the smallest non-negative integer with the property that there exists a $W(k)$ -submodule \mathfrak{m} of \mathfrak{g} which contains $p^{m_G} \mathfrak{g}$ and for which the pair (\mathfrak{m}, ϕ) is a Fontaine–Dieudonné p -divisible object over k in the sense of loc. cit. One has a direct sum decomposition $(\mathfrak{m}, \phi) = \bigoplus_{j \in J} (\mathfrak{m}_j, \phi)$ such that each pair (\mathfrak{m}_j, ϕ) is an elementary Fontaine–Dieudonné p -divisible object over k . The pair (\mathfrak{m}_j, ϕ) is a special type of isoclinic latticed F -isocrystals over k that are definable over \mathbb{F}_p ; let $\alpha_j \in \mathbb{Q}$ be the Newton polygon slope of (\mathfrak{m}_j, ϕ) . One basic property of (\mathfrak{m}_j, ϕ) is the following: if $\alpha_j > 0$ (resp. $\alpha_j = 0$ or $\alpha_j < 0$), then we have $\phi(\mathfrak{m}_j) \subseteq \mathfrak{m}_j$ (resp. $\phi(\mathfrak{m}_j) = \mathfrak{m}_j$ or $\phi^{-1}(\mathfrak{m}_j) \subseteq \mathfrak{m}_j$). Thus if $\alpha_j > 0$ (resp. $\alpha_j = 0$ or $\alpha_j < 0$), then we have $\mathfrak{m}_j \subseteq \mathfrak{g} \cap O_+$ (resp. $\mathfrak{m}_j \subseteq \mathfrak{g} \cap O_0$ or $\mathfrak{m}_j \subseteq \mathfrak{g} \cap O_-$). This implies that $\mathfrak{m} \subseteq O_G$. Therefore we have $\ell_G \leq m_G$ except in the case when $\mathfrak{g} = O_G = \mathfrak{m}$ and $\ell_G = 1$. This implies that $\ell_G \leq \max\{1, m_G\}$. In general, ℓ_G can be smaller than m_G (see Example 2.2).

1.2.2. Example. We assume that all Newton polygon slopes of (\mathfrak{g}, ϕ) are 0. Then we have $O_G = \mathfrak{g} \cap O_0$ and ℓ_G is the smallest non-negative integer such that we have inclusions $p^{\ell_G} \mathfrak{g} \subseteq O_G \subseteq \mathfrak{g}$. As the $W(k)$ -module \mathfrak{g} is a direct summand of $\text{End}(M)$, we have $O_G = \mathfrak{g} \cap O_0 = \mathfrak{g}[\frac{1}{p}] \cap O_0$. This implies that $\phi(O_G) = O_G$ and therefore O_G has a $W(k)$ -basis formed by elements of $\mathfrak{g} \cap A_0$. Thus (O_G, ϕ) is a Fontaine–Dieudonné p -divisible object over k ; therefore $\ell_G = m_G$.

Our first main goal is to prove (see Section 3) the following theorem.

1.3. Main Theorem A. *We recall that (M, ϕ, G) is a latticed F -isocrystal with a group over k and that we work under Assumption 1.1.1.*

- (a) *We have the inequality $n_G \leq \ell_G$.*
- (b) *Assume that (M, ϕ) is a direct sum of isoclinic latticed F -isocrystals over k . Then we have $n_{\text{GL}_M} = \ell_{\text{GL}_M}$.*

We neither know nor expect examples with $n_{\text{GL}_M} < \ell_{\text{GL}_M}$. Our second main goal is to apply Main Theorem A to study p -divisible groups over k .

1.4. First applications to p -divisible groups. Let D and n_D be as in the beginning paragraph of the paper. We say that D is isoclinic if its (contravariant) Dieudonné module is isoclinic. If (M, ϕ) is the Dieudonné module of D , then let

$$\ell_D := \ell_{\text{GL}_M} \in \mathbb{N} \cup \{0\}.$$

We call ℓ_D the *level torsion* of D . The following elementary lemma is our starting point for calculating and estimating n_D .

1.4.1. Lemma. *We assume that (M, ϕ) is the Dieudonné module of D . Then we have $n_D = n_{\text{GL}_M}$.*

See [Va1, Lemma 3.2.2 and Corollary 3.2.3] and [NV, Theorem 2.2 (a)] for two proofs of Lemma 1.4.1 (the second proof is not stated in the language of latticed F -isocrystals with a group). Accordingly, we call n_D the *i-number* (i.e., the isomorphism number) of D . Based on Lemma 1.4.1, we have the following corollary of Main Theorem A.

1.4.2. Basic Corollary. *For each non-trivial p -divisible group D over k we have $n_D \leq \ell_D$. If D is a direct sum of isoclinic p -divisible groups over k , then we have $n_D = \ell_D$.*

The inequality $n_D \leq \ell_D$ was first checked for the isoclinic case in [Va1, Example 3.3.5].

1.4.3. Proposition. *We assume that $D = \prod_{i \in I} D_i$ is a direct sum of at least two isoclinic p -divisible groups over k . Then we have the following basic estimate:*

$$n_D \leq \max\{1, n_{D_i}, n_{D_i} + n_{D_j} - 1 \mid i, j \in I, j \neq i\}.$$

Proposition 1.4.3 is proved in Section 4.5. Example 4.6.2 shows that in general, Proposition 1.4.3 is optimal. The next proposition (proved in Section 4.7) describes the possible range of variation of n_D and ℓ_D under isogenies.

1.4.4. Proposition. *Let $D \twoheadrightarrow \tilde{D}$ be an isogeny between non-trivial p -divisible groups over k . Let $\kappa \in \mathbb{N} \cup \{0\}$ be the smallest number such that p^κ annihilates the kernel of this isogeny. Then we have $n_D \leq \ell_D \leq \ell_{\tilde{D}} + 2\kappa$. Thus, if \tilde{D} is a direct sum of isoclinic p -divisible groups, then we have $n_D \leq \ell_D \leq n_{\tilde{D}} + 2\kappa$.*

In general, the constant 2κ of Proposition 1.4.4 is optimal (see Example 4.7.1).

1.5. Minimal and quasi-special types. Let $\mathcal{B} = \{e_1, \dots, e_r\}$ be a $W(k)$ -basis for M . Let π be an arbitrary permutation of the set $J_r := \{1, \dots, r\}$. Let (M, ϕ_π) be the Dieudonné module over k with the property that for each $s \in \{1, \dots, d\}$ we have $\phi_\pi(e_s) = pe_{\pi(s)}$ and for each $s \in \{d+1, \dots, d+c\}$ we have $\phi_\pi(e_s) = e_{\pi(s)}$. Let C_π be a p -divisible group over k whose Dieudonné module is (M, ϕ_π) . For a cycle $\pi_i = (e_{s_1}, \dots, e_{s_{r_i}})$ of π , let c_i and $d_i = r_i - c_i$ be the number of elements of the sets $\{s_1, \dots, s_{r_i}\} \cap \{d+1, \dots, d+c\}$ and $\{s_1, \dots, s_{r_i}\} \cap \{1, \dots, d\}$ (respectively), and let $\alpha_i := \frac{d_i}{r_i} \in \mathbb{Q} \cap [0, 1]$.

1.5.1. Definitions. We recall that c and d are non-negative integers such that $r := c + d > 0$, that D is a p -divisible group over k of codimension c and dimension d , and that $J_r = \{1, \dots, r\}$.

- (a) We say that D is F -cyclic (resp. F -circular), if there exists a permutation π (resp. an r -cycle permutation π) of J_r such that D is isomorphic to C_π .
- (b) We say that π is a *minimal permutation*, if the following condition holds:
 - (*) for each cycle $\pi_i = (e_{s_1}, \dots, e_{s_{r_i}})$ of π and for all $q \in \mathbb{N}$ and $u \in \{1, \dots, r_i\}$, we have $\phi_\pi^q(e_{s_u}) = p^{[q\alpha_i] + \varepsilon_q(s_u)} e_{\pi^q(s_u)}$ for some number $\varepsilon_q(s_u) \in \{0, 1\}$.
- (c) We say that D is *minimal*, if there exists a minimal permutation π of J_r such that D is isomorphic to C_π .
- (d) A non-trivial truncated Barsotti–Tate group B of level 1 over k is called *minimal*, if there exists a p -divisible group \tilde{D} over k such that $\tilde{D}[p]$ is isomorphic to B and $n_{\tilde{D}} \leq 1$.
- (e) Let $m := g.c.d.\{c, d\} \in \mathbb{N}$ and let $(d_1, r_1) := (\frac{d}{m}, \frac{r}{m})$. We say that D is *isoclinic quasi-special* (resp. *isoclinic special*), if we have $\phi^r(M) = p^d M$ (resp. we have $\phi^{r_1}(M) = p^{d_1} M$). We say that D is *quasi-special* (resp. *special*), if it is a direct sum of isoclinic quasi-special (resp. isoclinic special) p -divisible groups over k .

The terminology F -cyclic and F -circular is suggested by Definition 1.2.4 (c) in [Va2]. The terminology minimal p -divisible groups and minimal truncated Barsotti–Tate groups of level 1 is the one used in [Oo3] and [Oo4]. It is easy to check that the above definitions of minimal p -divisible groups over k and of minimal truncated Barsotti–Tate groups of level 1 over k are equivalent to the ones used in [Oo3, Section 1.1] (this also follows from Main Theorem B below). Moreover D is minimal if and only if $D[p]$ is minimal, cf. Main Theorem B below. The terminology special (see (e)) is as in [Ma, Chapter III, Section 2]. If D is F -cyclic, then it is also quasi-special but it is not necessarily special (see Lemma 4.2.4 (a) and Example 4.7.1). The class of isomorphism classes of quasi-special p -divisible groups of codimension c and dimension d over k , is a finite set (see Lemma 4.2.4 (b)); this result recovers and refines slightly [Ma, Chapter III, Section 3, Theorem 3.4].

A systematic approach to C_π 's was started in [Va2] and [Va3] using the language of Weyl groups (the role of a permutation π of J_r is that one of a representative of the Weyl group of GL_M with respect to its maximal torus that normalizes $W(k)e_s$ for all $s \in J_r$); for instance, we proved that for two permutations π_1, π_2 of J_r , the p -divisible groups C_{π_1} and C_{π_2} are isomorphic if and only if $C_{\pi_1}[p]$ and $C_{\pi_2}[p]$ are isomorphic (cf. [Va3, Theorem 1.3 (a) and Fact 4.3.1]). The p -divisible groups C_π are also studied in [Oo4] using the language of cyclic words in the variables ϕ and ϑ . We note down that in the condition (*), it suffices to consider natural numbers q which are at most equal to the order of π_i . Thus we view (b) and (d) as a more practical form of [Oo4, Section 4].

In Section 4.6 we prove the following theorem.

1.5.2. Theorem. *We assume that the non-trivial p -divisible group D is quasi-special (for instance, D is F -cyclic or special). Then D is a direct sum of isoclinic p -divisible groups over k and thus we have $n_D = \ell_D$. Moreover, we have an inequality $n_D \leq \min\{c, d\}$ i.e., the Traverso truncation conjecture holds for D .*

The proof of the inequality part of Theorem 1.5.2 relies on Proposition 1.4.3 and on an explicit formula for n_D (see property 4.6(ii); if D is F -cyclic, see also Scholium 4.6.1).

The importance of minimal p -divisible groups stems from the following theorem to be proved in Section 5.1.

1.6. Main Theorem B. *Let D be a non-trivial p -divisible group over k . Then the following three statements are equivalent:*

- (a) *we have $\ell_D \leq 1$;*
- (b) *we have $n_D \leq 1$ (equivalently, $D[p]$ is minimal);*
- (c) *the p -divisible group D over k is minimal.*

The implication (c) \Rightarrow (b) was first checked for the isoclinic case in [Va1, Example 3.3.6] and for the general case in [Oo3, Theorem 1.2]. A great part of [Oo4] is devoted to the proof of the equivalence between (b) and (c), cf. [Oo4, Theorem B].

2. Preliminaries

Let (M, ϕ) be a latticed F -isocrystal over k . In this section we include simple properties that pertain to (M, ϕ) . Let $M^* := \text{Hom}(M, W(k))$.

The notations $p, k, c, d, r = c + d, D, n_D, W(k), B(k), (M, \phi, G), \mathfrak{g}, M[\frac{1}{p}] = \bigoplus_{\alpha \in \mathbb{Q}} W(\alpha), L_+, L_0, L_-, O_+, A_0, O_0, O_-, O_G, \ell_G, \ell_D, J_r = \{1, \dots, r\}, (M, \phi_\pi)$, and C_π introduced in Section 1 will be used throughout the paper. Let D^\dagger be the p -divisible group over k which is the Cartier dual of D . For $m \in \mathbb{N}$, let $W_m(k) := W(k)/p^m W(k)$.

All finitely generated $W(k)$ -modules and all finite dimensional $B(k)$ -vector spaces are endowed with the p -adic topology. As in Section 1.2, in the whole paper we keep the following order: first $+$, next 0 , and last $-$.

2.1. Duals and homs. Let $\phi: M^*[\frac{1}{p}] \xrightarrow{\sim} M^*[\frac{1}{p}]$ be the σ -linear automorphism that takes $f \in M^*[\frac{1}{p}]$ to $\sigma \circ f \circ \phi^{-1} \in M^*[\frac{1}{p}]$. The latticed F -isocrystal (M^*, ϕ) is called the dual of (M, ϕ) , cf. [Va1, Section 2.1]. The canonical identification $\text{End}(M) = M \otimes_{W(k)} M^*$ defines an identification $(\text{End}(M), \phi) = (M, \phi) \otimes (M^*, \phi)$ of latticed F -isocrystals over k . If (M, ϕ) is the Dieudonné module of D , then $(M^*, p\phi)$ is the Dieudonné module of D^\dagger . Let (M_1, ϕ_1) and (M_2, ϕ_2) be two latticed F -isocrystals

over k . Let $\phi_{12}: \operatorname{Hom}(M_1, M_2)[\frac{1}{p}] \xrightarrow{\sim} \operatorname{Hom}(M_1, M_2)[\frac{1}{p}]$ be the σ -linear automorphism that takes $f \in \operatorname{Hom}(M_1[\frac{1}{p}], M_2[\frac{1}{p}])$ to $\phi_2 \circ f \circ \phi_1^{-1} \in \operatorname{Hom}(M_1[\frac{1}{p}], M_2[\frac{1}{p}])$. The latticed F -isocrystal $(\operatorname{Hom}(M_1, M_2), \phi_{12})$ over k is called the hom of (M_1, ϕ_1) and (M_2, ϕ_2) . Thus (M^*, ϕ) is the hom of (M, ϕ) and $(W(k), \sigma)$. The dual of $(\operatorname{Hom}(M_1, M_2), \phi_{12})$ is $(\operatorname{Hom}(M_2, M_1), \phi_{21})$ (here ϕ_{21} is defined similarly to ϕ_{12}). Thus the dual of $(\operatorname{End}(M), \phi)$ is $(\operatorname{End}(M), \phi)$ itself.

If \mathcal{B} is a $W(k)$ -basis for M , let $\mathcal{B}^* := \{x^* \mid x \in \mathcal{B}\}$ be the dual $W(k)$ -basis for M^* . Thus for $x, y \in \mathcal{B}$, we have $x^*(y) = \delta_{xy}$. For $q \in \mathbb{Z}$ and $x, y \in \mathcal{B}$, let $a_q(x, y) \in B(k)$ be such that we have $\phi^q(x) = \sum_{y \in \mathcal{B}} a_q(x, y)y$. We have $\phi^q(x^*) = \sum_{y \in \mathcal{B}} \sigma^q(a_{-q}(y, x))y^*$ and hence $\phi^{-q}(x^*) = \sum_{y \in \mathcal{B}} \sigma^{-q}(a_q(y, x))y^*$. This implies:

(*) If $s \in \mathbb{Z}$, then we have $p^s \phi^q(M) \subseteq M$ (i.e., $p^s a_q(x, y) \in W(k)$ for all $x, y \in \mathcal{B}$) if and only if we have $p^s \phi^{-q}(M^*) \subseteq M^*$ (i.e., $p^s a_q(y, x) \in W(k)$ for all $x, y \in \mathcal{B}$).

The set $\{x \otimes y^* \mid x, y \in \mathcal{B}\}$ is a $W(k)$ -basis for $\operatorname{End}(M) = M \otimes_{W(k)} M^*$.

2.2. Example. We assume that we have a direct sum decomposition $M = W(k) \oplus N$ such that ϕ acts on $W(k)$ as σ does, we have $\phi(N) \subseteq N$, and (N, ϕ) is isoclinic of Newton polygon slope $\gamma \in (\mathbb{Q} \cap (0, \infty)) \setminus \mathbb{Z}$. We have a direct sum decomposition of latticed F -isocrystals over k

$$(\operatorname{End}(M), \phi) = (\operatorname{End}(N), \phi) \oplus (N, \phi) \oplus (N^*, \phi) \oplus (W(k), \sigma).$$

The $W(k)$ -span of the product N^*N (taken inside $\operatorname{End}(M)$) is $\operatorname{End}(N)$. As $\phi(N) \subseteq N$ and $N^* \subseteq \phi^{-1}(N^*)$, we have $N \subseteq O_+$ and $N^* \subseteq O_-$. As $\gamma \notin \mathbb{Z}$, we have $O \cap \operatorname{End}(N) \subsetneq \operatorname{End}(N)$. Thus $O_+ O_- \not\subseteq O$. Therefore O is not a $W(k)$ -subalgebra of $\operatorname{End}(M)$.

We take G such that \mathfrak{g} is the $W(k)$ -subalgebra $N \oplus W(k)1_N$ of $\operatorname{End}(M)$. As $N \subseteq O_+$ and $1_M \in O_0$, we have $\mathfrak{g} = O_G = (\mathfrak{g} \cap O_+) \oplus (\mathfrak{g} \cap O_0)$ and $(\mathfrak{g} \cap O_+)$ is a nilpotent, two-sided ideal of the $W(k)$ -algebra \mathfrak{g} . Thus $\ell_G = 0$, cf. the rule 1.2 (a). If the pair (N, ϕ) is not a Dieudonné–Fontaine p -divisible object over k , then the Dieudonné–Fontaine torsion m_G of (\mathfrak{g}, ϕ) is positive (and in fact it can be any natural number).

2.3. Lemma. We assume that $n_{\operatorname{GL}_M} = 0$. Then there exists an integer s such that we have $\phi(M) = p^s M$. Thus $\phi(\operatorname{End}(M)) = \operatorname{End}(\phi(M)) = \operatorname{End}(M)$ and therefore we have $O_0 = \operatorname{End}(M)$ and $\ell_{\operatorname{GL}_M} = 0$.

Proof. Let $q \in \mathbb{N}$. By induction on q we show that the Lemma holds if the rank r of M is at most q . If $q = 1$ and $r = 1$, then the Lemma is obvious. The passage from q to $q + 1$ goes as follows. We can assume that $r = q + 1$. By multiplying ϕ with p^{-s} for some $s \in \mathbb{Z}$, we can assume that $\phi(M)$ is a $W(k)$ -submodule of

M that contains a direct summand of M of rank at least 1. Let $\tilde{x} \in M \setminus pM$ be such that $\phi(\tilde{x}) \in M \setminus pM$. Let $g_{\tilde{x}} \in \mathrm{GL}_M(W(k))$ be such that $g_{\tilde{x}}\phi(\tilde{x}) = \tilde{x}$. As $n_{\mathrm{GL}_M} = 0$, (M, ϕ) is isomorphic to $(M, g_{\tilde{x}}\phi)$. Thus there exists $x \in M \setminus pM$ such that $\phi(x) = x$. Let M_0 be the $W(k)$ -submodule of M generated by elements fixed by ϕ ; it is a direct summand of M which contains x .

If $M_0 = M$, then we are done as $\phi(M) = M$. Thus to end the proof it suffices to show that the assumption that $M_0 \neq M$ leads to a contradiction. Let $M_1 := M/M_0$ and let $\phi_1: M_1 \rightarrow M_1$ be the σ -linear endomorphism induced by ϕ . For each element $g_1 \in \mathrm{GL}_{M_1}(W(k))$ there exists an element $g \in \mathrm{GL}_M(W(k))$ that fixes M_0 and that maps naturally to g_1 . As $(M, g\phi)$ and (M, ϕ) are isomorphic and due to the definition of M_0 , we easily get that (M_1, ϕ_1) and $(M_1, g_1\phi_1)$ are isomorphic. Thus the i -number of $(M_1, \phi_1, \mathrm{GL}_{M_1})$ is 0. As the rank of M_1 is less than $q + 1$, by induction we get that there exists a natural number s_1 such that $\phi(M_1) = p^{s_1}M_1$ (we have $s_1 \neq 0$, due to the definition of M_0). Let $z_1 \in M_1 \setminus pM_1$ be such that $\phi_1(z_1) = p^{s_1}z_1$. Let $\tilde{z} \in M$ be such that it maps naturally to z_1 . We have $\phi(\tilde{z}) - p^{s_1}\tilde{z} \in M_0$. Let $\tilde{y} \in M_0$ be such that $\phi(\tilde{y}) - p^{s_1}\tilde{y} = -\phi(\tilde{z}) + p^{s_1}\tilde{z}$. If $z := \tilde{z} + \tilde{y}$, then we have $\phi(z) = p^{s_1}z$. As z maps naturally to $z_1 \in M_1 \setminus pM_1$, the $W(k)$ -module $M_0 \oplus W(k)z$ is a direct summand of M . Let $g_{xz} \in \mathrm{GL}_M(W(k))$ be such that it permutes x and z , it normalizes $M_0 \oplus W(k)z$, and it acts identically on $(M_0 \oplus W(k)z)/(W(k)x \oplus W(k)z)$ and on $M/(M_0 \oplus W(k)z)$. The Newton polygon slopes of $(M, g_{xz}\phi)$ are 0, $\frac{s_1}{2}$, and s_1 . As the Newton polygon slopes of (M, ϕ) are 0 and s_1 and as $s_1 \in \mathbb{N}$, we get that (M, ϕ) and $(M, g_{xz}\phi)$ are not isomorphic. This contradicts the equality $n_{\mathrm{GL}_M} = 0$. \square

2.4. Lemma. *Let $x \in \mathrm{End}(M)$ be such that for all $q \in \mathbb{N}$ (resp. for all $q \in -\mathbb{N}$) we have $\phi^q(x) \in \mathrm{End}(M)$. Then we have $x \in O_+ \oplus O_0$ (resp. we have $x \in O_0 \oplus O_-$).*

Proof. We will prove only the non-negative part of the Lemma as the non-positive part of it is proved in the same way. Thus we assume that we have $\phi^q(x) \in \mathrm{End}(M)$ for all $q \in \mathbb{N}$. We write $x = x_+ + x_0 + x_-$, where $x_+ \in L_+$, $x_0 \in L_0$, and $x_- \in L_-$. There exists a number $s \in \mathbb{N}$ such that $p^s x_+ \in O_+$ and $p^s x_0 \in O_0$. Thus $\phi^q(p^s x_+) \in O_+ \subseteq \mathrm{End}(M)$ and $\phi^q(p^s x_0) \in O_0 \subseteq \mathrm{End}(M)$. We easily get that we have $p^s \phi^q(x_-) \in \mathrm{End}(M)$ for all $q \in \mathbb{N}$. This implies that $x_- = 0$ (as all Newton polygon slopes of (L_-, ϕ) are negative). Thus $x = x_+ + x_0$. The sequence $(\phi^q(x_+))_{q \in \mathbb{N}}$ converges to 0 (as all Newton polygon slopes of (L_+, ϕ) are positive). Thus there exists $\tilde{q} \in \mathbb{N}$ such that $y_+ := \phi^{\tilde{q}}(x_+) \in O_+$. Let $y := \phi^{\tilde{q}}(x)$ and $y_0 := y - y_+ = \phi^{\tilde{q}}(x_0) \in \mathrm{End}(M) \cap L_0$. As for each $q \in \mathbb{N}$ we have $\phi^q(y_+) \in O_+ \subseteq \mathrm{End}(M)$ and $\phi^q(y) \in \mathrm{End}(M)$, we also have $\phi^q(y_0) \in \mathrm{End}(M)$. Thus $y_0 \in O_0$. Therefore $x_0 = \phi^{-\tilde{q}}(y_0) \in \phi^{-\tilde{q}}(O_0) = O_0$. This implies that for all $q \in \mathbb{N} \cup \{0\}$ we have $\phi^q(x_0) \in O_0 \subseteq \mathrm{End}(M)$. Thus for all $q \in \mathbb{N} \cup \{0\}$ we have $\phi^q(x_+) = \phi^q(x) - \phi^q(x_0) \in \mathrm{End}(M)$ i.e., $x_+ \in O_+$. Therefore $x = x_+ + x_0 \in O_+ \oplus O_0$. \square

2.5. Invertible elements. In this section we recall basic properties of invertible elements of $W(k)$ -subalgebras of $\text{End}(M)$. Let \mathfrak{h} be a $W(k)$ -subalgebra of $\text{End}(M)$.

- (i) If $x \in \mathfrak{h}$ has an inverse x^{-1} in $\text{End}(M)$ (i.e., if $x \in \mathfrak{h} \cap \text{GL}_M(W(k))$), then the determinant of x is an invertible element of $W(k)$ and therefore from the Cayley–Hamilton theorem we get that x^{-1} is a polynomial in x with coefficients in $W(k)$; thus $x^{-1} \in \mathfrak{h}$ (i.e., x is an invertible element of \mathfrak{h}).
- (ii) Each invertible element of \mathfrak{h} is also an invertible element of any other $W(k)$ -subalgebra of $\text{End}(M)$ that contains \mathfrak{h} .
- (iii) If we have a direct sum decomposition $\mathfrak{h} = \mathfrak{n} \oplus \mathfrak{h}_0$ such that \mathfrak{h}_0 is a $W(k)$ -subalgebra of \mathfrak{h} and \mathfrak{n} is a nilpotent, two-sided ideal of the $W(k)$ -algebra \mathfrak{h} , then we have a short exact sequence $1 \rightarrow 1_M + \mathfrak{n} \rightarrow \mathfrak{h} \cap \text{GL}_M(W(k)) \rightarrow \mathfrak{h}_0 \cap \text{GL}_M(W(k)) \rightarrow 1$ which splits and which is defined by the following rule: if $x \in \mathfrak{n}$ and $y \in \mathfrak{h}_0$ are such that $x + y \in \mathfrak{h} \cap \text{GL}_M(W(k))$, then the image of $x + y$ in $\mathfrak{h}_0 \cap \text{GL}_M(W(k))$ is y .
- (iv) We recall that a two-sided ideal \mathfrak{i} of the $W(k)$ -algebra \mathfrak{h} is called *topologically nilpotent* if for all $m \in \mathbb{N}$ there exists $\tilde{m} \in \mathbb{N}$ such that we have an inclusion $\mathfrak{i}^{\tilde{m}} \subseteq p^m \mathfrak{h}$ (this implies that $\bigcap_{m \in \mathbb{N}} \mathfrak{i}^m = 0$). If $x \in \mathfrak{i}$, then the element $1_M + \sum_{m=1}^{\infty} (-x)^m \in 1_M + \mathfrak{i}$ is well defined and is the inverse of $1_M + x$. This implies that an element of \mathfrak{h} is invertible if and only if its image in $\mathfrak{h}/\mathfrak{i}$ is an invertible element of $\mathfrak{h}/\mathfrak{i}$.

3. The proof of Main Theorem A

In this section we prove Main Theorem A (see Sections 3.4 and 3.5). We begin by introducing certain $W(k)$ -algebras and group schemes over $\text{Spec}(W(k))$ and by presenting basic properties of them (see Sections 3.1 and 3.2). In Section 3.3 we list simple properties of isomorphism classes of certain latticed F -isocrystals over k . All these properties play a key role in Section 3.4. In Sections 3.6 and 3.7 we include two remarks as well as a more general variant of Theorem 1.3 (b).

3.1. Group schemes of invertible elements. Let $\mathfrak{h}_+ := \mathfrak{g} \cap \mathcal{O}_+$ and $\mathfrak{h}_0 := \mathfrak{g} \cap \mathcal{O}_0$. Let $\mathfrak{h} := (\mathfrak{h}_+ \oplus \mathfrak{h}_0) + p^{\ell_G} \mathfrak{g}$. As \mathcal{O}_0 and $\mathcal{O}_+ \oplus \mathcal{O}_0$ are $W(k)$ -algebras and as $p^{\ell_G} \mathfrak{g}$ is a two-sided ideal of the $W(k)$ -algebra \mathfrak{g} , it is easy to see that \mathfrak{h} is a $W(k)$ -subalgebra of \mathfrak{g} . Let $\mathfrak{h}_- := \mathfrak{h} \cap \mathcal{O}_-$. As $p^{\ell_G} \mathfrak{g} \subseteq \mathcal{O}_G$ (see (1)), we have a direct sum decomposition

$$\mathfrak{h} = (\mathfrak{h}_+ \oplus \mathfrak{h}_0) + p^{\ell_G} \mathfrak{g} = \mathfrak{h}_+ \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-. \quad (2)$$

Let $\Pi_+ : \mathfrak{h} \rightarrow \mathfrak{h}$ be the projection on \mathfrak{h}_+ along $\mathfrak{h}_0 \oplus \mathfrak{h}_-$.

Let H , H_{+0} , and H_0 be the affine group scheme over $\text{Spec}(W(k))$ of invertible elements of \mathfrak{h} , $\mathfrak{h}_+ \oplus \mathfrak{h}_0$, and \mathfrak{h}_0 (respectively). Due to Section 2.5 (ii), we have a sequence

$$H_0(W(k)) \leq H_{+0}(W(k)) \leq H(W(k)) \leq G(W(k)) \quad (3a)$$

of subgroups. As \mathfrak{h}_+ and \mathfrak{h}_- are nilpotent subalgebras (without unit) of \mathfrak{h} , we have

$$1_M + \mathfrak{h}_+ \leq H_{+0}(W(k)) \quad \text{and} \quad 1_M + \mathfrak{h}_- \leq H(W(k)). \quad (3b)$$

From Section 2.5 (iii) we get that we have a natural split short exact sequence

$$1 \rightarrow 1_M + \mathfrak{h}_+ \rightarrow H_{+0}(W(k)) \rightarrow H_0(W(k)) \rightarrow 1. \quad (3c)$$

Based on (2), for each element $h \in H(W(k))$ we can write uniquely

$$h = 1_M + a(h) + b(h) + c(h),$$

where $a(h) \in \mathfrak{h}_+$, $b(h) \in \mathfrak{h}_0$, and $c(h) \in \mathfrak{h}_-$. We have $a(h) = \Pi_+(h)$.

3.1.1. The ideal \mathfrak{i} . If $\ell_G = 0$, let \mathfrak{i} be the two-sided ideal of the $W(k)$ -algebra $\mathfrak{h} = \mathfrak{g}$ generated by \mathfrak{h}_+ and \mathfrak{h}_- . If $\ell_G \geq 1$, let $\mathfrak{i} := \mathfrak{h}_+ + \mathfrak{h}_- + p^{\ell_G} \mathfrak{g}$. We check that \mathfrak{i} is a topologically nilpotent, two-sided ideal of \mathfrak{h} . If $\ell_G = 0$, this is so by the very definitions (see rules 1.2 (a) and (b)). We assume that $\ell_G \geq 1$. This implies that $p^{\ell_G} \mathfrak{g}$ is a topologically nilpotent, two-sided ideal of \mathfrak{h} . As \mathfrak{h}_+ is a nilpotent, two-sided ideal of $\mathfrak{h}_+ \oplus \mathfrak{h}_0$, its image in $\mathfrak{h}/p^{\ell_G} \mathfrak{g} = \mathfrak{h}_+ \oplus \mathfrak{h}_0/[(\mathfrak{h}_+ \oplus \mathfrak{h}_0) \cap p^{\ell_G} \mathfrak{g}]$ is a nilpotent, two-sided ideal. Thus $p^{\ell_G} \mathfrak{g} + \mathfrak{h}_+$ is a topologically nilpotent, two-sided ideal of \mathfrak{h} . As \mathfrak{h}_- is a nilpotent, two-sided ideal of $\mathfrak{h}_0 \oplus \mathfrak{h}_-$ and as $\mathfrak{h}_0 \oplus \mathfrak{h}_-$ surjects onto $\mathfrak{h}/(p^{\ell_G} \mathfrak{g} + \mathfrak{h}_+) = \mathfrak{h}_0/[\mathfrak{h}_0 \cap (p^{\ell_G} \mathfrak{g} + \mathfrak{h}_+)]$, the image of \mathfrak{h}_- in $\mathfrak{h}/(p^{\ell_G} \mathfrak{g} + \mathfrak{h}_+)$ is a nilpotent, two-sided ideal. From the last two sentences, we get that $\mathfrak{i} = p^{\ell_G} \mathfrak{g} + \mathfrak{h}_+ + \mathfrak{h}_-$ is a topologically nilpotent, two-sided ideal of \mathfrak{h} .

3.1.2. Fact. For each element $h = 1_M + a(h) + b(h) + c(h) \in H(W(k))$, we have $1_M + b(h) \in H_0(W(k))$. Therefore also $1_M + a(h) + b(h) \in H_{+0}(W(k))$.

Proof. As $1_M + b(h)$ and h are congruent modulo \mathfrak{i} , the first part of the fact follows from Section 2.5 (iv). The last part of the fact follows from (3c). \square

3.1.3. On H_0 . As $\mathfrak{h}_0 = \mathfrak{g} \cap \mathcal{O}_0$, we have $\phi(\mathfrak{h}_0) = \mathfrak{h}_0$ (see Example 1.2.2). Let $\mathfrak{h}_{0\mathbb{Z}_p}$ be the \mathbb{Z}_p -subalgebra of \mathfrak{h}_0 formed by elements fixed by ϕ . Let $H_{0\mathbb{Z}_p}$ be the affine group scheme over $\text{Spec}(\mathbb{Z}_p)$ of invertible elements of $\mathfrak{h}_{0\mathbb{Z}_p}$. The group scheme $H_{0\mathbb{Z}_p}$ is a \mathbb{Z}_p -structure of H_0 and thus the Frobenius automorphism σ acts naturally on $H_0(W(k)) = H_{0\mathbb{Z}_p}(W(k))$: for $*$ in $H_0(W(k))$ we have $\sigma(*) = \phi(*)$. The scheme $H_{0\mathbb{Z}_p}$ is an open subscheme of the vector group scheme over $\text{Spec}(\mathbb{Z}_p)$ defined by $\mathfrak{h}_{0\mathbb{Z}_p}$ (viewed only as a \mathbb{Z}_p -module). Thus the affine, smooth group scheme $H_{0\mathbb{Z}_p}$ has connected fibres.

3.1.4. Lemma. Let \mathfrak{f}_+ be a $W(k)$ -submodule of \mathfrak{h}_+ . Let $\mathfrak{j} := \mathfrak{f}_+ \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$. We consider the following three conditions:

- (i) we have $\Pi_+(\mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+) \subseteq \mathfrak{f}_+$;
- (ii) the $W(k)$ -module \mathfrak{f}_+ is a left and right \mathfrak{h}_0 -module;
- (iii) we have $\mathfrak{f}_+^2 \subseteq \mathfrak{f}_+$ (i.e., \mathfrak{f}_+ is an algebra).

Then the following three properties hold:

- (a) Conditions (i) and (ii) hold if and only if \mathfrak{j} is a left and right $\mathfrak{h}_0 \oplus \mathfrak{h}_-$ -module.
- (b) Conditions (ii) and (iii) hold if and only if $\mathfrak{f}_+ \oplus \mathfrak{h}_0$ is a $W(k)$ -subalgebra of $\mathfrak{h}_+ \oplus \mathfrak{h}_0$.
- (c) The three conditions (i) to (iii) hold if and only if \mathfrak{j} is a $W(k)$ -subalgebra of \mathfrak{h} .

Proof. As $\mathfrak{h}_0 \oplus \mathfrak{h}_-$ is a $W(k)$ -subalgebra of \mathfrak{h} , \mathfrak{j} is a left and right $\mathfrak{h}_0 \oplus \mathfrak{h}_-$ -module if and only if we have $\mathfrak{f}_+\mathfrak{h}_0 + \mathfrak{h}_0\mathfrak{f}_+ + \mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+ \subseteq \mathfrak{j}$. We have $\mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+ \subseteq \mathfrak{j}$ if and only if (i) holds. As \mathfrak{h}_+ is a left and right \mathfrak{h}_0 -module, we have $\mathfrak{f}_+\mathfrak{h}_0 + \mathfrak{h}_0\mathfrak{f}_+ \subseteq \mathfrak{j}$ if and only if $\mathfrak{f}_+\mathfrak{h}_0 + \mathfrak{h}_0\mathfrak{f}_+ \subseteq \mathfrak{f}_+$ and thus if and only if (ii) holds. Part (a) follows from the last three sentences. As \mathfrak{h}_+ is an algebra and a left and right \mathfrak{h}_0 -module, we have $(\mathfrak{f}_+ \oplus \mathfrak{h}_0)^2 \subseteq \mathfrak{f}_+ \oplus \mathfrak{h}_0$ if and only if $\mathfrak{f}_+^2 + \mathfrak{f}_+\mathfrak{h}_0 + \mathfrak{h}_0\mathfrak{f}_+ \subseteq \mathfrak{f}_+$ and thus if and only if conditions (ii) and (iii) hold. Thus (b) holds. Part (c) follows from (a) and (b). \square

3.2. Subalgebras. In this section we list several subalgebras of \mathfrak{h} .

3.2.1. Frobenius filtration of \mathfrak{h}_+ . For $i \in \mathbb{N} \cup \{0\}$ let

$$\mathfrak{h}_{+,i} := \mathfrak{h}_+ \cap \phi^i(\mathfrak{g}) = \mathfrak{h}_+ \cap \phi^i(\mathfrak{g} \cap L_+).$$

We have $\mathfrak{h}_{+,0} = \mathfrak{h}_+$, $\phi(\mathfrak{h}_{+,i}) \subseteq \mathfrak{h}_{+,i+1} \subseteq \mathfrak{h}_{+,i}$, and each $\mathfrak{h}_{+,i}$ is a $W(k)$ -module and a nilpotent algebra. As $\phi^i(\mathfrak{h}_0) = \mathfrak{h}_0$ (see Section 3.1.3) and as \mathfrak{h}_+ is a left and right \mathfrak{h}_0 -module, $\mathfrak{h}_{+,i}$ is also a left and right \mathfrak{h}_0 -module. As all Newton polygon slopes of (\mathfrak{h}_+, ϕ) are positive, we have $\bigcap_{i=0}^{\infty} \mathfrak{h}_{+,i} = 0$. Thus $(\mathfrak{h}_{+,i})_{i \in \mathbb{N} \cup \{0\}}$ is a decreasing, separated, and exhaustive filtration of \mathfrak{h}_+ to be called the Frobenius filtration.

3.2.2. The Theta operations. We assume that $\mathfrak{h}_+ \neq 0$. Let $\mathcal{M}(\mathfrak{h}_+)$ be the set of $W(k)$ -submodules of \mathfrak{h}_+ endowed with the pre-order relation defined by inclusions. We consider the increasing operators $\Theta, \Theta_a, \Theta_s: \mathcal{M}(\mathfrak{h}_+) \rightarrow \mathcal{M}(\mathfrak{h}_+)$ that take $\mathfrak{f}_+ \in \mathcal{M}(\mathfrak{h}_+)$ to

$$\Theta(\mathfrak{f}_+) := \mathfrak{f}_+^2 + \Pi_+(\mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+) + \phi(\mathfrak{f}_+) \in \mathcal{M}(\mathfrak{h}_+),$$

$\Theta_a(\mathfrak{f}_+) := \mathfrak{f}_+^2 + \Pi_+(\mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+)$, and $\Theta_s(\mathfrak{f}_+) := \Pi_+(\mathfrak{f}_+\mathfrak{h}_- + \mathfrak{h}_-\mathfrak{f}_+)$. We have identities $\Theta(\mathfrak{f}_+) = \Theta_a(\mathfrak{f}_+) + \phi(\mathfrak{f}_+)$ and $\Theta_a(\mathfrak{f}_+) = \mathfrak{f}_+^2 + \Theta_s(\mathfrak{f}_+)$. The lower right indices a and s stand for algebraic and slope module (respectively), as suggested by Lemma 3.1.4 (a) and (c). For $i \in \mathbb{N} \cup \{0\}$ let

$$\mathfrak{f}_i := \Theta^i(\mathfrak{h}_+).$$

As Θ is increasing and as $\Theta(\mathfrak{h}_+) \subseteq \mathfrak{h}_+$, we have $\mathfrak{f}_{i+1} = \Theta(\mathfrak{f}_i) \subseteq \mathfrak{f}_i \subseteq \mathfrak{h}_+$.

3.2.3. Lemma. *We assume that $\mathfrak{h}_+ \neq 0$. Let $i \in \mathbb{N} \cup \{0\}$. Then $e_i := \mathfrak{f}_i \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$ is a $W(k)$ -subalgebra of \mathfrak{h} .*

Proof. We use induction on i . For $i = 0$ we have $e_0 = \mathfrak{h}$ and thus the Lemma holds. The passage from i to $i + 1$ goes as follows. We check that the three conditions (i) to (iii) of Lemma 3.1.4 hold for $\mathfrak{f}_+ := \mathfrak{f}_{i+1}$. As $\mathfrak{f}_{i+1} \subseteq \mathfrak{f}_i$, we have $\Theta_s(\mathfrak{f}_{i+1}) \subseteq \Theta_s(\mathfrak{f}_i) \subseteq \Theta(\mathfrak{f}_i) = \mathfrak{f}_{i+1}$. Thus condition 3.1.4 (i) holds. To check that condition 3.1.4 (ii) holds, it suffices to show that each one of the following four elements \mathfrak{f}_i^2 , $\Pi_+(\mathfrak{f}_i \mathfrak{h}_-)$, $\Pi_+(\mathfrak{h}_- \mathfrak{f}_i)$, and $\phi(\mathfrak{f}_i)$ of $\mathcal{M}(\mathfrak{h}_+)$ are left and right \mathfrak{h}_0 -modules; we will only check that they are left \mathfrak{h}_0 -modules as the arguments for checking that they are right \mathfrak{h}_0 -modules are entirely the same. As \mathfrak{f}_i is a left \mathfrak{h}_0 -module, \mathfrak{f}_i^2 is also a left \mathfrak{h}_0 -module. We have $\mathfrak{h}_0 \Pi_+(\mathfrak{f}_i \mathfrak{h}_-) = \Pi_+(\mathfrak{h}_0 \mathfrak{f}_i \mathfrak{h}_-) = \Pi_+(\mathfrak{f}_i \mathfrak{h}_-)$ (the last equality as \mathfrak{f}_i is a left \mathfrak{h}_0 -module). We have $\mathfrak{h}_0 \Pi_+(\mathfrak{h}_- \mathfrak{f}_i) = \Pi_+(\mathfrak{h}_0 \mathfrak{h}_- \mathfrak{f}_i) = \Pi_+(\mathfrak{h}_- \mathfrak{f}_i)$ (the last equality as \mathfrak{h}_- is a left \mathfrak{h}_0 -module). We have $\mathfrak{h}_0 \phi(\mathfrak{f}_i) = \phi(\mathfrak{h}_0) \phi(\mathfrak{f}_i) = \phi(\mathfrak{h}_0 \mathfrak{f}_i) = \phi(\mathfrak{f}_i)$. Thus condition 3.1.4 (ii) holds. As $\mathfrak{f}_{i+1}^2 \subseteq \mathfrak{f}_i^2 \subseteq \mathfrak{f}_{i+1}$, condition 3.1.4 (iii) also holds. Thus e_{i+1} is a $W(k)$ -subalgebra of \mathfrak{h} , cf. Lemma 3.1.4 (c). This ends the induction. \square

3.2.4. Lemma. *We assume that $\mathfrak{h}_+ \neq 0$. Let $i \in \mathbb{N} \cup \{0\}$. Then $\Theta_s(\mathfrak{h}_{+,i}) \subseteq \mathfrak{h}_{+,i}$. Thus $\mathfrak{h}_{+,i} \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$ is a $W(k)$ -subalgebra of \mathfrak{h} .*

Proof. Let $x \in \mathfrak{h}_{+,i}$ and $y \in \mathfrak{h}_-$. As $z := -xy + \Pi_+(xy) \in \mathfrak{h}_0 \oplus \mathfrak{h}_-$, we have $\tilde{z} := \phi^{-i}(z) \in \mathfrak{g} \cap (O_0 \oplus O_-)$. As $x \in \mathfrak{h}_{+,i}$ and $y \in \mathfrak{h}_- \subseteq \mathfrak{g} \cap O_-$, we have $\tilde{x} := \phi^{-i}(x) \in \mathfrak{g} \cap L_+$ and $\tilde{y} := \phi^{-i}(y) \in \mathfrak{g} \cap L_-$. Thus $\Pi_+(xy) = z + xy = \phi^i(\tilde{z} + \tilde{x}\tilde{y}) \in \phi^i(\mathfrak{g})$ i.e., $\Pi_+(xy) \in \mathfrak{h}_+ \cap \phi^i(\mathfrak{g}) = \mathfrak{h}_{+,i}$. A similar argument shows that $\Pi_+(yx) \in \mathfrak{h}_{+,i}$. Thus $\Theta_s(\mathfrak{h}_{+,i}) \subseteq \mathfrak{h}_{+,i}$. As $\mathfrak{h}_{+,i}$ is an algebra and a left and right \mathfrak{h}_0 -module (see Section 3.2.1), from Lemma 3.1.4 (c) we get that $\mathfrak{h}_{+,i} \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$ is a $W(k)$ -subalgebra of \mathfrak{h} . \square

3.2.5. Lemma. *We assume that $\mathfrak{h}_+ \neq 0$. Then $\mathfrak{f}_{a,\infty} := \bigcap_{i=0}^{\infty} \Theta_a^i(\mathfrak{h}_+)$ is 0.*

Proof. Let i be as in Section 3.1.1. Let \mathfrak{n}_0 be the topologically nilpotent, two-sided ideal of the $W(k)$ -algebra \mathfrak{h}_0 such that we have $i = \mathfrak{h}_+ \oplus \mathfrak{n}_0 \oplus \mathfrak{h}_-$. We will check by induction on $q \in \mathbb{N}$ that $\mathfrak{f}_{a,\infty} \subseteq i^q + \mathfrak{n}_0 + \mathfrak{h}_-$. As $\mathfrak{f}_{a,\infty} \subseteq \mathfrak{h}_+ \subseteq i$, the basis of the induction holds. The passage from q to $q + 1$ goes as follows. Let $i \in \mathbb{N}$ be such that $\Theta_a^i(\mathfrak{h}_+) \subseteq i^q + \mathfrak{n}_0 + \mathfrak{h}_- \subseteq i$. We have

$$\begin{aligned} \Theta_a^{i+1}(\mathfrak{h}_+) &= (\Theta_a^i(\mathfrak{h}_+))^2 + \Theta_s(\Theta_a^i(\mathfrak{h}_+)) \subseteq (i^q + \mathfrak{n}_0 + \mathfrak{h}_-)^2 + \Theta_s(\mathfrak{h}_+ \cap (i^q + \mathfrak{n}_0 + \mathfrak{h}_-)) \\ &\subseteq i^{q+1} + \mathfrak{n}_0 + \mathfrak{h}_- + \Theta_s(\mathfrak{h}_+ \cap (i^q + \mathfrak{n}_0 + \mathfrak{h}_-)). \end{aligned}$$

Let $x \in \mathfrak{h}_+ \cap (i^q + \mathfrak{n}_0 + \mathfrak{h}_-) \subseteq i$ and $y \in \mathfrak{h}_- \subseteq i$. We have $\Pi_+(xy) - xy \in \mathfrak{n}_0 \oplus \mathfrak{h}_-$ and $xy \in i^{q+1} + (\mathfrak{n}_0 + \mathfrak{h}_-)\mathfrak{h}_- \subseteq i^{q+1} + \mathfrak{n}_0 + \mathfrak{h}_-$. Thus $\Pi_+(xy) = [\Pi_+(xy) - xy] +$

$xy \in i^{q+1} + n_0 + \mathfrak{h}_-$. A similar argument shows that $\Pi_+(yx) \in i^{q+1} + n_0 + \mathfrak{h}_-$. From the last two sentences we get that $\Theta_s(\mathfrak{h}_+ \cap (i^q + n_0 + \mathfrak{h}_-)) \subseteq i^{q+1} + n_0 + \mathfrak{h}_-$. We conclude that $\Theta_a^{i+1}(\mathfrak{h}_+) \subseteq i^{q+1} + n_0 + \mathfrak{h}_-$. This implies that $\mathfrak{f}_{a,\infty} \subseteq i^{q+1} + n_0 + \mathfrak{h}_-$. This ends the induction.

As i is topologically nilpotent, we have $\bigcap_{q \in \mathbb{N}} (i^q + n_0 + \mathfrak{h}_-) \subseteq \bigcap_{q \in \mathbb{N}} (p^q \mathfrak{h} + n_0 + \mathfrak{h}_-) = n_0 + \mathfrak{h}_-$. This implies that $\mathfrak{f}_{a,\infty} \subseteq n_0 + \mathfrak{h}_-$. Thus $\mathfrak{f}_{a,\infty} \subseteq \mathfrak{h}_+ \cap (n_0 + \mathfrak{h}_-) = 0$ i.e., $\mathfrak{f}_{a,\infty} = 0$. \square

3.2.6. Lemma. *We assume that $\mathfrak{h}_+ \neq 0$. Then $\mathfrak{f}_\infty := \bigcap_{i=0}^\infty \mathfrak{f}_i$ is 0.*

Proof. We show that the assumption that $\mathfrak{f}_\infty \neq 0$ leads to a contradiction. As we have inclusions $0 \subsetneq \mathfrak{f}_\infty \subseteq \mathfrak{f}_i \subseteq \mathfrak{h}_+ = \mathfrak{h}_{+,0}$ and as $\bigcap_{i=0}^\infty \mathfrak{h}_{+,i} = 0$, there exists a greatest number $i_0 \in \mathbb{N} \cup \{0\}$ for which there exists $i \in \mathbb{N}$ such that we have inclusions $\mathfrak{f}_\infty \subseteq \mathfrak{f}_i \subseteq \mathfrak{h}_{+,i_0}$.

As $\Theta_s(\mathfrak{h}_{+,i_0+1}) \subseteq \mathfrak{h}_{+,i_0+1}$ (cf. Lemma 3.2.4) and as $\mathfrak{h}_{+,i_0+1}^2 + \phi(\mathfrak{h}_{+,i_0+1}) \subseteq \mathfrak{h}_{+,i_0+1}$ (cf. Section 3.2.1), we have $\Theta(\mathfrak{h}_{+,i_0+1}) \subseteq \mathfrak{h}_{+,i_0+1}$. Based on this and the inclusion $\phi(\mathfrak{h}_{+,i_0}) \subseteq \mathfrak{h}_{+,i_0+1}$, an easy induction on $j \in \mathbb{N}$ shows that the images of $\mathfrak{f}_{i+j} = \Theta^j(\mathfrak{f}_i)$ and $\Theta_a^j(\mathfrak{f}_i)$ in $\mathfrak{h}_{+,i_0}/\mathfrak{h}_{+,i_0+1}$ coincide. Let $j_0 \in \mathbb{N}$ be such that we have $\Theta_a^{j_0}(\mathfrak{h}_+) \subseteq \mathfrak{h}_{+,i_0+1}$, cf. Lemma 3.2.5. Thus the image of \mathfrak{f}_{i+j_0} in $\mathfrak{h}_{+,i_0}/\mathfrak{h}_{+,i_0+1}$ is 0. Therefore $\mathfrak{f}_\infty \subseteq \mathfrak{f}_{i+j_0} \subseteq \mathfrak{h}_{+,i_0+1}$ and this contradicts the choice of i_0 . Thus $\mathfrak{f}_\infty = 0$. \square

3.3. Isomorphism properties. In this section we list properties of the isomorphism classes of those latticed F -isocrystals with a group over k which are of the form $(M, g\phi, G)$ with $g \in G(W(k))$. We recall that σ acts on $H_0(W(k))$ as ϕ does, cf. Section 3.1.3.

3.3.1. Lemma. (a) *We have $H_0(W(k)) = \{*\phi^{-1} \mid * \in H_0(W(k))\}$.*

(b) *If $m \in \mathbb{N}$, then*

$$\begin{aligned} \text{Ker}(H_0(W(k)) \rightarrow H_0(W_m(k))) \\ = \{*\phi^{-1} \mid * \in \text{Ker}(H_0(W(k)) \rightarrow H_0(W_m(k)))\}. \end{aligned}$$

(c) *For each $* \in H_{+0}(W(k))$, we have $\phi(*) \in H_{+0}(W(k))$.*

(d) *Let \mathfrak{f}_+ and \mathfrak{f}_- be two left and right \mathfrak{h}_0 -modules contained in \mathfrak{h}_+ and \mathfrak{h}_- (respectively). Let $g \in H(W(k))$ be such that $a(g) \in \mathfrak{f}_+$ and $c(g) \in \mathfrak{f}_-$. Then there exists an element $h_0 \in H_0(W(k))$ such that for $g_0 := h_0 g \phi(h_0)^{-1} \in H(W(k))$ we have $a(g_0) \in \mathfrak{f}_+$, $b(g_0) = 0$, and $c(g_0) \in \mathfrak{f}_-$.*

Proof. As $H_{0\mathbb{Z}_p}$ is an affine, smooth group scheme over $\text{Spec}(\mathbb{Z}_p)$ whose special fibre is connected (see Section 3.1.3), (a) and (b) are only the Witt vectors version of Lang theorem for affine, connected, smooth groups over \mathbb{F}_p ; see [NV, Proposition 2.1] and its proof for details. As $\phi(\mathfrak{h}_+) \subseteq \mathfrak{h}_+$ and $\phi(H_0(W(k))) = H_0(W(k))$, from (3c) we get that for each $* \in H_{+0}(W(k))$ we have $\phi(*) \in H_{+0}(W(k))$. Thus (c) holds.

We prove (d). We have $1_M + b(g) \in H_0(W(k))$, cf. Fact 3.1.2. Let $h_0 \in H_0(W(k))$ be such that $1_M + b(g) = h_0^{-1}\phi(h_0)$, cf. (a). We have

$$\begin{aligned} g_0 &= h_0 a(g)\phi(h_0)^{-1} + h_0[1_M + b(g)]\phi(h_0)^{-1} + h_0 c(g)\phi(h_0)^{-1} \\ &= 1_M + h_0 a(g)\phi(h_0)^{-1} + h_0 c(g)\phi(h_0)^{-1}. \end{aligned}$$

As \mathfrak{f}_+ and \mathfrak{f}_- are left and right \mathfrak{h}_0 -modules, we have $h_0 a(g)\phi(h_0)^{-1} \in \mathfrak{f}_+$ and $h_0 c(g)\phi(h_0)^{-1} \in \mathfrak{f}_-$. Therefore $a(g_0) = h_0 a(g)\phi(h_0)^{-1} \in \mathfrak{f}_+$ and $c(g_0) = h_0 c(g)\phi(h_0)^{-1} \in \mathfrak{f}_-$. Thus (d) holds. \square

3.3.2. Lemma. *Let $g = 1_M + c(g) \in 1_M + \mathfrak{h}_-$. Then there exists an element $h \in G(W(k)) \cap (1_M + \mathfrak{h}_-[\frac{1}{p}])$ such that we have $hg\phi(h)^{-1} = 1_M$.*

Proof. For $i \in \mathbb{N} \cup \{0\}$ let $g_i := \phi^{-i}(g) = 1_M + \phi^{-i}(c(g))$. As $\phi^{-i}(O_-) \subseteq O_-$, we have $\phi^{-i}(\mathfrak{h}_-) \subseteq \mathfrak{g} \cap O_- \subseteq \mathfrak{g} \cap \mathfrak{h}_-[\frac{1}{p}]$ and thus $\phi^{-i}(c(g))$ is a nilpotent element of \mathfrak{g} . This implies that $g_i = 1_M + \phi^{-i}(c(g))$ is an invertible element of \mathfrak{g} i.e., we have $g_i \in G(W(k))$. We have $g = g_0$. For $i \in \mathbb{N}$ we have $\phi(g_i) = g_{i-1}$. As all Newton polygon slopes of (L_-, ϕ) are negative, the sequence $(\phi^{-i}(c(g)))_{i \in \mathbb{N}}$ of elements of $\mathfrak{g} \cap O_-$ converges to 0. This implies that the element $h := \lim_{i \rightarrow \infty} g_i g_{i-1} \dots g_1 \in G(W(k))$ is well defined. We compute that

$$\begin{aligned} hg\phi(h)^{-1} &= \lim_{i \rightarrow \infty} g_i g_{i-1} \dots g_1 g\phi(g_1)^{-1} \dots \phi(g_{i-1})^{-1} \phi(g_i)^{-1} \\ &= \lim_{i \rightarrow \infty} g_i \dots g_0 g_0^{-1} \dots g_{i-1}^{-1} \end{aligned}$$

is equal to $\lim_{i \rightarrow \infty} g_i = 1_M$. \square

3.4. Proof of 1.3 (a). We prove Theorem 1.3 (a). Let $\tilde{g} \in G(W(k))$ be congruent to 1_M modulo p^{ℓ_G} . As $\tilde{g} - 1_M \in p^{\ell_G} \mathfrak{g} \subseteq \mathfrak{h}$, we have $\tilde{g} \in \mathfrak{h}$. As $\tilde{g} \in \mathrm{GL}_M(W(k))$, we have $\tilde{g} \in H(W(k))$ (cf. Section 2.5 (i)). Thus to prove Theorem 1.3 (a), it suffices to prove the following stronger statement:

(*) *for each element g in $H(W(k))$ there exists an element h_g in $G(W(k))$ such that $h_g g\phi(h_g)^{-1} = 1_M$.*

We will first prove the following lemma.

3.4.1. Lemma. *Let $g \in H(W(k))$. Then there exists an element $h_+ \in H_{+0}(W(k))$ such that $g_+ := h_+ g\phi(h_+)^{-1} \in H(W(k))$ has the property that $a(g_+) = 0$.*

Proof. We can assume that $\mathfrak{h}_+ \neq 0$. For $i \in \mathbb{N} \cup \{0\}$, let $e_i = \mathfrak{f}_i \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$ be the $W(k)$ -subalgebra of \mathfrak{h} constructed in Lemma 3.2.3. By induction on $i \in \mathbb{N} \cup \{0\}$ we show that there exists $h_i \in H_{+0}(W(k))$ such that $g_i := h_i g\phi(h_i)^{-1} \in H(W(k))$ has the property that $a(g_i) \in e_i$. Taking $h_0 = 1_M$, we have $g_0 = g \in \mathfrak{h} = e_0$. Thus the basis of the induction holds. The passage from i to $i + 1$ goes as follows.

We will take h_{i+1} to be a product of the form $h_{i,+}h_{i,0}h_i$. Let $\delta_i \in \mathbb{N} \cup \{0\}$ be the greatest number such that we have $b(g_i) \in p^{\delta_i}\mathfrak{h}_0$. Let $h_{i,0} \in \text{Ker}(H_0(W(k)) \rightarrow H_0(W_{\delta_i}(k)))$ be such that $h_{i,0}(1_M + b(g_i))\phi(h_{i,0})^{-1} = 1_M$, cf. Lemma 3.3.1 (b). The element $g_{i+1,0} := h_{i,0}g_i\phi(h_{i,0})^{-1} \in H(W(k))$ has the properties that $a(g_{i+1,0}) \in \mathfrak{f}_i$ and $b(g_{i+1,0}) = 0$, cf. proof of Lemma 3.3.1 (d) applied with $(\mathfrak{f}_+, \mathfrak{f}_-) = (\mathfrak{f}_i, \mathfrak{h}_-)$. Let $h_{i,+} := 1_M - a(g_{i+1,0}) \in 1_M + \mathfrak{h}_+ \leq H_{+0}(W(k))$. We compute that

$$\begin{aligned} g_{i+1} &= h_{i+1}g\phi(h_{i+1})^{-1} = h_{i,+}h_{i,0}g_i\phi(h_{i,0})^{-1}\phi(h_{i,+})^{-1} = h_{i,+}g_{i+1,0}\phi(h_{i,+})^{-1} \\ &= [1_M - a(g_{i+1,0})][1_M + a(g_{i+1,0}) + c(g_{i+1,0})][1_M - \phi(a(g_{i+1,0}))]^{-1} \\ &= [1_M - a(g_{i+1,0})^2 - a(g_{i+1,0})c(g_{i+1,0}) + c(g_{i+1,0})][1_M - \phi(a(g_{i+1,0}))]^{-1}. \end{aligned}$$

As $a(g_{i+1,0}) \in \mathfrak{f}_i$, the three elements $-a(g_{i+1,0})^2$, $\Pi_+(-a(g_{i+1,0})c(g_{i+1,0}))$, and $\phi(a(g_{i+1,0}))$ belong to $\Theta(\mathfrak{f}_i) = \mathfrak{f}_{i+1}$. As $\Pi_+(-a(g_{i+1,0})c(g_{i+1,0})) \in \mathfrak{f}_{i+1}$, we get that $-a(g_{i+1,0})c(g_{i+1,0}) \in e_{i+1}$. As e_{i+1} is a $W(k)$ -algebra, we conclude that both $1_M - \phi(a(g_{i+1,0}))$ and $[1_M - a(g_{i+1,0})][1_M + a(g_{i+1,0}) + c(g_{i+1,0})]$ belong to e_{i+1} . From Section 2.5 (i) we get that $[1_M - \phi(a(g_{i+1,0}))]^{-1} \in e_{i+1}$. Thus we have $g_{i+1} \in e_{i+1}$. This ends the induction.

Due to Lemma 3.2.6, the sequences $(a(g_i))_{i \in \mathbb{N} \cup \{0\}}$ and $(a(g_{i+1,0}))_{i \in \mathbb{N} \cup \{0\}}$ of elements of \mathfrak{h}_+ converge to 0. We have $b(g_{i+1}) = [1_M - a(g_{i+1,0})][1_M + a(g_{i+1,0}) + c(g_{i+1,0})][1_M - \phi(a(g_{i+1,0}))]^{-1} - 1_M - a(g_{i+1}) - c(g_{i+1}) \in \mathfrak{h}_0$. From the last two sentences we easily get that the sequence $(b(g_{i+1}))_{i \in \mathbb{N} \cup \{0\}}$ of elements of \mathfrak{h}_0 converges to 0. Thus the sequence $(\delta_i)_{i \in \mathbb{N} \cup \{0\}}$ of non-negative integers converges to ∞ . This implies that the sequence $(h_{i,0})_{i \in \mathbb{N} \cup \{0\}}$ of elements of $H_0(W(k))$ converges to 1_M . As $h_{i,+} = 1_M + a(g_{i+1,0})$, the sequence $(h_{i,+})_{i \in \mathbb{N} \cup \{0\}}$ converges to 1_M . Thus the sequence $(h_{i,+}h_{i,0})_{i \in \mathbb{N} \cup \{0\}}$ of elements of $H_{+0}(W(k))$ converges also to 1_M . As $h_{i+1} = h_{i,+}h_{i,0}h_i$, we get that the sequence $(h_i)_{i \in \mathbb{N} \cup \{0\}}$ of elements of $H_{+0}(W(k))$ converges to an element $h_+ \in H_{+0}(W(k))$. We have $g_+ = h_+g\phi(h_+)^{-1} = \lim_{i \rightarrow \infty} h_i g\phi(h_i)^{-1} = \lim_{i \rightarrow \infty} g_i \in \bigcap_{i=0}^{\infty} e_i = \bigcap_{i=0}^{\infty} \mathfrak{f}_i \oplus \mathfrak{h}_0 \oplus \mathfrak{h}_-$. Thus $g_+ \in \mathfrak{h}_0 \oplus \mathfrak{h}_-$, cf. Lemma 3.2.6. Therefore $a(g_+) = 0$. \square

3.4.2. End of the proof of 1.3(a). Let $g \in H(W(k))$. Let $h_+ \in H_{+0}(W(k))$ and $g_+ \in H(W(k))$ be as in Lemma 3.4.1. Let $h_0 \in H_0(W(k))$ be such that for $g_0 := h_0g_+\phi(h_0)^{-1} \in H(W(k))$ we have $a(g_0) = b(g_0) = 0$, cf. Lemma 3.3.1 (d) applied with $(\mathfrak{f}_+, \mathfrak{f}_-) = (0, \mathfrak{h}_-)$. Let $h_- \in G(W(k))$ be such that we have $g_0 = h_-^{-1}\phi(h_-)$, cf. Lemma 3.3.2. Due to (3a), the element $h_g := h_-h_0h_+$ belongs to $G(W(k))$. We have $h_gg\phi(h_g)^{-1} = h_-h_0h_+g\phi(h_+)^{-1}\phi(h_0)^{-1}\phi(h_-)^{-1} = h_-h_0g_+\phi(h_0)^{-1}\phi(h_-)^{-1} = h_-g_0\phi(h_-)^{-1} = 1_M$. Thus the statement 3.4 (*) holds. This ends the proof of Theorem 1.3 (a). \square

3.4.3. Remarks. (a) The proof of Theorem 1.3 (a) can be also worked out using $p^{\ell_G}\mathfrak{g} + \mathfrak{h}_0 + \mathfrak{h}_-$ instead of $\mathfrak{h} = p^{\ell_G}\mathfrak{g} + \mathfrak{h}_+ + \mathfrak{h}_0$.

(b) If $g \in H_{+0}(W(k))$, then $h_g = h_0 h_+ \in H_{+0}(W(k))$. Thus we have an identity $H_{+0}(W(k)) = \{ *^{-1} \phi(*) \mid * \in H_{+0}(W(k)) \}$ (to be compared with Lemma 3.3.1 (a)).

3.5. Proof of 1.3 (b). We prove Theorem 1.3 (b). We consider a direct sum decomposition

$$M = \bigoplus_{i \in I} M_i \quad (4a)$$

with the property that for all elements i of the finite set I we have $\phi(M_i[\frac{1}{p}]) = M_i[\frac{1}{p}]$ and (M_i, ϕ) is isoclinic. For instance, we can take I to be the set of Newton polygon slopes of (M, ϕ) and then as each M_i we can take $M \cap W(i)$ (see Section 1.1.2 for $W(\alpha)$ with $\alpha \in \mathbb{Q}$). For each $i \in I$, let $\alpha_i \in \mathbb{Q}$ be the unique Newton polygon slope of (M_i, ϕ) . In Section 3.5.1 we do not assume that the association $i \rightarrow \alpha_i$ is one-to-one.

3.5.1. Scholium. One computes ℓ_{GLM} as follows. For $i \in I$, let \mathcal{B}_i be a $W(k)$ -basis for M_i . Let $\mathcal{B} := \bigcup_{i \in I} \mathcal{B}_i$; it is a $W(k)$ -basis for M . Let $\mathcal{B}^* := \{x^* \mid x \in \mathcal{B}\}$ be the $W(k)$ -basis for M^* which is the dual of \mathcal{B} (see Section 2.1).

Due to (4a), we have direct sum decompositions

$$\begin{aligned} \text{End}(M) \cap L_+ &= \bigoplus_{\substack{i, j \in I \\ \alpha_i < \alpha_j}} \text{Hom}(M_i, M_j), \\ \text{End}(M) \cap L_0 &= \bigoplus_{i \in I} \text{End}(M_i), \end{aligned}$$

and

$$\text{End}(M) \cap L_- = \bigoplus_{\substack{i, j \in I \\ \alpha_i < \alpha_j}} \text{Hom}(M_j, M_i).$$

Thus $\text{End}(M) = (\text{End}(M) \cap L_+) \oplus (\text{End}(M) \cap L_0) \oplus (\text{End}(M) \cap L_-)$ and therefore

$$\begin{aligned} \text{End}(M)/O &= [(\text{End}(M) \cap L_+)/O_+] \oplus [(\text{End}(M) \cap L_0)/O_0] \\ &\quad \oplus [(\text{End}(M) \cap L_-)/O_-]. \end{aligned} \quad (4b)$$

For $i, j \in I$, $x \in \mathcal{B}_i$, and $y \in \mathcal{B}_j$, we define a number $\ell(x, y) \in \mathbb{N} \cup \{0\}$ via the following two rules:

- if $\alpha_i \geq \alpha_j$, let $\ell(x, y) \in \mathbb{N} \cup \{0\}$ be the smallest number such that we have $p^{\ell(x, y)} \phi^q(x \otimes y^*) \in \text{Hom}(M_j, M_i)$ for all $q \in \mathbb{N}$;
- if $\alpha_i < \alpha_j$, let $\ell(x, y) \in \mathbb{N} \cup \{0\}$ be the smallest number such that we have $p^{\ell(x, y)} \phi^{-q}(x \otimes y^*) \in \text{Hom}(M_j, M_i)$ for all $q \in \mathbb{N}$.

Let $\ell_+, \ell_0, \ell_- \in \mathbb{N} \cup \{0\}$ be the smallest numbers such that p^{ℓ_+} annihilates $(\text{End}(M) \cap L_+)/O_+$, p^{ℓ_0} annihilates $(\text{End}(M) \cap L_0)/O_0$, and p^{ℓ_-} annihilates $(\text{End}(M) \cap L_-)/O_-$. As

$$O_+ = \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^{-q}(\text{End}(M) \cap L_+) = \bigcap_{q \in \mathbb{N} \cup \{0\}} \text{End}(\phi^{-q}(M) \cap L_+),$$

ℓ_+ is the smallest non-negative integer with the property that we have $p^{\ell_+}(\text{End}(M) \cap L_+) \subseteq \phi^{-q}(M) \cap L_+$ for all $q \in \mathbb{N}$ (i.e., we have $p^{\ell_+} \phi^q(\text{End}(M) \cap L_+) \subseteq \text{End}(M)$ for all $q \in \mathbb{N}$). As $\{x \otimes y^* \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j, i, j \in I, \alpha_i > \alpha_j\}$ is a $W(k)$ -basis for $\text{End}(M) \cap L_+$, we get that ℓ_+ is the smallest non-negative integer such that we have $p^{\ell_+} \phi^q(x \otimes y^*) \in \text{Hom}(M_j, M_i)$ for all $q \in \mathbb{N}$, all $i, j \in I$ with $\alpha_i > \alpha_j$, and all $x \in \mathcal{B}_i$ and $y \in \mathcal{B}_j$. Therefore

$$\ell_+ := \max\{\ell(x, y) \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j, i, j \in I, \alpha_i > \alpha_j\}. \quad (5a)$$

Similar arguments show that

$$\ell_0 = \max\{\ell(x, y) \mid x, y \in \mathcal{B}_i, i \in I\} \quad (5b)$$

and that

$$\ell_- := \max\{\ell(x, y) \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j, i, j \in I, \alpha_i < \alpha_j\}. \quad (5c)$$

From (4b) and the very definitions of ℓ_+, ℓ_0 , and ℓ_- we get that $\max\{\ell_+, \ell_0, \ell_-\} \in \mathbb{N} \cup \{0\}$ is the smallest number such that $p^{\max\{\ell_+, \ell_0, \ell_-\}}$ annihilates $\text{End}(M)/O$.

Next we define a number $\varepsilon_{\text{GL}_M} \in \{0, 1\}$ via the following rules. If $O = \text{End}(M)$, let $\varepsilon_{\text{GL}_M} := \ell_{\text{GL}_M}$ (cf. rules 1.2 (a) and (b)); we have $\ell_+ = \ell_0 = \ell_- = 0$ and thus $\ell_{\text{GL}_M} = \max\{\varepsilon_{\text{GL}_M}, \ell_+, \ell_0, \ell_-\}$. If $O \neq \text{End}(M)$, let $\varepsilon_{\text{GL}_M} := 0$; we have $\ell_{\text{GL}_M} = \max\{\ell_+, \ell_0, \ell_-\}$ (cf. rule 1.2 (b)). From the last two sentences and the formulas (5a), (5b), and (5c) we get that, regardless of what O is, we have

$$\ell_{\text{GL}_M} = \max\{\varepsilon_{\text{GL}_M}, \ell_+, \ell_0, \ell_-\} = \max\{\varepsilon_{\text{GL}_M}, \ell(x, y) \mid x, y \in \mathcal{B}\}. \quad (6a)$$

The latticed F -isocrystals $(\text{Hom}(M_j, M_i), \phi)$ and $(\text{Hom}(M_i, M_j), \phi)$ are dual to each other (cf. Section 2.1) and the dual of the $W(k)$ -basis $\{x \otimes y^* \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j\}$ of $\text{Hom}(M_j, M_i)$ is the $W(k)$ -basis $\{y \otimes x^* \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j\}$ of $\text{Hom}(M_i, M_j)$. Based on this, from the property 2.1 (*) we get that for all $i, j \in I$ we have an equality

$$\max\{\ell(x, y) \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j\} = \max\{\ell(y, x) \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j\}. \quad (6b)$$

3.5.2. Reduction steps and notations. Let $\ell := \ell_{\text{GL}_M}$. Based on Theorem 1.3 (a), we have $n_{\text{GL}_M} \leq \ell$. Thus to prove that $n_{\text{GL}_M} = \ell$, it suffices to show that $n_{\text{GL}_M} > \ell - 1$. If $n_{\text{GL}_M} = 0$, then $\ell = 0$ (see Lemma 2.3) and therefore $n_{\text{GL}_M} > \ell - 1$. Thus to prove that $n_{\text{GL}_M} = \ell$, it suffices to show that for $\ell \geq 2$ we have $n_{\text{GL}_M} > \ell - 1$. To check this we can assume that the map $I \rightarrow \mathbb{Q}$ that takes $l \in I$ to $\alpha_l \in \mathbb{Q}$ is injective (i.e., for each element $l \in I$ we have $M_l = M \cap W(\alpha_l)$).

Let q be the smallest positive integer for which the following two properties hold:

- (i) there exists a $W(k)$ -basis $\mathcal{B} = \cup_{l \in I} \mathcal{B}_l$ for M which is contained in $\cup_{l \in I} M_l$ and for which there exist elements $i, j \in I$, $x \in \mathcal{B}_i \subseteq M_i$, and $y \in \mathcal{B}_j \subseteq M_j$ such that (cf. (6a) and (6b)) we have $\ell(x, y) = \ell$ and $\alpha_j \leq \alpha_i$;
- (ii) we have $e_{q,x,y} := p^{\ell(x,y)} \phi^q(x \otimes y^*) = p^\ell \phi^q(x \otimes y^*) \in \text{Hom}(M_j, M_i) \setminus p \text{Hom}(M_j, M_i)$.

The existence of q follows from (6a), (6b), and the very definition of the numbers $\ell(x, y)$.

For $z \in M$, let $a_{z,q}$ be the unique integer such that we have $\phi^q(z) \in p^{a_{z,q}} M \setminus p^{a_{z,q}+1} M$. We can choose the $W(k)$ -basis $\mathcal{B} = \cup_{l \in I} \mathcal{B}_l$ such that we have a direct sum decomposition $M = \bigoplus_{z \in \mathcal{B}} W(k) p^{-a_{z,q}} \phi^q(z)$ i.e., we have $\phi^{-q}(M) = \bigoplus_{z \in \mathcal{B}} W(k) p^{-a_{z,q}} z$. Let

$$a_{i,q} := \min\{a_{z,q} \mid z \in \mathcal{B}_i\} \quad \text{and} \quad b_{j,q} := \max\{a_{z,q} \mid z \in \mathcal{B}_j\}.$$

Therefore $a_{i,q}$ is the greatest integer such that we have $\phi^q(M_i) \subseteq p^{a_{i,q}} M_i$ and $b_{j,q}$ is the smallest integer such that we have $p^{b_{j,q}} M_j \subseteq \phi^q(M_j)$. The smallest number $s \in \mathbb{N} \cup \{0\}$ with the property that $p^s \phi^q(\text{Hom}(M_j, M_i)) = p^s \text{Hom}(\phi^q(M_j), \phi^q(M_i))$ is contained in $\text{Hom}(M_j, M_i)$, equals $\max\{0, b_{j,q} - a_{i,q}\}$; as $e_{q,x,y} \in \text{Hom}(M_j, M_i) \setminus p \text{Hom}(M_j, M_i)$, we have $s \geq \ell(x, y) = \ell \geq 2$. As $s \leq \max\{\ell_+, \ell_0\} \leq \ell$, we conclude that $2 \leq \ell = s = b_{j,q} - a_{i,q}$. It is easy to see that we have $\max\{\ell_+, \ell_0\} \geq a_{y,q} - a_{x,q} \geq \ell(x, y)$, cf. property (ii) for the second inequality. From the last two sentences we get that $a_{x,q} = a_{i,q}$ and $a_{y,q} = b_{j,q}$. Thus we have $\ell = \ell(x, y) = a_{y,q} - a_{x,q} = b_{j,q} - a_{i,q}$. As $\ell = a_{y,q} - a_{x,q} > 0$, we have $x \neq y$.

3.5.3. The set Λ . Let $\Lambda := \{w \in M_i \setminus pM_i \mid a_{w,q} = a_{x,q}\}$; it is the set of those elements $w \in M_i$ for which $p^{-a_{x,q}} \phi^q(w)$ is a direct summand of M . Obviously the set Λ is stable under multiplication by invertible elements of $W(k)$. For $w \in \Lambda$ let

$$g_w := 1_M + p^{\ell-1} w \otimes y^* \in \text{End}(M);$$

it is the endomorphism of M that fixes each element $z \in \mathcal{B} \setminus \{y\}$ and that takes y to $y + p^{\ell-1} w$. As $\ell \geq 2$, we have $g_w \in \text{GL}_M(W(k))$. As each g_w is congruent to 1_M modulo $p^{\ell-1}$, to prove that $n_{\text{GL}_M} > \ell - 1$ it suffices to show that there exists an element $w \in \Lambda$ such that the latticed F -isocrystals $(M, g_w \phi)$ and (M, ϕ) are not isomorphic. We show that the assumption that this is not true leads to a contradiction. This assumption implies that for each element $w \in \Lambda$ there exists an element $h_w \in \text{GL}_M(W(k))$ which is an isomorphism between $(M, g_w \phi)$ and (M, ϕ) . Thus we have $h_w g_w \phi h_w^{-1} = \phi$ i.e., we have

$$h_w g_w = \phi(h_w). \tag{7a}$$

We write $h_w = 1_M + u_w$, where $u_w \in \text{End}(M)$. Substituting the expressions of h_w and g_w in (7a), we come across the following identity

$$u_w + p^{\ell-1}w \otimes y^* + p^{\ell-1}u_w(w \otimes y^*) = u_w + p^{\ell-1}[w + u_w(w)] \otimes y^* = \phi(u_w) \quad (7b)$$

(here $u_w(w \otimes y^*)$ is the product inside $\text{End}(M)$ of u_w and $w \otimes y^*$). In other words, if $v_w := w + u_w(w)$ then the pair (u_w, v_w) is a solution of the following equation

$$U + p^{\ell-1}V \otimes y^* = \Phi(U) \quad (7c)$$

in variables U and V that can take values in $\text{End}(M)[\frac{1}{p}]$ and M (respectively).

3.5.4. Fact. *There exists an isomorphism between $(M, g_w\phi)$ and (M, ϕ) defined by an element \tilde{h}_w of $\text{GL}_M(W(k))$ which has the following two properties:*

- (i) *it acts identically on each M_l with $l \in I \setminus \{i, j\}$ and leaves invariant M_i ;*
- (ii) *if $i \neq j$, then it acts identically on M_i , leaves invariant $M_i \oplus M_j$, and acts identically on $(M_i + M_j)/M_i$.*

Proof. We will prove this only in the case when $i \neq j$ (as the case $i = j$ is even simpler). We know that g_w acts identically on each M_l with $l \in I \setminus \{j\}$ and on $(M_i \oplus M_j)/M_i$. This implies that each M_l with $l \in I \setminus \{j\}$ is the maximal direct summand of M such that all Newton polygon slopes of $(M_l, g_w\phi)$ are equal to α_l and that $M_i \oplus M_j$ is the maximal direct summand of M such that all Newton polygon slopes of $(M_i \oplus M_j, g_w\phi)$ are equal to either α_i or α_j . From this and the fact that $h_w \in \text{GL}_M(W(k))$ is an isomorphism between $(M, g_w\phi)$ and (M, ϕ) , we get that h_w leaves invariant each M_l with $l \in I \setminus \{j\}$ as well as $M_i \oplus M_j$. Even more, from the second sentence of this proof we get that h_w restricted to each M_l with $l \in I \setminus \{j\}$ is an automorphism h_{lw} of (M_l, ϕ) and moreover h_w induces an automorphism of $((M_i \oplus M_j)/M_i, \phi)$ and thus an automorphism h_{jw} of (M_j, ϕ) .

Let $h_{0w} := \prod_{l \in I} h_{lw} \in \prod_{l \in I} \text{GL}_{M_l}(W(k)) \leq \text{GL}_M(W(k))$; it is an automorphism of (M, ϕ) . The element $\tilde{h}_w := h_{0w}^{-1}h_w \in \text{GL}_M(W(k))$ has all the desired properties. \square

To reach the desired contradiction we can assume that we have $h_w = \tilde{h}_w$, where \tilde{h}_w is as in Fact 3.5.4. We first consider the case when $i \neq j$.

3.5.5. The case $i \neq j$. We assume that $i \neq j$ (i.e., $\alpha_j < \alpha_i$). As $h_w = \tilde{h}_w$, we have $u_w \in \text{Hom}(M_j, M_i)$. From this and the relation $i \neq j$ we get that $u_w(w) = 0$. As $\alpha_j < \alpha_i$, all Newton polygon slopes of $(\text{Hom}(M_j, M_i), \phi)$ are positive. Therefore for each V in M_i the sequence $(\phi^m(p^{\ell-1}V \otimes y^*))_{m \geq 0}$ converges to 0 and thus all the solutions of the equation (7c) in $\text{Hom}(M_j, M_i)[\frac{1}{p}] \times M_i$ are of the form $(-\sum_{m=0}^{\infty} \phi^m(p^{\ell-1}V \otimes y^*), V)$. From this and the relation $u_w(w) = 0$ we get the

following identity

$$u_w = - \sum_{m=0}^{\infty} \phi^m(p^{\ell-1}w \otimes y^*). \quad (7d)$$

We have the following two properties of the terms of the sum (7d).

(i) All the terms of the sum of (7d) belong to $\frac{1}{p} \text{Hom}(M_j, M_i)$ (this is so as w and y belong to a $W(k)$ -basis for M formed by elements of $\cup_{l \in I} M_l$ and therefore the element $\ell(w, y)$ can be defined as in Section 3.5.1 and it is equal to $\ell(x, y) = \ell$). Moreover, all but a finite number of these terms belong to $\text{Hom}(M_j, M_i)$.

(ii) The term $\phi^q(p^{\ell-1}w \otimes y^*)$ of the sum of (7d) belongs to $\frac{1}{p} \text{Hom}(M_j, M_i) \setminus \text{Hom}(M_j, M_i)$ (cf. property 3.5.2 (ii) and the fact that $a_{w,q} = a_{x,q}$).

Let γ be an invertible element of $W(k)$. Let $\bar{\gamma} \in k \setminus \{0\}$ be its reduction modulo p . Based on properties (i) and (ii), the condition that the element $u_{\gamma w}$ obtained as in (7d) belongs to $\text{Hom}(M_j, M_i)$ is expressed by $\bar{\gamma}$ being a solution of a system of polynomial equations in one variable which have coefficients in k and which contain at least one polynomial of degree at least p^q . Therefore there exist such elements γ with the property that we have $u_{\gamma w} \in \frac{1}{p} \text{Hom}(M_j, M_i) \setminus \text{Hom}(M_j, M_i)$. Thus for such an element γ we have $\gamma w \in \Lambda$ and $h_{\gamma w} = 1_M + u_{\gamma w} \notin \text{GL}_M(W(k))$. Contradiction.

3.5.6. Extra reduction steps. To reach the desired contradiction we can assume that $i = j$ (i.e., $\alpha_i = \alpha_j$), cf. Section 3.5.5. As $h_w = \tilde{h}_w$ and $i = j$, to reach a contradiction we can assume based on Fact 3.5.4 (i) that $M_i = M$ (i.e., that $I = \{i\}$). Thus (M, ϕ) is isoclinic and we have $O = O_0 = A_0 \otimes_{\mathbb{Z}_p} W(k)$, cf. Section 1.2.

3.5.7. Lemma. *We recall that $v_w = w + u_w(w)$. Then we have $v_w \in \Lambda$.*

Proof. Due to the definition of q , we have $\phi^s(g_w) \in \text{GL}_M(W(k))$ for all $s \in \{1, \dots, q-1\}$ but $\phi^q(g_w) \notin \text{GL}_M(W(k))$. From this and the equation (7a) we get that $\phi^s(h_w) \in \text{GL}_M(W(k))$ for all $s \in \{1, \dots, q\}$ but $\phi^{q+1}(h_w) \notin \text{GL}_M(W(k))$. Thus we have $\phi^s(u_w) \in \text{End}(M)$ for all $s \in \{1, \dots, q\}$ but $\phi^{q+1}(u_w) \notin \text{End}(M)$.

Due to this and the identity (7b) we get that $\phi^q(p^{\ell-1}v_w \otimes y^*) \notin \text{End}(M)$. If $\phi^q(p^{\ell-1}v_w \otimes y^*) \notin \frac{1}{p} \text{End}(M)$ or if $v_w \in pM$, then we have $\ell_0 \geq \ell + 1$ and this contradicts (6a). Thus we have $\phi^q(p^{\ell-1}v_w \otimes y^*) \in \frac{1}{p} \text{End}(M) \setminus \text{End}(M)$ and $v_w \in M \setminus pM$. Therefore $\phi^q(p^{\ell}v_w \otimes y^*) \in \text{End}(M) \setminus p \text{End}(M)$ and $v_w \in M \setminus pM$. But we also have $\phi^q(p^{\ell}x \otimes y^*) \in \text{End}(M) \setminus p \text{End}(M)$, cf. property 3.5.2 (ii). From the last two sentences and the very definitions of $a_{w,q}$ and $a_{x,q}$, we get that $a_{v_w,q} = a_{x,q}$. From this and the relation $v_w \in M \setminus pM$ we conclude that $v_w \in \Lambda$. \square

3.5.8. Lemma. *Let $(u, v) \in \text{End}(M) \times M$ be a solution of the equation (7c).*

(a) *Then we have $\{u, p^{\ell-1}v \otimes y^*\} \subset \text{End}(M) \cap \frac{1}{p}O$.*

(b) Let $v_1 \in pM$. Then there exists a solution $(u + u_1, v + v_1)$ of the equation (7c) with $u_1 \in O$.

Proof. We have $pp^{\ell-1}v \otimes y^* \in p^\ell \text{End}(M) \subseteq O$. It is easy to see that for each element $\tilde{w} \in O$, the equation $* + \tilde{w} = \phi(*)$ in $*$ has a solution in $\phi(O) = O = O_0$ and thus also in $\text{End}(M)$. Let $\tilde{u} \in O$ be such that we have $\tilde{u} + pp^{\ell-1}v \otimes y^* = \phi(\tilde{u})$. Thus $pu - \tilde{u} = \phi(pu - \tilde{u})$ belongs to $A_0[\frac{1}{p}] \cap \text{End}(M) = A_0 \subseteq O$. Therefore we have $u \in \text{End}(M) \cap \frac{1}{p}O$. Thus (a) holds. Part (b) follows from the fact that there exists $u_1 \in O$ such that $u_1 + p^\ell v_1 \otimes y^* = \phi(u_1)$. \square

3.5.9. Morphisms between k -schemes. Let \mathcal{M} be the affine space (scheme) over k defined naturally by the k -vector space M/pM . Let $\varphi_q: \mathcal{M} \rightarrow \mathcal{M}$ be the morphism of k -schemes that takes $\bar{*} \in \mathcal{M}(k) = M/pM$ to the element of $\mathcal{M}(k)$ which is the reduction modulo p of $p^{-a_{x,q}}\phi^q(*) \in M$, where $*$ $\in M$ is an arbitrary lift of \bar{m} .

The set $\text{Im}(\Lambda \rightarrow M/pM)$ is the set of k -valued points of the open, non-empty subscheme $\mathcal{S} := \varphi_q^{-1}(\mathcal{M} \setminus \{0\})$ of \mathcal{M} . For each solution $(u, v) \in \text{End}(M) \times M$ of the equation (7c), a similar argument to the proof of Lemma 3.5.8 shows that v modulo p determines u modulo A_0 up to a finite number of possibilities. From this and the identity $w = v_w - u_w$, we get that the association that takes (w, u_w) modulo p to v_w modulo p has finite fibres. This association can be viewed as the one defined naturally (at the level of k -valued points) by a morphism of k -schemes whose codomain is \mathcal{S} and whose domain has the same dimension r as \mathcal{S} . By reasons of dimensions, we get the following:

(i) There exists an open, non-empty subscheme \mathcal{V} of \mathcal{S} which has the property that each k -valued point \bar{v} of \mathcal{V} is of the form v_w modulo p for some elements $w \in \Lambda$ and $u_w \in \text{End}(M)$ such that $(w, v_w) := (w, w + u_w(w))$ is a solution of the equation (7c).

Let $\bar{O} := \frac{1}{p}O/O$ and let \bar{E} be the image of $\text{End}(M) \cap \frac{1}{p}O$ in \bar{O} . Both \bar{O} and \bar{E} are k -vector spaces. Let \mathcal{O} and \mathcal{E} be the affine spaces (schemes) over k defined naturally by the k -vector spaces \bar{O} and \bar{E} (respectively). Let $\bar{\phi}: \mathcal{O} \rightarrow \mathcal{O}$ be the morphism which takes a k -valued point of \mathcal{O} defined by some element $o \in \frac{1}{p}O$ to the k -valued point of \mathcal{O} defined by the element $\phi(o) - o \in \frac{1}{p}O$ (we think of $\bar{\phi}$ as a finite, surjective endomorphism of $\mathbb{G}_a^{r^2}$). Let $\mathcal{F} := \mathcal{E} \cap \bar{\phi}^{-1}(\mathcal{E})$. Thus \mathcal{F} is a closed subscheme of \mathcal{O} equipped with a morphism $m_1: \mathcal{F} \rightarrow \mathcal{E}$ induced from $\bar{\phi}$ (we think of m_1 as a homomorphism between closed subgroup schemes of $\mathbb{G}_a^{r^2}$). Based on Lemma 3.5.8 (a) we can speak about the natural images \bar{u}_w and $\bar{v}_{w,y}$ of u_w and $p^{\ell-1}v_w \otimes y^*$ (respectively) in \bar{O} and thus about k -valued points (denoted in the same way) $\bar{u}_w \in \mathcal{F}(k)$ and $\bar{v}_{w,y} \in \mathcal{E}(k)$ with the property that m_1 maps \bar{u}_w to $\bar{v}_{w,y}$.

We have a natural morphism of k -schemes $m_2: \mathcal{V} \rightarrow \mathcal{E}$ which at the level of k -valued points maps a k -valued point of \mathcal{V} represented by an element $v \in \Lambda$ to the k -valued point of \mathcal{E} defined by the image of $p^{\ell-1}v \otimes y^* \in p^{\ell-1} \text{End}(M) \subseteq \frac{1}{p}O$ in \bar{O} .

From the property (i) and the previous paragraph we get that the natural morphism

$$\iota: \mathcal{V} \times_{\mathcal{E}} \mathcal{F} \rightarrow \mathcal{V}$$

associated to the fibre product of m_1 and m_2 , is surjective. As the morphism $\bar{\phi}: \mathcal{O} \rightarrow \mathcal{O}$ has finite fibres and it is of finite type, the finite type morphism ι is quasi-finite and therefore it is generically finite. From this, the property (i), and Lemma 3.5.8 (b) we get the following:

(ii) There exists an element $v \in \Lambda$ that defines naturally a k -valued point of \mathcal{V} and there exists a finite subset $\bar{\Gamma}$ of k such that for each algebraically closed field k_1 that contains k and for every invertible element γ of $W(k_1)$ whose reduction modulo p does not belong to $\bar{\Gamma}$, the following equation in U ,

$$U + p^{\ell-1} \gamma v \otimes y^* = (\phi \otimes \sigma_{k_1})(U), \quad (7e)$$

obtained from (7c) by replacing (V, ϕ) with $(\gamma v, \phi \otimes \sigma_{k_1})$, possesses a solution in $\text{End}(M \otimes_{W(k)} W(k_1))$. Here σ_{k_1} is the Frobenius automorphism of the ring $W(k_1)$ of Witt vectors with coefficients in k_1 .

3.5.10. Good choice of γ . We will take k_1 to be an algebraic closure of $k((X))$, where X is an independent variable. We identify $W(k)[[X]]$ with a $W(k)$ -subalgebra of $W(k_1)$ that contains the invertible element $X = (X, 0, \dots)$ of $W(k_1)$. We will take $\gamma := \tau X$, where τ is an invertible element of $W(k)$. We have $\sigma_{k_1}(X) = X^p$. For this choice of γ , the equation (7e) has (up to addition of elements in the free \mathbb{Z}_p -module $\frac{1}{p}A_0$ of rank r^2) a unique solution

$$u_{\tau X} = - \sum_{m=0}^{\infty} X^{p^m} \phi^m(p^{\ell-1} \tau v \otimes y^*) \quad (7f)$$

in $\frac{1}{p}O \otimes_{W(k)} W(k_1)$. In fact we have $u_{\tau X} \in \frac{1}{p} \text{End}(M) \otimes_{W(k)} W(k)[[X]]$. As $\tau v \in \Lambda$, from the property 3.5.2 (ii) we get that the term $X^{p^q} \phi^q(p^{\ell-1} \tau v \otimes y^*)$ of (7f) does not belong to $\text{End}(M) \otimes_{W(k)} W(k_1)$. The last two sentences imply that the intersection $(u_{\tau X} + \frac{1}{p}A_0) \cap [\text{End}(M) \otimes_{W(k)} W(k_1)]$ is empty and therefore we reached the desired contradiction.

3.5.11. End of the proof. The contradiction we reached implies that $n_{\text{GL}_M} > \ell - 1$. Thus $n_{\text{GL}_M} = \ell = \ell_{\text{GL}_M}$. This ends the proof of Theorem 1.3 (b) and therefore also of Main Theorem A. \square

3.6. Remarks. Suppose (M, ϕ) is a direct sum of isoclinic latticed F -isocrystals over k .

(a) We have a direct sum decomposition $\mathfrak{g} = (\mathfrak{g} \cap L_+) \oplus (\mathfrak{g} \cap L_0) \oplus (\mathfrak{g} \cap L_-)$ of $W(k)$ -modules. Thus $O_G = \mathfrak{g} \cap O$ and therefore $\mathfrak{g}/O_G \subseteq \text{End}(M)/O$. From

this inclusion we easily get the following *monotony properties*: we have $\ell_G \leq \ell_{\mathrm{GL}_M}$ and therefore (cf. Main Theorem A) we also have $n_G \leq n_{\mathrm{GL}_M}$.

(b) We assume that (M, ϕ) is the Dieudonné module of D ; thus $\ell_{\mathrm{GL}_M} = \ell_D$. We will use the notations of Section 3.5. We also assume that there exist elements $x, y \in \mathcal{B}$ such that $\ell(x, y) \geq 2$ and we have $x \in \mathcal{B}_i$ and $y \in \mathcal{B}_j$ with $\alpha_j < \alpha_i$. Let $g \in \mathrm{GL}_M(W(k))$ be the element that fixes each $z \in \mathcal{B} \setminus \{y\}$ and that takes y to $y + p^{\ell(x,y)-1}x$. Let D_g be a p -divisible group over k whose Dieudonné module is isomorphic to $(M, g\phi)$. Then $D_g[p^{\ell(x,y)-1}]$ is isomorphic to $D[p^{\ell(x,y)-1}]$ and D_g has the same Newton polygon as D (as $\alpha_j \neq \alpha_i$). If by chance we also have an identity $\ell_D = \ell(x, y)$, then Section 3.5 can be easily adapted to give us that, up to a replacement of $x \in \mathcal{B}_i$ by a multiple of it with an invertible element of $W(k)$, we can assume that D_g is not isomorphic to D .

3.7. Variant of 1.3 (b). Let (M, ϕ, G) be a latticed F -isocrystal with a group over k such that Assumption 1.1.1 holds. We assume that the following two conditions hold:

- (i) we have $n_G \geq 1$ and a direct sum decomposition $\mathfrak{g} = (\mathfrak{g} \cap L_+) \oplus (\mathfrak{g} \cap L_0) \oplus (\mathfrak{g} \cap L_-)$ (or $\mathfrak{g} = (\mathfrak{g} \cap L_+) \oplus [\mathfrak{g} \cap (L_0 \oplus L_-)]$ or $\mathfrak{g} = [\mathfrak{g} \cap (L_+ \oplus L_0)] \oplus (\mathfrak{g} \cap L_-)$) of $W(k)$ -modules;
- (ii) for all $q \in \mathbb{N}$, there exists a $W(k)$ -basis \mathcal{B} for M and a sequence of integers $(a_{z,q})_{z \in \mathcal{B}}$ such that certain subsets of $\{x \otimes y^* \mid x, y \in \mathcal{B}\}$ are $W(k)$ -bases for all direct summands of \mathfrak{g} listed in (i) and moreover we have $M = \bigoplus_{z \in \mathcal{B}} W(k) p^{-a_{z,q}} \phi^q(z)$.

Then the proof of Theorem 1.3 (b) (see Section 3.5) can be entirely adapted to give us that $n_G = \ell_G$. We only add here two things. First, if by chance in Section 3.5.2 we have $\ell(x, y) = \ell_{\mathrm{GL}_M}$ with $x \in \mathcal{B}_i$ and $y \in \mathcal{B}_j$ such that $\alpha_j > \alpha_i$, then one needs to use ϕ^{-q} (instead of ϕ^q) with $q \in \mathbb{N}$ in order to reach the desired contradiction. Second, if we have $\mathfrak{g} = (\mathfrak{g} \cap L_+) \oplus [\mathfrak{g} \cap (L_0 \oplus L_-)]$ (resp. $\mathfrak{g} = [\mathfrak{g} \cap (L_+ \oplus L_0)] \oplus (\mathfrak{g} \cap L_-)$), then one needs to use Lemma 2.4 in order to be able to treat $\mathfrak{g} \cap (L_0 \oplus L_-)$ (resp. $\mathfrak{g} \cap (L_+ \oplus L_0)$) in the same manner as $\mathfrak{g} \cap L_-$ (resp. as $\mathfrak{g} \cap L_+$).

4. Direct applications to p -divisible groups

In this section we prove the results stated in Sections 1.4.2 to 1.4.4 (see Sections 4.5 to 4.7). In Sections 4.1 to 4.4 we introduce basis invariants of p -divisible groups over k and we present basic properties of them that are needed in Sections 4.5 to 4.7. Until the end we will assume that (M, ϕ) is the Dieudonné module of D .

4.1. Definitions. (a) Let $q \in \mathbb{N}$. Let $\alpha_D(q) \in \mathbb{N} \cup \{0\}$ be the greatest number such that we have $\phi^q(M) \subseteq p^{\alpha_D(q)}M$. Let $\beta_D(q) \in \mathbb{N} \cup \{0\}$ be the smallest number such that we have $p^{\beta_D(q)}M \subseteq \phi^q(M)$. Let $\delta_D(q) := \beta_D(q) - \alpha_D(q)$; as $p^{\beta_D(q)}M \subseteq \phi^q(M) \subseteq p^{\alpha_D(q)}M$, we have $\delta_D(q) \in \mathbb{N} \cup \{0\}$.

(b) We assume that D is isoclinic. Let $m := \text{g.c.d.}\{c, d\}$. Let $(c_1, d_1, r_1) := (\frac{c}{m}, \frac{d}{m}, \frac{r}{m})$. Let $u_D := \sup\{0, \beta_D(r_1n) - d_1n \mid n \in \mathbb{N}\}$. Let $\tilde{u}_D := \sup\{0, d_1n - \alpha_D(r_1n) \mid n \in \mathbb{N}\}$. Let $v_D := \sup\{0, \beta_D(rn) - dn \mid n \in \mathbb{N}\}$. Let $\tilde{v}_D := \sup\{0, dn - \alpha_D(rn) \mid n \in \mathbb{N}\}$. Proposition 4.3 (c) and (b) below will imply that u_D , \tilde{u}_D , v_D , and \tilde{v}_D are non-negative integers and therefore that in their definition we can replace \sup by \max .

(c) Let (M_1, ϕ_1) and (M_2, ϕ_2) be the Dieudonné modules of two isoclinic p -divisible groups D_1 and D_2 (respectively) over k . Let α_1 and α_2 be the unique Newton polygon slopes of D_1 and D_2 (respectively). Let $\ell_{D_1, D_2} \in \mathbb{N} \cup \{0\}$ be the smallest number that has the following property:

- (i) if $\alpha_1 \leq \alpha_2$, then for all $q \in \mathbb{N}$ the $W(k)$ -module $\phi^q(p^{\ell_{D_1, D_2}} \text{Hom}(M_1, M_2)) = p^{\ell_{D_1, D_2}} \text{Hom}(\phi^q(M_1), \phi^q(M_2))$ is included in $\text{Hom}(M_1, M_2)$;
- (ii) if $\alpha_1 > \alpha_2$, then for all $q \in \mathbb{N}$ the $W(k)$ -module $\phi^{-q}(p^{\ell_{D_1, D_2}} \text{Hom}(M_1, M_2)) = p^{\ell_{D_1, D_2}} \text{Hom}(\phi^{-q}(M_1), \phi^{-q}(M_2))$ is included in $\text{Hom}(M_1, M_2)$.

(d) Let $\varepsilon_D := \varepsilon_{\text{GL}_M}$, where the number $\varepsilon_{\text{GL}_M} \in \{0, 1\}$ is as in Scholium 3.5.1.

4.1.1. Remark. If $cd = 0$, then $O_0 = \text{End}(M)$ and therefore $\ell_D = \varepsilon_D = 0$. If $c, d \geq 1$ and D is ordinary (i.e., isomorphic to $(\mathbb{Q}_p/\mathbb{Z}_p)^c \oplus (\mu_{p^\infty})^d$), then $O = \text{End}(M)$ and the two-sided ideal of the $W(k)$ -algebra $\text{End}(M)$ generated by $O_+ \oplus O_-$ is $\text{End}(M)$; thus $\ell_D = \varepsilon_D = 1$. If $c, d \geq 1$ and D is not isomorphic to $(\mathbb{Q}_p/\mathbb{Z}_p)^c \oplus (\mu_{p^\infty})^d$, then $\text{End}(M) \neq O$ and therefore $\ell_D > 0$ and $\varepsilon_D = 0$; moreover $\ell_D \in \mathbb{N}$ is the smallest number such that we have $p^{\ell_D} \text{End}(M) \subseteq O$ (cf. rule 1.2 (b)).

4.2. Simple properties. In this section we list few simple properties of the invariants we have introduced so far.

4.2.1. Fact. We have $n_D = n_{D^\dagger}$ and $\ell_D = \ell_{D^\dagger}$.

Proof. We show that $n_D \leq n_{D^\dagger}$. Let C be a p -divisible group of codimension c and dimension d over k . If $C[p^{n_{D^\dagger}}]$ is isomorphic to $D[p^{n_{D^\dagger}}]$, then taking Cartier duals we get that $C^\dagger[p^{n_{D^\dagger}}]$ is isomorphic to $D^\dagger[p^{n_{D^\dagger}}]$ and thus that C^\dagger is isomorphic to D^\dagger . Taking Cartier duals, we get that C is isomorphic to D . This implies that $n_D \leq n_{D^\dagger}$. As D is the Cartier dual of D^\dagger , we also have $n_{D^\dagger} \leq n_D$. Thus $n_D = n_{D^\dagger}$.

As $(M^*, p\phi)$ is the Dieudonné module of D^\dagger , under the natural identification $\text{End}(M^*) = \text{End}(M)$, the level module of $(M^*, p\phi)$ gets identified with O . Thus we have $\ell_D = \ell_{D^\dagger}$. \square

4.2.2. Fact. *The following three properties hold:*

(a) *for all $q \in \mathbb{N}$, we have inclusions*

$$M \subseteq p^{-\beta_D(q)} \phi^q(M) \subseteq p^{-\delta_D(q)} M \quad (8a)$$

which are optimal in the sense that we also have

$$M \not\subseteq p^{-\beta_D(q)+1} \phi^q(M) \quad \text{and} \quad p^{-\beta_D(q)} \phi^q(M) \not\subseteq p^{-\delta_D(q)+1} M; \quad (8b)$$

(b) *for all $q \in \mathbb{N}$, we have $\alpha_{D^t}(q) = q - \beta_D(q)$ and $\beta_{D^t}(q) = q - \alpha_D(q)$;*

(c) *if D is isoclinic, then we have $u_D = \tilde{u}_{D^t}$, $\tilde{u}_D = u_{D^t}$, $v_D = \tilde{v}_{D^t}$, and $\tilde{v}_D = v_{D^t}$.*

Proof. Part (a) follows from the very Definition 4.1 (a). As $p^{\beta_D(q)} M \subseteq \phi^q(M) \subseteq p^{\alpha_D(q)} M$, we have $p^{-\alpha_D(q)} M^* \subseteq \phi^q(M^*) \subseteq p^{-\beta_D(q)} M^*$ i.e., $p^{q-\alpha_D(q)} M^* \subseteq (p\phi)^q(M^*) \subseteq p^{q-\beta_D(q)} M^*$. As $(M^*, p\phi)$ is the Dieudonné module of D^t and due to (8a) and (8b), we get that (b) holds. We prove (c). Due to (b) we have an equality $\beta_D(r_1 n) - d_1 n = r_1 n - \alpha_{D^t}(r_1 n) - d_1 n = c_1 n - \alpha_{D^t}(r_1 n)$. This implies that $u_D = \tilde{u}_{D^t}$. By replacing D with D^t , we get that $u_{D^t} = \tilde{u}_D$. Similar arguments show that $v_D = \tilde{v}_{D^t}$ and $\tilde{v}_D = v_{D^t}$. Thus (c) holds. \square

4.2.3. Lemma. *We assume that D is isoclinic. Let $\alpha := \frac{d}{r} \in \mathbb{Q} \cap [0, 1]$ be its unique Newton polygon slope. Then the following two properties hold:*

(a) *we have $\alpha_D(q) \leq q\alpha \leq \beta_D(q)$;*

(b) *if $\alpha_D(q) = q\alpha$ (or if $\beta_D(q) = q\alpha$), then we have $\alpha_D(q) = \beta_D(q) = q\alpha$.*

Proof. As $\varphi_q := p^{-\alpha_D(q)} \phi^q$ is a σ^q -linear endomorphism of M , the Newton polygon slopes of φ_q are on one hand non-negative and on the other hand are all equal to $q\alpha - \alpha_D(q)$. Thus $\alpha_D(q) \leq q\alpha$. If $q\alpha = \alpha_D(q)$, then all the Newton polygon slopes of $\varphi_q: M \rightarrow M$ are 0 and therefore we have $\varphi_q(M) = M$. This implies that $\beta_D(q) = \alpha_D(q) = q\alpha$. The part involving $\beta_D(q)$ is proved in the same way but working with $p^{\beta_D(q)} \phi^{-q}$. \square

4.2.4. Lemma. (a) *If D is either F -cyclic or special, then D is also quasi-special.*

(b) *The class $\mathcal{Q}_{c,d}$ of isomorphism classes of quasi-special p -divisible groups of codimension c and dimension d over k , is a finite set.*

Proof. Each isoclinic special p -divisible group over k is isoclinic quasi-special. Each F -cyclic p -divisible group over k is a direct sum of F -circular p -divisible groups over k . Based on the last two sentences, it suffices to prove (a) in the case when D is F -circular. Let π be an r -cycle of J_r such that D is isomorphic to C_π . We have $\phi_\pi^r(M) = p^d M$ and therefore C_π is isoclinic quasi-special of Newton polygon slope $\frac{d}{r}$. Thus (a) holds.

To prove (b) it suffices to show that for all pairs $(c, d) \in (\mathbb{N} \cup \{0\})^2$ with $c + d > 0$, the class $\mathcal{J}_{c,d}$ of isomorphism classes of isoclinic quasi-special p -divisible groups of codimension c and dimension d over k , is a finite set. We assume that D is isoclinic quasi-special. Then we have $\phi^r(M) = p^d M$. Therefore $\varphi_r := p^{-d} \phi^r: M \rightarrow M$ is a σ^r -linear automorphism of M . Let $M_{W(\mathbb{F}_{p^r})} := \{x \in M \mid \varphi_r(x) = x\}$. We have $M_{W(\mathbb{F}_{p^r})} \otimes_{W(\mathbb{F}_{p^r})} W(k) = M$. Moreover $\phi(M_{W(\mathbb{F}_{p^r})}) \subseteq M_{W(\mathbb{F}_{p^r})}$. Therefore the Dieudonné module (M, ϕ) is definable over the finite field \mathbb{F}_{p^r} . Thus every isoclinic quasi-special p -divisible group of codimension c and dimension d over k has a Dieudonné module over k which (i) is isomorphic to $(M, g\phi)$ for a suitable element $g \in \mathrm{GL}_M(W(k))$, and (ii) it is definable over \mathbb{F}_{p^r} .

Let $M = F^1 \oplus F^0$ be a direct decomposition such that F^1/pF^1 is the kernel of ϕ modulo p . We have $\phi(\frac{1}{p}F^1 + F^0) = M$. Thus the cocharacter $\mu: \mathbb{G}_m \rightarrow \mathrm{GL}_M$ that fixes F^0 and that acts on F^1 via the inverse of the identity character of \mathbb{G}_m , is a Hodge cocharacter of (M, ϕ, GL_M) in the sense of [Va1, Section 2.2.1 (d)]. Thus the triple (M, ϕ, GL_M) is a latticed F -isocrystal with a group over k for which the W -condition of loc. cit. holds. From the Atlas Principle applied to (M, ϕ, GL_M) and to an emphasized family of tensors indexed by the empty set (see [Va1, Theorem 5.2.3]), we get that the set of isomorphism classes of Dieudonné modules over k which are of the form $(M, g\phi)$ with $g \in \mathrm{GL}_M(W(k))$ and which are definable over the finite field \mathbb{F}_{p^r} is finite. From this and the classical Dieudonné theory, we get that the class $\mathcal{J}_{c,d}$ is a finite set. \square

4.3. Proposition. *We assume that D is isoclinic. Then the following six properties hold:*

- (a) *we have $\ell_D = \max\{\delta_D(q) \mid q \in \mathbb{N}\}$;*
- (b) *if $\alpha := \frac{d}{r}$, then we have $\lim_{q \rightarrow \infty} \frac{\beta_D(q)}{q} = \lim_{q \rightarrow \infty} \frac{\alpha_D(q)}{q} = \alpha$;*
- (c) *if M_0 (resp. \tilde{M}_0) is the $W(k)$ -submodule of M generated by elements fixed by $\varphi_{r_1} := p^{-d_1} \phi^{r_1}$ (resp. by $\varphi_r := p^{-d} \phi^r$), then u_D (resp. v_D) is finite and it is the smallest non-negative integer such that p^{u_D} (resp. p^{v_D}) annihilates M/M_0 (resp. M/\tilde{M}_0);*
- (d) *if M_1 (resp. \tilde{M}_1) is the smallest $W(k)$ -submodule of $M[\frac{1}{p}]$ which is generated by elements fixed by φ_{r_1} (resp. by φ_r) and which contains \tilde{M} , then \tilde{u}_D (resp. \tilde{v}_D) is finite and it is the smallest non-negative integer such that $p^{\tilde{u}_D}$ (resp. $p^{\tilde{v}_D}$) annihilates M_1/M (resp. \tilde{M}_1/M);*
- (e) *we have $u_D = \tilde{u}_D$ (resp. $v_D = \tilde{v}_D$);*
- (f) *we have $u_D \leq \ell_D$.*

Proof. We prove (a). The $W(k)$ -span of endomorphisms of (M, ϕ) is $O = O_0$. The number ℓ_D is the smallest number such that $p^{\ell_D} \mathrm{End}(M) \subseteq O \subseteq \mathrm{End}(M)$,

cf. Example 1.2.2. As $O = \bigcap_{q \in \mathbb{N} \cup \{0\}} \phi^q(\text{End}(M)) = \bigcap_{q \in \mathbb{N} \cup \{0\}} \text{End}(\phi^q(M)) = \bigcap_{q \in \mathbb{N} \cup \{0\}} \text{End}(p^{-\beta_D(q)} \phi^q(M))$, ℓ_D is the smallest (non-negative) integer such that we have $p^{\ell_D} \text{End}(M) \subseteq \text{End}(p^{-\beta_D(q)} \phi^q(M))$ for all $q \in \mathbb{N}$. Thus from (8a) and (8b) we get that ℓ_D is the smallest integer which is greater than or equal to $\delta_D(q)$ for all $q \in \mathbb{N}$. From this (a) follows.

We prove (b). From (a) we get that $\delta_D(q) = \beta_D(q) - \alpha_D(q) \leq \ell_D$. Thus $\frac{\alpha_D}{q} \leq \alpha \leq \frac{\beta_D}{q} \leq \frac{\alpha_D}{q} + \frac{\ell_D}{q}$, cf. Lemma 4.2.3 (a). From these inequalities we get that (b) holds.

We will prove (c) only for M_0 as the case of \tilde{M}_0 is argued in the same manner. We have $M_0 = \bigcap_{n \in \mathbb{N} \cup \{0\}} \varphi_{r_1}^n(M)$. As $\varphi_{r_1}^n(M) = p^{-d_1 n} \phi^{r_1 n}(M)$, from (8a) and (8b) we get that $p^{\beta_D(r_1 n) - d_1 n} M \subseteq \varphi_{r_1}^n(M)$ and $p^{\beta_D(r_1 n) - d_1 n - 1} M \not\subseteq \varphi_{r_1}^n(M)$. Thus the smallest non-negative number s such that we have $p^s M \subseteq \varphi_{r_1}^n(M)$ for all $n \in \mathbb{N}$ is $\sup\{0, \beta_D(r_1 n) - d_1 n \mid n \in \mathbb{N}\}$ and therefore it is u_D . Thus (c) holds.

We will prove (d) only for M_1 as the case of \tilde{M}_1 is argued in the same manner. The $W(k)$ -submodule M_1^* of M^* is the largest $W(k)$ -submodule of M^* generated by elements fixed by $p^{-d_1} \phi^{r_1}$. As $(M^*, p\phi)$ is the Dieudonné module of D^t , the analogue of $p^{-d_1} \phi^{r_1}$ for D^t is the σ^{r_1} -linear automorphism $p^{-c_1} (p\phi)^{r_1} = p^{-d_1} \phi^{r_1}$ of $M^*[\frac{1}{p}]$. Thus from (c) applied to D^t , we get that u_{D^t} is the smallest non-negative integer with the property that $p^{u_{D^t}}$ annihilates M^*/M_1^* . As $\tilde{u}_D = u_{D^t}$ (see Fact 4.2.2 (c)) and as the $W(k)$ -modules M_1/M and M^*/M_1^* are isomorphic, we get that \tilde{u}_D is the smallest non-negative integer such that $p^{\tilde{u}_D}$ annihilates M_1/M . Thus (d) holds.

We will prove (e) for u_D and \tilde{u}_D as the case of v_D and \tilde{v}_D is argued in the same manner. As $p^{\tilde{u}_D} M_1 \subseteq M$ and as $p^{\tilde{u}_D} M_1$ is $W(k)$ -generated by elements fixed by φ_{r_1} , we have $p^{\tilde{u}_D} M_1 \subseteq M_0$. Thus $p^{\tilde{u}_D}$ annihilates M/M_0 and therefore $\tilde{u}_D \geq u_D$, cf. (c). A similar argument shows that $M_1 \subseteq p^{-u_D} M_0$ and that $\tilde{u}_D \leq u_D$. Thus $u_D = \tilde{u}_D$ i.e., (e) holds.

We prove (f). Each endomorphism of (M, ϕ) maps M_0 to M_0 . Thus $O = O_0 \subseteq \text{End}(M_0)$. But due to (c), the smallest number $s \in \mathbb{N} \cup \{0\}$ such that $p^s \text{End}(M) \subseteq \text{End}(M_0)$ is u_D . As $p^{\ell_D} \text{End}(M) \subseteq O \subseteq \text{End}(M_0)$, we get that $u_D \leq \ell_D$. Thus (f) holds. \square

4.3.1. Remark. We have $M_0 \subseteq \tilde{M}_0 \subseteq M$ and $M \subseteq \tilde{M}_1 \subseteq M_1$. Thus $v_D = \tilde{v}_D \leq u_D = \tilde{u}_D$. The $W(k)$ -submodule M_0 (resp. \tilde{M}_0) of M is the largest one with the property that (M_0, ϕ) (resp. (\tilde{M}_0, ϕ)) is the Dieudonné module of an isoclinic special (resp. isoclinic quasi-special) p -divisible group over k . Thus we call $u_D = \tilde{u}_D$ (resp. $v_D = \tilde{v}_D$) the *Manin height* (resp. the *Manin quasi-height*) of D , cf. [Ma, Chapter III, Section 2]. Similarly, the $W(k)$ -submodule M_1 (resp. \tilde{M}_1) of $M[\frac{1}{p}]$ is the smallest one with the properties that it contains M and that (M_1, ϕ) (resp. (\tilde{M}_1, ϕ)) is the Dieudonné module of an isoclinic special (resp. isoclinic quasi-special) p -divisible group over k .

4.4. Proposition. *We assume that $D = \prod_{i \in I} D_i$ is a product of at least two non-trivial isoclinic p -divisible groups over k . Then the following three properties hold:*

- (a) *for all $i, j \in I$ with $i \neq j$, we have $\ell_{D_i, D_j} = \ell_{D_j, D_i}$;*
- (b) *we have $\ell_{D_i} = \ell_{D_i, D_i}$ and $\ell_D = \max\{\varepsilon_D, \ell_{D_i}, \ell_{D_i, D_j} \mid i \in I, j \in I \setminus \{i\}\}$;*
- (c) *if $i, j \in I$ with $i \neq j$ and if the Newton polygon slope α_i of D_i is less than or equal to the Newton polygon slope α_j of D_j , then we have $\ell_{D_i, D_j} = \max\{0, \beta_{D_i}(q) - \alpha_{D_j}(q) \mid q \in \mathbb{N}\}$.*

Proof. Let $M = \bigoplus_{i \in I} M_i$ be the direct sum decomposition such that (M_i, ϕ) is the Dieudonné module of D_i . Let \mathcal{B}_i be a $W(k)$ -basis for M_i . Let $\mathcal{B} := \cup_{i \in I} \mathcal{B}_i$. For $x, y \in \mathcal{B}$, let $\ell(x, y) \in \mathbb{N} \cup \{0\}$ be defined as in Scholium 3.5.1. We have $\ell_{D_i, D_j} = \max\{\ell(y, x) \mid x \in \mathcal{B}_i, y \in \mathcal{B}_j\}$, cf. the very definitions. Thus (a) is a particular case of Formula (6b). As D_i is isoclinic, we have $\varepsilon_{D_i} = 0$. Thus (b) is a particular case of Formulas (6a), (5a), (5b), and (5c). We prove (c). Due to (8a), for all $q \in \mathbb{N}$ we have

$$\begin{aligned} \text{Hom}(\phi^q(M_i), \phi^q(M_j)) &\subseteq \text{Hom}(p^{\beta_{D_i}(q)} M_i, p^{\alpha_{D_j}(q)} M_j) \\ &= p^{\alpha_{D_j}(q) - \beta_{D_i}(q)} \text{Hom}(M_i, M_j). \end{aligned}$$

From this and making use of (8a) and (8b) we get that there exist a direct summand of $\text{Hom}(\phi^q(M_i), \phi^q(M_j))$ which is a direct summand of $p^{\alpha_{D_j}(q) - \beta_{D_i}(q)} \text{Hom}(M_i, M_j)$ as well. Thus the smallest number $s \in \mathbb{N} \cup \{0\}$ with the property that for all $q \in \mathbb{N}$ the $W(k)$ -module $p^s \phi^q(\text{Hom}(M_i, M_j))$ is included in $\text{Hom}(M_i, M_j)$, is $\max\{0, \beta_{D_i}(q) - \alpha_{D_j}(q) \mid q \in \mathbb{N}\}$. From this and the rule (i) of the Definition 4.1 (c), we get that (c) holds. \square

4.4.1. Example. We assume that D is isoclinic and that $d < r \leq 2d$. Thus $\alpha := \frac{d}{r} \in \mathbb{Q} \cap [\frac{1}{2}, 1)$. For $q \in \mathbb{N}$ we have $\beta_{D^\vee}(q) - \alpha_D(q) = q - 2\alpha_D(q)$, cf. Fact 4.2.2 (b). From this and Proposition 4.4 (c) we get that $\ell_{D^\vee, D} = \max\{0, q - 2\alpha_D(q) \mid q \in \mathbb{N}\}$. As $\varepsilon_D = 0$, from Proposition 4.4 (a) and (b) we get that $\ell_{D \oplus D^\vee} = \max\{\ell_D, \ell_{D^\vee}, \ell_{D^\vee, D}\}$. As $\ell_D = \ell_{D^\vee} = \max\{\delta_D(q) \mid q \in \mathbb{N}\}$ (cf. Fact 4.2.1 and Proposition 4.3 (a)), we conclude that

$$\ell_{D \oplus D^\vee} = \max\{\delta_D(q), q - 2\alpha_D(q) \mid q \in \mathbb{N}\}.$$

4.5. Proof of 1.4.3. We assume that $D = \prod_{i \in I} D_i$ is a product of at least two non-trivial isoclinic p -divisible groups over k . Let $M = \bigoplus_{i \in I} M_i$ be the direct sum decomposition defined by the product decomposition $D = \prod_{i \in I} D_i$. As $n_D = \ell_D$ and as for $i \in I$ we have $n_{D_i} = \ell_{D_i}$ (cf. Corollary 1.4.2), based on Proposition 4.4 (a) and (b), to prove Proposition 1.4.3 it suffices to show that for all $i, j \in I$ with $i \neq j$ we have

$$\ell_{D_i, D_j} \leq \max\{0, \ell_{D_i} + \ell_{D_j} - 1\}. \quad (9)$$

As $\ell_{D_i, D_j} = \ell_{D_j, D_i}$ (see Proposition 4.4 (a)), to check the inequality (9) we can assume that $\alpha_i \leq \alpha_j$. We have $\alpha_{D_i}(q) \leq q\alpha_i \leq q\alpha_j \leq \beta_{D_j}(q)$, cf. Lemma 4.2.3 (a). Thus $\alpha_{D_i}(q) \leq \beta_{D_j}(q)$. Based on Proposition 4.4 (c), to prove the inequality (9) it suffices to show that for all $q \in \mathbb{N}$ we have $\beta_{D_i}(q) - \alpha_{D_j}(q) \leq \max\{0, \ell_{D_i} + \ell_{D_j} - 1\}$. We have $\delta_{D_i}(q) + \delta_{D_j}(q) \leq \ell_{D_i} + \ell_{D_j}$, cf. Proposition 4.3 (a). From this and the inequality $\alpha_{D_i}(q) \leq \beta_{D_j}(q)$ we get:

$$\begin{aligned} \beta_{D_i}(q) - \alpha_{D_j}(q) &= \delta_{D_i}(q) + \alpha_{D_i}(q) + \delta_{D_j}(q) - \beta_{D_j}(q) \\ &\leq \ell_{D_i} + \ell_{D_j} + \alpha_{D_i}(q) - \beta_{D_j}(q) \leq \ell_{D_i} + \ell_{D_j}. \end{aligned} \quad (10)$$

If we have an equality $\beta_{D_i}(q) - \alpha_{D_j}(q) = \ell_{D_i} + \ell_{D_j}$, then $\beta_{D_j}(q) = \alpha_{D_i}(q) = q\alpha_i = q\alpha_j$ and therefore also $\beta_{D_i}(q) = q\alpha_i$ and $\alpha_{D_j}(q) = q\alpha_j = q\alpha_i$ (cf. Lemma 4.2.3 (b)). Thus the assumption that $\beta_{D_i}(q) - \alpha_{D_j}(q) = \ell_{D_i} + \ell_{D_j}$ implies that $\beta_{D_i}(q) - \alpha_{D_j}(q) = 0 = \ell_{D_i} + \ell_{D_j} \leq \max\{0, \ell_{D_i} + \ell_{D_j} - 1\}$. From this and (10) we get that the inequality $\beta_{D_i}(q) - \alpha_{D_j}(q) \leq \max\{0, \ell_{D_i} + \ell_{D_j} - 1\}$ always holds; therefore the inequality (9) holds. This ends the proof of Proposition 1.4.3. \square

4.6. Proof of 1.5.2. Let $D = \prod_{i \in I} D_i$ be a product decomposition into isoclinic quasi-special p -divisible groups over k . For $i \in I$, let c_i and d_i be the codimension and the dimension (respectively) of D_i , and let $\alpha_i := \frac{d_i}{r_i}$. Let $M = \bigoplus_{i \in I} M_i$ be the direct sum decomposition such that (M_i, ϕ) is the Dieudonné module of D_i . As each D_i is isoclinic, we have $n_{D_i} = \ell_{D_i}$ and $n_D = \ell_D$ (cf. Corollary 1.4.2). For $i \in I$ we have $\phi^{r_i}(M_i) = p^{d_i} M_i$, cf. Definition 1.5.1 (e). Let $m'_i \in \mathbb{N}$ be the greatest divisor of $\text{g.c.d.}\{c_i, d_i\}$ such that for $(c_{i2}, d_{i2}, r_{i2}) := (\frac{c_i}{m'_i}, \frac{d_i}{m'_i}, \frac{r_i}{m'_i})$ we have $\phi^{r_{i2}}(M_i) = p^{d_{i2}} M_i$. This identity implies that

(i) we have $\alpha_{D_i}(r_{i2}) = \beta_{D_i}(r_{i2}) = d_{i2}$ and for all $q \in \mathbb{N}$ we have $\alpha_{D_i}(q + r_{i2}) = \alpha_{D_i}(q) + d_{i2}$ and $\beta_{D_i}(q + r_{i2}) = \beta_{D_i}(q) + d_{i2}$.

From (i) we get that for all $q \in \mathbb{N}$ we have $\delta_{D_i}(q + r_{i2}) = \delta_{D_i}(q)$. From this and Proposition 4.3 (a) applied to D_i , we get that

(ii) $n_{D_i} = \ell_{D_i} = \max\{\delta_{D_i}(q) \mid q \in \mathbb{N}\} = \max\{\delta_{D_i}(q) \mid q \in \{1, \dots, r_{i2}\}\}$.

As the function $\beta_{D_i}(\ast)$ defined for $\ast \in \mathbb{N}$ is increasing, for all $q \in \{1, \dots, r_{i2}\}$ we have $\delta_{D_i}(q) \leq \beta_{D_i}(q) \leq \beta_{D_i}(r_{i2}) = d_{i2}$. From this and (ii) we get that $n_{D_i} = \ell_{D_i} \leq d_{i2}$. It is easy to see that the p -divisible group D_i^t is isoclinic quasi-special and that the analogue of the triple (r_{i2}, d_{i2}, c_{i2}) for it is (r_{i2}, c_{i2}, d_{i2}) . Thus we have $n_{D_i^t} \leq c_{i2}$. As $n_{D_i} = n_{D_i^t}$ (see Fact 4.2.1), we have $n_{D_i} \leq c_{i2}$. Thus

$$n_{D_i} = \ell_{D_i} \leq \min\{c_{i2}, d_{i2}\} \leq \min\{c_i, d_i\}. \quad (11a)$$

This proves Theorem 1.5.2 if $D = D_i$ i.e., if $I = \{i\}$.

We assume that I has at least two elements. From Proposition 1.4.3 we get that $n_D = \ell_D \leq \max\{1, n_{D_i} + n_{D_j} \mid i \in I, j \in I \setminus \{i\}\}$. From this and (11a) we get that

$$n_D \leq \max\{1, \min\{c_{i2} + c_{j2}, d_{i2} + d_{j2}\} \mid i, j \in I, i \neq j\}. \quad (11b)$$

As $c_{i2} + c_{j2} \leq c$ and $d_{i2} + d_{j2} \leq d$, we have $\min\{c_{i2} + c_{j2}, d_{i2} + d_{j2}\} \leq \min\{c, d\}$. From this and (11b) we get that $n_D = \ell_D \leq \max\{1, \min\{c, d\}\}$. But if $\min\{c, d\} = 0$ (i.e., if $cd = 0$), then $n_D = \ell_D = 0$. Thus, regardless of what the product cd is, we have $n_D = \ell_D \leq \min\{c, d\}$. This ends the proof of Theorem 1.5.2. \square

4.6.1. Scholium. Let π be a permutation of $J_r = \{1, \dots, r\}$. Let o be the order of π . We assume that (D, ϕ) is (C_π, ϕ_π) ; thus D is F -cyclic and therefore (cf. Lemma 4.2.4 (a)) quasi-special. We will translate the property 4.6 (ii) and Proposition 4.4 (a) and (b) in terms only of the permutation π . Let $\pi = \prod_{i \in I} \pi_i$ be the product decomposition of the permutation π into cycles. As in Section 1.5, we write $\pi_i = (e_{s_1}, \dots, e_{s_{r_i}})$ for some number $r_i \in \mathbb{N}$ (which can be 1). Let M_i be the $W(k)$ -span of $\{e_{s_1}, \dots, e_{s_{r_i}}\}$. We have $pM_i \subseteq \phi_\pi(M_i) \subseteq M_i$, cf. the definition of ϕ_π . Thus we have a direct sum decomposition $(M, \phi_\pi) = \bigoplus_{i \in I} (M_i, \phi_\pi)$ of Dieudonné modules. Let $D = \prod_{i \in I} D_i$ be the product decomposition that corresponds to the direct sum decomposition $(M, \phi_\pi) = \bigoplus_{i \in I} (M_i, \phi_\pi)$. Each p -divisible group D_i is F -circular and quasi-special. Let $c_i, d_i = r_i - c_i, \alpha_i = \frac{d_i}{r_i} \in \mathbb{N} \cup \{0\}$ be as in Section 1.5.

Due to the property 4.6 (i), the difference $\delta_{D_i}(q) = \beta_{D_i}(q) - \alpha_{D_i}(q)$ depends only on q modulo o . For $s \in J_r$ and $q \in \{1, \dots, o\}$, let $\eta_q(s) \in \mathbb{N} \cup \{0\}$ be such that we have $\phi_\pi(e_s) = p^{\eta_q(s)} e_{\pi^q(s)}$. Thus $\eta_q(s)$ is the number of elements of the sequence $e_s, \pi(e_s), \dots, \pi^{q-1}(e_s)$ that belong to the set $\{e_1, \dots, e_d\}$. We have

$$\alpha_{D_i}(q) = \min\{\eta_q(s_j) \mid j \in \{1, \dots, r_i\}\} \quad (12a)$$

and

$$\beta_{D_i}(q) = \max\{\eta_q(s_j) \mid j \in \{1, \dots, r_i\}\}. \quad (12b)$$

The $W(k)$ -basis $\mathcal{B} = \{e_1, \dots, e_r\}$ for M is a disjoint union of $W(k)$ -basis for M_i 's. We consider the standard $W(k)$ -basis $\{e_s \otimes e_t^* \mid s, t \in J_r\}$ for $\text{End}(M)$ defined by \mathcal{B} . We have $\phi_\pi^q(e_s \otimes e_t^*) = p^{\eta_q(s) - \eta_q(t)} e_{\pi^q(s)} \otimes e_{\pi^q(t)}^*$. If $\eta_o(s) > \eta_o(t)$ (resp. $\eta_o(s) = \eta_o(t)$ or $\eta_o(s) < \eta_o(t)$), then $e_s \otimes e_t^*$ belongs to L_+ (resp. to L_0 or L_-) and therefore the number $\ell(e_s, e_t)$ defined in Scholium 3.5.1 is $\max\{0, \eta_q(t) - \eta_q(s) \mid q \in \{1, \dots, o\}\}$ (resp. is $\max\{0, \eta_q(s) - \eta_q(t) \mid q \in \{1, \dots, o\}\}$). From Formula (6a) we get that

$$\ell_{C_\pi} = \max\{\varepsilon_{C_\pi}, \ell(e_s, e_t) \mid s, t \in J_r\}. \quad (12c)$$

4.6.2. Example. We assume that $c = d = 8$; thus $r = 16$. Let $\pi = \pi_1 \pi_2$, where $\pi_1 = (9 \ 10 \ 5 \ 11 \ 12 \ 6 \ 7 \ 8)$ and $\pi_2 = (1 \ 2 \ 13 \ 3 \ 4 \ 14 \ 15 \ 16)$ are

8-cycles. We have $o = 8$. Let $D = C_\pi = D_1 \oplus D_2$ be the product decomposition corresponding to the cycle decomposition $\pi = \pi_1 \pi_2$. All Newton polygon slopes of D_1 and D_2 are $\frac{1}{2}$ and it is easy to see that D_1^t is isomorphic to D_1 . We have $(\delta_{D_1}(1), \dots, \delta_{D_1}(8)) = (1, 2, 2, 2, 2, 2, 1, 0)$; thus $n_{D_1} = 2$ (cf. property 4.6(ii)). From Fact 4.2.1 we get that $n_{D_2} = 2$. We have $(\eta_1(9), \dots, \eta_8(9)) = (0, 0, 1, 1, 1, 2, 3, 4)$ and $(\eta_1(1), \dots, \eta_8(1)) = (1, 2, 2, 3, 4, 4, 4, 4)$. Therefore $(\eta_1(9) - \eta_1(1), \dots, \eta_8(9) - \eta_8(1)) = (-1, -2, -1, -2, -3, -2, -1, 0)$ and thus $\ell(e_9, e_1) = \ell(e_1, e_9) = 3$. This implies that $n_D = \ell_D \geq 3$. From Proposition 1.4.3 we get that $n_D \leq 3$. Thus $n_{D_1 \oplus D_2} = n_D = 3 = n_{D_1} + n_{D_2} - 1$.

Plenty of similar examples can be constructed in which the identity $n_{D_1 \oplus D_2} = n_{D_1} + n_{D_2} - 1$ holds and D_1 and D_2 are isoclinic of equal height and different dimension.

4.7. Proof of 1.4.4. The Dieudonné module of \tilde{D} is (\tilde{M}, ϕ) , where \tilde{M} is a $W(k)$ -submodule of M which contains $p^\kappa M$. Let $\tilde{O} = \tilde{O}_+ \oplus \tilde{O}_0 \oplus \tilde{O}_-$ be the level module of (\tilde{M}, ϕ) . If D and \tilde{D} are ordinary, then Proposition 1.4.4 is trivial. Thus to prove Proposition 1.4.4, we can assume that D and \tilde{D} are not ordinary; thus from Remark 4.1.1 we get that ℓ_D (resp. $\ell_{\tilde{D}}$) is the smallest natural number such that we have $p^{\ell_D} \text{End}(M) \subseteq O$ (resp. we have $p^{\ell_{\tilde{D}}} \text{End}(\tilde{M}) \subseteq \tilde{O}$). As $p^\kappa M \subseteq \tilde{M} \subseteq M$, we have

$$p^{2\kappa} \text{End}(M) \subseteq p^\kappa \text{End}(\tilde{M}) \subseteq \text{End}(M). \quad (13)$$

For $q \in \mathbb{N}$ we have $\phi^q(p^\kappa \tilde{O}_+) \subseteq p^\kappa \tilde{O}_+ \subseteq \text{End}(\tilde{M}) \cap L_+$. As $p^\kappa \tilde{O}_+ \subseteq \text{End}(M)$ (cf. (13)), we get that $p^\kappa \tilde{O}_+ \subseteq O_+$. A similar argument shows that $p^\kappa \tilde{O}_0 \subseteq O_0$ and $p^\kappa \tilde{O}_- \subseteq O_-$. Thus $p^\kappa \tilde{O} \subseteq O$. From this, the inclusion $p^{\ell_{\tilde{D}}} \text{End}(\tilde{M}) \subseteq \tilde{O}$, and (13) we get that

$$p^{2\kappa + \ell_{\tilde{D}}} \text{End}(M) \subseteq p^{\kappa + \ell_{\tilde{D}}} \text{End}(\tilde{M}) \subseteq p^\kappa \tilde{O} \subseteq O \subseteq \text{End}(M).$$

Thus $\ell_D \leq 2\kappa + \ell_{\tilde{D}}$. Based on this inequality, Proposition 1.4.4 follows from Corollary 1.4.2. This ends the proof of Proposition 1.4.4. \square

4.7.1. Example. We assume that $c = d$. We have $r = 2d$. Let $\pi := (12 \dots r)$; its cyclic decomposition is $\pi = \pi_i$ (with i as an index). As $\phi_\pi^r(M) = p^{\frac{r}{2}}(M)$, the F -circular p -divisible group C_π is supersingular. If $d \geq 2$, then $\phi^2(M) \neq pM$ and therefore C_π is not special. As $\phi_\pi^d(e_1) = p^d e_{d+1}$ and $\phi_\pi^d(e_{d+1}) = e_1$, we have $\alpha_D(d) = 0$ and $\beta_D(d) = d$. This implies $\delta_D(d) = d$ and therefore from Proposition 4.3(a) we get that $n_{C_\pi} = \ell_{C_\pi} \geq d$. As $n_{C_\pi} \leq d$ (cf. Theorem 1.5.2), we have $n_{C_\pi} = d$. See [NV, Example 3.3] for a simpler proof that $n_{C_\pi} = d$ (in loc. cit. C_π is denoted as C_d). Let E be a supersingular p -divisible group over k of height 2. From [NV, Remark 2.6 and Example 3.3] we get that the smallest number $\kappa \in \mathbb{N} \cup \{0\}$ such that we have an isogeny $C_\pi \twoheadrightarrow E^d$ is $\kappa := \lceil \frac{d-1}{2} \rceil$. It is well known that E^d is uniquely determined up to isomorphism by $E^d[p]$ (for instance,

see [NV, Scholium 2.3] or see Formula (12c) applied to the minimal permutation $(1\ d+1)\dots(d\ r)$ of J_r). Thus $n_{E^d} = 1$. If d is odd, then $\kappa = \frac{d-1}{2}$ and therefore $n_{C_\pi} = d = n_{E^d} + 2\kappa$. This implies that in general, Proposition 1.4.4 is optimal.

4.7.2. Example. We assume that $r > 0$ and that D is isoclinic. Let M_0 and \tilde{M}_0 be as in Proposition 4.3 (c). Let D_0 and \tilde{D}_0 be the p -divisible groups over k whose Dieudonné modules are isomorphic to (M_0, ϕ) and (\tilde{M}_0, ϕ) (respectively), cf. Remark 4.3.1. To the inclusions $M_0 \subseteq M$ and $\tilde{M}_0 \subseteq M$ correspond isogenies $D \twoheadrightarrow D_0$ and $D \twoheadrightarrow \tilde{D}_0$ whose kernels are annihilated by $p^{u_D} = p^{\tilde{u}_D}$ and $p^{v_D} = p^{\tilde{v}_D}$ (respectively), cf. Proposition 4.3 (c). Let $j_D := n_{D_0}$ and $\tilde{j}_D := n_{\tilde{D}_0}$. From Propositions 1.4.4 and 4.3 (f) we get that

$$u_D \leq n_D = \ell_D \leq \min\{j_D + 2u_D, \tilde{j}_D + 2v_D\}. \quad (14a)$$

If (c_1, d_1, r_1) is as in Definition 4.1 (b), then $j_D \leq \min\{c_1, d_1\}$ (cf. (11a)). From this and (14a) we get

$$u_D \leq n_D \leq 2u_D + \min\{c_1, d_1\}. \quad (14b)$$

5. On Main Theorem B

In Section 5.1 we prove Main Theorem B. Sections 5.2 and 5.3 present two applications of Main Theorem B. For instance, Theorem 5.3 presents applications to extensions between two minimal p -divisible groups over k . We recall that (M, ϕ) is the Dieudonné module of D .

5.1. The proof of Main Theorem B. If $n_D \leq 1$, then $D[p]$ is minimal (cf. Definition 1.5.1 (d)). If $D[p]$ is minimal, then there exists a p -divisible group \tilde{D} over k such that $n_{\tilde{D}} \leq 1$ and $\tilde{D}[p]$ is isomorphic to $D[p]$; the codimension and the dimension of \tilde{D} are c and d (respectively) and thus from the very definition of $n_{\tilde{D}}$ we get that D is isomorphic to \tilde{D} and therefore that we have $n_D = n_{\tilde{D}} \leq 1$. Thus we have $n_D \leq 1$ if and only if $D[p]$ is minimal. As $n_D \leq \ell_D$ (see Corollary 1.4.2), 1.6 (a) implies 1.6 (b). Thus to end the proof of Main Theorem B, it suffices to show that 1.6 (b) implies 1.6 (c) and that 1.6 (c) implies 1.6 (a).

5.1.1. On 1.6 (b) \Rightarrow 1.6 (c). Let $\phi_1, \vartheta_1: M/pM \rightarrow M/pM$ be the reductions modulo p of $\phi, \vartheta: M \rightarrow M$. In [Kr] (see also [Oo1, Section (2.3) and Lemma (2.4)] and [Mo, Section 2.1]) it is shown that there exists a k -basis $\{b_1, \dots, b_r\}$ for M/pM and a permutation π of $J_r = \{1, \dots, r\}$ such that the following two properties hold:

- (i) if $s \in \{1, \dots, d\}$, then $\phi_1(b_s) = 0$ and $\vartheta_1(b_{\pi(s)}) = b_s$, and
- (ii) if $s \in \{d+1, \dots, r\}$, then $\phi_1(b_s) = b_{\pi(s)}$ and $\vartheta_1(b_{\pi(s)}) = 0$.

Let $\vartheta_\pi := p\phi_\pi^{-1}: M \rightarrow M$; if $s \in \{1, \dots, d\}$, then $\vartheta_\pi(e_{\pi(s)}) = e_s$, and if $s \in \{d+1, \dots, r\}$, then $\vartheta_\pi(e_{\pi(s)}) = pe_s$. Properties (i) and (ii) imply that the k -linear map $M/pM \rightarrow M/pM$ that takes b_s to e_s modulo p , is an isomorphism between $(M/pM, \phi_1, \vartheta_1)$ and the reduction modulo p of $(M, \phi_\pi, \vartheta_\pi)$. This means that $D[p]$ is isomorphic to $C_\pi[p]$, cf. the classical Dieudonné theory. As $n_D \leq 1$, we get that D is isomorphic to C_π .

We check that π is a minimal permutation in the sense of Definition 1.5.1 (b). Let $\pi = \prod_{i \in I} \pi_i$ be the product decomposition of π into cycles. We write $\pi_i = (e_{s_1}, \dots, e_{s_{r_i}})$, where $r_i \in \mathbb{N}$. Let $c_i, d_i = r_i - c_i$, and $\alpha_i = \frac{d_i}{r_i}$ be as in Section 1.5. Let $M_i := \bigoplus_{u=1}^{r_i} W(k)e_{s_u}$. Let $D = \prod_{i \in I} D_i$ be the product decomposition defined by the direct sum decomposition $(M, \phi_\pi) = \bigoplus_{i \in I} (M_i, \phi_\pi)$. Each D_i is an F -circular p -divisible group over k and therefore isoclinic. From Proposition 4.4 (b) we get that $n_{D_i} \leq n_D \leq 1$. But $n_{D_i} = \ell_{D_i} = \max\{\delta_{D_i}(q) \mid q \in \mathbb{N}\}$, cf. Corollary 1.4.2 and Proposition 4.3 (a). From the last two sentences we get that for all $q \in \mathbb{N}$ we have $\delta_{D_i}(q) \in \{0, 1\}$. Thus either $\alpha_{D_i}(q) = \beta_{D_i}(q)$ or $\alpha_{D_i}(q) + 1 = \beta_{D_i}(q)$. If $\alpha_{D_i}(q) = \beta_{D_i}(q)$, then from Lemma 4.2.3 (a) we get that $\alpha_{D_i}(q) = \beta_{D_i}(q) = q\alpha_i$. If $\alpha_{D_i}(q) + 1 = \beta_{D_i}(q)$, then from Lemma 4.2.3 (a) we get that either $(\alpha_{D_i}(q), \beta_{D_i}(q)) = ([q\alpha_i], [q\alpha_i] + 1)$ or $(\alpha_{D_i}(q), \beta_{D_i}(q)) = (q\alpha_i - 1, q\alpha_i)$. But the second possibility is excluded by Lemma 4.2.3 (b). We conclude that in all cases we have $\alpha_{D_i}(q), \beta_{D_i}(q) \in \{[q\alpha_i], [q\alpha_i] + 1\}$. Therefore $p^{[q\alpha_i]+1}M_i \subseteq \phi^q(M_i) \subseteq p^{[q\alpha_i]}M_i$. Thus for each $u \in \{1, \dots, r_i\}$, we have $\phi_\pi^q(e_{s_u}) = p^{[q\alpha_i]+\varepsilon_q(s_u)}e_{\pi^q(s_u)}$ for some number $\varepsilon_q(s_u) \in \{0, 1\}$. As this property holds for all pairs $(q, i) \in \mathbb{N} \times I$, π is a minimal permutation. As D is isomorphic to C_π , we get that D is minimal. Thus 1.6 (b) implies 1.6 (c).

5.1.2. On 1.6 (c) \Rightarrow 1.6 (a). To prove that 1.6 (c) implies 1.6 (a), we can assume that π is a minimal permutation of J_r , that $D = C_\pi$, and that $\phi = \phi_\pi$. Let $\pi = \prod_{i \in I} \pi_i$, $M = \bigoplus_{i \in I} M_i$, and $D = \prod_{i \in I} D_i$ be the decompositions obtained as in Section 5.1.1. For $i \in I$, let $\pi_i = (e_{s_1}, \dots, e_{s_{r_i}})$, $c_i, d_i = r_i - c_i$, and $\alpha_i = \frac{d_i}{r_i}$ be as in Section 1.5. As the permutation π is minimal, for all $u \in \{1, \dots, r_i\}$ we have $\phi_\pi^q(e_{s_u}) = p^{[q\alpha_i]+\varepsilon_q(s_u)}e_{\pi^q(s_u)}$ for some number $\varepsilon_q(s_u) \in \{0, 1\}$. This implies that $p^{[q\alpha_i]+1}M_i \subseteq \phi_\pi^q(M_i) \subseteq p^{[q\alpha_i]}M_i$. Thus

$$\alpha_{D_i}(q), \beta_{D_i}(q) \in \{[q\alpha_i], [q\alpha_i] + 1\}. \quad (15)$$

From (15) and the fact that $\delta_{D_i}(q) \geq 0$, we get that $\delta_{D_i}(q) \in \{0, 1\}$. From this and Proposition 4.3 (a), we get that $n_{D_i} = \ell_{D_i} \leq 1$. If $D = D_i$ (i.e., if $I = \{i\}$), then $\ell_D \leq 1$ and thus 1.6 (a) holds. If I has at least two elements, then from Proposition 1.4.3 we get that $\ell_D = n_D \leq 1$. Thus regardless of what I is, we have $\ell_D \leq 1$. This ends the argument that the implication 1.6 (c) \Rightarrow 1.6 (a) holds. This ends the proof of Main Theorem B. \square

5.2. Corollary. We assume that $\ell_D \leq 2$. Then $n_D = \ell_D$.

Proof. If $n_D \leq 1$, then D is minimal (cf. Main Theorem B) and therefore F -cyclic; thus $n_D = \ell_D$ (cf. Theorem 1.5.2). As $n_D \leq \ell_D \leq 2$, we have $n_D = \ell_D$ even if $n_D = 2$. \square

The next theorem generalizes and refines [Va1, Proposition 4.5.1].

5.3. Theorem. *We assume that we have a short exact sequence $0 \rightarrow D_1 \rightarrow D \rightarrow D_2 \rightarrow 0$ of p -divisible groups over k , with D_1 and D_2 as minimal p -divisible groups.*

- (a) *Then we have $n_D \leq \ell_D \leq 3$.*
- (b) *We assume that $d = c \geq 3$ and that D_1 and D_2 are isoclinic of Newton polygon slopes $\frac{1}{d}$ and $\frac{d-1}{d}$ (respectively). Then $n_D = \ell_D \leq 2$.*

Proof. Let $0 \rightarrow D_1 \rightarrow \tilde{D} \rightarrow D_2 \rightarrow 0$ be the pull forward of our initial short exact sequence via the multiplication by p isogeny $D_1 \twoheadrightarrow D_1$. The kernel of the resulting isogeny $D \twoheadrightarrow \tilde{D}$ is annihilated by p . The p -divisible group $D_{12} := D_1 \oplus D_2$ is minimal. As $\tilde{D}[p]$ is isomorphic to $D_{12}[p]$, from the equivalence between 1.6 (b) and (c) we get that \tilde{D} is isomorphic to D_{12} and that $n_{\tilde{D}} \leq 1$. As \tilde{D} is F -cyclic and thus a direct sum of isoclinic p -divisible groups, we have $n_D \leq \ell_D \leq n_{\tilde{D}} + 2 \leq 3$ (cf. Proposition 1.4.4). Thus (a) holds.

We prove (b). We know that there exists an isogeny $D_{12} \twoheadrightarrow D$ whose kernel K is annihilated by p . We will choose such an isogeny of the smallest degree possible. It is well known that up to isomorphisms, there exists a unique p -divisible group over k of height d and Newton polygon slope $\frac{*}{d}$, where $* \in \{1, d-1\}$ (see [De, Chapter IV, Section 8]). Thus if K has a proper subgroup scheme K_1 (resp. K_2) whose image in D_1 (resp. in D_2) is trivial, then $D'_{12} := D_{12}/K_1$ (resp. $D'_{12} := D_{12}/K_2$) is isomorphic to D_{12} and thus we would get an isogeny $D_{12} \xrightarrow{\sim} D'_{12} \twoheadrightarrow D$ of smaller degree. This implies that the projections of K on D_1 and D_2 are monomorphisms. Thus the codimension and the dimension of K are both at most 1. Based on the last two sentences, as $d \geq 3$ we easily get that K is either trivial or isomorphic to α_p . If K is trivial, then D is minimal and therefore we have $n_D \leq 1$ (in fact we have $n_D = 1$). Thus to prove (b), we can assume that K is isomorphic to α_p . We reached the case when we have isogenies

$$D_{12} \twoheadrightarrow D_{12}/\alpha_p \xrightarrow{\sim} D \twoheadrightarrow D_{12}/(\alpha_p \times_k \alpha_p) = D_1/\alpha_p \times_k D_2/\alpha_p.$$

At the level of Dieudonné modules, this means the following things. Let $N_{12} := \bigoplus_{s=1}^{2d} W(k)e_s$ be a free $W(k)$ -module of rank $r = 2d$. Let $\phi: N_{12} \rightarrow N_{12}$ be the σ -linear endomorphism such that it takes (e_1, \dots, e_d) and (e_{d+1}, \dots, e_{2d}) to $(pe_2, e_3, \dots, e_d, e_1)$ and $(pe_{d+2}, \dots, pe_{2d}, e_{d+1})$ (respectively). We can assume that (N_{12}, ϕ) is the Dieudonné module of $D_1/\alpha_p \times_k D_2/\alpha_p$ (cf. the mentioned uniqueness property). As $D \twoheadrightarrow D_{12}/(\alpha_p \times_k \alpha_p) = D_1/\alpha_p \times_k D_2/\alpha_p$ is an isogeny of kernel α_p and as K maps monomorphically to both D_1 and D_2 , there exists an

invertible element $\gamma \in W(k)$ such that we can identify M with $N_{12} + W(k)(\frac{\gamma}{p}e_1 + \frac{1}{p}e_{d+1})$. Moreover, if $M_{12} := N_{12} + W(k)\frac{1}{p}e_1 + W(k)\frac{1}{p}e_{d+1}$, then (M_{12}, ϕ) is the Dieudonné module of D_{12} .

We check that $\ell_D \leq 2$. We have

$$p^2 \operatorname{End}(M) \subseteq p \operatorname{Hom}(M_{12}, N_{12}) + W(k)[(p\gamma e_1 + pe_{d+1}) \otimes e_1^*] \subseteq \operatorname{End}(M). \quad (16)$$

The latticed F -isocrystal $(\operatorname{Hom}(M_{12}, N_{12}), \phi)$ is isomorphic to $\operatorname{End}(M_{12}, \phi)$ and moreover we have $\ell_{D_{12}} \leq 1$. From this and the fact that $\operatorname{Hom}(M_{12}, N_{12}) \subseteq \operatorname{End}(M)$, we get that O contains $p \operatorname{Hom}(M_{12}, N_{12})$. It is easy to see that for all $q \in \mathbb{N}$ we have $\phi^q((p\gamma e_1 + pe_{d+1}) \otimes e_1^*) \in \operatorname{End}(M)$; like $\phi((p\gamma e_1 + pe_{d+1}) \otimes e_1^*) = (p\sigma(\gamma)e_2 + pe_{d+2}) \otimes e_2^*$, $\phi^2((p\gamma e_1 + pe_{d+1}) \otimes e_1^*) = (p\sigma^2(\gamma)e_3 + p^2e_{d+3}) \otimes e_3^*$, etc. Thus $(p\gamma e_1 + pe_{d+1}) \otimes e_1^* \in O_+ \oplus O_0$, cf. Lemma 2.4. Based on (16) we conclude that $p^2 \operatorname{End}(M) \subseteq O$. Thus $\ell_D \leq 2$. From Corollary 5.2 we get that $n_D = \ell_D \leq 2$. Thus (b) holds. \square

Acknowledgments. We would like to thank the University of Arizona and Binghamton University for providing good conditions during the writing of this paper. We would like to thank M.-H. Nicole for introducing us to Traverso's work on February 2006, for many valuable comments, and for suggesting the ideas of the second paragraph of the proof of Theorem 5.3 (cf. [Ni, Theorem 1.5.2]). We would like to thank the referee for many valuable comments and suggestions.

References

- [De] M. Demazure, *Lectures on p -divisible groups*. Lecture Notes in Math. 302, Springer-Verlag, Berlin 1972. [Zbl 0247.14010](#) [MR 0344261](#)
- [Di] J. Dieudonné, Groupes de Lie et hyperalgèbres de Lie sur un corps de caractéristique $p > 0$. VII. *Math. Ann.* **134** (1957), 114–133. [Zbl 0086.02605](#) [MR 0098146](#)
- [Kr] H. Kraft, *Kommutative algebraische p -Gruppen (mit Anwendungen auf p -divisible Gruppen und abelsche Varietäten)*. Manuscript, Universität Bonn, 1975.
- [Ma] J. I. Manin, The theory of formal commutative groups in finite characteristic. *Russian Math. Surveys* **18** (6) (1963), 1–83. [Zbl 0128.15603](#) [MR 0157972](#)
- [Mo] B. Moonen, Group schemes with additional structures and Weyl group cosets. In *Moduli of abelian varieties* (Texel Island, 1999), Progr. Math. 195, Birkhäuser, Basel, 2001, 255–298. [Zbl 1084.14523](#) [MR 1827024](#)
- [Ni] M.-H. Nicole, Superspecial abelian varieties, theta series and the Jacquet-Langlands correspondence. Ph.D. thesis, McGill University, 2005.
- [NV] M.-H. Nicole and A. Vasiu, Minimal truncations of supersingular p -divisible groups. *Indiana Univ. Math. J.* **56** (6) (2007), 2887–2897. [Zbl 05254004](#) [MR 2375706](#)

- [Oo1] F. Oort, A stratification of a moduli space of abelian varieties. In *Moduli of abelian varieties* (Texel Island, 1999), Progr. Math. 195, Birkhäuser, Basel 2001, 345–416. [Zbl 1052.14047](#) [MR 1827027](#)
- [Oo2] F. Oort, Foliations in moduli spaces of abelian varieties. *J. Amer. Math. Soc.* **17** (2) (2004), 267–296. [Zbl 1041.14018](#) [MR 2051612](#)
- [Oo3] F. Oort, Minimal p -divisible groups. *Ann. of Math. (2)* **161** (2) (2005), 1021–1036. [Zbl 1081.14065](#) [MR 2153405](#)
- [Oo4] F. Oort, Simple p -kernels of p -divisible groups. *Adv. Math.* **198** (1) (2005), 275–310. [Zbl 1093.14064](#) [MR 2183256](#)
- [Tr1] C. Traverso, Sulla classificazione dei gruppi analitici commutativi di caratteristica positiva. *Ann. Scuola Norm. Sup. Pisa (3)* **23** (3) (1969), 481–507. [Zbl 0214.48301](#) [MR 0260757](#)
- [Tr2] C. Traverso, p -divisible groups over fields. *Symposia Math.* **XI** (Convegno di Algebra Commutativa, INDAM, Rome, 1971), Academic Press, London 1973, 45–65. [Zbl 0293.14011](#) [MR 0344262](#)
- [Tr3] C. Traverso, Specializations of Barsotti–Tate groups. *Symposia Math.* **XXIV** (Sympos., INDAM, Rome, 1979), Academic Press, London 1981, 1–21. [Zbl 0466.14016](#) [MR 0619238](#)
- [Va1] A. Vasiu, Crystalline boundedness principle. *Ann. Sci. École Norm. Sup.* **39** (2) (2006), 245–300. [Zbl 1143.14037](#) [MR 2245533](#)
- [Va2] A. Vasiu, Generalized Serre–Tate ordinary theory. Preprint 2008. [arXiv:math/0208216](#)
- [Va3] A. Vasiu, Mod p classification of Shimura F -crystals. Preprint 2006. [arXiv:math/0304030](#)

Received February 15, 2008

Adrian Vasiu, Department of Mathematical Sciences, Binghamton University, Binghamton, NY 13902-6000, U.S.A.

E-mail: adrian@math.binghamton.edu