

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 85 (2010)

Artikel: The $K(1)$ conjecture for a class of Artin groups
Autor: Ellis, Graham / Sköldbberg, Emil
DOI: <https://doi.org/10.5169/seals-130668>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The $K(\pi, 1)$ conjecture for a class of Artin groups

Graham Ellis and Emil Sköldberg*

Abstract. Salvetti constructed a cellular space B_D for any Artin group A_D defined by a Coxeter graph D . We show that B_D is an Eilenberg–Mac Lane space if $B_{D'}$ is an Eilenberg–Mac Lane space for every subgraph D' of D involving no ∞ -edges.

Mathematics Subject Classification (2000). 55P20, 20F36.

Keywords. Artin group, Eilenberg–Mac Lane space, cohomology groups.

1. Introduction

A *Coxeter matrix* is a symmetric $n \times n$ matrix whose entries $m(i, j)$ are either positive integers or the symbol ∞ , with $m(i, j) = 1$ if and only if $i = j$. Such a matrix is represented by an n -vertex labelled graph D (called a *Coxeter graph*) with edge joining vertices i and j if and only if $m(i, j) \geq 3$; the edge is labelled by $m(i, j)$. The *Artin group* A_D is defined to be the group generated by the set of symbols $S = \{x_1, \dots, x_n\}$ subject to relations $(x_i x_j)_{m(i, j)} = (x_j x_i)_{m(i, j)}$ for all $i \neq j$, where $(xy)_m$ denotes the word $xyxyx \dots$ of length m . The *Coxeter group* W_D is the quotient of A_D obtained by imposing additional relations $x^2 = 1$ for $x \in S$.

For each Coxeter graph D there is an interesting finite CW-space B_D arising as a quotient of a union of certain convex polytopes (see Section 2 for precise details). It has fundamental group $\pi_1(B_D) = A_D$ and we have the following.

Conjecture 1. The space B_D is an Eilenberg–Mac Lane space $K(A_D, 1)$.

From work of Squier in the 1980s (published posthumously [16]) one can deduce that the conjecture holds whenever the Coxeter group W_D is finite. (Squier established a free $\mathbb{Z}A_D$ -resolution R_*^D of \mathbb{Z} having the same number of free generators in each degree as the cellular chain complex $C_*(\tilde{B}_D)$. It is clear that R_*^D coincides with $C_*(\tilde{B}_D)$ in degrees ≤ 2 and hence R_*^D is the cellular chain complex of the universal cover of some $K(A_D, 1)$. A detailed analysis suggests that R_*^D is in fact the cellular

*Supported by Marie Curie fellowship HPMD-CT-2001-00079.

chain complex of \tilde{B}_D .) Also, it follows immediately from a result of Appel and Schupp [1, Lemma 6] that the conjecture holds if, for every triple of generators $a, b, c \in S$, the three Artin relators $(ab)_k = (ba)_k$, $(bc)_l = (cb)_l$, $(ac)_m = (ca)_m$ are such that $\frac{1}{k} + \frac{1}{l} + \frac{1}{m} \leq 1$. (In this case B_D is just the standard 2-dimensional CW-space associated to the presentation and Lemma 6 in [1] implies that any element of $\pi_2(B_D)$ would have to be represented by a non-positively curved piecewise euclidean 2-sphere.)

Given a Coxeter graph D we shall say that a subgraph D' is an ∞ -free subgraph if (1) D' is a connected and full subgraph of D ; (2) no edge of D' is labelled by ∞ . (In a full subgraph an edge must be included if its two boundary vertices are present.) Our main result is obtained using a technique of D. E. Cohen [8] and is the following.

Theorem 2. *An Artin group A_D satisfies Conjecture 1 if $A_{D'}$ satisfies Conjecture 1 for every ∞ -free subgraph D' in D .*

For Artin groups satisfying Conjecture 1 the cellular chains of the universal cover \tilde{B}_D yield an explicit small free $\mathbb{Z}A_D$ -resolution from which cohomology calculations can be made. Section 4 gives such a cohomology calculation based on Theorem 2.

To place Theorem 2 in context we mention that there is an alternative statement of Conjecture 1. Every Coxeter group W_D acts canonically as a linear group generated by “reflections” on a real vector space V and properly discontinuously on an open cone $I \subset V$ called the *Tits cone*. Denote by A the set of reflecting hyperplanes of W_D and consider the following subspace of $\mathbb{C} \otimes V = V \oplus \mathbf{i}V$:

$$M(W_D) = I \oplus \mathbf{i}V \setminus \left(\bigcup_{H \in A} H \oplus \mathbf{i}H \right).$$

The group W_D acts freely and properly discontinuously on $M(W)$ and the quotient $N(W_D) = M(W_D)/W_D$ has fundamental group equal to A_D .

Conjecture 3. The space $N(W_D)$ is an Eilenberg–Mac Lane space $K(A_D, 1)$.

Conjecture 3 is known as the $K(\pi, 1)$ -conjecture for Artin groups and is attributed to Arnold, Pham and Thom in [5]. It has been proved in many cases: Deligne [10] proved it for finite W_D ; Hendriks [12] proved it for W_D of large type; Charney and Davis [5] proved it when W_D is 2-dimensional and when W_D is of FC type; Charney and Peifer [7] proved it for W_D of affine type \tilde{A}_n ; Callegaro, Moroni and Salvetti [4] have recently proved it for W_D of affine type \tilde{B}_n .

Salvetti [15] showed that the space B_D is homotopy equivalent to $N(W_D)$ for finite W_D . This homotopy equivalence was extended to arbitrary W_D by Charney and Davis [6]. Conjectures 1 and 3 are thus equivalent and so Theorem 2 can consequently be viewed as a generalisation of the solution to the $K(\pi, 1)$ -conjecture for Artin groups of FC type provided in [5]. (Recall that A_D is said to be of *FC type* if $W_{D'}$ is finite for every ∞ -free subgraph D' in D .)

We would like to thank the referees for helpful comments and references.

2. The space B_D

Let D be a Coxeter graph. Let \bar{x} be the image in W_D of the generator $x \in S \subset A_D$ and set $\bar{S} = \{\bar{x} : x \in S\}$. We say that D is of *finite type* if the Coxeter group W_D is finite.

Assume for the moment that D is of finite type and let $n = |S|$. Then W_D can be realized as a group of orthogonal transformations of \mathbb{R}^n with generators \bar{x} equal to reflections [9]. Let A be the set of hyperplanes corresponding to all the reflections in W_D . For any point e in $\mathbb{R}^n \setminus A$ we denote by P_D the convex hull of the orbit of e under the action of W_D . The face lattice of the n -dimensional convex polytope P_D depends only on the graph D . (To see this, first note that the vertices of P_D are the points $w \cdot e$ for $w \in W_D$ and that there is an edge between $w \cdot e$ and $w' \cdot e$ if and only if $w^{-1}w' \in \bar{S}$. Thus the combinatorial type of the 1-skeleton of P_D does not depend on the choice of point e . Furthermore, each vertex of the n -dimensional polytope P_D is incident with precisely n edges; hence P_D is simple and the face lattice of the polytope is determined by the combinatorial type of the 1-skeleton [2].)

Label each edge in P_D by the generating reflection $\bar{x} = w^{-1}w' \in \bar{S}$ determined by the edge's boundary vertices $w \cdot e, w' \cdot e$. Define the *length* of an element g in W_D to be the shortest length of a word in the generators representing it. It is possible to orient each edge in P_D so that its initial vertex gv and final vertex $g'v$ are such that the length of g is less than the length of g' . With this edge orientation the 1-skeleton coincides with the Hasse diagram for the weak Bruhat order on W_D . Each k -face in P_D has a least vertex in the weak Bruhat order. Reading the edge labels along the boundary of any 2-face, starting at the least vertex and using edge orientations to determine exponents ± 1 , yields a relator $(xy)_{m(i,j)}(yx)_{m(i,j)}^{-1}$ of the Artin group A_D . Furthermore, if F is any k -face of P_D , then $V_F = \{w \in W_D : w \cdot e \in F\}$ is a left coset of the *parabolic subgroup* $\langle T \rangle$ of W_D generated by some subset $T \subset \bar{S}$ of size $|T| = k$; this induces an isomorphism between the face lattice of P_D and the poset of cosets $\{w \cdot \langle T \rangle : T \subset \bar{S}, w \in W_D\}$ ordered by inclusion.

The above description of the polytope P_D is well known. (We note that many authors prefer to deal with the dual polytope: since P_D is simple the dual is simplicial.)


The space B_D is obtained from the polytope P_D by isometrically identifying any two cells with similarly labelled 1-skeleta. More precisely, the group W_D acts cellularly on P_D . If a k -face F is mapped to a k -face F' under the action of $w \in W_D$, then there is a unique $w_0 \in W_D$ which maps F to F' in such a way that the least vertex of F maps to the least vertex of F' ; we identify $w_0 \cdot f$ with f for each point $f \in F$. Thus the face lattice of B_D is isomorphic to the poset of subsets of \bar{S} .

Suppose now that D is not of finite type. We define a subgraph D_i of D to be *maximal finite* if D_i is a full subgraph of D of finite type that is not contained in any larger subgraph of finite type. Let D_1, \dots, D_k be the list of maximal finite subgraphs of D . We denote by $D_i \cap D_j$ the full subgraph of D with vertices common to D_i and

D_j . There is a canonical embedding of the polytope $P_{D_i \cap D_j}$ into the polytope P_{D_i} ; such embeddings allow us to define P_D as the amalgamated sum of the polytopes P_{D_1}, \dots, P_{D_k} . The space B_D is the connected space obtained from P_D by isometrically identifying any two cells with similarly labelled 1-skeleta; the identification is the unique one which respects orientations of edges. The face lattice of the space B_D is isomorphic to the poset $S^f = \{T \subset \bar{S} : |\langle T \rangle| < \infty\}$ ordered by inclusion.

Note that if a Coxeter graph D with vertex set S is a disjoint union of two Coxeter graphs D', D'' with vertex sets S', S'' respectively, then there is a poset isomorphism $S^f = S'^f \times S''^f$. It is not difficult to see that this poset isomorphism extends to a CW-homeomorphism $B_D = B_{D'} \times B_{D''}$.


Example. Consider the graph



where edges whose label is not indicated are assumed to have edge label 3. Letting vertices correspond to generators w, x, y, z (starting at the top left corner and working clockwise) the associated Artin group is

$$A_D = \langle w, x, y, z : wxw = xwx, wy = yw, \\ wz w = zwz, xz = zx, yzyz = zyzy \rangle.$$

The 3-dimensional space B_D is obtained from the following two 3-dimensional polytopes by identifying similarly labelled faces and edges.



The space B_D contains four 1-cells, five 2-cells and two 3-cells. \square

3. Proof of Theorem 2

Suppose that A_D satisfies the hypothesis of the Theorem 2. Let X_D denote the universal covering space of B_D . We shall use induction on the number of infinity edges in D and the number of connected components in D to show that X_D is contractible.

If there are no infinity edges and the graph D is connected then X_D is contractible by hypothesis.

If D is not connected then A_D is a direct product $A_D = A_{D'} \times A_{D''}$ of two non-trivial Artin groups $A_{D'}$ and $A_{D''}$ where the graph D is the disjoint union of D' and D'' . The space B_D is the direct product $B_{D'} \times B_{D''}$. Thus X_D is contractible if and only if both $X_{D'}$ and $X_{D''}$ are contractible. Hence, by induction on the number of connected components in D , it suffices to prove the theorem in the case where the graph D is connected.

Suppose that the Coxeter graph D is connected. Suppose that there is an infinity edge in D whose endpoints correspond to the generators $a, b \in S = \{x_1, \dots, x_n\}$. Let $A_{\hat{a}}$ be the subgroup of A_D generated by $S \setminus \{a\}$, and $A_{\hat{a}, \hat{b}}$ the subgroup generated by $S \setminus \{a, b\}$. Let $D \setminus \{a\}$ denote the graph obtained from D by removing vertex a and all edges incident with a . Let $D \setminus \{a, b\}$ be the subgraph obtained by removing vertices a, b and all edges incident with them. There are clearly surjective homomorphisms $A_{D \setminus \{a\}} \rightarrow A_{\hat{a}}$ and $A_{D \setminus \{a, b\}} \rightarrow A_{\hat{a}, \hat{b}}$. A result of H. van der Lek [13] (see also [14]) shows that these surjections are in fact isomorphisms. Note that each of the groups $A_{D \setminus \{a\}}$, $A_{D \setminus \{b\}}$, $A_{D \setminus \{a, b\}}$ is an Artin group satisfying the hypothesis of the theorem and with Coxeter graph involving fewer infinity edges than are in D .

Suppose that D has $n \geq 1$ infinity edges. As an inductive hypothesis assume that the theorem holds for all Artin groups satisfying its hypothesis and having Coxeter graph with fewer than n infinity edges. Thus we can assume that $B_{D \setminus \{a\}}$, $B_{D \setminus \{b\}}$, $B_{D \setminus \{a, b\}}$ are classifying spaces for the subgroups $A_{\hat{a}}$, $A_{\hat{b}}$, $A_{\hat{a}, \hat{b}}$. Consider the homotopy pushout

$$\begin{array}{ccc} B_{D \setminus \{a, b\}} & \longrightarrow & B_{D \setminus \{a\}} \\ \downarrow & & \downarrow \\ B_{D \setminus \{b\}} & \longrightarrow & W. \end{array}$$


The space $W = B_{D \setminus \{a\}} \cup B_{D \setminus \{b\}}$ is precisely the space $W = B_D$. Now by a theorem of J. H. C. Whitehead (see for example [3], Chapter II-7) the space W is a classifying space. Hence its universal cover X_D is contractible. \square

An argument similar to the above was used in [8] to study properties of graph products of groups. Also, a version of this proof for Artin groups of FC type can be found in [5] as a remark following Lemma 4.3.7.

4. An application

The cellular chain complex $C_*(X_D)$ has been implemented in the computational algebra package HAP [11]. In cases where the $K(\pi, 1)$ conjecture is known to hold this chain complex is a free $\mathbb{Z}A_D$ -resolution of \mathbb{Z} and can be used to compute the cohomology of the Artin group A_D . The following was obtained in this way.

Proposition 4. *The Artin group A_D defined by the Coxeter graph*



has integral cohomology groups

$$\begin{aligned} H^0(A_D, \mathbb{Z}) &\cong \mathbb{Z}, & H^1(A_D, \mathbb{Z}) &\cong \mathbb{Z}^5, & H^2(A_D, \mathbb{Z}) &\cong \mathbb{Z}^{11}, \\ H^3(A_D, \mathbb{Z}) &\cong \mathbb{Z}_2 \oplus \mathbb{Z}^{14}, & H^4(A_D, \mathbb{Z}) &\cong \mathbb{Z}_2^2 \oplus \mathbb{Z}^{12}, & H^5(A_D, \mathbb{Z}) &\cong \mathbb{Z}_2 \oplus \mathbb{Z}^6, \\ H^6(A_D, \mathbb{Z}) &\cong \mathbb{Z}, & H^n(A_D, \mathbb{Z}) &= 0 \quad (n \geq 7). \end{aligned}$$

Proof. The graph D is such that for every ∞ -free subgraph D' the Artin group $A_{D'}$ satisfies the $K(\pi, 1)$ conjecture by results mentioned in Section 1. By Theorem 2 the group A_D itself satisfies the $K(\pi, 1)$ conjecture. We can thus use the computer implementation of $C_*(X_D)$ in [11] to make the cohomology calculations. The space X_D is 6-dimensional in this example. \square

References

- [1] K. J. Appel and P. E. Schupp, Artin groups and infinite Coxeter groups. *Invent. Math.* **72** (1983), 201–220. [Zbl 0536.20019](#) [MR 0700768](#) [410](#)
- [2] R. Blind and P. Mani-Levitska, Puzzles and polytope isomorphisms. *Aequationes Math.* **34** (1987), 287–297. [Zbl 0634.52005](#) [MR 0921106](#) [411](#)
- [3] K. S. Brown, *Cohomology of groups*. Grad. Texts in Math. 87, Springer-Verlag, New York 1982. [Zbl 0584.20036](#) [MR 0672956](#) [413](#)
- [4] F. Callegaro, D. Moroni and M. Salvetti, The $K(\pi, 1)$ problem for the affine Artin group of type \tilde{B}_n and its cohomology. Preprint, 2007. [arXiv:0705.2830](#) [410](#)

- [5] R. Charney and M. W. Davis, The $K(\pi, 1)$ problem for hyperplane complements associated to infinite reflection groups. *J. Amer. Math. Soc.* **8** (1995), no. 3, 597–627. [Zbl 0833.51006](#) [MR 1303028](#) 410, 414
- [6] R. Charney and M. W. Davis, Finite $K(\pi, 1)$ s for Artin groups. In *Prospects in topology* (Princeton, NJ, 1994), Ann. of Math. Stud. 138, Princeton University Press, Princeton, NJ, 1995, 110–124. [Zbl 0930.55006](#) [MR 1368655](#) 410
- [7] R. Charney and D. Peifer, The $K(\pi, 1)$ -conjecture for the affine braid groups. *Comment. Math. Helv.* **78** (2003), no. 3, 584–600. [Zbl 1066.20043](#) [MR 1998395](#) 410
- [8] D. E. Cohen, Projective resolutions for graph products of groups. *Proc. Edinburgh Math. Soc.* **38** (1995) 185–188. [Zbl 0836.20074](#) [MR 1317337](#) 410, 414
- [9] H. S. M. Coxeter, Discrete groups generated by reflections. *Ann. of Math.* (2) **35** (1934), no. 3, 588–621. [Zbl 60.0898.02](#) [MR 1503182](#) 411
- [10] P. Deligne, Les immeubles des groupes de tresses généralisés. *Invent. Math.* **17** (1972), 273–302. [Zbl 0238.20034](#) [MR 0422673](#) 410
- [11] G. Ellis, Homological algebra programming. A GAP package for computational homological algebra. <http://www.gap-system.org/Packages/hap.html> 414
- [12] H. Hendriks, Hyperplane complements of large type. *Invent. Math.* **79** (1985), 375–381. [Zbl 0564.57016](#) [MR 0778133](#) 410
- [13] H. van der Lek, The homotopy type of complex hyperplane complements. Ph.D. Thesis, Nijmegen, 1983. 413
- [14] L. Paris, Parabolic subgroups of Artin groups. *J. Algebra* **196** (1997), no. 2, 369–399. [Zbl 0926.20022](#) [MR 1475116](#) 413
- [15] M. Salvetti, The homotopy type of Artin groups. *Math. Res. Lett.* **1** (1994), no. 5, 565–577. [Zbl 0847.55011](#) [MR 1295551](#) 410
- [16] C. C. Squier, The homological algebra of Artin groups. *Math. Scand.* **75** (1994), no. 1, 5–43. [Zbl 0839.20065](#) [MR 1308935](#) 409

Received April 18, 2008

Graham Ellis, Mathematics Department, National University of Ireland, Galway

E-mail: graham.ellis@nuigalway.ie

Emil Sköldbberg, Mathematics Department, National University of Ireland, Galway, Ireland

E-mail: emil.skoldberg@nuigalway.ie