

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 85 (2010)

Artikel: Small exotic Stein manifolds
Autor: Akbulut, Selman / Yasui, Kouichi
DOI: <https://doi.org/10.5169/seals-130676>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 16.02.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Small exotic Stein manifolds

Selman Akbulut* and Kouichi Yasui

Abstract. It is known that the only Stein filling of the standard contact structure on S^3 is B^4 . In this paper, we construct pairs of homeomorphic but not diffeomorphic simply connected compact Stein 4-manifolds, for any Betti number $b_2 \geq 1$; we do this by enlarging corks and plugs.

Mathematics Subject Classification (2010). 57R55, 57R17, 57R65.

Keywords. 4-manifold, Stein manifold, handlebody, cork, plug.

1. Introduction

A properly imbedded complex submanifold of an affine space $X \subset \mathbb{C}^N$ is called a *Stein manifold*. A compact smooth submanifold $M \subset X$ is called a *compact Stein manifold* if it is cut out from X by $f \leq c$, where $f: X \rightarrow \mathbb{R}$ is a strictly plurisubharmonic (proper) Morse function, and c is a regular value. In particular M is a symplectic manifold with convex boundary and the symplectic form $\omega = \frac{1}{2}\partial\bar{\partial}f$. The form ω induces a contact structure ξ on the boundary ∂M . We call (M, ω) a *Stein filling* of the boundary contact manifold $(\partial M, \xi)$. Stein manifolds have been a useful tools for studying exotic smooth structures on 4-manifolds, since smooth 4-manifolds can be decomposed into codimension zero Stein pieces (e.g. [3], [5]).

In [8], [9] Eliashberg characterized the topology of Stein manifolds and proved that S^3 with the standard contact structure has a unique Stein filling B^4 (for more uniqueness results see [11]). However, Ozbagci–Stipsicz [12] and Smith [13] constructed infinitely many different Stein fillings of some contact 3-manifolds up to diffeomorphism. Their Stein fillings are distinguished by their first homology groups, so in particular they are not homeomorphic to each other. It is thus interesting to find pairs of homeomorphic but not diffeomorphic compact Stein 4-manifolds. Recently, by using Fintushel–Stern’s knot surgery, Akhmedov–Etnyre–Mark–Smith [6] constructed infinitely many simply connected compact Stein 4-manifolds that are

*The first author is partially supported by NSF, and the second author is partially supported by JSPS Research Fellowships for Young Scientists.

mutually homeomorphic but not diffeomorphic, and the induced contact structures on their boundaries are isomorphic. Here we give various examples of small simply connected compact Stein manifold pairs, that are homeomorphic but not diffeomorphic to each other, and describe their handlebodies concretely. In the light of the Eliashberg's theorem it is interesting to seek such exotic pairs with the smallest second Betti number b_2 . Our examples provide the case $b_2 = 1$. Note that the previously constructed exotic contractible manifold of [1] is only exotic rel boundary, and the exotic manifold pairs of [2] have $b_2 = 1$ but are not both Stein.

Corks and plugs are fundamental objects detecting exotic smooth structures on 4-manifolds [5]. In [1] the first example of a cork was found. In [2] an exotic pair of simply connected 4-manifolds with boundaries was constructed, by enlarging a cork. In [5] examples of infinitely many corks and plugs were found. In this paper, we construct examples of exotic pairs of compact Stein 4-manifolds, by enlarging corks and plugs. We also give handlebody pictures of these Stein manifolds.

Theorem 1.1. *For every second Betti number $b_2 \geq 1$, there exist pairs of simply connected compact Stein 4-manifolds which are homeomorphic but not diffeomorphic to each other.*

Theorem 1.2. *For every $n \geq 0$, there exist pairs of simply connected compact Stein 4-manifolds with the following properties:*

- (i) *They are homeomorphic but not diffeomorphic to each other;*
- (ii) *The first homology groups of their boundaries are $\mathbb{Z}/n\mathbb{Z}$.*

Theorem 1.3. *There exist pairs of simply connected compact Stein 4-manifolds with the following properties:*

- (i) *Their boundaries are diffeomorphic;*
- (ii) *Their integer coefficient homology groups are isomorphic;*
- (iii) *Their intersection forms are not isomorphic. In particular, they are not homeomorphic.*

Let X be a simply connected compact smooth 4-manifold with $H_2(X; \mathbb{Z}) \cong \mathbb{Z}$. Define the genus $G(X)$ of X as the minimal number of the genera of surfaces which represent a generator of $H_2(X; \mathbb{Z})$. In this paper, by enlarging corks we also construct the examples of exotic pairs below. These examples show that the distance of genera of two homeomorphic (Stein) 4-manifolds can be arbitrarily large.

Theorem 1.4. *For every $n \geq 0$, there exist simply connected compact Stein 4-manifolds X_n and Y_n with the following properties:*

- (i) *They are homeomorphic but not diffeomorphic to each other;*

- (ii) $H_2(X_n; \mathbb{Z}) \cong H_2(Y_n; \mathbb{Z}) \cong \mathbb{Z};$
- (iii) $G(X_n) - G(Y_n) \geq n.$

Acknowledgment. The second author would like to thank his adviser Hisaaki Endo for his heartfelt encouragement. This work was partially done during the second author's stay at Michigan State University. The second author is grateful for their hospitality. Finally, the authors would like to thank the referee for his/her useful comments.

2. Corks and plugs

In this section, we briefly recall corks and plugs. For details, see [5]. For basics of handlebody pictures, see [10].

Definition 2.1. Let C be a compact Stein 4-manifold with boundary, and let $\tau : \partial C \rightarrow \partial C$ be an involution on the boundary. We call (C, τ) a *Cork* if τ extends to a self-homeomorphism of C , but cannot extend to any self-diffeomorphism of C . A cork (C, τ) is called a cork of a smooth 4-manifold X , if $C \subset X$ and X changes its diffeomorphism type when removing C and regluing it via τ . Note that this operation does not change the homeomorphism type of X .

Definition 2.2. Let P be a compact Stein 4-manifold with boundary, and let $\tau : \partial P \rightarrow \partial P$ be an involution on the boundary, which cannot extend to any self-homeomorphism of P . We call (P, τ) a *Plug* of X , if $P \subset X$ and X keeps its homeomorphism type and changes its diffeomorphism type when removing P and gluing it via τ . We call (P, τ) a *Plug* if there exists a smooth 4-manifold X such that (P, τ) is a plug of X .

Definition 2.3. Let W_n and $W_{m,n}$ be the smooth 4-manifolds in Figure 1. Let $f_n : \partial W_n \rightarrow \partial W_n$ and $f_{m,n} : \partial W_{m,n} \rightarrow \partial W_{m,n}$ be the obvious involutions obtained




Figure 1

by first surgering $S^1 \times B^3$ to $B^2 \times S^2$ in the interiors of W_n and $W_{m,n}$, then surgering

the other imbedded $B^2 \times S^2$ back to $S^1 \times B^3$ (i.e., replacing the dots in Figure 1), respectively. Note that the diagrams of W_n and $W_{m,n}$ are symmetric links.

Theorem 2.4 ([1], [5]). (1) For $n \geq 1$, the pair (W_n, f_n) is a cork.

(2) For $m \geq 1$ and $n \geq 2$, the pair $(W_{m,n}, f_{m,n})$ is a plug.

3. Proofs

3.1. Enlarging corks. In this subsection, we construct exotic smooth pairs of compact Stein 4-manifolds by enlarging corks W_n .

Definition 3.1. Let $C_1(m, n, p, q)$ be the simply connected compact 4-manifold in Figure 2. Let $C_2(m, n, p, q)$ be the simply connected compact 4-manifold in Figure 3. Here each m -framed knot of Figure 2 and 3 is the $(p, p-1)$ -torus knot.




Figure 2. $C_1(m, n, p, q)$.


Lemma 3.2. (1) $C_1(m, n, p, q)$ and $C_2(m, n, p, q)$ are homeomorphic.

(2) $C_1(m, n, p, q)$ and $C_2(m, n, p, q)$ are compact Stein 4-manifolds for $n \geq 1$, $q \geq 0$ and $m \leq p^2 - 3p$.


(3) $H_1(\partial C_1(m, n, p, 0); \mathbb{Z}) \cong H_1(\partial C_2(m, n, p, 0); \mathbb{Z}) \cong \mathbb{Z}/m\mathbb{Z}$.


(4) $H_2(C_1(m, n, p, q); \mathbb{Z}) \cong H_2(C_2(m, n, p, q); \mathbb{Z}) \cong \bigoplus_{q+1} \mathbb{Z}$.

Proof. (1) $C_2(m, n, p, q)$ is obviously obtained from $C_1(m, n, p, q)$ by removing the contractible 4-manifold W_n and regluing it via the involution f_n . The claim now follows from Freedman's theorem.

Figure 3. $C_2(m, n, p, q)$.

(2) By isotoping the diagrams of $C_1(m, n, p, q)$ and $C_2(m, n, p, q)$ into Legendrian diagrams (and changing the 1-handle notation) as in Figure 4 and 5, respectively, we can easily see that these manifolds are Stein, by checking the Eliashberg's criterium: The framing of each 2-handle is less than the Thurston–Bennequin number.

Figure 4. $C_1(m, n, p, q)$.

Figure 5. $C_2(m, n, p, q)$.

(3) The claim follows from the fact that the intersection forms of $C_1(m, n, p, 0)$ and $C_2(m, n, p, 0)$ are isomorphic to $\langle m \rangle$.

(4) This clearly follows from the definition. \square

We first prove the theorem below. This theorem together with Lemma 3.2 clearly gives Theorem 1.1 and 1.2.


Theorem 3.3. (1) $C_1(m, n, p, 0)$ and $C_2(m, n, p, 0)$ are homeomorphic but not diffeomorphic to each other, for $n \geq 4$, $p \geq 1$ and $m \leq p^2 - 3p + 1$.

(2) $C_1(m, n, p, 0)$ and $C_2(m, n, p, 0)$ are homeomorphic but not diffeomorphic to each other, for $1 \leq n \leq 3$, $p \geq 3$ and $m \leq p^2 - 3p + 1$.


(3) $C_1(m, n, p, q)$ and $C_2(m, n, p, q)$ are homeomorphic but not diffeomorphic to each other, for $n \geq 1$, $p, q \geq 1$ and $0 \leq m \leq p^2 - 3p + 1$.

Let $E(n)$ be the relatively minimal simply connected elliptic surface with Euler characteristic $12n$. We begin with the theorem below.

Theorem 3.4 (Gompf–Stipsicz [10]). *For $n \geq 1$, the elliptic surface $E(n)$ has the handle decomposition in Figure 6. The obvious cusp neighborhood (i.e. the dotted circle, -1 -framed meridian of the dotted circle, and the left most 0 -framed unknot) is isotopic to the regular neighborhood of a cusp fiber of $E(n)$.*

Figure 6. $E(n)$.

Corollary 3.5. For $n \geq 1$, the elliptic surface $E(n)$ has a handle decomposition as in Figure 7. The obvious cusp neighborhood is isotopic to the regular neighborhood of a cusp fiber of $E(n)$.

Figure 7. $E(n)$.

Proof. In Figure 6, pull off the leftmost 0-framed unknot from the dotted circle by sliding over -1 -framed knot. \square

Proposition 3.6. For $p, q, r \geq 1$, the elliptic surface $E(p + q + 2r + 1)$ has a handle decomposition as in Figure 8, where $k = 3r - 2, 3r - 1, 3r$. The obvious cusp neighborhood is isotopic to the regular neighborhood of a cusp fiber of

$E(p + q + 2r + 1)$. The symbol F denotes the class of a regular fiber of $E(p + q + 2r + 1)$ in $H_2(E(p + q + 2r + 1); \mathbb{Z})$.




Figure 8. $E(p + q + 2r + 1)$ ($k = 3r - 2, 3r - 1, 3r$).


Proof. We give a proof for $k = 3r - 2$. The other cases are only slightly different. Figure 7 shows that $E(p + q + 2r + 1)$ has a handle decomposition as in Figure 9. We slide handles as in Figure 10. Notice that we ignored many 2-handles in the process. We next introduce a 2-handle/3-handle pair and slide the new 2-handle as in the second and third diagrams of Figure 11. We repeat a handle slide as in the third, fourth and fifth diagrams. (For more details, see [14, Figure 15–17] and [15, Proposition 3.1(1)].) We now get the sixth diagram by an isotopy. By sliding the p^2 -framed knot over -1 -framed knots, we obtain the last diagram. This procedure gives a diagram of $E(p + q + 2r + 1)$ as in Figure 8. \square

By blowing ups, we get the following corollary.

Corollary 3.7. *For $p, q, r \geq 1$, the 4-manifold $E(p + q + 2r + 1) \# (2r - 1)\overline{\mathbb{CP}^2}$ has a handle decomposition as in Figure 12. The symbol F denotes the class of a regular fiber of $E(p + q + 2r + 1)$ in $H_2(E(p + q + 2r + 1); \mathbb{Z})$. The symbols $E_1, E_2, \dots, E_{2r-1}$ denote a standard basis of $H_2((2r - 1)\overline{\mathbb{CP}^2}; \mathbb{Z})$.*

We can now prove Theorem 3.3.

Proof of Theorem 3.3. Let α be a generator of $H_2(C_1(m, n, p, 0); \mathbb{Z}) \cong \mathbb{Z}$. Let β be a generator of $H_2(C_2(m, n, p, 0); \mathbb{Z}) \cong \mathbb{Z}$. We can easily check that β is represented by a surface with its genus $\frac{1}{2}(p^2 - 3p + 2)$.

Figure 9. $E(p+q+2r+1)$.

(1) (i) The case $m = p^2 - 3p + 1$ and $n = 3r - 2$ ($r \in \mathbb{Z}_{\geq 2}$): Since $\text{PD}((p+2r-1)F)$ is a Seiberg–Witten basic class of $E(p+2r+1)$ (see [10].), the blow up formula shows that $K := \text{PD}((p+2r-1)F + E_1 + E_2 + \dots + E_{2r-1})$ is a Seiberg–Witten basic class of $E(p+2r+1) \# (2r-1)\overline{\mathbb{CP}^2}$. Here the symbol PD denotes the Poincaré dual. Let α' be the element of $H_2(E(p+2r+1) \# (2r-1)\overline{\mathbb{CP}^2}; \mathbb{Z})$ induced from α by the inclusion of $C_1(m, n, p, 0)$ as shown in Figure 12. Put g as the genus of a surface which represents α . Then the adjunction inequality for the symplectic 4-manifold $E(p+2r+1) \# (2r-1)\overline{\mathbb{CP}^2}$ shows the following inequality:

$$|K(\alpha')| + \alpha'^2 \leq \begin{cases} 2g - 2 & \text{if } g \geq 2, \\ 0 & \text{if } g = 0, 1. \end{cases}$$

This inequality together with Figure 12 shows that

$$g \geq \frac{1}{2}(p^2 - 3p + 2r + 2).$$

Note that the right side of the inequality above is greater than or equal to 2. Since β is represented by a surface with its genus $\frac{1}{2}(p^2 - 3p + 2)$, the inequality above shows that $C_1(m, n, p, 0)$ and $C_2(m, n, p, 0)$ are not diffeomorphic.

(ii) The other cases: We first blow up Figure 12 and proceed similarly as above by using the adjunction inequality.

- (2) The same argument as the proof of (1) shows the claim.
- (3) Suppose that $C_1(m, n, p, q)$ and $C_2(m, n, p, q)$ are diffeomorphic. Then there exists a genus $\frac{1}{2}(p^2 - 3p + 2)$ surface with self-intersection number m in




Figure 10. Handle slides.




Figure 11. Handle slide.




Figure 12. $E(p + q + 2r + 1) \# (2r - 1)\overline{\mathbb{CP}^2}$, ($k = 3r - 2, 3r - 1, 3r$).

$C_1(m, n, p, q)$, since $C_2(m, n, p, q)$ has such a surface. Let γ be an element of $H_2(C_1(m, n, p, q); \mathbb{Z})$ with self-intersection number m . Put g as the genus of a surface which represent γ .

(i) The case $m = p^2 - 3p + 1$ and $n = 3r - 2$ ($r \in \mathbb{Z}_{\geq 2}$): Let γ' be the element of $H_2(E(p + q + 2r + 1) \# (2r - 1)\overline{\mathbb{CP}^2}; \mathbb{Z})$ induced from γ by the inclusion of $C_1(m, n, p, q)$ as shown in Figure 12. The fact $\gamma'^2 = m \geq 0$ and Figure 2 shows the existence of some element x of $H_2(E(p + q + 2r + 1) \# (2r - 1)\overline{\mathbb{CP}^2}; \mathbb{Z})$ such that $x \cdot ((p + q + 2r - 1)F + E_1 + E_2 + \dots + E_{2r-1}) = 0$ and $\gamma' = a\alpha' + x$ ($a \geq 1$). Since $K := \text{PD}((p + q + 2r - 1)F + E_1 + E_2 + \dots + E_{2r-1})$ is a Seiberg–Witten basic class of $E(p + q + 2r + 1) \# (2r - 1)\overline{\mathbb{CP}^2}$, the adjunction inequality gives the following inequality:

$$|K(\gamma')| + \gamma'^2 \leq \begin{cases} 2g - 2 & \text{if } g \geq 2, \\ 0 & \text{if } g = 0, 1. \end{cases}$$

This inequality together with Figure 12 shows

$$g \geq \frac{1}{2}(p^2 - 3p + 3 + a(2r - 1)).$$

Note that the right side of the inequality above is greater than or equal to 2. On the

other hand, the assumption shows that γ is represented by a genus $\frac{1}{2}(p^2 - 3p + 2)$ surface. This is a contradiction.

(ii) The other cases: We first blow up Figure 12 and proceed similarly as above by using the adjunction inequality. \square

Remark 3.8. In Theorem 3.3, we split the cases into (1) and (2) to avoid the $n \leq 3$, $p = 1, 2, q = 0$ case. This is because, in this case, the adjunction inequalities in the proof above cause difficulties with the genus $g = 0$ case.

The proof above of Theorem 3.3(1) shows the corollary below. This corollary together with Lemma 3.2 clearly gives Theorem 1.4.

Corollary 3.9. *If $r \geq 2$, $p \geq 1$ and $m \leq p^2 - 3p + 1$, then*

$$G(C_1(m, 3r - 2, p, 0)) - G(C_2(m, 3r - 2, p, 0)) \geq r$$

3.2. Enlarging plugs. In this subsection, by enlarging plugs $W_{m,n}$, we construct exotic smooth pairs of compact Stein 4-manifolds. We also construct non-homeomorphic pairs of compact Stein 4-manifolds with the same boundary 3-manifold.

Definition 3.10. Let $P_1(m, n)$ and $P_2(m, n)$ be the simply connected compact smooth 4-manifolds in Figure 13.





Figure 13

We can easily prove the lemma below, similarly to the proof of Lemma 3.2. See also the equivalent diagrams of $P_1(m, n)$ and $P_2(m, n)$ in Figure 14.


Lemma 3.11. *$P_1(m, n)$ and $P_2(m, n)$ are simply connected compact Stein 4-manifolds for $m, n \geq 1$. The boundaries $\partial P_1(m, n)$ and $\partial P_2(m, n)$ are diffeomorphic.*

We first prove the following theorem.


Theorem 3.12. *Simply connected compact Stein 4-manifolds $P_1(1, 3)$ and $P_2(1, 3)$ are homeomorphic but not diffeomorphic to each other.*

Figure 14. $P_1(m,n)$ and $P_2(m,n)$.

Proposition 3.13. $E(2) \# 2\overline{\mathbb{CP}^2}$ has a handle decomposition as in Figure 15. Here E_1 and E_2 denote a standard basis of $H_2(2\overline{\mathbb{CP}^2}; \mathbb{Z})$.

Figure 15. $E(2) \# 2\overline{\mathbb{CP}^2}$.

Proof. The $n = 2$ case of Figure 7 obviously gives the first diagram of $E(2)$ in Figure 16. We get the second diagram of Figure 16 by handle moves as in the $p = 3$ case of Figure 11. Handle slides as in Figure 17 gives the third diagram of Figure 16. Finally blow up twice. Then we get Figure 15. \square

Figure 16. Handle moves of $E(2)$.

Proof of Theorem 3.12. We can easily check that the intersection forms of both $P_1(1, 3)$ and $P_2(1, 3)$ are $\langle 1 \rangle \oplus \langle -1 \rangle$. Boyer's theorem [7] thus shows that $P_1(1, 3)$ and $P_2(1, 3)$ are homeomorphic.

It follows from Figure 15 that the subspace $H_2(P_1(1, 3); \mathbb{Z})$ of $H_2(E(2) \# 2\overline{\mathbb{CP}^2}; \mathbb{Z})$ is spanned by elements x_1 and x_2 such that $E_1 \cdot x_1 = 1$, $E_2 \cdot x_1 = 0$, $E_1 \cdot x_2 = 0$ and $E_2 \cdot x_2 = 1$. The blow-up formula shows that the Seiberg–Witten basic classes of $E(2) \# 2\overline{\mathbb{CP}^2}$ are $\pm \text{PD}(E_1 \pm E_2)$. Using the adjunction inequality for $E(2) \# 2\overline{\mathbb{CP}^2}$, we can now easily prove that there exists no non-zero element of $H_2(P_1(1, 3); \mathbb{Z})$ such that the square of the element is zero and that the element is represented by a torus.




Figure 17. Handle slide.

On the other hand, we can easily check that there exists a non-zero element of $H_2(P_2(1, 3); \mathbb{Z})$ such that the square of the element is zero and that the element is represented by a torus. Therefore, the claim follows. \square

Remark 3.14. We can easily extend Theorem 3.12 as in Theorem 3.3.

We finally give the theorem below. This shows Theorem 1.3.

Theorem 3.15. *If $m \geq 1$ is odd, and $n \geq 1$ is even, then simply connected compact Stein 4-manifolds $P_1(m, n)$ and $P_2(m, n)$ have non-isomorphic intersection forms, whereas they have diffeomorphic boundaries and isomorphic integer homology groups.*

Proof. We can easily check that the intersection form of $P_1(m, n)$ is odd and that the one of $P_2(m, n)$ is even, under the assumption. \square

References

- [1] S. Akbulut, A fake compact contractible 4-manifold. *J. Differential Geom.* **33** (1991), no. 2, 335–356. [Zbl 0839.57015](#) [MR 1094459](#)
- [2] S. Akbulut, An exotic 4-manifold. *J. Differential Geom.* **33** (1991), 357–361. [Zbl 0839.57016](#) [MR 1094460](#)
- [3] S. Akbulut and R. Matveyev, A convex decomposition of 4-manifolds. *Internat. Math. Res. Notices* **1998** (1998), no. 7, 371–381. [Zbl 0911.57025](#) [MR 1623402](#)
- [4] S. Akbulut and R. Matveyev, Exotic structures and adjunction inequality. *Turkish J. Math.* **21** (1997), no. 1, 47–53. [Zbl 0885.57011](#) [MR 1456158](#)
- [5] S. Akbulut and K. Yasui, Corks, plugs and exotic structures. *J. Gökova Geom. Topol.* **2** (2008), 40–82. [Zbl 05660539](#) [MR 2466001](#)
- [6] A. Akhmedov, J. B. Etnyre, T. E. Mark, and I. Smith, A note on Stein fillings of contact manifolds. *Math. Res. Lett.* **15** (2008), no. 6, 1127–1132. [Zbl 1156.57019](#) [MR 2470389](#)
- [7] S. Boyer, Simply-connected 4-manifolds with a given boundary. *Trans. Amer. Math. Soc.* **298** (1986), no. 1, 331–357. [Zbl 0615.57008](#) [MR 0857447](#)
- [8] Y. Eliashberg, Topological characterization of Stein manifolds of dimension > 2 . *Internat. J. Math.* **1** (1990), no. 1, 29–46. [Zbl 0699.58002](#) [MR 1044658](#)
- [9] Y. Eliashberg, Filling by holomorphic discs and its applications. In *Geometry of low-dimensional manifolds*, 2 (Durham, 1989), London Math. Soc. Lecture Note Ser. 151, Cambridge University Press, Cambridge 1990, 45–67. [Zbl 0731.53036](#) [MR 1171908](#)
- [10] R. E. Gompf and A. I. Stipsicz, *4-manifolds and Kirby calculus*. Grad. Stud. Math. 20, Amer. Math. Soc., Providence, R.I., 1999. [Zbl 0933.57020](#) [MR 1707327](#)
- [11] B. Ozbagci and A. I. Stipsicz, *Surgery on contact 3-manifolds and Stein surfaces*. Bolyai Soc. Math. Stud. 13, Springer-Verlag, Berlin; Janos Bolyai Mathematical Society, Budapest 2004. [Zbl 1067.57024](#) [MR 2114165](#)

- [12] B. Ozbagci and A.I. Stipsicz, Contact 3-manifolds with infinitely many Stein fillings. *Proc. Amer. Math. Soc.* **132** (2004), no. 5, 1549–1558. [Zbl 1045.57014](#) [MR 2053364](#)
- [13] I. Smith, Torus fibrations on symplectic four-manifolds. *Turkish J. Math.* **25** (2001), no. 1, 69–95. [Zbl 0996.53055](#) [MR 1829080](#)
- [14] K. Yasui, Exotic rational elliptic surfaces without 1-handles. *Algebr. Geom. Topol.* **8** (2008), 971–996. [Zbl 1161.57017](#) [MR 2443105](#)
- [15] K. Yasui, Small exotic rational surfaces without 1- and 3-handles. *Trans. Amer. Math. Soc.*, to appear; preprint 2008, [arXiv:0807.0373v2](#) [math.GT].

Received July 24, 2008

Selman Akbulut, Department of Mathematics, Michigan State University, East Lansing, MI 48824, U.S.A.

E-mail: akbulut@math.msu.edu

Kouichi Yasui, Department of Mathematics, Graduate School of Science, Kyoto University, Kyoto 606-8502 Japan

E-mail: kyasui@math.kyoto-u.ac.jp