

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 86 (2011)

**Artikel:** Comparison of dynamical degrees for semi-conjugate meromorphic maps  
**Autor:** Dinh, Tien-Cuong / Nguyễn, Việt-Anh  
**DOI:** <https://doi.org/10.5169/seals-283468>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 15.04.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Comparison of dynamical degrees for semi-conjugate meromorphic maps

Tien-Cuong Dinh and Viêt-Anh Nguyễn

**Abstract.** Let  $f : X \rightarrow X$  be a dominant meromorphic map on a projective manifold  $X$  which preserves a meromorphic fibration  $\pi : X \rightarrow Y$  of  $X$  over a projective manifold  $Y$ . We establish formulas relating the dynamical degrees of  $f$ , the dynamical degrees of  $f$  relative to the fibration and the dynamical degrees of the map  $g : Y \rightarrow Y$  induced by  $f$ . Applications are given.

**Mathematics Subject Classification (2010).** Primary 37F; Secondary 32U40, 32H50.

**Keywords.** Semi-conjugate maps, dynamical degree, relative dynamical degree.

### 1. Introduction

Let  $(X, \omega_X)$  be a compact Kähler manifold of dimension  $k$  and let  $f : X \rightarrow X$  be a meromorphic map. We assume that  $f$  is *dominant*, i.e. the image of  $f$  contains an open subset of  $X$ . Let  $\pi : X \rightarrow Y$  be a dominant meromorphic map from  $X$  onto a compact Kähler manifold  $(Y, \omega_Y)$  of dimension  $l \leq k$ . The fibers of  $\pi$  define a fibration on  $X$  which might be singular. If  $f$  preserves this fibration, i.e.  $f$  sends generic fibers of  $\pi$  to fibers of  $\pi$ , it induces a dominant meromorphic map  $g : Y \rightarrow Y$  such that  $\pi \circ f = g \circ \pi$ . In that case, we say that  $f$  is *semi-conjugate* to  $g$ . For simplicity, we assume that  $\omega_Y$  is normalized so that  $\omega_Y^l$  is a probability measure.

A natural question is how the dynamical system defined by  $f$  is similar to the one defined by  $g$  when  $f$  is semi-conjugate to  $g$  as above. One of the first steps towards understanding this question should be to find out the relations between some invariants associated to  $f$  and  $g$ . In this paper, we will compare their dynamical degrees.

Let  $f^n := f \circ \dots \circ f$ ,  $n$  times, denote the iterate of order  $n$  of  $f$ . The dynamical degree  $d_p(f)$  of order  $p$  is the quantity which measures the growth of the norms of  $(f^n)^*$  acting on the Hodge cohomology group  $H^{p,p}(X, \mathbb{R})$  when  $n$  tends to infinity. By Poincaré duality, it also measures the growth of the norms of  $(f^n)_*$  acting on  $H^{k-p,k-p}(X, \mathbb{R})$ . If  $X$  is a projective manifold,  $d_p(f)$  represents the volume growth of  $f^n(V)$  for  $p$ -dimensional (closed complex) submanifolds  $V$  of  $X$ .

It was shown by Sibony and the first author in [6], [7] that dynamical degrees are bi-meromorphic invariants, that is, if  $f$  and  $g$  are bi-meromorphically conjugate, they have the same dynamical degrees. Dynamical degrees capture important dynamical information, in particular, in the computation of the topological entropy or in the construction of Green currents and of measures of maximal entropy. We refer the reader to the above references and to [8], [10], [15], [20] for more results on this matter.

When  $f$  preserves a fibration  $\pi : X \rightarrow Y$  as above, the dynamical degree  $d_p(f|\pi)$  of order  $p$  of  $f$  relative to  $\pi$  measures the growth of  $(f^n)^*$  acting on the subspace  $H_\pi^{l+p, l+p}(X, \mathbb{R})$  of classes in  $H^{l+p, l+p}(X, \mathbb{R})$  which can be supported by a generic fiber of  $\pi$ . It also measures the growth of  $(f^n)_*$  acting on  $H_\pi^{k-p, k-p}(X, \mathbb{R})$  and represents the volume growth of  $f^n(V)$  for  $p$ -dimensional submanifolds  $V$  of a generic fiber of  $\pi$  when  $X$  is projective. Precise definitions and properties will be given in Section 3. Here is our main result.

**Theorem 1.1.** *Let  $X$  and  $Y$  be projective manifolds of dimension  $k$  and  $l$  respectively with  $k \geq l$ . Let  $f : X \rightarrow X$ ,  $g : Y \rightarrow Y$  and  $\pi : X \rightarrow Y$  be dominant meromorphic maps such that  $\pi \circ f = g \circ \pi$ . Then the dynamical degrees  $d_p(f)$  of  $f$  are related to the dynamical degrees  $d_p(g)$  of  $g$  and the relative dynamical degrees  $d_p(f|\pi)$  by the formulas*

$$d_p(f) = \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} d_j(g) d_{p-j}(f|\pi)$$

for  $0 \leq p \leq k$ .

Note that the condition  $\max\{0, p-k+l\} \leq j \leq \min\{p, l\}$  is equivalent to  $0 \leq j \leq l$  and  $0 \leq p-j \leq k-l$ . It guarantees that  $d_j(g)$  and  $d_{p-j}(f|\pi)$  are meaningful<sup>1</sup>. We deduce from the above result that  $\max d_p(f) \geq \max d_p(g)$ . This gives an affirmative answer to the problem 9.3 in Hasselblatt–Propp [12]. When  $X$  and  $Y$  have the same dimension, generic fibers of  $\pi$  are finite and have the same cardinality. Moreover,  $f$  defines bijections between generic fibers of  $\pi$ . We deduce from the proof of Theorem 1.1 the following corollary which generalizes a result in [6], [7]. It was proved by Nakayama–Zhang for holomorphic maps in [14].

**Corollary 1.2.** *Let  $X$  and  $Y$  be compact Kähler manifolds of same dimension  $k$ . Let  $f : X \rightarrow X$ ,  $g : Y \rightarrow Y$  and  $\pi : X \rightarrow Y$  be dominant meromorphic maps such that  $\pi \circ f = g \circ \pi$ . Then the dynamical degrees of  $f$  are equal to the dynamical degrees of  $g$ .*

<sup>1</sup>We will find later analogous conditions, essentially for the same reason but also to avoid expressions which always vanish, e.g.  $\omega_Y^{l+1} = 0$ .

Recall that by a theorem of Khovanskii [13], Teissier [17] and Gromov [9], the dynamical degrees of  $f$  are log-concave, i.e.  $p \mapsto \log d_p(f)$  is concave. Therefore, there are integers  $p \leq p'$  such that

$$1 = d_0(f) < \cdots < d_p(f) = \cdots = d_{p'}(f) > \cdots > d_k(f).$$

An instructive example with  $p \neq p'$  is a map  $f(x_1, x_2) = (h(x_1), x_2)$  on a product  $X_1 \times X_2$  of projective manifolds. A natural problem is to find dynamically interesting examples of maps on projective manifolds. Therefore, it would be interesting to see construction of maps with distinct consecutive dynamical degrees, i.e. with  $p = p'$ . Somehow, this condition insures that there is no trivial direction in the associated dynamical systems. We have the following useful results.

**Corollary 1.3.** *Let  $f, \pi, g$  be as in Theorem 1.1. If the consecutive dynamical degrees of  $f$  are distinct, then the same property holds for  $g$  and for the consecutive dynamical degrees of  $f$  relative to  $\pi$ .*

The following result is obtained using the Iitaka fibrations of  $X$ .

**Corollary 1.4.** *Let  $X$  be a projective manifold admitting a dominant meromorphic map with distinct consecutive dynamical degrees. Then the Kodaira dimension of  $X$  is either equal to 0 or  $-\infty$ .*

Note that the same result was proved for compact Kähler surfaces by Cantat in [3] and Guedj in [11], and for holomorphic maps on compact Kähler manifolds by Nakayama and Zhang in [14], [21]. We also refer to Amerik–Campana [1] and Nakayama–Zhang [14], [22] for other invariant fibrations for which Theorem 1.1 may be applied in order to compute dynamical degrees.

**Acknowledgment.** The paper was written while the second author was visiting the Abdus Salam International Centre for Theoretical Physics in Trieste and the Korea Institute for Advanced Study in Seoul. He wishes to express his gratitude to these organizations.

## 2. Positive closed currents

The proof of our main result uses a delicate calculus on positive closed currents on compact Kähler manifolds<sup>2</sup>. In this section, we prove some useful results which can be applied to currents of integration on varieties and may have independent interest. The reader will find in Demailly [4] and Voisin [19] the basic facts on currents and on Kähler geometry.

---

<sup>2</sup>In this paper, we only consider the strong positivity, see e.g. [8], A.2, for the terminology.

Let  $(X, \omega_X)$  be a compact Kähler manifold of dimension  $k$ . Let  $\mathcal{K}^p(X)$  denote the cone of classes of strictly positive closed  $(p, p)$ -forms in  $H^{p,p}(X, \mathbb{R})$ . This is an open cone which is salient, i.e.  $\overline{\mathcal{K}^p(X)} \cap -\overline{\mathcal{K}^p(X)} = \{0\}$ . If  $c, c'$  are two classes in  $H^{p,p}(X, \mathbb{R})$ , we write  $c \leq c'$  and  $c' \geq c$  when  $c' - c$  is in  $\mathcal{K}^p(X) \cup \{0\}$ .

If  $T$  is a real closed  $(p, p)$ -current, denote by  $\{T\}$  its class in  $H^{p,p}(X, \mathbb{R})$ . If moreover  $T$  is positive, the mass of  $T$  is defined by  $\|T\| := \langle T, \omega_X^{k-p} \rangle$ . We often use the properties that  $\|T\|$  depends only on the class of  $T$  and  $\{T\} \leq A\|T\|\{\omega_X^p\}$  for some constant  $A > 0$  independent of  $T$ . The following semi-regularization of currents was proved by Sibony and the first author in [6], [7].

**Proposition 2.1.** *Let  $T$  be a positive closed  $(p, p)$ -current on a compact Kähler manifold  $(X, \omega_X)$ . Then there is a sequence of smooth positive closed  $(p, p)$ -forms  $T_n$  on  $X$  which converges weakly to a positive closed  $(p, p)$ -current  $T'$  such that  $T' \geq T$ , i.e.  $T' - T \geq 0$ ,  $\|T_n\| \leq A\|T\|$  and  $\{T_n\} \leq A\|T\|\{\omega_X^p\}$ , where  $A > 0$  is a constant independent of  $T$ . Moreover, if  $T$  is smooth on an open set  $U$ , then for every compact set  $K \subset U$ , we have  $T_n \geq T$  on  $K$  when  $n$  is large enough.*

We need the following lemma.

**Lemma 2.2.** *Let  $T$  and  $S$  be positive closed currents on  $X$  of bidegree  $(p, p)$  and  $(q, q)$  respectively with  $p + q \leq k$ . Assume that  $T$  is smooth on a dense Zariski open set  $U$  of  $X$ . Then  $T|_U \wedge S|_U$  has a finite mass. More precisely, there is a constant  $A > 0$  independent of  $T, S$  and  $U$  such that*

$$\|T|_U \wedge S|_U\| := \langle T|_U \wedge S|_U, \omega_X^{k-p-q} \rangle \leq A\|T\|\|S\|.$$

*Proof.* Let  $T_n$  and  $K$  be as in Proposition 2.1. Since  $\|T_n \wedge S\|$  can be computed cohomologically, we have

$$\|T|_K \wedge S|_K\| \leq \liminf_{n \rightarrow \infty} \|T_n \wedge S\| \leq A\|T\|\|\omega_X^p \wedge S\| = A\|T\|\|S\|.$$

This property holds for every compact subset  $K$  of  $U$ . Therefore,

$$\|T|_U \wedge S|_U\| \leq A\|T\|\|S\|.$$

The lemma follows. □

Consider currents  $T$  and  $S$  as in Lemma 2.2. So,  $T|_U \wedge S|_U$  has a finite mass. Therefore, by Skoda’s theorem [16], its trivial extension defines a positive closed current on  $X$ . We denote by  $T \overset{\circ}{\wedge} S$  this current obtained for the maximal Zariski open set  $U$  on which  $T$  is smooth (in that case  $T|_U$  is the regular part of  $T$ ). Observe that when  $S$  has no mass on proper analytic subsets of  $X$ , the current obtained in this

way does not change if we replace  $U$  with another dense Zariski open set. We often use this property in the sequence. By Lemma 2.2, we always have

$$\|T \overset{\circ}{\wedge} S\| \leq A\|T\|\|S\|.$$

We will be interested in positive closed currents  $T$  on  $Y \times \mathbb{P}^m$ , where  $(Y, \omega_Y)$  is a compact Kähler manifold of dimension  $l$  and  $\mathbb{P}^m$  is the projective space of dimension  $m$  endowed with the standard Fubini–Study form  $\omega_{\text{FS}}$ . We assume that  $\omega_{\text{FS}}$  is normalized so that  $\omega_{\text{FS}}^m$  is a probability measure. In practice, we will take  $m := k - l = \dim X - \dim Y$ . In order to simplify the notation, the pull-back of  $\omega_Y$  and  $\omega_{\text{FS}}$  to  $Y \times \mathbb{P}^m$  under the canonical projections are also denoted by  $\omega_Y$  and  $\omega_{\text{FS}}$ . Consider on  $Y \times \mathbb{P}^m$  the Kähler form  $\omega := \omega_Y + \omega_{\text{FS}}$ . The pull-back of a class  $c$  in  $H^*(Y, \mathbb{C})$  or  $H^*(\mathbb{P}^m, \mathbb{C})$  to  $H^*(Y \times \mathbb{P}^m, \mathbb{C})$  under the canonical projections is also denoted by  $c$ .

If  $T$  is a positive closed  $(p, p)$ -current on  $Y \times \mathbb{P}^m$ , define for  $\max\{0, p - m\} \leq j \leq \min\{l, p\}$  (or equivalently, for  $0 \leq j \leq l$  and  $0 \leq p - j \leq m$ )

$$\alpha_j(T) := \langle T, \omega_Y^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle. \tag{1}$$

Observe that  $\alpha_j(T)$  depends only on the class  $\{T\}$  of  $T$ . Denote by  $\cup$  the cup-product on Hodge cohomology groups.

**Proposition 2.3.** *Let  $T$  be a positive closed  $(p, p)$ -current on  $Y \times \mathbb{P}^m$  as above. Then*

$$\{T\} \leq A \sum_{\max\{0, p-m\} \leq j \leq \min\{l, p\}} \alpha_j(T) \{\omega_Y^j\} \cup \{\omega_{\text{FS}}^{p-j}\},$$

where  $A > 0$  is a constant independent of  $T$ .

*Proof.* By the Künneth formula (see e.g. [19], p. 266) we have

$$H^*(Y \times \mathbb{P}^m, \mathbb{C}) = H^*(Y, \mathbb{C}) \otimes H^*(\mathbb{P}^m, \mathbb{C}).$$

Therefore, there are classes  $c_j \in H^{j,j}(Y, \mathbb{R})$  such that

$$\{T\} = \sum_{\max\{0, p-m\} \leq j \leq \min\{l, p\}} c_j \cup \{\omega_{\text{FS}}^{p-j}\}.$$

Let  $S$  be a smooth positive closed  $(l - j, l - j)$ -form on  $Y$  and  $S'$  its canonical pull-back to  $Y \times \mathbb{P}^m$ . Recall that  $c_j$  denotes also the pull-back of  $c_j$  to  $Y \times \mathbb{P}^m$ . Since  $\omega_{\text{FS}}^m$  is a probability measure on  $\mathbb{P}^m$ , a simple computation on bidegree gives

$$c_j \cup \{S\} = c_j \cup \{S'\} \cup \{\omega_{\text{FS}}^m\} = \langle T, S' \wedge \omega_{\text{FS}}^{m-p+j} \rangle \geq 0.$$

So,  $c_j$  belongs to the convex closed cone  $\mathcal{K}$  of classes  $c$  in  $H^{j,j}(Y, \mathbb{R})$  with  $c \cup c' \geq 0$  for  $c' \in \mathcal{K}^{l-j}(Y)$ . Since  $\mathcal{K}^{l-j}(Y)$  is open and since  $\cup$  is non-degenerate,  $\mathcal{K}$  is salient, i.e.  $\mathcal{K} \cap -\mathcal{K} = \{0\}$ . The fact that  $\{\omega_Y^{l-j}\}$  is in the interior of  $\mathcal{K}^{l-j}(Y)$  implies that  $c_j \cup \{\omega_Y^{l-j}\} = 0$  only when  $c_j = 0$ . Moreover, we have

$$\|c_j\| \leq A' c_j \cup \{\omega_Y^{l-j}\} = A' \langle T, \omega_Y^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle = A' \alpha_j(T)$$

for a fixed norm  $\| \cdot \|$  on  $H^{j,j}(Y, \mathbb{R})$  and for some constant  $A' > 0$ . It follows that

$$c_j \leq A \alpha_j(T) \{\omega_Y^j\}$$

for some constant  $A > 0$ . The result follows. □

**Proposition 2.4.** *Let  $T$  be a positive closed  $(p, p)$ -current on  $Y \times \mathbb{P}^m$  as above. Assume that  $Y$  is a projective manifold. Then there is a sequence of smooth positive closed  $(p, p)$ -forms  $T_n$  on  $Y \times \mathbb{P}^m$  which converges weakly to a current  $T' \geq T$  such that  $\alpha_j(T_n) \leq A \alpha_j(T)$  for all  $j$ , where  $A > 0$  is a constant independent of  $T$ . Moreover, if  $T$  is smooth on an open set  $U$ , then for every compact subset  $K$  of  $U$  and every  $\epsilon > 0$ , we have  $T_n \geq T - \epsilon \omega^p$  on  $K$  when  $n$  is large enough.*

*Proof.* We first consider the case where  $Y = \mathbb{P}^l$  and  $\omega_Y$  is the Fubini–Study form normalized so that  $\omega_Y^l$  is a probability measure. The Künneth formula applied to this particular case says that  $T$  is cohomologous to

$$\sum_{\max\{0, p-m\} \leq j \leq \min\{l, p\}} \alpha_j(T) \{\omega_Y^j\} \cup \{\omega_{\text{FS}}^{p-j}\}.$$

Since  $Y \times \mathbb{P}^m$  is homogeneous, we can regularize  $T$  using the automorphisms of  $Y \times \mathbb{P}^m$  which are close to the identity.

More precisely, let  $\nu_n$  be a sequence of smooth probability measures on the group of automorphisms  $\text{Aut}(Y \times \mathbb{P}^m)$  of  $Y \times \mathbb{P}^m$  whose supports converge to the identity  $\text{id} \in \text{Aut}(Y \times \mathbb{P}^m)$ . Define

$$T_n := \int_{\tau \in \text{Aut}(Y \times \mathbb{P}^m)} \tau_*(T) d\nu_n(\tau).$$

Then, the  $T_n$  are smooth positive closed  $(p, p)$ -forms and converge weakly to  $T$ . We also have  $\{T_n\} = \{T\}$  and hence  $\alpha_j(T_n) = \alpha_j(T)$ . This gives the first assertion for  $Y = \mathbb{P}^l$ .

For the second assertion, we can prove a stronger property. Let  $\Phi$  be a smooth positive  $(p, p)$ -form on  $U$  such that  $\Phi \leq T$ . We do not assume that  $T$  is smooth nor that  $\Phi$  is closed on  $U$ . Then

$$\Phi_n := \int_{\tau \in \text{Aut}(Y \times \mathbb{P}^m)} \tau_*(\Phi) d\nu_n(\tau)$$

converges uniformly to  $\Phi$  on  $K$ . Since  $\Phi_n \leq T_n$ , we have  $T_n \geq \Phi - \epsilon \omega^p$  on  $K$  for  $n$  large enough. With our hypothesis,  $T$  is smooth on  $U$  and we can replace  $\Phi$  with  $T$ .

Assume now that  $Y$  is a general projective manifold. We may find a finite family of open holomorphic maps  $\Psi_i$ ,  $1 \leq i \leq s$ , from  $Y$  onto  $\mathbb{P}^l$  such that for every point  $y \in Y$  at least one map  $\Psi_i$  is of maximal rank at  $y$ . To do this it suffices to embed  $Y$  into a projective space and take a family of central projections. Let  $\Pi_i: Y \times \mathbb{P}^m \rightarrow \mathbb{P}^l \times \mathbb{P}^m$  be defined by

$$\Pi_i(y, z) := (\Psi_i(y), z), \quad (y, z) \in Y \times \mathbb{P}^m.$$

We apply the first case to the currents  $T^{(i)} := (\Pi_i)_*(T)$ .

We construct as above smooth positive closed  $(p, p)$ -forms  $T_n^{(i)}$  on  $\mathbb{P}^l \times \mathbb{P}^m$  converging to  $T^{(i)}$  such that  $\{T_n^{(i)}\} = \{T^{(i)}\}$ . Define  $T_n := \sum_i \Pi_i^*(T_n^{(i)})$ . Since the cohomology classes of  $T_n^{(i)}$  are bounded, the classes of  $T_n$  are also bounded. Therefore, the masses of  $T_n$  are bounded. Up to extracting a subsequence, we can assume that  $\Pi_i^*(T_n^{(i)})$  converges and hence the  $T_n$  converge to a positive closed current  $T'$ . If  $(y, z)$  is a point in  $Y \times \mathbb{P}^m$  and  $\Psi_i$  has maximal rank at  $y$ , then  $\Pi_i$  defines a local bi-holomorphic map on a neighbourhood of  $(y, z)$ . In this neighbourhood, we have

$$T \leq \Pi_i^*(\Pi_i)_*(T) = \Pi_i^*(T^{(i)}) \leq \lim_{n \rightarrow \infty} \Pi_i^*(T_n^{(i)}) \leq T'.$$

The choice of  $\Psi_i$  implies that  $T \leq T'$  on  $Y \times \mathbb{P}^m$ . The second assertion of the proposition is a local property. So, it is also easy to check.

It remains to prove the estimate on  $\alpha_j(T_n)$ . Let  $\tilde{\omega}_{\text{FS}}$  denote the Fubini–Study form of  $\mathbb{P}^l$  normalized so that  $\tilde{\omega}_{\text{FS}}^l$  is a probability measure. Since  $\tilde{\omega}_{\text{FS}}$  is strictly positive, there is a constant  $A_1 > 0$  such that  $(\Psi_i)_*(\omega_Y^{l-j}) \leq A_1 \{\tilde{\omega}_{\text{FS}}^{l-j}\}$ . We also have  $(\Psi_i)^*(\tilde{\omega}_{\text{FS}}^{l-j}) \leq A_2 \omega_Y^{l-j}$  for some constant  $A_2 > 0$ . For simplicity, we will also denote by  $\omega_Y$ ,  $\omega_{\text{FS}}$  and  $\tilde{\omega}_{\text{FS}}$  the pull-backs of these forms to  $Y \times \mathbb{P}^m$  or to  $\mathbb{P}^l \times \mathbb{P}^m$ . In particular,  $(\Pi_i)_*(\omega_Y^{l-j} \wedge \omega_{\text{FS}}^{m-p+j})$  and  $(\Psi_i)_*(\omega_Y^{l-j}) \wedge \omega_{\text{FS}}^{m-p+j}$  represent the same form on  $\mathbb{P}^l \times \mathbb{P}^m$ . Since the  $T_n^{(i)}$  are smooth and since the following integrals can be computed cohomologically, we have

$$\begin{aligned} \langle \Pi_i^*(T_n^{(i)}), \omega_Y^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle &= \langle T_n^{(i)}, (\Psi_i)_*(\omega_Y^{l-j}) \wedge \omega_{\text{FS}}^{m-p+j} \rangle \\ &\leq A_1 \langle T_n^{(i)}, \tilde{\omega}_{\text{FS}}^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle \\ &= A_1 \langle T^{(i)}, \tilde{\omega}_{\text{FS}}^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle \\ &= A_1 \langle T, \Psi_i^*(\tilde{\omega}_{\text{FS}}^{l-j}) \wedge \omega_{\text{FS}}^{m-p+j} \rangle \\ &\leq A_1 A_2 \langle T, \omega_Y^{l-j} \wedge \omega_{\text{FS}}^{m-p+j} \rangle. \end{aligned}$$

It follows that  $\alpha_j(\Pi_i^*(T_n^{(i)})) \leq A_1 A_2 \alpha_j(T)$  and hence  $\alpha_j(T_n) \leq A \alpha_j(T)$  for some constant  $A > 0$ . □

### 3. Dynamical degrees

Let  $\pi: (X, \omega_X) \rightarrow (Y, \omega_Y)$  be a dominant meromorphic map between compact Kähler manifolds of dimension  $k$  and  $l$  respectively. The map  $\pi$  is holomorphic outside the indeterminacy set  $I_\pi$  which is an analytic subset of  $X$  of codimension at least 2. The closure  $\Gamma$  of its graph over  $X \setminus I_\pi$  is an irreducible analytic subset of dimension  $k$  of  $X \times Y$ . If,  $\tau_X$  and  $\tau_Y$  denote the projections from  $X \times Y$  onto its factors, then  $\tau_X$  defines a bi-holomorphic map between  $\Gamma \setminus \tau_X^{-1}(I_\pi)$  and  $X \setminus I_\pi$ . The fibers of  $\tau_X|_\Gamma$  over  $I_\pi$  have positive dimension. One can identify  $\pi$  with  $\tau_Y \circ (\tau_X|_\Gamma)^{-1}$ . For  $A \subset X$  and  $B \subset Y$ , define  $\pi(A) := \tau_Y(\tau_X|_\Gamma)^{-1}(A)$  and  $\pi^{-1}(B) := \tau_X(\tau_Y|_\Gamma)^{-1}(B)$ .

The map  $\pi$  induces linear operators on currents. If  $\Phi$  is a smooth  $(p, q)$ -form on  $Y$ , then  $\pi^*(\Phi)$  is the  $(p, q)$ -current defined by

$$\pi^*(\Phi) := (\tau_X)_*(\tau_Y^*(\Phi) \wedge [\Gamma]),$$

where  $[\Gamma]$  is the current of integration on  $\Gamma$ . It is not difficult to see that  $\pi^*(\Phi)$  is an  $L^1$  form smooth outside  $I_\pi$ . If  $\Psi$  is a smooth  $(p, q)$ -form on  $X$  with  $p, q \geq k - l$ , then  $\pi_*(\Psi)$  is the  $(p - k + l, q - k + l)$ -current defined by

$$\pi_*(\Psi) := (\tau_Y)_*(\tau_X^*(\Psi) \wedge [\Gamma]).$$

If  $\Phi$  and  $\Psi$  are closed or positive, so are  $\pi^*(\Phi)$  and  $\pi_*(\Psi)$ . Therefore,  $\pi^*$  and  $\pi_*$  induce linear operators on the Hodge cohomology groups of  $X$  and  $Y$ .

In general, the above operators do not extend continuously to positive closed currents. We will use instead the strict transforms of currents  $\pi^\bullet$  and  $\pi_\bullet$  which coincide with  $\pi^*$  and  $\pi_*$  on smooth positive closed forms. In this paper, we only need these operators in the case where  $X$  and  $Y$  have the same dimension  $k$ .

Let  $U$  be the maximal Zariski open set in  $X \setminus I_\pi$  such that  $\pi: U \rightarrow \pi(U)$  is locally invertible. The complement of  $U$  in  $X$  is called the *critical set* of  $\pi$ . If  $T$  is a positive closed  $(p, p)$ -current on  $Y$ ,  $(\pi|_U)^*(T)$  is well-defined and is a positive closed  $(p, p)$ -current on  $U$ . Proposition 2.1 allows us to show that this current has finite mass. By Skoda's Theorem [16], its trivial extension to  $X$  is a positive closed  $(p, p)$ -current that we denote by  $\pi^\bullet(T)$ .

Let  $V$  be the maximal Zariski open set in  $Y \setminus \pi(I_\pi)$  such that  $\pi: \pi^{-1}(V) \rightarrow V$  is a non-ramified covering. The complement of  $V$  in  $Y$  is called the *set of critical values* of  $\pi$ . If  $S$  is a positive closed  $(p, p)$ -current on  $X$ , then  $\pi_\bullet(S)$  is the trivial extension of  $(\pi|_{\pi^{-1}(V)})_*(S)$  to  $Y$ . This is also a positive closed  $(p, p)$ -current. We will use the properties that  $\|\pi^\bullet(T)\| \leq A\|T\|$  and  $\|\pi_\bullet(S)\| \leq A\|S\|$  for some constant  $A > 0$  independent of  $T, S$ , see [6], [7] for details.

Consider now a dominant meromorphic self-map  $f: X \rightarrow X$ . The iterate of order  $n$  of  $f$  is defined by  $f^n := f \circ \dots \circ f$  ( $n$  times) on a dense Zariski open set and extends to a dominant meromorphic map on  $X$ . Define for  $0 \leq p \leq k$

$$\lambda_p(f^n) := \|(f^n)^*(\omega_X^p)\| = \langle (f^n)^*(\omega_X^p), \omega_X^{k-p} \rangle.$$

It is not difficult to see that

$$\lambda_p(f^n) = \|(f^n)_*(\omega_X^{k-p})\| = \langle (f^n)_*(\omega_X^{k-p}), \omega_X^p \rangle.$$

It was shown in [6], [7] that  $[\lambda_p(f^n)]^{1/n}$  converges to a constant  $d_p(f)$  which is the *dynamical degree of order p* of  $f$ . Note that the main difficulty here is that in general we do not have  $(f^{n+s})^* = (f^n)^* \circ (f^s)^*$  on cohomology classes.

Let  $\|\cdot\|_{H^{p,p}}$  denote the norm of an operator acting on  $H^{p,p}(X, \mathbb{R})$  with respect to a fixed norm on that space. Since the mass of a positive closed current depends only on its cohomology class, we deduce from the above discussion that

$$A^{-1}\lambda_p(f^n) \leq \|(f^n)^*\|_{H^{p,p}} \leq A\lambda_p(f^n),$$

for some constant  $A > 0$ . It follows that

$$d_p(f) = \lim_{n \rightarrow \infty} \|(f^n)^*\|_{H^{p,p}}^{1/n}.$$

Note that we also have  $d_p(f^n) = d_p(f)^n$  for  $n \geq 1$ . The last dynamical degree  $d_k(f)$  is also called the *topological degree* of  $f$ . It is equal to the number of points in a generic fiber of  $f$  and we have  $\lambda_k(f^n)\|\omega_X^k\|^{-1} = d_k(f^n) = d_k(f)^n$ .

**Proposition 3.1.** *Let  $T$  be a positive closed  $(p, p)$ -current and  $S$  a positive closed  $(k - p, k - p)$ -current on  $X$ . Then*

$$\|(f^n)^\bullet(T)\| \leq A\|T\|\lambda_p(f^n) \quad \text{and} \quad \|(f^n)_\bullet(S)\| \leq A\|S\|\lambda_p(f^n)$$

for some constant  $A > 0$  independent of  $T, S$  and  $n$ . In particular, we have

$$\limsup_{n \rightarrow \infty} \|(f^n)^\bullet(T)\|^{1/n} \leq d_p(f) \quad \text{and} \quad \limsup_{n \rightarrow \infty} \|(f^n)_\bullet(S)\|^{1/n} \leq d_p(f).$$

*Proof.* We show the first inequality. The second one is proved in the same way. Let  $T_i$  be smooth positive closed forms as in Proposition 2.1. It follows from the definition of  $(f^n)^\bullet$  that any limit value of  $(f^n)^*(T_i)$  is larger than or equal to  $(f^n)^\bullet(T)$ . So, it is enough to bound the mass of  $(f^n)^*(T_i)$ . Since this mass can be computed cohomologically and since  $\{T_i\} \leq A\|T\|\{\omega_X^p\}$ , we obtain that  $\|(f^n)^*(T_i)\| \leq A\|T\|\lambda_p(f^n)$  for some constant  $A > 0$ . This completes the proof.  $\square$

The above proposition can be applied to currents of integration on submanifolds  $V$  of dimension  $k - p$  or  $p$  of  $X$  and gives an upper bound for the volume growth of the preimage or image of  $V$  by  $f^n$ .

It was shown in [6], [7] that dynamical degrees are bi-meromorphic invariants, i.e. conjugate maps have the same dynamical degrees. This property allows us to define

dynamical degrees for maps on singular manifolds having a kählerian desingularization. We will use the same argument in order to define dynamical degrees relative to an invariant meromorphic fibration.

Let  $f, g, \pi$  be as in Theorem 1.1. So,  $\pi$  defines a fibration and  $f$  preserves this fibration. Let us assume first that  $\pi$  is a *holomorphic* map. By the Bertini–Sard theorem, the set  $Z$  of critical values of  $\pi$  is a proper analytic subset of  $Y$ . Therefore,  $\pi: X \setminus \pi^{-1}(Z) \rightarrow Y \setminus Z$  defines a regular holomorphic fibration. Its fibers form a continuous family of smooth submanifolds of dimension  $k - l$  of  $X$ .

Let  $P_f$  and  $P_g$  denote the union of the critical set and the set of critical values of  $f$  and  $g$  respectively. They contain the indeterminacy sets of  $f$  and of  $g$ . A fiber  $L_y := \pi^{-1}(y)$  with  $y \in Y \setminus Z$  is called *generic* if for every  $n \geq 0$

- (a)  $g^n(y)$  and  $g^{-n}(y)$  do not intersect  $P_g$ ;
- (b) For every point  $b$  in  $g^n(y) \cup g^{-n}(y)$ , no component of  $L_b$  is contained in  $P_f$ .

Denote by  $\Sigma$  the set of  $y$  such that  $L_y$  is generic. Observe that  $Y \setminus \Sigma$  is contained in a finite or countable union of proper analytic subsets of  $Y$ . So,  $\Sigma$  is connected. We also have  $g(\Sigma) = g^{-1}(\Sigma) = \Sigma$ . We will use the following lemma for  $\nu = \omega_Y^l$  and for  $\nu = [d_l(g)]^{-n}(g^n)^*(\omega_Y^l)$ .

**Lemma 3.2.** *Let  $L_y$  be a generic fiber as above. Let  $\nu$  be a probability measure on  $Y$  which has no mass on proper analytic subsets of  $Y$ . Then, for  $0 \leq p \leq k - l$  and for  $n \geq 0$ , the 6 positive closed currents*

$$d_l(g)^{-n}(f^n)^\bullet(\omega_X^p \wedge [L_y]), \quad (f^n)^\bullet(\omega_X^p) \overset{\circ}{\wedge} [L_y], \quad (f^n)^\bullet(\omega_X^p) \overset{\circ}{\wedge} \pi^*(\nu)$$

and

$$(f^n)_\bullet(\omega_X^{k-l-p} \wedge [L_y]), \quad d_l(g)^{-n}(f^n)_*(\omega_X^{k-l-p}) \overset{\circ}{\wedge} [L_y], \\ d_l(g)^{-l}(f^n)_*(\omega_X^{k-l-p}) \overset{\circ}{\wedge} \pi^*(\nu)$$

have the same mass. In particular, their mass does not depend on  $y \in \Sigma$ .

*Proof.* For  $y \in \Sigma$ , define

$$\varphi(y) := d_l(g)^{-n} \|(f^n)^\bullet(\omega_X^p \wedge [L_y])\| \quad \text{and} \quad \psi(y) := \|(f^n)_\bullet(\omega_X^{k-l-p} \wedge [L_y])\|.$$

It is not difficult to see that these functions are continuous on  $\Sigma$ . We have

$$\begin{aligned} \varphi(y) &= d_l(g)^{-n} \langle (f^n)^\bullet(\omega_X^p \wedge [L_y]), \omega_X^{k-l-p} \rangle \\ &= d_l(g)^{-n} \langle \omega_X^p, [L_y] \overset{\circ}{\wedge} (f^n)_*(\omega_X^{k-l-p}) \rangle. \end{aligned}$$

It follows that

$$\varphi = d_l(g)^{-n} \pi_*(\omega_X^p \wedge (f^n)_*(\omega_X^{k-l-p}))$$

in the sense of currents on  $Y$ . Therefore,  $\varphi$  defines a closed 0-current on  $Y$  and it should be constant on  $\Sigma$ .

We also deduce from the above computation that

$$\varphi(y) = d_l(g)^{-n} \|(f^n)_*(\omega_X^{k-l-p}) \overset{\circ}{\wedge} [L_y]\|.$$

Since  $\nu$  has no mass on  $Y \setminus \Sigma$ , we obtain

$$\varphi = \int \varphi(y) d\nu = d_l(g)^{-n} \|(f^n)_*(\omega_X^{k-l-p}) \overset{\circ}{\wedge} \pi^*(\nu)\|.$$

In the same way, we prove that  $\psi$  is constant on  $\Sigma$  and

$$\psi = \|(f^n)^*(\omega_X^p) \overset{\circ}{\wedge} [L_y]\| = \|(f^n)^*(\omega_X^p) \overset{\circ}{\wedge} \pi^*(\nu)\|.$$

It remains to check that  $\varphi = \psi$ . Using that  $\psi$  is constant and  $\#g^{-n}(y) = d_l(g)^n$ , we have

$$\varphi = d_l(g)^{-n} \|(f^n)^*(\omega_X^p) \overset{\circ}{\wedge} [L_y]\| = d_l(g)^{-n} \sum_{b \in g^{-n}(y)} \|(f^n)^*(\omega_X^p) \overset{\circ}{\wedge} [L_b]\| = \psi.$$

This completes the proof. □

Define  $\lambda_p(f^n|\pi)$  the mass of the currents in Lemma 3.2. We have in particular

$$\lambda_p(f^n|\pi) = \|(f^n)^*(\omega_X^p) \wedge \pi^*(\omega_Y^l)\|.$$

Recall that  $\pi$  is holomorphic and then  $\pi^*(\omega_Y^l)$  is smooth.

**Proposition 3.3.** *The sequence  $\lambda_p(f^n|\pi)^{1/n}$  converges to a constant  $d_p(f|\pi)$ . Let  $T$  be a positive closed  $(p+l, p+l)$ -current and  $S$  a positive closed  $(k-p, k-p)$ -current on  $X$  which are supported on a generic fiber  $L_y$ . Then*

$$\|(f^n)^\bullet(T)\| \leq A_y \|T\| d_l(g)^n \lambda_p(f^n|\pi) \quad \text{and} \quad \|(f^n)_\bullet(S)\| \leq A_y \|S\| \lambda_p(f^n|\pi)$$

for some constant  $A_y > 0$  independent of  $T$  and  $S$ . In particular, we have

$$\limsup_{n \rightarrow \infty} \|(f^n)^\bullet(T)\|^{1/n} \leq d_l(g) d_p(f|\pi)$$

and

$$\limsup_{n \rightarrow \infty} \|(f^n)_\bullet(S)\|^{1/n} \leq d_p(f|\pi).$$

*Proof.* Fix a generic fiber  $L_y$  with  $y \in \Sigma$ . We will show that

$$\lambda_p(f^{n+m}|\pi) \leq A_y \lambda_p(f^n|\pi) \lambda_p(f^m|\pi)$$

for some constant  $A_y > 0$  and for all  $n, m \geq 0$ . This will imply the first assertion because the sequence  $A_y \lambda_p(f^n|\pi)$  is sub-multiplicative.

Since  $L_y$  is a compact Kähler manifold, we can apply Proposition 2.1 to  $L_y$ . Let  $b$  be a point in  $\Sigma$  such that  $g^m(b) = y$ . Define  $R := (f^m)_\bullet(\omega_X^{k-l-p} \wedge [L_b])$ . This is a positive closed  $(k-p, k-p)$ -current on  $X$  which is also a  $(k-l-p, k-l-p)$ -current on  $L_y$ . By Lemma 3.2, we have  $\|R\| = \lambda_p(f^m|\pi)$ . Therefore, there are smooth positive closed  $(k-l-p, k-l-p)$ -forms  $\Theta_i$  on  $L_y$  which converge to a current  $\Theta \geq R$ . Moreover, we have  $\{\Theta_i\} \leq A_y \lambda_p(f^m|\pi) \{\omega_{X|L_y}^{k-l-p}\}$  for some constant  $A_y > 0$ , where the inequality is considered in  $H^*(L_y, \mathbb{R})$ .

Let  $h$  denote the restriction of  $f^n$  to  $L_y$ . It defines a meromorphic map from  $L_y$  to  $L_{g^n(y)}$ . Since the mass of a positive closed current can be computed cohomologically, we obtain

$$\begin{aligned} \lambda_p(f^{n+m}|\pi) &= \|(f^n)_\bullet(R)\| \leq \liminf_{i \rightarrow \infty} \|h_\star(\Theta_i)\| \leq A_y \lambda_p(f^m|\pi) \|h_\star(\omega_{X|L_y}^{k-l-p})\| \\ &= A_y \lambda_p(f^m|\pi) \|(f^n)_\bullet(\omega_X^{k-l-p} \wedge [L_y])\| \\ &= A_y \lambda_p(f^m|\pi) \lambda_p(f^n|\pi). \end{aligned}$$

This implies the first assertion in the proposition. The rest is proved in the same way using the semi-regularization result for  $T$  and  $S$  on  $L_y$ , see also Proposition 3.1.  $\square$

We call  $d_p(f|\pi)$  the *dynamical degree of order  $p$  of  $f$  relative to  $\pi$* . The convergence in Proposition 3.3 implies that  $d_p(f^n|\pi) = d_p(f|\pi)^n$ .

**Remark 3.4.** Our choice of  $\Sigma$  simplifies the calculus on currents but several properties above still hold for some  $y \notin \Sigma$ . For example, if  $y$  is a fixed point of  $g$  which is not a critical value of  $\pi$  and if no component of  $L_y$  is contained in the critical set of  $f$ , then  $d_p(f|\pi) = d_p(f|_{L_y})$ . The proof is left to the reader.

The next result shows that the relative dynamical degrees are bi-meromorphic invariants. Consider a bi-meromorphic map  $\tau : (\tilde{X}, \omega_{\tilde{X}}) \rightarrow (X, \omega_X)$  between compact Kähler manifolds. Define  $\tilde{\pi} := \pi \circ \tau$  and  $\tilde{f} := \tau^{-1} \circ f \circ \tau$ . Then,  $\tilde{f}$  is a dominant meromorphic map conjugate to  $f$  and  $\tilde{\pi} \circ \tilde{f} = g \circ \tilde{\pi}$ .

**Proposition 3.5.** *Assume that  $\tilde{\pi}$  is holomorphic. Then*

$$d_p(f|\pi) = d_p(\tilde{f}|\tilde{\pi})$$

for  $0 \leq p \leq k-l$ .

*Proof.* Since  $\tau$  is bi-meromorphic,  $\tau_*\tilde{\pi}^*(\omega_Y^l) = \pi^*(\omega_Y^l)$  and  $\tilde{f}^n = \tau^{-1} \circ f^n \circ \tau$ , we have

$$\begin{aligned} \lambda_p(\tilde{f}^n|\tilde{\pi}) &= \langle (\tilde{f}^n)^*(\omega_{\tilde{X}}^p) \wedge \tilde{\pi}^*(\omega_Y^l), \omega_{\tilde{X}}^{k-l-p} \rangle \\ &= \langle \tau^*(f^n)^*\tau_*(\omega_X^p) \wedge \omega_{\tilde{X}}^{k-l-p}, \tilde{\pi}^*(\omega_Y^l) \rangle \\ &= \langle (f^n)^*\tau_*(\omega_X^p) \wedge \tau_*(\omega_{\tilde{X}}^{k-l-p}), \pi^*(\omega_Y^l) \rangle. \end{aligned}$$

Using the semi-regularization result for  $\tau_*(\omega_{\tilde{X}}^{k-l-p})$ , we deduce that

$$\lambda_p(\tilde{f}^n|\tilde{\pi}) \leq A \langle (f^n)^*\tau_*(\omega_X^p) \wedge \omega_X^{k-l-p}, \pi^*(\omega_Y^l) \rangle$$

for some constant  $A > 0$ . Then, using a semi-regularization of  $\tau_*(\omega_X^p)$ , we obtain

$$\lambda_p(\tilde{f}^n|\tilde{\pi}) \leq A' \langle (f^n)^*(\omega_X^p) \wedge \omega_X^{k-l-p}, \pi^*(\omega_Y^l) \rangle = A' \lambda_p(f^n|\pi)$$

for some constant  $A' > 0$ . It follows that  $d_p(\tilde{f}|\tilde{\pi}) \leq d_p(f|\pi)$ . The converse inequality is proved in the same way.  $\square$

The last proposition allows us to define relative dynamical degrees in the general case. Assume now that  $f$  preserves a meromorphic fibration  $\pi : X \rightarrow Y$ , i.e.  $\pi \circ f = g \circ \pi$  as in Theorem 1.1. Let  $\Gamma$  denote the closure of the graph of  $\pi$  in  $X \times Y$ . Then  $\Gamma$  is an irreducible analytic set of dimension  $k$  which is bi-meromorphic to  $X$ . Let  $\sigma : \tilde{X} \rightarrow \Gamma$  be a desingularization of  $\Gamma$  which can be constructed using a blow-up along the singularities. By Blanchard's theorem [2],  $\tilde{X}$  is a compact Kähler manifold. Then,  $\tau := \tau_X \circ \sigma$  is a bi-meromorphic map from  $\tilde{X}$  to  $X$ . Define also  $\tilde{\pi} := \tau_Y \circ \sigma$  and  $\tilde{f} := \tau^{-1} \circ f \circ \tau$ . The map  $\tilde{\pi}$  is holomorphic and  $\tilde{\pi} \circ \tilde{f} = g \circ \tilde{\pi}$ . Define the dynamical degree of order  $p$  of  $f$  relative to  $\pi$  by

$$d_p(f|\pi) := d_p(\tilde{f}|\tilde{\pi}).$$

Proposition 3.5 implies that the definition does not depend on the choice of  $\sigma$ . The following result is a consequence of a theorem by Khovanskii, Teissier and Gromov.

**Proposition 3.6.** *The function  $p \mapsto \log d_p(f|\pi)$  is concave for  $0 \leq p \leq k - l$ . In particular,  $d_p(f|\pi) \geq 1$  for  $0 \leq p \leq k - l$ .*

*Proof.* We can assume that  $\pi$  is holomorphic. We have to show that

$$d_{p-1}(f|\pi)d_{p+1}(f|\pi) \leq d_p(f|\pi)^2.$$

For this purpose, it is enough to check that

$$\lambda_{p-1}(f^n|\pi)\lambda_{p+1}(f^n|\pi) \leq \lambda_p(f^n|\pi)^2.$$

Observe that for non-critical values  $y$  of  $\pi$ , the fibers  $L_y$  are not necessarily connected but they contain the same number  $s$  of components. The family of these components is connected since  $X$  is connected. It defines a covering of degree  $s$  over the set of non-critical values of  $\pi$ . Let  $\Sigma'$  denote the parameter space for the components  $L'_y$  of  $L_y$  with  $y \in \Sigma$ . We may think of  $\Sigma'$  as a covering of degree  $s$  over  $\Sigma$ . We can then prove as in Lemma 3.2 that the function  $L'_y \mapsto \|(f^n)^*(\omega_X^p) \frown [L'_y]\|$  is constant on  $\Sigma'$ . Therefore, it is equal to  $s^{-1} \|(f^n)^*(\omega_X^p) \frown [L_y]\|$  and then to  $s^{-1} \lambda_p(f^n|\pi)$ .

Let  $h$  be the restriction of  $f^n$  to  $L_1 := L'_y$  and define  $L_2 := h(L_1)$ . Let  $\Gamma$  denote the graph of  $h$  in  $L_1 \times L_2$  and  $\tau: \hat{\Gamma} \rightarrow \Gamma$  a desingularization of  $\Gamma$  using some blow-up along the singularities. By Blanchard's theorem [2],  $\hat{\Gamma}$  is a compact Kähler manifold. Denote by  $\tau_1: \hat{\Gamma} \rightarrow L_1$  and  $\tau_2: \hat{\Gamma} \rightarrow L_2$  the canonical projections. We have  $h = \tau_2 \circ \tau_1^{-1}$ . Define  $\omega_1 := \tau_1^*(\omega_X)$  and  $\omega_2 := \tau_2^*(\omega_X)$ . We deduce from the above discussion that

$$s^{-1} \lambda_p(f^n|\pi) = \|(h^n)^*(\omega_X^p)\| = \int_{\hat{\Gamma}} \omega_1^{k-l-p} \wedge \omega_2^p.$$

If  $\gamma_p$  denotes the last integral, Gromov proved in [9] that  $p \mapsto \log \gamma_p$  is concave, i.e.  $\gamma_{p-1} \gamma_{p+1} \leq \gamma_p^2$ , when  $\omega_1$  and  $\omega_2$  are Kähler forms. By continuity, this still holds in our case where these forms are only smooth positive and closed. Hence,  $p \mapsto \log d_p(f|\pi)$  is concave.

In order to deduce the second assertion of the proposition, it is enough to show that  $d_0(f|\pi) = 1$  and  $d_{k-l}(f|\pi) \geq 1$ . For  $y$  generic, we have

$$\lambda_0(f^n|\pi) = d_l(g)^{-n} \|(f^n)^\bullet [L_y]\| = d_l(g)^{-n} \sum_{b \in g^{-n}(y)} \|[L_b]\|.$$

Hence,  $\lambda_0(f^n|\pi)$  is independent of  $n$  since  $\#g^{-n}(y) = d_l(g)^n$  and the mass of  $[L_b]$ , with  $b \in \Sigma$ , is independent of  $b$ . It follows that  $d_0(f|\pi) = 1$ .

We also have for  $y$  generic and  $b \in g^{-n}(y)$

$$\lambda_{k-l}(f^n|\pi) = \|(f^n)^\bullet [L_b]\| \geq \|[L_y]\|.$$

So, the sequence  $\lambda_{k-l}(f^n|\pi)$  is bounded from below by a positive constant. Therefore,  $d_{k-l}(f|\pi) \geq 1$ . This completes the proof of the lemma. Note that we can show that  $d_{k-l}(f|\pi)$  is the number of points in a generic fiber of the restriction of  $f$  to  $L_y$ . □

Consider now some examples, see also [1], [14], [21], [22].

**Example 3.7.** Let  $X = Y \times Z$  be the product of two compact Kähler manifolds and  $\pi: X \rightarrow Y$  the canonical projection. Consider  $f(y, z) := (g(y), h(z))$  where

$g: Y \rightarrow Y$  and  $h: Z \rightarrow Z$  are dominant meromorphic maps. So,  $f$  is semi-conjugate to  $g$ . The relative dynamical degree  $d_p(f|\pi)$  is equal to  $d_p(h)$ . We easily deduce from the definition of dynamical degrees that

$$d_p(f) = \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} d_j(g) d_{p-j}(h).$$

There are more interesting examples of maps on the product  $Y \times Z$ . Let  $F$  be a compact Kähler manifold. Assume that  $F$  is also the parameter space of a meromorphic family of meromorphic self-maps of  $Z$ . Let  $\tau: Y \rightarrow F$  be a meromorphic map. Then  $f(y, z) := (g(y), \tau(y)(z))$  is a meromorphic self-map of  $Y \times Z$  which preserves the fibration  $\pi$ . The example is also interesting when  $\tau(y)$  is holomorphic for generic  $y$  or when a Zariski open set  $G$  of  $F$  is a Lie group and  $\tau$  is a morphism from  $G$  to the group of bi-meromorphic maps of  $Z$ .

**Example 3.8.** Let  $g: Y \rightarrow Y$  be a dominant meromorphic map on a compact Kähler manifold  $Y$ . It induces a meromorphic self-map  $f$  on the projectivization  $X := \mathbb{P}T_Y$  of the holomorphic tangent bundle of  $Y$ . The map  $f$  preserves the fibration associated to the canonical projection from  $X$  onto  $Y$  and is semi-conjugate to  $g$ . This example and some applications were considered in [5].

#### 4. Proofs of the results

We first prove Theorem 1.1. Since the dynamical degrees are bi-meromorphic invariants, Proposition 3.5 allows us to assume that  $\pi$  is a holomorphic map. Since  $X$  is projective, we can construct a dominant meromorphic map  $v: X \rightarrow \mathbb{P}^{k-l}$ . Indeed, it is enough to embed  $X$  in a projective space and choose a generic central projection on  $\mathbb{P}^{k-l}$ . Replacing  $X$  with a desingularization of the graph of  $v$  allows to assume that  $v$  is holomorphic. Consider the holomorphic map  $\Pi: X \rightarrow Y \times \mathbb{P}^{k-l}$  defined by

$$\Pi(x) := (\pi(x), v(x)).$$

Since the chosen central projection is generic, the intersection of a generic fiber of  $\pi$  and a generic fiber of  $v$  is finite. Therefore,  $\Pi$  is dominant.

Our proof is based on a delicate calculus on currents. If  $X = Y \times \mathbb{P}^{k-l}$  and  $\pi$  is the canonical projection onto  $Y$ , the proof is simpler and the properties obtained in Section 2 can be directly applied. A rough idea is to reduce the general case to the particular case using the map  $\Pi$ . In other words, we use the fact that  $f$  is, in some sense, “semi-conjugate” to the multi-valued map  $\Pi \circ f \circ \Pi^{-1}$  which is defined on  $Y \times \mathbb{P}^{k-l}$ .

Let  $\omega_{\text{FS}}$  denote the Fubini–Study form on  $\mathbb{P}^{k-l}$ . For simplicity, the canonical pull-back of  $\omega_Y$  and  $\omega_{\text{FS}}$  to  $Y \times \mathbb{P}^{k-l}$  are still denoted by  $\omega_Y$  and  $\omega_{\text{FS}}$ . In particular,

$\Pi^*(\omega_Y)$  and  $\pi^*(\omega_Y)$  represent the same form on  $X$ . We consider on  $Y \times \mathbb{P}^{k-l}$  the Kähler form  $\omega := \omega_Y + \omega_{\text{FS}}$ . Our calculus will involve the quantities  $a_{q,p}(n)$  defined for  $n \geq 0$ ,  $0 \leq q \leq k-l$  and  $q \leq p \leq l+q$  by

$$\begin{aligned} a_{q,p}(n) &:= \|\Pi_*(f^n)^* \Pi^*(\omega^p) \wedge \omega_Y^{l-p+q}\| \\ &= \langle \Pi_*(f^n)^* \Pi^*(\omega^p), \omega_Y^{l-p+q} \wedge \omega^{k-l-q} \rangle \\ &= \langle (f^n)^* \Pi^*(\omega^p), \Pi^*(\omega_Y^{l-p+q} \wedge \omega^{k-l-q}) \rangle \\ &= \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^{l-p+q}), \Pi^*(\omega^{k-l-q}) \rangle. \end{aligned}$$

Observe that

$$a_{q,p}(n) \geq \alpha_{p-q}(\Pi_*(f^n)^* \Pi^*(\omega^p)),$$

where  $\alpha_{p-q}(\cdot)$  is defined in (1).

**Lemma 4.1.** *There is a constant  $A > 0$  independent of  $p, n$  such that*

$$A^{-1} \lambda_p(f^n | \pi) \leq a_{p,p}(n) \leq A \lambda_p(f^n | \pi)$$

*In particular,  $[a_{p,p}(n)]^{1/n}$  converges to  $d_p(f | \pi)$ .*

*Proof.* Since the pull-back of a smooth form under  $\Pi$  is smooth, we have

$$\begin{aligned} a_{p,p}(n) &= \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l), \Pi^*(\omega^{k-l-p}) \rangle \\ &\leq A \langle (f^n)^*(\omega_X^p) \wedge \pi^*(\omega_Y^l), \omega_X^{k-l-p} \rangle = A \lambda_p(f^n | \pi) \end{aligned}$$

for some constant  $A > 0$ . This gives the second inequality in the lemma.

Define  $T := \Pi_*(\omega_X^p)$ . Since  $\Pi^\bullet(T) \geq \omega_X^p$ , we have

$$\lambda_p(f | \pi) = \|(f^n)^*(\omega_X^p) \wedge \pi^*(\omega_Y^l)\| \leq \|(f^n)^\bullet \Pi^\bullet(T) \wedge \pi^*(\omega_Y^l)\|.$$

We apply Proposition 2.1 to the current  $T$  on  $Y \times \mathbb{P}^{k-l}$  which is an  $L^1$  form smooth on a Zariski open set. Let  $T_i$  be as in that proposition with  $\{T_i\} \leq A\{\omega^p\}$  for some constant  $A > 0$ . If  $S := \Pi_*(\omega_X^{k-l-p})$ , we have  $\Pi^\bullet(S) \geq \omega_X^{k-l-p}$  and hence

$$\begin{aligned} \lambda_p(f | \pi) &\leq \liminf_{i \rightarrow \infty} \|(f^n)^* \Pi^*(T_i) \wedge \pi^*(\omega_Y^l)\| \\ &\leq A \|(f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l)\| \\ &= A \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l), \omega_X^{k-l-p} \rangle \\ &\leq A \|(f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l) \overset{\circ}{\wedge} \Pi^\bullet(S)\|. \end{aligned}$$

Now, we apply again Proposition 2.1, in particular its last assertion, to the current  $S$  which is an  $L^1$  form smooth on a Zariski open set. If  $S_i$  are smooth forms satisfying that proposition, the latter expression is bounded from above by

$$\begin{aligned} \liminf_{i \rightarrow \infty} \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l), \Pi^*(S_i) \rangle \\ \lesssim \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^l), \Pi^*(\omega^{k-l-p}) \rangle. \end{aligned}$$

The last integral is equal to  $a_{p,p}(n)$ . The first inequality in the lemma follows.  $\square$

Define for  $0 \leq p \leq k$

$$b_p(n) := \sum_{\max\{0, p-l\} \leq q \leq \min\{p, k-l\}} a_{q,p}(n).$$

We have the following lemma.

**Lemma 4.2.** *The sequence  $b_p(n)^{1/n}$  converges to  $d_p(f)$ .*

*Proof.* Since  $\Pi^*(\omega^p)$ ,  $\pi^*(\omega_Y^{l-p+q})$  and  $\Pi^*(\omega^{k-l-q})$  are smooth on  $X$ , we have

$$\begin{aligned} a_{q,p}(n) &= \langle (f^n)^* \Pi^*(\omega^p) \wedge \pi^*(\omega_Y^{l-p+q}), \Pi^*(\omega^{k-l-q}) \rangle \\ &\leq A \|(f^n)^*(\omega_X^p)\| = A \lambda_p(f^n) \end{aligned}$$

for some constant  $A > 0$ . We deduce that  $\limsup b_p(n)^{1/n} \leq d_p(f)$ .

It remains to check that  $\liminf b_p(n)^{1/n} \geq d_p(f)$ . For this purpose, we only need to show that  $\lambda_p(f^n) \leq A b_p(n)$  for some constant  $A > 0$ . Define  $T := \Pi_*(f^n)^* \Pi^*(\omega^p)$ . We prove that  $\lambda_p(f^n) \lesssim \|T\| \lesssim b_p(n)$  which will imply the result.

Define  $S := \Pi_*(\omega_X^p)$ . We have  $\Pi^\bullet(S) \geq \omega_X^p$ . Therefore,

$$\lambda_p(f^n) = \langle (f^n)^*(\omega_X^p), \omega_X^{k-p} \rangle \leq \langle (f^n)^\bullet \Pi^\bullet(S), \omega_X^{k-p} \rangle.$$

Using a semi-regularization of  $S$ , we deduce that

$$\lambda_p(f^n) \lesssim \langle (f^n)^* \Pi^*(\omega^p), \omega_X^{k-p} \rangle.$$

Define  $R := \Pi_*(\omega_X^{k-p})$ . We also have  $\Pi^\bullet(R) \geq \omega_X^{k-p}$ . We obtain as above using a semi-regularization of  $R$  that

$$\begin{aligned} \lambda_p(f^n) &\lesssim \|(f^n)^* \Pi^*(\omega^p) \overset{\circ}{\wedge} \Pi^\bullet(R)\| \lesssim \|(f^n)^* \Pi^*(\omega^p) \wedge \Pi^*(\omega^{k-p})\| \\ &= \langle \Pi_*(f^n)^* \Pi^*(\omega^p), \omega^{k-p} \rangle = \|T\|. \end{aligned}$$

Now, since  $\omega_Y^{l+1} = 0$  and  $\omega_{\text{FS}}^{k-l+1} = 0$ , we have

$$\begin{aligned} \|T\| &= \langle T, (\omega_Y + \omega_{\text{FS}})^{k-p} \rangle \lesssim \sum_{\max\{0, p-l\} \leq q \leq \min\{p, k-l\}} \langle T, \omega_Y^{l-p+q} \wedge \omega_{\text{FS}}^{k-l-q} \rangle \\ &\leq \sum_{\max\{0, p-l\} \leq q \leq \min\{p, k-l\}} a_{q,p}(n) = b_p(n). \end{aligned}$$

This completes the proof of the lemma.  $\square$

For every  $n \geq 0$  and  $0 \leq p \leq l$  define

$$c_p(n) := \lambda_p(g^n) = \|(g^n)^*(\omega_Y^p)\| = \langle (g^n)^*(\omega_Y^p), \omega_Y^{l-p} \rangle.$$

We have the following lemma.

**Lemma 4.3.** *There is a constant  $A > 0$  such that*

$$\langle \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q} \wedge \omega^q), \omega_Y^{l-p+p_0} \wedge \omega^{k-l-p_0} \rangle \leq A a_{p_0,q}(n) c_{p-q}(n)$$

for  $0 \leq p_0 \leq k-l$ ,  $p_0 \leq p \leq l+p_0$ ,  $p_0 \leq q \leq p$  and  $n \geq 0$ . Moreover, the above integral vanishes when  $q < p_0$ .

*Proof.* Observe that by definition of  $\Pi_*$

$$\begin{aligned} \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q} \wedge \omega^q) &= \Pi_*[(f^n)^* \Pi^*(\omega_Y^{p-q}) \overset{\circ}{\wedge} (f^n)^* \Pi^*(\omega^q)] \\ &\leq \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q}) \overset{\circ}{\wedge} \Pi_*(f^n)^* \Pi^*(\omega^q). \end{aligned}$$

Hence, the left hand side of the inequality in the lemma is smaller than or equal to

$$\langle \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q}) \overset{\circ}{\wedge} \Pi_*(f^n)^* \Pi^*(\omega^q), \omega_Y^{l-p+p_0} \wedge \omega^{k-l-p_0} \rangle.$$

Define  $T := \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q}) \wedge \omega_Y^{l-p+p_0}$  and  $S := \Pi_*(f^n)^* \Pi^*(\omega^q) \wedge \omega^{k-l-p_0}$ . Note that  $T$  and  $S$  are of bidegree  $(l-q+p_0, l-q+p_0)$  and  $(k-l+q-p_0, k-l+q-p_0)$  respectively. The quantity considered above is equal to the mass of the measure  $T \overset{\circ}{\wedge} S$ .

We first show that  $\alpha_j(T) = 0$  when  $j < l-q+p_0$  and  $\alpha_{l-q+p_0}(T) \leq A c_{p-q}(n)$  for some constant  $A > 0$ . Since  $\pi \circ f^n = g^n \circ \pi$ , we have

$$T = \Pi_*(f^n)^* \pi^*(\omega_Y^{p-q}) \wedge \omega_Y^{l-p+p_0} = \Pi_* \pi^\bullet(g^n)^*(\omega_Y^{p-q}) \wedge \omega_Y^{l-p+p_0}.$$

Hence,

$$\begin{aligned} \alpha_j(T) &= \langle \Pi_* \pi^\bullet(g^n)^*(\omega_Y^{p-q}) \wedge \omega_Y^{l-p+p_0}, \omega_Y^{l-j} \wedge \omega_{\text{FS}}^{k-2l+q-p_0+j} \rangle \\ &= \langle \pi^\bullet(g^n)^*(\omega_Y^{p-q}) \wedge \pi^*(\omega_Y^{l-p+p_0}), \pi^*(\omega_Y^{l-j}) \wedge \Pi^*(\omega_{\text{FS}}^{k-2l+q-p_0+j}) \rangle \\ &= \langle \pi^\bullet[(g^n)^*(\omega_Y^{p-q}) \wedge \omega_Y^{2l-p+p_0-j}], \Pi^*(\omega_{\text{FS}}^{k-2l+q-p_0+j}) \rangle. \end{aligned}$$

When  $j < l - q + p_0$ , the form in the brackets has bidegree  $\geq (l + 1, l + 1)$  and should vanish because  $\dim Y = l$ . Therefore,  $\alpha_j(T) = 0$  in that case. When  $j = l - q + p_0$ , this form defines a positive measure of mass  $\lambda_{p-q}(g^n)$ . Its cohomology class is equal to  $\lambda_{p-q}(g^n)\{\omega_Y^l\}$ . Therefore, using a semi-regularization as above, we obtain

$$\alpha_{l-q+p_0}(T) \lesssim \lambda_{p-q}(g^n)\langle \pi^*(\omega_Y^l), \Pi^*(\omega_{FS}^{k-l}) \rangle \leq Ac_{p-q}(n)$$

for some constant  $A > 0$ .

We deduce from Proposition 2.3 that  $\{T\} \lesssim c_{p-q}(n)\{\omega_Y^{l-q+p_0}\}$ . Using the semi-regularization in Proposition 2.4 for  $T$ , we obtain

$$\{T \overset{\circ}{\wedge} S\} \lesssim c_{p-q}(n)\|\omega_Y^{l-q+p_0} \wedge S\| = c_{p-q}(n)a_{p_0,q}(n).$$

This completes the proof of the first assertion in the lemma. For the second one, it is enough to observe that when  $q < p_0$ , we have  $\alpha_j(T) = 0$  for every  $j$  and hence  $T = 0$ . □

The following lemma is crucial in our proof.

**Lemma 4.4.** *There exists a constant  $A > 0$  such that for all  $0 \leq p_0 \leq k - l$ ,  $p_0 \leq p \leq l + p_0$  and all  $n, r \geq 1$*

$$a_{p_0,p}(nr) \leq A^r \sum \prod_{s=1}^r a_{p_{s-1},p_s}(n)c_{p-p_s}(n),$$

where the sum is taken over  $(p_1, \dots, p_r)$  with  $p_0 \leq p_1 \leq p_2 \leq \dots \leq p_r \leq p$  and  $p_{r-1} \leq k - l$ .

*Proof.* We proceed by induction on  $r$ . Clearly, the lemma is true for  $r = 1$ . Suppose the lemma true for  $r$ , we need to prove it for  $r + 1$ . In what follows, the constants  $A_i$  depend only on the geometry of  $X$  and  $Y$ .

Define  $T^{(r)} := \Pi_*(f^{nr})^*\Pi^*(\omega^p)$ . This is a positive closed  $L^1$  form, smooth on a dense Zariski open set. Observe that  $\Pi^\bullet\Pi_* \geq \text{id}$  on positive closed currents having no mass on proper analytic subsets of  $X$ . Therefore,

$$T^{(r+1)} \leq \Pi_*(f^n)^\bullet\Pi^\bullet\Pi_*(f^{nr})^\bullet\Pi^*(\omega^p) = \Pi_*(f^n)^\bullet\Pi^\bullet(T^{(r)}).$$

On the other hand, by Proposition 2.4, we can find a sequence of smooth positive closed  $(p, p)$ -forms  $T_i^{(r)}$  converging weakly to a positive closed current  $\tilde{T}^{(r)} \geq T^{(r)}$  such that

$$\alpha_{p-q}(T_i^{(r)}) \leq A_1\alpha_{p-q}(T^{(r)}) \leq A_1a_{q,p}(nr)$$

for  $\max\{0, p - l\} \leq q \leq \min\{p, k - l\}$  and  $A_1 > 0$  a constant. By Proposition 2.3, there is a constant  $A_2 > 0$  such that

$$\{T_i^{(r)}\} \leq A_2 \sum_{\max\{0, p-l\} \leq q \leq \min\{p, k-l\}} a_{q,p}(nr) \{\omega_Y^{p-q}\} \cup \{\omega_{FS}^q\}.$$

We deduce from the above discussion and Lemma 4.3 that

$$\begin{aligned} a_{p_0,p}(n(r + 1)) &= \langle T^{(r+1)}, \omega_Y^{l-p+p_0} \wedge \omega^{k-l-p_0} \rangle \\ &\leq \liminf_{i \rightarrow \infty} \langle \Pi_*(f^n)^* \Pi^*(T_i^{(r)}), \omega_Y^{l-p+p_0} \wedge \omega^{k-l-p_0} \rangle \\ &\leq A_2 \sum_{\substack{\max\{0, p-l\} \leq q \\ \leq \min\{p, k-l\}}} a_{q,p}(nr) \langle \Pi_*(f^n)^* \Pi^*(\omega_Y^{p-q} \wedge \omega_{FS}^q), \omega_Y^{l-p+p_0} \wedge \omega^{k-l-p_0} \rangle \\ &\leq A_3 \sum_{p_0 \leq q \leq \min\{p, k-l\}} a_{q,p}(nr) a_{p_0,q}(n) c_{p-q}(n) \end{aligned}$$

for some constant  $A_3 > 0$ . Consequently, the induction hypothesis implies the result. □

Theorem 1.1 is a consequence of the next two propositions.

**Proposition 4.5.** *We have*

$$d_p(f) \geq \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} d_j(g) d_{p-j}(f|\pi)$$

for  $0 \leq p \leq k$ .

*Proof.* Since  $\Pi^*(\omega_Y^j \wedge \omega^{p-j})$  is a smooth form, we have for some constant  $A > 0$

$$\|(f^n)^* \Pi^*(\omega_Y^j \wedge \omega^{p-j})\| \leq A \lambda_p(f^n).$$

So, by definition of dynamical degrees and Lemma 4.1, it is enough to bound  $\|(f^n)^* \Pi^*(\omega_Y^j \wedge \omega^{p-j})\|$  from below by a constant times  $\lambda_j(g^n) a_{p-j, p-j}(n)$ .

Fix a constant  $A > 0$  large enough. Using the identity  $\pi \circ f^n = g^n \circ \pi$  and that  $\Pi^*(\omega_Y^{l-j} \wedge \omega^{k-l-p+j})$  is smooth, we obtain

$$\begin{aligned} &A \|(f^n)^* \Pi^*(\omega_Y^j \wedge \omega^{p-j})\| \\ &\geq \langle (f^n)^* \Pi^*(\omega_Y^j \wedge \omega^{p-j}), \Pi^*(\omega_Y^{l-j} \wedge \omega^{k-l-p+j}) \rangle \\ &= \langle (f^n)^* \pi^*(\omega_Y^j) \overset{\circ}{\wedge} (f^n)^* \Pi^*(\omega^{p-j}), \pi^*(\omega_Y^{l-j}) \wedge \Pi^*(\omega^{k-l-p+j}) \rangle \\ &= \|(f^n)^* \pi^*(\omega_Y^j) \wedge \pi^*(\omega_Y^{l-j}) \overset{\circ}{\wedge} (f^n)^* \Pi^*(\omega^{p-j}) \wedge \Pi^*(\omega^{k-l-p+j})\| \\ &= \|\pi^*[(g^n)^*(\omega_Y^j) \wedge \omega_Y^{l-j}] \overset{\circ}{\wedge} (f^n)^* \Pi^*(\omega^{p-j}) \wedge \Pi^*(\omega^{k-l-p+j})\|. \end{aligned}$$

Observe that  $(g^n)^*(\omega_Y^j) \wedge \omega_Y^{l-j}$  is a positive measure of mass  $\lambda_j(g^n)$ . As in Lemma 3.2, we show that the last expression in the previous identities is equal to  $\lambda_j(g^n)$  times the mass of the restriction of  $(f^n)^* \Pi^*(\omega^{p-j}) \wedge \Pi^*(\omega^{k-l-p+j})$  to a generic fiber  $L_y$  of  $\pi$ . Therefore, it is also equal to

$$\lambda_j(g^n) \langle \pi^*(\omega_Y^l), (f^n)^* \Pi^*(\omega^{p-j}) \wedge \Pi^*(\omega^{k-l-p+j}) \rangle = \lambda_j(g^n) a_{p-j, p-j}(n).$$

This completes the proof. □

**Proposition 4.6.** *We have*

$$d_p(f) \leq \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} d_j(g) d_{p-j}(f|\pi)$$

for  $0 \leq p \leq k$ .

*Proof.* For every  $0 \leq p \leq k$  and  $n \geq 0$  let

$$\mu_p(n) := \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} c_j(n) a_{p-j, p-j}(n).$$

Observe that for  $r > p$ , in Lemma 4.4, there are at most  $p$  indices  $s$  such that  $p_{s-1} < p_s$ . Moreover, the sum in that lemma contains at most  $(k+1)^r$  terms and the sum in the definition of  $b_p(n)$  contains at most  $p+1$  terms. We infer the following estimate

$$b_p(rn) \leq \left[ (p+1)(k+1)^r A^r b_0(n) \cdots b_p(n) \prod_{j=0}^l c_j(n) \right] \mu_p(n)^r.$$

We deduce that

$$\begin{aligned} [b_p(rn)]^{1/rn} &\leq (p+1)^{1/nr} (k+1)^{1/n} A^{1/n} [b_0(n)^{1/n} \cdots b_p(n)^{1/n}]^{1/r} \\ &\quad \cdot \left[ \prod_{j=0}^l c_j(n)^{1/n} \right]^{1/r} \mu_p(n)^{1/n}. \end{aligned}$$

Letting  $n$  tend to infinity, we obtain using Lemma 4.2 that

$$d_p(f) \leq [d_0(f) \cdots d_p(f)]^{1/r} \left[ \prod_{j=0}^l d_j(g) \right]^{1/r} \liminf_{n \rightarrow \infty} \mu_p(n)^{1/n}.$$

Now, letting  $r \rightarrow \infty$ , the first two factors in the right hand side tend to 1. Therefore, using Lemma 4.1, we obtain

$$d_p(f) \leq \liminf_{n \rightarrow \infty} \mu_p(n)^{1/n} = \max_{\max\{0, p-k+l\} \leq j \leq \min\{p, l\}} d_j(g) d_{p-j}(f|\pi).$$

This completes the proof. □

*Proof of Corollary 1.2.* When  $X$  and  $Y$  are projective, the corollary is a direct consequence of Theorem 1.1. We only used the projectivity in Proposition 2.4 applied to  $m := k - l$  and for the existence of  $v: X \rightarrow \mathbb{P}^{k-l}$ . This is superfluous when  $X$  and  $Y$  have the same dimension, i.e.  $k = l$ .  $\square$

*Proof of Corollary 1.3.* Let  $j, p$  be such that  $d_j(g) = \max_q d_q(g)$  and  $d_{p-j}(f|\pi) = \max_q d_q(f|\pi)$ . We have  $0 \leq j \leq l$  and  $0 \leq p - j \leq k - l$ . By Theorem 1.1,  $d_p(f)$  is the maximal dynamical degree of  $f$  and  $d_p(f) = d_j(g)d_{p-j}(f|\pi)$ . We have  $d_{p-1}(f) < d_p(f) < d_{p+1}(f)$ . Theorem 1.1 implies that

$$d_{j-1}(g) < d_j(g) < d_{j+1}(g) \quad \text{and} \quad d_{p-j-1}(f|\pi) < d_{p-j}(f|\pi) < d_{p-j+1}(f|\pi).$$

The log-concavity of  $d_q(g)$  and  $d_q(f|\pi)$  implies the result. Note that when  $j = 0, l$  or  $p - j = 0, k - l$ , in the above inequalities, one has to remove the expressions which are not meaningful.  $\square$

In the rest of the paper, we prove Corollary 1.4. Let  $K_X$  denote the canonical lines bundle of  $X$ . Let  $H^0(X, K_X^n)$  denote the space of holomorphic sections of  $K_X^n$  and  $H^0(X, K_X^n)^*$  its dual space. Assume that  $H^0(X, K_X^n)$  has positive dimension. If  $x$  is a generic point in  $X$ , the family  $H_x$  of sections which vanish at  $x$  is a hyperplane of  $H^0(X, K_X^n)$  passing through 0. So, the correspondence  $x \mapsto H_x$  defines a meromorphic map

$$\pi_n: X \rightarrow \mathbb{P} H^0(X, K_X^n)^*$$

from  $X$  to the projectivization of  $H^0(X, K_X^n)^*$  which is called an *Iitaka fibration* of  $X$ . Let  $Y_n$  denote the image of  $X$  by  $\pi_n$ . The *Kodaira dimension* of  $X$  is  $\kappa_X := \max_{n \geq 1} \dim Y_n$ . When  $H^0(X, K_X^n) = 0$  for every  $n \geq 1$ , the Kodaira dimension of  $X$  is  $-\infty$ . We have the following result.

**Theorem 4.7** ([14], [18]). *Let  $f: X \rightarrow X$  be a dominant meromorphic map. Assume that  $\kappa_X \geq 1$ . Then  $f$  preserves the Iitaka fibration  $\pi_n: X \rightarrow Y_n$ . Moreover, the map  $g: Y_n \rightarrow Y_n$  induced by  $f$  is periodic, i.e.  $g^N = \text{id}$  for some integer  $N \geq 1$ .*

*Proof of Corollary 1.4.* Assume in order to get a contradiction that  $\kappa_X \geq 1$ . Let  $n \geq 1$  be such that  $l := \dim Y_n \geq 1$ . Replacing  $f$  with an iterate, we can assume that  $g = \text{id}$ . A priori,  $Y_n$  may be singular, but we can use a blow-up and assume that  $Y_n$  is smooth. We have  $d_j(g) = 1$  for  $0 \leq j \leq l$ . This contradicts Corollary 1.3. Note that in order to prove that  $d_j(g) = 1$ , instead of Theorem 4.7, it is enough to use the weaker result that  $g$  is induced by a linear endomorphism of  $\mathbb{P} H^0(X, K_X^n)^*$ .  $\square$

## References

- [1] E. Amerik and F. Campana, Fibrations méromorphes sur certaines variétés à fibré canonique trivial. *Pure Appl. Math. Q.* **4** (2008), no. 2, part 1, 509–545. [Zbl 1143.14035](#) [MR 1143.14035](#)
- [2] A. Blanchard, Sur les variétés analytiques complexes. *Ann. Sci. École Norm. Sup. (3)* **73** (1956), 157–202. [Zbl 0073.37503](#) [MR 0087184](#)
- [3] S. Cantat, Dynamique des automorphismes des surfaces projectives complexes. *C. R. Acad. Sci. Paris Sér. I Math.* **328** (10) (1999), 901–906. [Zbl 0943.37021](#) [MR 1689873](#)
- [4] J.-P. Demailly, Complex analytic and differential geometry. Available at <http://www-fourier.ujf-grenoble.fr/~demailly/manuscripts/agbook.pdf>.
- [5] T.-C. Dinh, Suites d'applications méromorphes multivaluées et courants laminaires. *J. Geom. Anal.* **15** (2005), no. 2, 207–227. [Zbl 1085.37039](#) [MR 2152480](#)
- [6] T.-C. Dinh and N. Sibony, Une borne supérieure pour l'entropie topologique d'une application rationnelle. *Ann. of Math. (2)* **161** (2005), 1637–1644. [Zbl 1084.54013](#) [MR 2180409](#)
- [7] T.-C. Dinh and N. Sibony, Regularization of currents and entropy. *Ann. Sci. École Norm. Sup. (4)* **37** (2004), no. 6, 959–971. [Zbl 1074.53058](#) [MR 2119243](#)
- [8] T.-C. Dinh and N. Sibony, Dynamics in several complex variables: endomorphisms of projective spaces and polynomial-like mappings. In *Holomorphic dynamical systems*, Lecture Notes in Math. 1998, Springer-Verlag, Berlin 2010, 165–294. [Zbl 05879453](#) [MR 2648690](#)
- [9] M. Gromov, Convex sets and Kähler manifolds. In *Advances in differential geometry and topology*, World Scientific Publishing Co., Inc., Teaneck, NJ, 1998, 1–38. [Zbl 0770.53042](#) [MR 1095529](#)
- [10] M. Gromov, On the entropy of holomorphic maps. *Enseign. Math.* **49** (2003), 217–235. [Zbl 1080.37051](#) [MR 2026895](#)
- [11] V. Guedj, Propriétés ergodiques des applications rationnelles. *Panoramas & Synthèses*, to appear, [arXiv:math/0611302](https://arxiv.org/abs/math/0611302) [math.CV].
- [12] B. Hasselblatt and J. Propp, Degree-growth of monomial maps. *Ergodic Theory Dynam. Systems* **27** (2007), no. 5, 1375–1397. [Zbl 1143.37032](#) [MR 2358970](#)
- [13] A. G. Khovanskii, The geometry of convex polyhedra and algebraic geometry. *Uspehi Mat. Nauk.* **34** (1979), no. 4, 160–161.
- [14] N. Nakayama and D.-Q. Zhang, Building blocks of étale endomorphisms of complex projective manifolds. *Proc. London Math. Soc.* **99** (2009), 725–756. [Zbl 1185.14012](#) [MR 2551469](#)
- [15] N. Sibony, Dynamique des applications rationnelles de  $\mathbb{P}^k$ . In *Dynamique et géométrie complexes* (Lyon, 1997), Panoramas et Synthèses 8, Soc. Math. France, Paris 1999, 97–185. [Zbl 1020.37026](#) [MR 1760844](#)
- [16] H. Skoda, Prolongement des courants positifs, fermés de masse finie. *Invent. Math.* **66** (1982), 361–376. [Zbl 0488.58002](#) [MR 0662596](#)
- [17] B. Teissier, Du théorème de l'index de Hodge aux inégalités isopérimétriques. *C. R. Acad. Sci. Paris Sér. A-B* **288** (1979), no. 4, 287–289. [Zbl 0406.14011](#) [MR 0524795](#)

- [18] K. Ueno, *Classification theory of algebraic varieties and compact complex spaces*. Notes written in collaboration with P. Cherenack, Lecture Notes in Math. 439, Springer-Verlag, Berlin 1975. [Zbl 0299.14007](#) [MR 0506253](#)
- [19] C. Voisin, *Hodge theory and complex algebraic geometry. I*. Cambridge Stud. Adv. Math. 76, Cambridge University Press, Cambridge 2002. [Zbl 1005.14002](#) [MR 2451566](#)
- [20] Y. Yomdin, Volume growth and entropy. *Israel J. Math.*, **57** (1987) 285–300. [Zbl 0641.54036](#) [MR 0889979](#)
- [21] D.-Q. Zhang, Cohomologically hyperbolic endomorphisms of complex manifolds. *Internat. J. Math.* **20** (2009), 803–816. [Zbl 1172.14009](#) [MR 2548399](#)
- [22] D.-Q. Zhang, Polarized endomorphisms of uniruled varieties (with an Appendix by Y. Fujimoto and N. Nakayama). *Compos. Math.* **146** (2010), no. 1, 145–168. [Zbl 05706184](#) [MR 2581245](#)

Received March 29, 2009

Tien-Cuong Dinh, UPMC, Université Paris 06, UMR 7586, Institut de Mathématiques de Jussieu, 4 place Jussieu, 75005 Paris, France

E-mail: [dinh@math.jussieu.fr](mailto:dinh@math.jussieu.fr)

<http://www.math.jussieu.fr/~dinh>

Viêt-Anh Nguyễn, Équipe Analyse Harmonique, Mathématiques, Bâtiment 425, Université Paris-Sud 11, 91405 Orsay Cedex, France

E-mail: [vietanh.nguyen@math.u-psud.fr](mailto:vietanh.nguyen@math.u-psud.fr)

<http://www.math.u-psud.fr/~vietanh>