

Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	86 (2011)
Artikel:	Actions of automorphism groups of free groups on homology spheres and acyclic manifolds
Autor:	Bridson, Martin R. / Vogtmann, Karen
DOI:	https://doi.org/10.5169/seals-283445

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Actions of automorphism groups of free groups on homology spheres and acyclic manifolds

Martin R. Bridson and Karen Vogtmann*

Abstract. For $n \geq 3$, let $\mathrm{SAut}(F_n)$ denote the unique subgroup of index two in the automorphism group of a free group. The standard linear action of $\mathrm{SL}(n, \mathbb{Z})$ on \mathbb{R}^n induces non-trivial actions of $\mathrm{SAut}(F_n)$ on \mathbb{R}^n and on \mathbb{S}^{n-1} . We prove that $\mathrm{SAut}(F_n)$ admits no non-trivial actions by homeomorphisms on acyclic manifolds or spheres of smaller dimension. Indeed, $\mathrm{SAut}(F_n)$ cannot act non-trivially on any generalized \mathbb{Z}_2 -homology sphere of dimension less than $n-1$, nor on any \mathbb{Z}_2 -acyclic \mathbb{Z}_2 -homology manifold of dimension less than n . It follows that $\mathrm{SL}(n, \mathbb{Z})$ cannot act non-trivially on such spaces either. When n is even, we obtain similar results with \mathbb{Z}_3 coefficients.

Mathematics Subject Classification (2010). 20F65, 20F28, 53C24, 57S25.

Keywords. Automorphism groups of free groups, rigidity, group actions, Smith theory, generalized manifolds.

1. Introduction

In geometric group theory one attempts to elucidate the algebraic properties of a group by studying its actions on spaces with good geometric properties. For irreducible lattices in higher-rank semisimple Lie groups, versions of Margulis superrigidity place severe restrictions on the spaces that are useful for this purpose. Our focus in this article is on the rigidity properties of the group $\mathrm{Aut}(F_n)$ of automorphisms of a free group, which is not a lattice but nevertheless enjoys many similar properties.

In [5] we exhibited strong constraints on homomorphisms from $\mathrm{Aut}(F_n)$ and pointed out that such constraints restrict the way in which $\mathrm{Aut}(F_n)$ can act on various spaces. We illustrated this point by showing that if $n \geq 3$ then any action of $\mathrm{Aut}(F_n)$ on the circle by homeomorphisms must factor through the determinant homomorphism $\det: \mathrm{Aut}(F_n) \rightarrow \mathbb{Z}_2$. We now show that similar restrictions apply much more

*Bridson's work was supported by an EPSRC Senior Fellowship and a Royal Society Nuffield Research Merit Award. Vogtmann is supported by NSF grant DMS-0204185. Bridson thanks the University of Geneva and the EPFL (Lausanne) for their hospitality during the writing of this paper, and the Swiss National Research Foundation for funding his stay.

generally, to actions on higher-dimensional generalized homology spheres over \mathbb{Z}_p and to generalized manifolds that are \mathbb{Z}_p -acyclic, for $p = 2, 3$.

For $n \geq 3$ we denote by $\text{SAut}(F_n)$ the unique subgroup of index two in $\text{Aut}(F_n)$. The action of $\text{Aut}(F_n)$ on the abelianization of the free group F_n gives a natural map $\text{Aut}(F_n) \rightarrow \text{GL}(n, \mathbb{Z})$, sending $\text{SAut}(F_n)$ onto $\text{SL}(n, \mathbb{Z})$. Thus the standard linear action of $\text{SL}(n, \mathbb{Z})$ on \mathbb{R}^n induces non-trivial actions of $\text{SAut}(F_n)$ on \mathbb{R}^n and on the sphere \mathbb{S}^{n-1} . However, we will prove that $\text{SAut}(F_n)$ cannot act non-trivially on spheres or contractible manifolds of any smaller dimension. For linear actions, elementary results in the representation theory of finite groups can be combined with an understanding of the torsion in $\text{SAut}(F_n)$ to prove this statement; the real challenge lies with non-linear actions.

Smooth actions are considerably easier to handle than topological ones. Thus we begin by proving, in Section 2, that for $n \geq 3$, $\text{SAut}(F_n)$ cannot act non-trivially by diffeomorphisms on a \mathbb{Z}_2 -acyclic smooth manifold of dimension less than n . The proof we present is deliberately constructed so as to point out the difficulties encountered in the purely topological setting. In particular, the proof requires understanding the fixed point sets of involutions. This immediately creates a problem in the topological setting because the fixed point sets of involutions are not in general manifolds, but only homology manifolds over \mathbb{Z}_2 . A second difficulty arises because there is no tangent space in the topological setting; in the smooth case the tangent space allows one to use linear algebra to transport information about the action near fixed point sets to information about the action on the ambient manifold.

These are well-known difficulties that lie at the heart of the theory of transformation groups and much effort has gone into confronting them [4], [3]. They are overcome using (local and global) *Smith theory*, but one has to accept the necessity of working with generalized manifolds rather than classical manifolds. (See Section 4 for definitions concerning generalized manifolds.)

We shall prove the following results by following the architecture of the proof we give in the smooth setting, combining Smith theory with an analysis of the torsion in $\text{SAut}(F_n)$ to overcome the technical problems that arise.

Theorem 1.1. *If $n \geq 3$ and $d < n - 1$, then any action of $\text{SAut}(F_n)$ by homeomorphisms on a generalized d -sphere over \mathbb{Z}_2 is trivial, and hence $\text{Aut}(F_n)$ can act only via the determinant map.*

Theorem 1.2. *If $n \geq 3$ and $d < n$, then any action of $\text{SAut}(F_n)$ by homeomorphisms on a d -dimensional \mathbb{Z}_2 -acyclic homology manifold over \mathbb{Z}_2 is trivial, and hence $\text{Aut}(F_n)$ can act only via the determinant map.*

As special cases we obtain the desired minimality result for the standard linear action of $\text{SAut}(F_n)$ on \mathbb{R}^n and \mathbb{S}^{n-1} .

Corollary 1.3. *If $n \geq 3$, then $\text{SAut}(F_n)$ cannot act non-trivially by homeomorphisms on any contractible manifold of dimension less than n , or on any sphere of dimension less than $n - 1$.*

We also note that these theorems have as immediate corollaries the analogous statements for $\text{SL}(n, \mathbb{Z})$ and $\text{GL}(n, \mathbb{Z})$.

Corollary 1.4. *If $n \geq 3$ and $d < n$, then $\text{SL}(n, \mathbb{Z})$ cannot act non-trivially by homeomorphisms on any generalized $(d - 1)$ -sphere over \mathbb{Z}_2 , or on any d -dimensional homology manifold over \mathbb{Z}_2 that is \mathbb{Z}_2 -acyclic. Hence $\text{GL}(n, \mathbb{Z})$ can act on such spaces only via the determinant map.*

Corollary 1.4 was conjectured by Parwani [13]; see Remark 4.16.

In Section 3 we describe a subgroup $T \subset \text{SAut}(F_{2m})$ isomorphic to $(\mathbb{Z}_3)^m$ that intersects every proper normal subgroup of $\text{SAut}(F_{2m})$ trivially. This provides a stronger degree of rigidity than is offered by the 2-torsion in $\text{SAut}(F_n)$ and consequently one can deduce the following theorems from Smith theory more readily than is possible in the case of \mathbb{Z}_2 (see Section 4.3).

Theorem 1.5. *If $n > 3$ is even and $d < n - 1$, then any action of $\text{SAut}(F_n)$ by homeomorphisms on a generalized d -sphere over \mathbb{Z}_3 is trivial.*

Theorem 1.6. *If $n > 3$ is even and $d < n$, then any action of $\text{SAut}(F_n)$ by homeomorphisms on a d -dimensional \mathbb{Z}_3 -acyclic homology manifold over \mathbb{Z}_3 is trivial.*

We expect that our results concerning $\text{SL}(n, \mathbb{Z})$ should be true for other lattices in $\text{SL}(n, \mathbb{R})$, but our techniques do not apply because we make essential use of the torsion in $\text{SL}(n, \mathbb{Z})$. What happens for subgroups of finite index in $\text{SAut}(F_n)$ is less clear: there are subgroups of finite index in $\text{SAut}(F_n)$ that map non-trivially to $\text{SL}(n - 1, \mathbb{R})$ and hence act non-trivially on \mathbb{R}^{n-1} , but one does not know if such subgroups can act non-trivially on contractible manifolds of dimension less than $n - 1$.

In a brief final section we explain how our results concerning torsion in $\text{Aut}(F_n)$, together with the application of Smith theory in [12], imply the following result.

Theorem 1.7. *Let p be a prime and let M be a compact d -dimensional homology manifold over \mathbb{Z}_p . There exists an integer $\eta(p, d, B)$, depending only on p , d and the sum B of the mod p Betti numbers of M , so that $\text{Aut}(F_n)$ cannot act non-trivially by homeomorphisms on M if $n > \eta(p, d, B)$.*

Acknowledgements. We would like to thank the colleagues who helped us struggle with the technicalities of generalized manifolds and Smith theory over the past year, including in particular Mladen Bestvina, Mike Davis, Ian Hambleton and Shmuel Weinberger. We also thank Linus Kramer and Olga Varghese for their comments concerning Theorem 4.5.

2. Smooth actions

In this section we indicate how to prove Theorem 1.2 for smooth actions. Our intent here is to explain the structure of the proof of our general results without the technical difficulties that occur in the topological setting.

Theorem 2.1. *Let X be a k -dimensional differentiable manifold that is \mathbb{Z}_2 -acyclic (i.e., has the \mathbb{Z}_2 -homology of a point). If $n \geq 3$ and $k < n$ then any action of $\mathrm{SAut}(F_n)$ by diffeomorphisms on X is trivial.*

Proof. The proof proceeds by induction on n . We omit the cases $n \leq 4$, where ad hoc arguments apply (cf. subsection 4.5). Suppose, then, that $n \geq 5$, fix a basis a_1, \dots, a_n for F_n and consider the involutions ε_{ij} of F_n defined as follows:

$$\varepsilon_{ij} : \begin{cases} a_i \mapsto a_i^{-1}, \\ a_j \mapsto a_j^{-1}, \\ a_k \mapsto a_k, \quad k \neq i, j. \end{cases}$$

These involutions are all conjugate in $\mathrm{SAut}(F_n)$, and the quotient of $\mathrm{SAut}(F_n)$ by the normal closure of any ε_{ij} is $\mathrm{SL}(n, \mathbb{Z}_2)$, which is a simple group (cf. Proposition 3.1). Thus to prove that an action of $\mathrm{SAut}(F_n)$ is trivial it suffices to show first that some ε_{ij} acts trivially, so that the action factors through $\mathrm{SL}(n, \mathbb{Z}_2)$, and then that some non-trivial element of $\mathrm{SL}(n, \mathbb{Z}_2)$ acts trivially.

Since X is \mathbb{Z}_2 -acyclic it must be orientable, and since $\mathrm{SAut}(F_n)$ is perfect it must act by orientation-preserving diffeomorphisms. Therefore either the action of ε_{12} is trivial or the fixed point set \mathcal{F}_{12} of ε_{12} is a smooth submanifold of codimension at least 2, and Smith theory [22] tells us that this fixed point set will itself be \mathbb{Z}_2 -acyclic.

The centralizer of ε_{12} contains an obvious copy of $\mathrm{SAut}(F_{n-2})$, corresponding to the sub-basis a_3, \dots, a_n , and by induction this must act trivially on \mathcal{F}_{12} . In particular, the automorphism ε_{45} acts trivially, so its fixed point set \mathcal{F}_{45} contains \mathcal{F}_{12} . But ε_{12} and ε_{45} are conjugate, so in fact $\mathcal{F}_{12} = \mathcal{F}_{45}$, i.e., we have two commuting involutions with the same (non-empty) fixed point set. On the tangent space at a common fixed point these induce commuting linear involutions of \mathbb{R}^k with the same fixed vectors, which must be identical by basic linear algebra. But the action of a finite group on a connected smooth manifold is determined by its action on the tangent space of a fixed point, so the actions themselves must be identical. Thus the product $\varepsilon_{12}\varepsilon_{45}$ acts trivially. A similar argument shows that $\varepsilon_{23}\varepsilon_{45}$ acts trivially, and we conclude that the product $\varepsilon_{12}\varepsilon_{45}\varepsilon_{23}\varepsilon_{45} = \varepsilon_{13}$ acts trivially.

Now look at the induced action of $\mathrm{SL}(n, \mathbb{Z}_2)$ on X , and consider the elementary matrices E_{1j} . These generate a subgroup isomorphic to \mathbb{Z}_2^{n-1} , and we claim that any such group acting by orientation-preserving homeomorphisms on X must contain an element which acts trivially. To see this, choose an element of \mathbb{Z}_2^{n-1} whose fixed point

set \mathcal{F} has the largest dimension. By induction (starting with the trivial case $n = 3$), some other element of the group must act trivially on \mathcal{F} , and one thus obtains two commuting involutions that have the same fixed point set, as in the previous paragraph. As before, the involutions must be the same and the product acts trivially. \square

3. Concerning the quotients of $\text{Aut}(F_n)$

3.1. Notation. Fix a generating set $\{a_1, \dots, a_n\}$ for F_n . The right and left Nielsen automorphisms ρ_{ij} and λ_{ij} are defined by

$$\rho_{ij} : \begin{cases} a_i \mapsto a_i a_j, \\ a_k \mapsto a_k, \quad k \neq i; \end{cases} \quad \lambda_{ij} : \begin{cases} a_i \mapsto a_j a_i, \\ a_k \mapsto a_k, \quad k \neq i. \end{cases}$$

We denote by e_i the automorphism which inverts the generator a_i . Elements of the subgroup Σ_n of automorphisms which permute the generators a_i will be denoted using standard cycle notation; for example (ij) is the automorphism interchanging a_i and a_j :

$$e_i : \begin{cases} a_i \mapsto a_i^{-1}, \\ a_k \mapsto a_k, \quad k \neq i; \end{cases} \quad (ij) : \begin{cases} a_i \mapsto a_j, \\ a_j \mapsto a_i, \\ a_k \mapsto a_k, \quad k \neq i, j. \end{cases}$$

W_n is the subgroup of $\text{Aut}(F_n)$ generated by Σ_n and the inversions e_i , and SW_n is the intersection of W_n with $\text{SAut}(F_n)$. The subgroup of W_n generated by the e_i is a normal subgroup $N \cong (\mathbb{Z}_2)^n$, and W_n decomposes as the semidirect product $N \rtimes \Sigma_n$. The intersection of N with $\text{SAut}(F_n)$ is denoted SN . Note that the central element $\Delta = e_1 e_2 \dots e_n$ of W_n is in SN if and only if n is even.

Although it seems awkward at first glance, it is convenient to work with the right action of $\text{Aut}(F_n)$ on F_n : so $\alpha\beta$ acts as α followed by β . An advantage of this is the neatness of the formula $[\lambda_{ij}, \lambda_{jk}] = \lambda_{ik}$, where our commutator convention is $[a, b] = aba^{-1}b^{-1}$.

3.2. How kernels can intersect SW_n . The following variation on Proposition 9 of [5] will be useful here.

Proposition 3.1. *Suppose $n \geq 3$ and let ϕ be a homomorphism from $\text{SAut}(F_n)$ to a group G . If $\phi|_{SW_n}$ has non-trivial kernel K , then one of the following holds:*

1. n is even, $K = \langle \Delta \rangle$ and ϕ factors through $\text{PSL}(n, \mathbb{Z})$,
2. $K = SN$ and the image of ϕ is isomorphic to $\text{SL}(n, \mathbb{Z}_2)$, or
3. ϕ is the trivial map.

Proof. In $\text{Aut}(F_n)$ one has the semidirect product decomposition $W_n = N \rtimes \Sigma_n$ and accordingly we write elements of SW_n as $\alpha\sigma$, with $\alpha = e_1^{\epsilon_1} e_2^{\epsilon_2} \dots e_n^{\epsilon_n} \in N$ and $\sigma \in \Sigma_n$. (Note that it may be that neither α nor σ is itself in $\text{SAut}(F_n)$.)

Using exponential notation to denote conjugation, we have

$$\lambda_{ij}^\alpha = \begin{cases} \lambda_{ij} & \text{if } \epsilon_i = \epsilon_j = 0, \\ \rho_{ij} & \text{if } \epsilon_i = \epsilon_j = 1, \\ \lambda_{ij}^{-1} & \text{if } \epsilon_i = 0 \text{ and } \epsilon_j = 1, \\ \rho_{ij}^{-1} & \text{if } \epsilon_i = 1 \text{ and } \epsilon_j = 0. \end{cases} \quad (1)$$

Also, for $\theta \in \{\lambda, \rho\}$, we have $\theta_{ij}^\sigma = \theta_{\sigma(i)\sigma(j)}$. Hence $\lambda_{ij}^{\alpha\sigma} = \theta_{\sigma(i)\sigma(j)}^{\pm 1}$ for some $\theta \in \{\lambda, \rho\}$.

If K contains the center $\langle \Delta \rangle$ of W_n then n must be even and the relations $\Delta \lambda_{ij} \Delta = \rho_{ij}$ imply that the map ϕ factors through $\text{SL}(n, \mathbb{Z})$, since by [9] adding the relations $\lambda_{ij} = \rho_{ij}$ to a presentation for $\text{SAut}(F_n)$ gives a presentation for $\text{SL}(n, \mathbb{Z})$. Since Δ maps to the center of $\text{SL}(n, \mathbb{Z})$, the map in fact factors through $\text{PSL}(n, \mathbb{Z})$.

If K contains an element $\alpha \in SN$ which is not central in W_n , then we can write $\alpha = e_1^{\epsilon_1} e_2^{\epsilon_2} \dots e_n^{\epsilon_n}$ with $\sum \epsilon_i$ even and some $\epsilon_k = 0$. Given any indices i and j we can conjugate α by an element of the alternating group $A_n \leq SW_n$ to obtain elements in the kernel of ϕ with any desired values of $\epsilon_i, \epsilon_j \in \{0, 1\}$. Conjugating λ_{ij} by these elements, we see from (1) that $\lambda_{ij}, \rho_{ij}, \lambda_{ij}^{-1}$ and ρ_{ij}^{-1} all have the same image under ϕ . This implies not only that ϕ factors through $\text{SL}(n, \mathbb{Z})$, but also that the images of all Nielsen automorphisms have order 2, and so ϕ factors through $\text{SL}(n, \mathbb{Z}_2)$. The image of SN is trivial under this map, i.e. $K \supseteq SN$. Since $\text{SL}(n, \mathbb{Z}_2)$ is simple, the image of ϕ is either trivial or isomorphic to $\text{SL}(n, \mathbb{Z}_2)$.

Finally, suppose that K contains an element $\alpha\sigma$ which is not in SN , i.e. $\sigma \neq 1$. If σ is not an involution, then for some i, j, k with $i \neq k$ we have $\sigma(i) = j$ and $\sigma(j) = k$, hence $\lambda_{ij}^\sigma = \theta_{jk}$ with $\theta \in \{\lambda, \rho\}$. By combining the relations $[\lambda_{ij}, \lambda_{jk}] = \lambda_{ik}$ and $[\theta_{ij}^{\pm 1}, \lambda_{ij}] = 1$ with the fact that $\phi(x^y) = \phi(x)$ for all $x \in \text{SAut}(F_n)$ and $y \in K$, we deduce:

$$\begin{aligned} \phi(\lambda_{ik}) &= [\phi(\lambda_{ij}), \phi(\lambda_{jk})] = [\phi(\lambda_{ij}^{\alpha\sigma}), \phi(\lambda_{jk})] \\ &= [\phi(\theta_{jk}^{\pm 1}), \phi(\lambda_{jk})] = \phi([\theta_{jk}^{\pm 1}, \lambda_{jk}]) = 1. \end{aligned}$$

Since all Nielsen automorphisms are conjugate in $\text{SAut}(F_n)$ and they together generate $\text{SAut}(F_n)$, we conclude that ϕ is trivial (and $K = SW_n$).

Finally, if σ is an involution interchanging $j \neq k$, then a similar calculation produces the conclusion that

$$\phi(\lambda_{ik}) = [\phi(\lambda_{ij}), \phi(\lambda_{jk}^{\alpha\sigma})] = \phi([\lambda_{ij}, \theta_{jk}^{\pm 1}]) = 1$$

so that ϕ is again trivial. \square

3.3. All non-trivial quotients of $\text{SAut}(F_{2m})$ contain $(\mathbb{Z}_3)^m$. In this subsection we are only interested in free groups of even rank. It is convenient to switch notation: if $n = 2m$ we fix a basis $\{a_1, b_1, \dots, a_m, b_m\}$ for F_n ; we write $\lambda_{a_i b_i}$ and $\rho_{a_i b_i}$ for the Nielsen transformations that send a_i to $b_i a_i$ and $a_i b_i$, respectively; we write $(a_i \ b_i)$ for the automorphism that interchanges a_i and b_i , fixing the other basis elements; we write e_{a_1} instead of e_1 , and so on.

Let T be the subgroup of $\text{SAut}(F_n)$ generated by $\{R_i \mid i = 1, \dots, m\}$ where

$$R_i : \begin{cases} a_i \mapsto b_i^{-1}, \\ b_i \mapsto b_i^{-1} a_i, \\ a_j \mapsto a_j, & j \neq i, \\ b_j \mapsto b_j, & j \neq i. \end{cases}$$

Lemma 3.2. $T \cong (\mathbb{Z}_3)^m$.

Proof. One can verify this by direct calculation but the nature of T is most naturally described in terms of the labelled graph \mathcal{T}_m depicted in Figure 1.




Figure 1. Graph realizing the subgroup T .

\mathcal{T}_m has $m + 1$ vertices v_0, v_1, \dots, v_m and 3 edges joining v_0 to each of the other vertices. A maximal tree is obtained by choosing an (unlabelled) edge joining v_0 to each of the other vertices. For each i , the remaining two edges incident at v_i are oriented towards v_0 and labelled a_i and b_i .

This labelling identifies $\pi_1(\mathcal{T}_m, v_0)$ with $F_{2m} = F(a_1, b_1, \dots, a_m, b_m)$ and defines an injective homomorphism $\psi : \text{Sym}(\mathcal{T}_m, v_0) \rightarrow \text{Aut}(F_n)$ whose image contains T . Indeed R_i is the image under ψ of the symmetry of order 3 that cyclically permutes the edges joining v_i to v_0 , sending the edge labelled a_i to that labelled b_i and sending the edge labelled b_i to the unlabelled edge. \square

A routine calculation yields:

Lemma 3.3. For $i = 1, \dots, m$, let $\beta_i \in \text{SAut}(F_n)$ be the automorphism that sends a_i to a_i^{-1} and b_i to $a_i^{-1} b_i^{-1} a_i$ while fixing the other basis elements.

- (1) $R_i e_{a_i} e_{b_i} R_i^{-1} = \beta_i$.
- (2) $[R_j, e_{a_i}] = [R_j, e_{b_i}] = 1$ if $j \neq i$.
- (3) $R_i e_{b_i} R_i^{-1} e_{a_i} = \lambda_{b_i a_i}^2$.
- (4) $R_i^{-1} e_{a_i} R_i e_{b_i} = \lambda_{a_i b_i}^2$.

Proposition 3.4. *For $m \geq 2$ and any group G , let $\phi: \mathrm{SAut}(F_{2m}) \rightarrow G$ be a homomorphism. If $\phi|_T$ is not injective, then ϕ is trivial.*

Proof. Let $t \in T$ be a non-trivial element of the kernel of ϕ . Replacing t by t^{-1} if necessary, we may write $t = R_i u$ where u is a word in the R_j with $j \neq i$. Since each R_j commutes with e_{a_i} and e_{b_i} , we have $t e_{a_i} e_{b_i} t^{-1} = R_i e_{a_i} e_{b_i} R_i^{-1} = \beta_i$. Since $\phi(t) = 1$, applying ϕ to this equation gives $\phi(e_{a_i} e_{b_i}) = \phi(\beta_i)$.

We now note that $e_{a_i} e_{b_i}$ conjugates $\lambda_{a_i b_i}$ to $\rho_{a_i b_i}$, whereas β_i commutes with $\lambda_{a_i b_i}$. Since the images of $e_{a_i} e_{b_i}$ and β_i under ϕ are the same, this gives

$$\phi(\rho_{b_i a_i}) = \phi(\lambda_{b_i a_i}).$$

As in the proof of Proposition 3.1, we appeal to [9] to deduce that ϕ factors through $\mathrm{SAut}(F_n) \rightarrow \mathrm{SL}(n, \mathbb{Z})$.

Next we consider the effect of the relations (3) and (4) from Lemma 3.3. Unfortunately, these are relations in $\mathrm{Aut}(F_n)$ not $\mathrm{SAut}(F_n)$. But since R_i commutes with e_{a_j} when $j \neq i$ we have the following relation in $\mathrm{SAut}(F_n)$,

$$R_i^{-1} e_{a_i} e_{a_j} R_i e_{b_i} e_{a_j} = \lambda_{a_i b_i}^2.$$

If $t = R_i$ then applying ϕ to this equation gives $\phi(e_{a_i} e_{b_i}) = \phi(\lambda_{a_i b_i})^2$. Conjugating both sides by the permutation $(a_i \ a_j)(b_i \ b_j)$, we get the same equality with j subscripts. Since all the automorphisms with i subscripts commute with those that have j subscripts, we deduce

$$(*) \quad \phi(e_{b_i} e_{b_j} e_{a_i} e_{a_j}) = \phi(\lambda_{a_i b_i}^2 \lambda_{a_j b_j}^2).$$

If $t = R_i R_j v$ for some $j \neq i$ and v a (possibly empty) word in the R_k with $k \neq i, j$, then combining relation (4) for i and j gives

$$t^{-1} e_{a_i} e_{a_j} t e_{b_i} e_{b_j} = R_i^{-1} R_j^{-1} e_{a_i} e_{a_j} R_j R_i e_{b_i} e_{b_j} = \lambda_{a_i b_i}^2 \lambda_{a_j b_j}^2.$$

Applying ϕ to this equation gives equation $(*)$ in this case as well.

If $t = R_i R_j^{-1} v$, then relation (3) for i and relation (4) for j give

$$t e_{b_i} e_{a_j} t^{-1} e_{a_i} e_{b_j} = R_i R_j^{-1} e_{b_i} e_{a_j} R_i^{-1} R_j e_{a_i} e_{b_j} = \lambda_{a_i b_i}^2 \lambda_{b_j a_j}^2.$$

Applying ϕ to this equation gives $\phi(e_{a_i} e_{b_j} e_{b_i} e_{a_j}) = \phi(\bar{\lambda}_{a_i b_i}^2 \bar{\lambda}_{b_j a_j}^2)$. Conjugating both sides by $(a_j b_j) e_{a_k}$ for some $k \neq i, j$ gives equation $(*)$ once again.

Next we claim that equation $(*)$ forces ϕ to factor not only through $\text{SAut}(F_n) \rightarrow \text{SL}(n, \mathbb{Z})$ but also through $\text{SAut}(F_n) \rightarrow \text{SL}(n, \mathbb{Z}_2)$. In order to prove this, it suffices to argue that the image under ϕ of some Nielsen transformation has order at most 2.

Let $\bar{\alpha}$ denote the image of $\alpha \in \text{SAut}(F_n)$ in $\text{SL}(n, \mathbb{Z})$. Consider the subgroup $\text{SL}(4, \mathbb{Z}) \subset \text{SL}(n, \mathbb{Z})$ corresponding to the sub-basis $\{a_i, b_i, a_j, b_j\}$. Equation $(*)$ tells us that $\bar{\lambda}_{a_i b_i}^2 \bar{\lambda}_{a_j b_j}^2$ becomes central in the image of $\text{SL}(4, \mathbb{Z})$ under ϕ . But in this copy of $\text{SL}(4, \mathbb{Z})$ one has the relations $[\bar{\lambda}_{a_i b_i}^2, \bar{\lambda}_{b_i b_j}] = \bar{\lambda}_{a_i b_j}^2$ and $[\bar{\lambda}_{a_j b_j}^2, \bar{\lambda}_{b_i b_j}] = 1$. So forcing $\bar{\lambda}_{a_i b_i}^2 \bar{\lambda}_{a_j b_j}^2$ to become central implies that $\phi(\bar{\lambda}_{a_i b_j})^2 = 1$, as required.

We have proved that ϕ factors through $\text{SAut}(F_n) \rightarrow \text{SL}(n, \mathbb{Z}_2)$. The final point to observe is that the restriction to T of this last map is injective; in particular the image of t is non-trivial, and hence so is the image of $\ker \phi$. Thus the image of ϕ in G is a proper quotient of the *simple* group $\text{SL}(n, \mathbb{Z}_2)$, and therefore is trivial. \square

4. Actions on generalized spheres and acyclic homology manifolds

Because the fixed point set of a finite-period homeomorphism of a sphere or contractible manifold need not be a manifold, we must expand the category we are working in to that of generalized manifolds. We follow the exposition in Bredon's book on sheaf theory [3]. All homology groups in this section are Borel–Moore homology with compact supports and coefficients in a sheaf \mathcal{A} of modules over a principle ideal domain L . The homology groups of X are denoted $H_*^c(X; \mathcal{A})$. If X is a locally finite CW-complex and \mathcal{A} is the constant sheaf $X \times L$ (which we will denote simply by L), then $H_*^c(X; L)$ is isomorphic to singular homology with coefficients in L (see [3], p. 279).

All cohomology groups are sheaf cohomology with compact supports, denoted $H_c^*(X; \mathcal{A})$. If \mathcal{A} is the constant sheaf, this is isomorphic to Čech cohomology with compact supports. If F is a closed subset of X , then sheaf cohomology satisfies $H_c^k(X, F; \mathcal{A}) \cong H_c^k(X \setminus F; \mathcal{A})$.

In fact, the only sheaves we will consider other than the constant sheaf are the sheaves \mathcal{O}_k associated to the pre-sheaves $U \mapsto H_k^c(X, X \setminus U; L)$.

4.1. Homology manifolds. Let L be one of \mathbb{Z} or \mathbb{Z}_p (the integers mod p , where p is a prime).

Definition 4.1 ([3], p. 329). An m -dimensional homology manifold over L (denoted $m\text{-hm}_L$) is a locally compact Hausdorff space X with finite homological dimension over L , that has the local homology properties of a manifold. Specifically, the sheaves

\mathcal{O}_k are locally constant with stalk 0 if $k \neq m$ and L if $k = m$. The sheaf $\mathcal{O} = \mathcal{O}_m$ is called the *orientation sheaf*.

We will further assume that our homology manifolds are first-countable.

Definition 4.2. If X is an m -hm $_L$ and $H_*^c(X; L) \cong H_*^c(\mathbb{S}^m; L)$ then X is called a *generalized m -sphere* over L .

Definition 4.3. If X is an m -hm $_L$ with $H_0^c(X; L) = L$ and $H_k^c(X; L) = 0$ for $k > 0$, then X is said to be *L -acyclic*.

There is a similar notion of *cohomology manifold over L* , denoted m -cm $_L$ (see [3], p. 373). If $L = \mathbb{Z}_p$, a connected space X is an n -cm $_L$ if and only if it is an n -hm $_L$ and is locally connected ([3], p. 375 Theorem 16.8 and footnote). If X is a locally connected homology manifold over \mathbb{Z}_p , then the fixed point set of any homeomorphism of order p is also locally connected (see [4], Theorem 1.6, p. 72, where there is a stronger connectivity statement (clc_L), but the proof, which relies on Proposition 1.4, p. 68, also applies to local connectivity). These remarks show that the theorems we state below for homology manifolds are also valid for cohomology manifolds.

Finally, we note that homology manifolds satisfy Poincaré duality between Borel–Moore homology and sheaf cohomology ([3], Theorem 9.2), i.e., if X is an m -hm $_L$ then

$$H_k^c(X; L) \cong H_{m-k}^c(X; \mathcal{O}).$$

4.2. Elements of Smith theory. There are two types of Smith theorems, usually referred to as “global” and “local” Smith theorems. The global theorems require only that X be a locally compact Hausdorff space with the homology of a sphere or a point, while the local theorems concern homology manifolds. These were originally proved by P. A. Smith ([14], [15]), but we follow the exposition in Bredon’s book and Borel’s Seminar on Transformation groups [4].

Theorem 4.4 (The Local Smith Theorem, [3], Theorem 20.1, Proposition 20.2, pp. 409–410). *Let p be a prime and $L = \mathbb{Z}_p$. The fixed point set of any action of \mathbb{Z}_p on an n -hm $_L$ is the disjoint union of (open and closed) components each of which is an r -hm $_L$ with $r \leq m$. If p is odd then each component of the fixed point set has even codimension.*

By invariance of domain for homology manifolds ([3], Corollary 16.19, p. 383) the fixed point set of any *non-trivial* action of \mathbb{Z}_p on a connected, locally connected m -hm $_{\mathbb{Z}_p}$ is a (locally connected) r -hm $_{\mathbb{Z}_p}$ with $r \leq m - 1$.

Theorem 4.5 (Global Smith Theorems, [3], Corollaries 19.8 and 19.9, p. 144). *Let p be a prime and X a locally compact Hausdorff space of finite dimension over \mathbb{Z}_p . Suppose that \mathbb{Z}_p acts on X with fixed point set F .*

- *If $H_*^c(X; \mathbb{Z}_p) \cong H_*^c(\mathbb{S}^m; \mathbb{Z}_p)$, then $H_*^c(F; \mathbb{Z}_p) \cong H_*^c(\mathbb{S}^r; \mathbb{Z}_p)$ for some r with $-1 \leq r \leq m$. If p is odd, then $m - r$ is even.*
- *If X is \mathbb{Z}_p -acyclic, then F is \mathbb{Z}_p -acyclic (in particular non-empty and connected).*

In Section 19 of [3] the Global Smith Theorem is stated for cohomology; the homology version above follows using the Smith theory sequence (132) on page 408 of [3]. The details of this translation have been worked out by Olga Varghese in [21].

Together these theorems imply

Corollary 4.6. *Let X be an m -hm $_{\mathbb{Z}_p}$.*

- *If X is a generalized m -sphere over \mathbb{Z}_p , the fixed point set of any homeomorphism of order p is a (possibly empty) generalized r -sphere, with $r \leq m - 1$. If p is odd, $r \leq m - 2$.*
- *If X is \mathbb{Z}_p -acyclic, the fixed point set of any homeomorphism of order p is a (non-empty) \mathbb{Z}_p -acyclic r -hm $_{\mathbb{Z}_p}$, for some $r \leq m - 1$. If p is odd, $r \leq m - 2$.*

We want to use this corollary as the basis for an induction that bounds the dimensions in which elementary p -groups can act effectively on generalized spheres and acyclic homology manifolds. But in the case of spheres we need an additional result that guarantees the existence of fixed points. This is provided by P. A. Smith's theorem that $\mathbb{Z}_p \times \mathbb{Z}_p$ cannot act freely on a generalized sphere over \mathbb{Z}_p (see [16]; cf Theorem 4.8 below).

The proof of the following theorem is again due to P. A. Smith [16]. (In [16] he only gave the proof for generalized spheres, but the acyclic case is similar.)

Theorem 4.7. *If $m < d - 1$, the group $(\mathbb{Z}_2)^d$ cannot act effectively on a generalized m -sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic $(m + 1)$ -dimensional homology manifold over \mathbb{Z}_2 .*

If $m < 2d - 1$ and p is odd, then $(\mathbb{Z}_p)^d$ cannot act effectively on a generalized m -sphere or a \mathbb{Z}_p -acyclic $(m + 1)$ -dimensional homology manifold over \mathbb{Z}_p .

Proof. The cases that arise when $d = 1$ are vacuous or trivial except when p is odd and the putative action is on a 1-hm $_{\mathbb{Z}_p}$, in which case one needs to recall that a 1-hm $_{\mathbb{Z}_p}$ is an actual manifold.

We assume $d \geq 2$ and proceed by induction. Let X be one of the spaces that the theorem asserts $G := (\mathbb{Z}_p)^d$ cannot act effectively on.

Among the non-trivial elements of G we choose one, a say, whose fixed point set F_a is maximal with respect to inclusion. We also choose a complement $G_0 \cong$

$(\mathbb{Z}_p)^{d-1}$ to $\langle a \rangle$ in G . From Theorem 4.5 (in the acyclic case) and Smith's theorem for $\mathbb{Z}_p \times \mathbb{Z}_p$ (in the case of spheres), we know that F_a is non-empty. We shall prove that $F_a = X$ by assuming it false and obtaining a contradiction.

If F_a is not the whole of X then it has codimension at least 1 if $p = 2$ and codimension at least 2 if p is odd. In the light of Corollary 4.6, we may apply induction to the action of G_0 on F_a and hence conclude that some non-trivial element $b \in G_0$ fixes F_a pointwise; in other words $F_a \subseteq F_b$. But F_a is maximal, so $F_b = F_a$, which implies $F_a = \text{Fix}(A)$ for $A = \langle a, b \rangle$. Thus for any non-trivial element x of A we have $F_a \subseteq F_x$, so again maximality tells us that $F_a = F_x$.

Theorem 4.3 on page 182 of [4] (which requires us to know that $\text{Fix}(A)$ is non-empty) provides a formula relating the dimensions of the fixed point sets of elements of A : writing $n = \dim_p(X)$, $r = \dim_p(\text{Fix}(A))$, and $r_C = \dim_p(\text{Fix}(C))$ for each cyclic subgroup $C < A$, we have

$$n - r = \sum (r_C - r),$$

where the sum is taken over the non-trivial cyclic subgroups of A . We have just argued that $F_a = \text{Fix}(C) = \text{Fix}(A)$ for all non-trivial $C < A$, so each summand on the right is 0 and hence $n = r = \dim_p(F_a)$. Since X is connected, invariance of domain gives $X = F_a$, i.e., a acts trivially. This contradiction completes the induction. \square

We need one more result from Smith theory:

Theorem 4.8. *Let X be a generalized sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic $\text{hm}_{\mathbb{Z}_2}$, and let a and b be commuting homeomorphisms of X , each of order 2, with fixed point sets F_a and F_b . If $F_a = F_b$ then $a = b$.*

Proof. For actions on generalized spheres, this is explicit in [16], so we consider only the acyclic case.

If $a \neq b$ then the subgroup $A \leq \text{Homeo}(X)$ generated by a and b is isomorphic to $\mathbb{Z}_2 \times \mathbb{Z}_2$ and $\text{Fix}(a) = \text{Fix}(b) = \text{Fix}(A)$. Thus in the formula $n - r = \sum (r_C - r)$ displayed in the preceding proof, the only non-zero summand on the right is the one for $\langle ab \rangle$. Hence $n = r_C$, that is, $\dim_p(X) = \dim_p(\text{Fix}(ab))$. Since X is connected, invariance of domain gives $X = \text{Fix}(ab)$, which means that $a = b$. \square

4.3. Actions on generalized spheres and \mathbb{Z}_3 -acyclic homology manifolds over \mathbb{Z}_3 , for n even. The results we have developed to this point easily yield the following theorem, for n even.

Theorem 4.9. *Let X be a generalized m -sphere over \mathbb{Z}_3 or a \mathbb{Z}_3 -acyclic $(m+1)$ -dimensional homology manifold over \mathbb{Z}_3 , and let $\phi: \text{SAut}(F_n) \rightarrow \text{Homeo}(X)$ be an action. If n is even and $m < n-1$, then ϕ is trivial.*

Proof. Write $n = 2d$ and let $T \subset \text{SAut}(F_{2d})$ be as in Lemma 3.2. Since $T \cong (\mathbb{Z}_3)^d$ and $m < n - 1 = 2d - 1$, Theorem 4.7 tells us that T cannot act effectively on X , so $\phi(t) = 1$ for some $t \in T \setminus \{1\}$. We proved in Proposition 3.4 that this forces ϕ to be the trivial map. \square

4.4. Actions on generalized spheres and \mathbb{Z}_2 -acyclic homology manifolds over \mathbb{Z}_2 . The proof of Theorems 1.1 and 1.2 is considerably more involved than that of the preceding result. This is largely due to the fact that Corollary 4.6 yields a weaker conclusion for $p = 2$ than for odd primes. Lemma 4.12 will allow us to circumvent this difficulty. It relies on the separation property of codimension 1 fixed point sets that is established in Lemma 4.11 using Poincaré duality and the following theorem about sheaf cohomology.

Theorem 4.10 (Theorem 16.16, [3]). *If X is a connected m -cm L with orientation sheaf \mathcal{O} , and F is a proper closed subset, then for any non-empty open subset U*

- (1) $H_c^m(U; \mathcal{O})$ is the free L -module on the components of U ;
- (2) $H_c^m(F; L) = 0$.

Lemma 4.11. *Let X be a generalized m -sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic m -hm \mathbb{Z}_2 , and let τ be an involution of X . If $\text{Fix}(\tau)$ has dimension $m - 1$, then $X \setminus \text{Fix}(\tau)$ has two \mathbb{Z}_2 -acyclic components and τ interchanges them.*

Proof. If $m = 1$, then X is a circle or a line, $\text{Fix}(\tau)$ is two points or one, and the theorem is clear, so we may assume $m \geq 2$. Let $F = \text{Fix}(\tau)$, and set $L = \mathbb{Z}_2$. Since F is closed, the long exact sequence in sheaf cohomology for the pair (X, F) reads

$$\begin{aligned} \cdots &\rightarrow H_c^{m-2}(F; L) \rightarrow H_c^{m-1}(X \setminus F; L) \rightarrow H_c^{m-1}(X; L) \rightarrow H_c^{m-1}(F; L) \\ &\rightarrow H_c^m(X \setminus F; L) \rightarrow H_c^m(X; L) \rightarrow H_c^m(F; L) \rightarrow 0 \end{aligned}$$

By (2) above, the last term $H_c^m(F; L)$ is 0. Since $L = \mathbb{Z}_2$, the orientation sheaf \mathcal{O} is actually constant, and by (1), we get $H_c^m(X; L) = H_c^{m-1}(F; L) = L$.

Poincaré duality says $H_c^k(X; L) \cong H_{m-k}^c(X; L)$; in particular, $H_c^{m-1}(X; L) \cong H_1^c(X; L) = 0$ (since $m \geq 2$), and the end of the sequence is

$$0 \rightarrow L \rightarrow H_c^m(X \setminus F; L) \rightarrow L \rightarrow 0$$

Thus $H_c^m(X \setminus F; L) \cong L \oplus L$, and another application of (1) shows that $X \setminus F$ has two components. (This is the argument in [3], Corollary 16.26.)

Suppose X is L -acyclic. Applying Poincaré duality to each remaining term in the long exact sequence (F has dimension $m - 1$) gives

$$\cdots \rightarrow H_k^c(F; L) \rightarrow H_k^c(X \setminus F; L) \rightarrow H_k^c(X; L) \rightarrow \cdots$$

for $k \geq 1$. Since F and X are L -acyclic, this shows that each component of $X \setminus F$ is also L -acyclic.

If X is a generalized m -sphere then F is a generalized $(m-1)$ -sphere, and the above argument shows that most of the homology of $X \setminus F$ vanishes as in the acyclic case. In dimensions m and $m-1$ we have

$$0 \rightarrow H_m^c(X \setminus F; L) \rightarrow H_m^c(X; L) \rightarrow H_{m-1}^c(F; L) \rightarrow H_{m-1}^c(X \setminus F; L) \rightarrow 0$$

which becomes

$$0 \rightarrow H_m^c(X \setminus F; L) \rightarrow L \cong L \rightarrow H_{m-1}^c(X \setminus F; L) \rightarrow 0,$$

so again the homology of $X \setminus F$ vanishes in positive degrees, and each component of $X \setminus F$ is acyclic.

In both situations the complement of F has two \mathbb{Z}_2 -acyclic components. Since the involution acts freely on this complement, it cannot preserve either component, by the Global Smith Theorem. \square

Lemma 4.12. *Let X be a generalized m -sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic $(m+1)$ - $\text{hm}_{\mathbb{Z}_2}$, and let G be a group acting by homeomorphisms on X . Suppose G contains a subgroup $P \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ all of whose non-trivial elements are conjugate in G . If P acts non-trivially, then the fixed point sets of its non-trivial elements have codimension at least 2, and $m \geq 2$.*

Proof. Since the non-trivial elements of P are all conjugate, they must all act non-trivially.

Let a and b be generators of P . If $\text{Fix}(a)$ had codimension 1, then by Lemma 4.11, its complement in X would have two components and the action of a would interchange these. Consider the action of b : since it commutes with a it leaves $\text{Fix}(a)$ invariant, so it either interchanges the components of the complement or leaves them invariant. Reversing the roles of b and ab if necessary, we may assume that it interchanges them and hence that $\text{Fix}(b) \subset \text{Fix}(a)$. Since a and b are conjugate, invariance of domain for homology manifolds implies that $\text{Fix}(a) = \text{Fix}(b)$ and hence, by Theorem 4.8, that the actions of a and b on X are identical. Thus ab acts trivially, contradicting the assumption that the action of P is non-trivial.

Thus the fixed point set of any non-trivial element of P has codimension at least 2. If $m = 1$ and X is a generalized sphere this says $\text{Fix}(a) = \text{Fix}(b) = \emptyset$. If X is 2-dimensional and acyclic, then $\text{Fix}(a)$ and $\text{Fix}(b)$ are 0-dimensional acyclic homology manifolds, i.e. points, so $\text{Fix}(a) \subset \text{Fix}(b)$ implies $\text{Fix}(a) = \text{Fix}(b)$. In either case, Theorem 4.8 again implies that ab acts trivially, contradicting our assumptions. \square

Proposition 4.13. *Let X be a generalized m -sphere or \mathbb{Z}_2 -acyclic $(m+1)$ - $\text{hm}_{\mathbb{Z}_2}$. If $m < n-1$ and $n \geq 3$, then any action of $\text{SL}(n, \mathbb{Z}_2)$ on X is trivial.*

Proof. Since $\text{SL}(n, \mathbb{Z}_2)$ is simple, it is enough to find a subgroup of $\text{SL}(n, \mathbb{Z}_2)$ that cannot act effectively on X .

The elementary matrices E_{j1} , $j \neq 1$, generate an elementary 2-group $Q \cong (\mathbb{Z}_2)^{n-1}$. All elementary matrices are conjugate in $\text{SL}(n, \mathbb{Z}_2)$. Furthermore, $E_{32}E_{21}E_{32} = E_{31}E_{21}$. Thus we are in the situation of Lemma 4.12 with $a = E_{21}$ and $b = E_{31}$. An appeal to that lemma completes the proof in the case $n = 3$.

If $n \geq 4$ then $\text{SL}(n, \mathbb{Z}_2)$ contains a larger elementary 2-group than Q , namely that generated by the elementary matrices E_{ij} with $i \leq n/2$ and $j > n/2$. This has rank at least n , so Theorem 4.7 tells us it cannot act effectively X . \square

Proposition 3.1 and Proposition 4.13 together give:

Corollary 4.14. *Let X be a generalized m -sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic $(m+1)$ -hm $_{\mathbb{Z}_2}$, with $m < n-1$. If a non-central element of W_n is in the kernel of an action of $\text{SAut}(F_n)$ on X , then the action is trivial.*

4.5. Proof of Theorems 1.1 and 1.2. We retain the notation introduced at the beginning of Section 3.

Let X be a generalized m -sphere or a \mathbb{Z}_2 -acyclic $(m+1)$ -dimensional homology manifold over \mathbb{Z}_2 , with $m < n-1$. Let $\Phi: \text{SAut}(F_n) \rightarrow \text{Homeo}(X)$ be an action of $\text{SAut}(F_n)$ on X . In the light of the preceding corollary, we will be done if we can prove that the kernel of Φ contains an element of $SN \cong (\mathbb{Z}_2)^{n-1}$ other than $\Delta = e_1 \dots e_n$.

If $n = 3$, then conjugating $a := e_1e_2$ by $(1\ 3)e_2$ and $(2\ 3)e_1$, respectively, yields $b := e_2e_3$ and $ab = e_1e_3$. Thus we may appeal to Lemma 4.12, to see that the action of SN on X is trivial if $m < 2$.

If $n = 4$, then SN is generated by a , b , and $c := e_2e_4$, which are conjugate in SW_n to each other and to each of the products ab , ac and bc . If the action of SN on X is trivial then we are done. Suppose that this is not the case. We know from Lemma 4.12 that $\text{Fix}(a)$ is a generalized d -sphere over \mathbb{Z}_2 or a \mathbb{Z}_2 -acyclic $(d+1)$ -hm $_{\mathbb{Z}_2}$ with $d < 1$. Since b and c commute with a , the group $\langle b, c \rangle \cong \mathbb{Z}_2^2$ acts on $\text{Fix}(a)$, so by Theorem 4.7 some element acts trivially, say g . Then $\text{Fix}(g) \supset \text{Fix}(a)$. But since a and g are conjugate in SW_n , this implies $\text{Fix}(a) = \text{Fix}(g)$, and then by Theorem 4.8, ag acts trivially on X . If $g = b$ or $g = c$, we have found a non-central element of the kernel of $\Phi|_{SW_n}$, so Φ is trivial by Corollary 4.14. If $g = bc$, the action factors through $\text{PSL}(4, \mathbb{Z})$, which contains a subgroup isomorphic to $(\mathbb{Z}_2)^4$ generated by e_1e_2 , e_2e_3 , $\sigma = (12)(34)$, and $\tau = (13)(24)$. By Theorem 4.7, some nontrivial element of this subgroup must map trivially to $\text{Homeo}(X)$. Pulling back these elements to SW_n , we see that some element of the form $e_i e_j$ or γ or $e_i e_j \gamma$, with $\gamma \in \langle \sigma, \tau \rangle$, is in the kernel. Corollary 4.14 again shows that Φ is trivial.

Now we suppose $n > 4$ and proceed by induction. If e_1e_2 acts trivially then we are done by Corollary 4.14. If not then, appealing to Lemma 4.12 once more, we may

suppose that the fixed point set of e_1e_2 in X is a generalized r -sphere or \mathbb{Z}_2 -acyclic $(r+1)$ -homology manifold over \mathbb{Z}_2 with $r < n-3$; call it Y . The centralizer of e_1e_2 will act on Y . This centralizer contains a copy of $\text{SAut}(F_{n-2})$ corresponding to the sub-basis a_3, \dots, a_n , and by induction this acts trivially on Y . In particular, the fixed point set of e_3e_4 contains that of e_1e_2 . Similarly, the reverse inclusion holds. But then by Theorem 4.8 the actions of e_3e_4 and e_1e_2 on X must be the same. Thus the kernel of any homomorphism $\text{SAut}(F_n) \rightarrow \text{Homeo}(X)$ intersects N in $e_1e_2e_3e_4 \neq \Delta$, and Corollary 4.14 says that the action is therefore trivial. \square

Remark 4.15. For $n = 3$, Theorem 1.1 also follows from the results of [5] because a generalized \mathbb{Z}_2 -sphere of dimension one is just a circle and it is shown in [5] that any action of $\text{SAut}(F_n)$ by homeomorphisms on a circle is trivial for $n \geq 3$.

Remark 4.16. This work was stimulated in part by the proof of Corollary 1.4 suggested by Zimmermann in [22]. His proof relied on earlier work of Parwani [13] which sets forth a good strategy but contains a flaw: it is assumed in [13] that if X is a homology manifold over \mathbb{Z} with the \mathbb{Z}_2 -homology of a sphere, then the fixed point set of any involution of X will again be such a space; this is false (see the following remark). It is also assumed in [13] that such a fixed point set will be an ENR, and this is also false.

Remark 4.17. In [11], L. Jones showed that almost any PL homology manifold over \mathbb{Z}_2 satisfying the Smith conditions can arise as the fixed point set of an involution of a genuine sphere. In particular, the fixed point set of an involution of a sphere need not be a \mathbb{Z} -homology manifold.

There are also involutions of spheres for which the fixed point set is not locally 1-connected, so in particular is not an ENR. Indeed Ancel and Guilbault [1] proved that if $\bar{M} = M \cup \Sigma$ is any Z -set compactification of a contractible n -manifold M , with $n > 4$, then the double of \bar{M} along Σ is homeomorphic to the n -sphere. One can realize Σ as the fixed point set of the involution that interchanges the two copies of M in this double, and Σ need not be locally 1-connected. To obtain a concrete example, we can take M to be the universal cover of one of the aspherical manifolds constructed by Davis [6] and take Σ to be its ideal boundary (cf. [7]).

5. Actions on arbitrary compact homology manifolds

In [12], Mann and Su use Smith theory and a spectral sequence developed by Swan [18] to prove that for every prime p and every compact d -dimensional homology manifold X over \mathbb{Z}_p , the sum of whose mod p Betti numbers is B , there exists an integer $v(d, B)$, depending only on d and B , so that \mathbb{Z}_p^r cannot act effectively by homeomorphisms on X if $r > v(d, B)$. (An explicit bound on v is given.)

If n is sufficiently large then the alternating group A_n will be simple and contain a copy of \mathbb{Z}_p^ν . Hence it will admit no non-trivial action on X . Theorem 1.7 stated in the introduction is an immediate consequence of this result and Proposition 3.1, since $SW_n \subset \text{SAut}(F_n)$ contains a copy of A_n . \square

The preceding argument allows one to bound the constant $\eta(p, d, B)$ in Theorem 1.7 by a multiple (depending on p) of $\nu(d, B)$. In the cases $p = 2$ and $p = 3$ one can sharpen this estimate by appealing directly to Propositions 3.1 and 3.4 instead of using A_n .

Remark 5.1. Various of the Higman–Thompson groups, including Richard Thompson’s *vagabond group* V , are finitely presented, simple, and contain an isomorphic copy of every finite group [10]. Given any class of objects each of which has the property that some finite group cannot act effectively on it, groups such as V cannot act non-trivially on any object in the class. In particular, it follows from the Mann and Su result that V cannot act non-trivially by homeomorphisms on any compact manifold. And Theorem 4.7 above implies that V cannot act non-trivially by homeomorphisms on any finite-dimensional \mathbb{Z}_p -acyclic homology manifold over \mathbb{Z}_p for any prime p (cf. [2] and [8]).

References

- [1] F. D. Ancel and C. R. Guilbault, Z -compactifications of open manifolds. *Topology* **38** (1999), 1265–1280. [Zbl 0951.57011](#) [MR 1690157](#)
- [2] G. Arzhantseva, M. R. Bridson, T. Januszkiewicz, I. J. Leary, A. Minasyan, and J. Świątkowski, Infinite groups with fixed point properties. *Geom. Topol.* **13** (2009), no. 3, 1229–1263. [Zbl 1197.20034](#) [MR 2496045](#)
- [3] G. E. Bredon, *Sheaf theory*. Second edition, Grad. Texts in Math. 170, Springer-Verlag, New York 1997. [Zbl 0874.55001](#) [MR 1481706](#)
- [4] A. Borel, *Seminar on transformation groups*. Ann. of Math. Stud. 46, Princeton University Press, Princeton, N.J., 1960. [Zbl 0091.37202](#) [MR 0116341](#)
- [5] M. R. Bridson and K. Vogtmann, Homomorphisms from automorphism groups of free groups. *Bull. London Math. Soc.* **35** (2003), 785–792. [Zbl 1049.20021](#) [MR 2000025](#)
- [6] M. W. Davis, Groups generated by reflections and aspherical manifolds not covered by Euclidean space. *Ann. of Math.* (2) **117** (1983), 293–325. [Zbl 0531.57041](#) [MR 0690848](#)
- [7] H. Fischer, Boundaries of right-angled Coxeter groups with manifold nerves. *Topology* **42** (2003), 423 – 446. [Zbl 1025.57040](#) [MR 1941443](#)
- [8] D. Fisher and L. Silberman, Groups not acting on manifolds. *Internat. Math. Res. Notices IMRN* **16** (2008), Art. ID rnn060, 11pp. [Zbl 1154.22024](#) [MR 2435750](#)
- [9] S. M. Gersten, A presentation for the special automorphism group of a free group. *J. Pure Appl. Algebra* **33** (1984), 269–279. [Zbl 0542.20021](#) [MR 0761633](#)

- [10] G. Higman, *Finitely presented infinite simple groups*. Notes on Pure Math. 8, Department of Pure Mathematics, Department of Mathematics, I.A.S. Australian National University, Canberra 1974. [Zbl](#) [MR](#)
- [11] L. Jones, The converse to the fixed point theorem of P. A. Smith. I. *Ann. of Math.* (2) **94** (1971), 52–68. [Zbl](#) [0229.55006](#) [MR](#) [0295361](#)
- [12] L. N. Mann and J. C. Su, Actions of elementary p -groups on manifolds. *Trans. Amer. Math. Soc.* **106** (1963), 115–126. [Zbl](#) [0114.39104](#) [MR](#) [0143840](#)
- [13] K. Parwani, Actions of $SL(n, \mathbb{Z})$ on homology spheres. *Geom. Dedicata* **112** (2005), 215–223. [Zbl](#) [1080.57034](#) [MR](#) [2163900](#)
- [14] P. A. Smith, Transformations of finite period. *Ann. of Math.* (2) **39** (1938), no. 1, 127–164. [Zbl](#) [0018.33204](#) [MR](#) [1503393](#)
- [15] P. A. Smith, Transformations of finite period II. *Ann. of Math.* (2) **40** (1939), 690–711. [Zbl](#) [0021.43002](#) [MR](#) [0000177](#)
- [16] P. A. Smith, *Permutable periodic transformations*, Proc. Nat. Acad. Sci. U. S. A. 30, (1944). 105–108. [Zbl](#) [0063.07092](#) [MR](#) [0010278](#)
- [17] P. A. Smith, *New results and old problems in finite transformation groups*, Bull. Amer. Math. Soc., 66 (1960), 401–415. [Zbl](#) [0096.37501](#) [MR](#) [0125581](#)
- [18] R. G. Swan, A new method in fixed point theory. *Comment. Math. Helv.* 34 (1960), 1–16. [Zbl](#) [0144.22602](#) [MR](#) [0115176](#)
- [19] R. J. Thompson, Embeddings into finitely generated simple groups which preserve the word problem. In *Word Problems II*, ed. by S. I. Adian, W. W. Boone and G. Higman, North-Holland, Amsterdam 1980. [Zbl](#) [0431.20030](#) [MR](#) [0579955](#)
- [20] J. H. C. Whitehead, On involutions of spheres. *Ann. of Math.* (2) **66** (1957), 27–29. [Zbl](#) [0080.16101](#) [MR](#) [0088732](#)
- [21] O. Varghese, Wirkungen von $\text{Aut}(F_n)$. Diplomarbeit, Westfälische Wilhelms-Universität Münster, 2010.
- [22] B. P. Zimmermann, $SL(n, \mathbb{Z})$ cannot act on small spheres. *Topology Appl.* **156** (2009), no. 6, 1167–1169. [Zbl](#) [1165.57024](#) [MR](#) [2493377](#)

Received September 22, 2008

Martin R. Bridson, Mathematical Institute, 24-29 St Giles', Oxford OX1 3LB, U.K.

E-mail: bridson@maths.ox.ac.uk

Karen Vogtmann, Department of Mathematics, Cornell University, Ithaca, NY 14853, U.S.A.

E-mail: vogtmann@math.cornell.edu