

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 87 (2012)  
  
**Artikel:** Spherical pairs over close local fields  
**Autor:** Aizenbud, Avraham / Avni, Nir / Gourevitch, Dmitry  
**DOI:** <https://doi.org/10.5169/seals-323267>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 12.12.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Spherical pairs over close local fields

Avraham Aizenbud, Nir Avni and Dmitry Gourevitch

**Abstract.** Extending results of [Kaz86] to the relative case, we relate harmonic analysis over some spherical spaces  $G(F)/H(F)$ , where  $F$  is a field of positive characteristic, to harmonic analysis over the spherical spaces  $G(E)/H(E)$ , where  $E$  is a suitably chosen field of characteristic 0.

We apply our results to show that the pair  $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$  is a strong Gelfand pair for all local fields of arbitrary characteristic, and that the pair  $(\mathrm{GL}_{n+k}(F), \mathrm{GL}_n(F) \times \mathrm{GL}_k(F))$  is a Gelfand pair for local fields of any characteristic different from 2. We also give a criterion for finite generation of the space of  $K$ -invariant compactly supported functions on  $G(E)/H(E)$  as a module over the Hecke algebra.

**Mathematics Subject Classification (2010).** 20G05, 20G25, 14L35.

**Keywords.** Multiplicity, Gelfand pair, reductive group, uniqueness of linear periods.

### Contents

0	Introduction . . . . .	929
1	Preliminaries and notation . . . . .	933
2	Finite generation of Hecke modules . . . . .	934
3	Uniform spherical pairs . . . . .	947
4	Applications . . . . .	955
	References . . . . .	961

### 0. Introduction

Local fields of positive characteristic can be approximated by local fields of characteristic zero. If  $F$  and  $E$  are local fields, we say that they are  $m$ -close if  $\mathcal{O}_F/\mathcal{P}_F^m \cong \mathcal{O}_E/\mathcal{P}_E^m$ , where  $\mathcal{O}_F, \mathcal{O}_E$  are the rings of integers of  $F$  and  $E$ , and  $\mathcal{P}_F, \mathcal{P}_E$  are their maximal ideals. For example,  $F_p((t))$  is  $m$ -close to  $\mathbb{Q}_p(\sqrt[m]{p})$ . More generally, for any local field  $F$  of positive characteristic  $p$  and any  $m$  there exists a (sufficiently ramified) extension of  $\mathbb{Q}_p$  that is  $m$ -close to  $F$ .

Let  $G$  be a reductive group defined over  $\mathbb{Z}$ . For any local field  $F$  and conductor  $\ell \in \mathbb{Z}_{\geq 0}$ , the Hecke algebra  $\mathcal{H}_\ell(G(F))$  is finitely generated and finitely presented. Based on this fact, Kazhdan showed in [Kaz86] that for any  $\ell$  there exists  $m \geq \ell$  such that the algebras  $\mathcal{H}_\ell(G(F))$  and  $\mathcal{H}_\ell(G(E))$  are isomorphic for any  $m$ -close fields  $F$  and  $E$ . This allows one to transfer certain results in representation theory of reductive groups from local fields of zero characteristic to local fields of positive characteristic.

In this paper we investigate a relative version of this technique. Let  $G$  be a reductive group and  $H$  be a spherical subgroup. Suppose for simplicity that both are defined over  $\mathbb{Z}$ .

In the first part of the paper we consider the space  $\mathcal{S}(G(F)/H(F))^K$  of compactly supported functions on  $G(F)/H(F)$  which are invariant with respect to a compact open subgroup  $K$ . We prove under certain assumption on the pair  $(G, H)$  that this space is finitely generated (and hence finitely presented) over the Hecke algebra  $\mathcal{H}_K(G(F))$ .

**Theorem A** (see Theorem 2.19). *Let  $F$  be a (non-Archimedean) local field. Let  $G$  be a reductive group and  $H < G$  be an algebraic subgroup both defined over  $F$ . Suppose that for any parabolic subgroup  $P \subset G$ , there is a finite number of double cosets  $P(F) \backslash G(F)/H(F)$ . Suppose also that for any irreducible smooth representation  $\rho$  of  $G(F)$  we have*

$$\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) < \infty. \quad (1)$$

*Then for any compact open subgroup  $K < G(F)$ , the space  $\mathcal{S}(G(F)/H(F))^K$  is a finitely generated module over the Hecke algebra  $\mathcal{H}_K(G(F))$ .*

Assumption (1) is rather weak in light of the results of [Del], [SV]. In particular, it holds for all symmetric pairs over fields of characteristic different from 2. One can easily show that the converse is also true. Namely, that if  $\mathcal{S}(G(F)/H(F))^K$  is a finitely generated module over the Hecke algebra  $\mathcal{H}_K(G(F))$  for any compact open subgroup  $K < G(F)$ , then (1) holds.

**Remark.** Theorem A implies that, if  $\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C})$  is finite, then it is bounded on every Bernstein component.

In the second part of the paper we introduce the notion of a uniform spherical pair and prove for them the following analog of Kazhdan's theorem.

**Theorem B** (see Theorem 3.12). *Let  $H < G$  be reductive groups defined over  $\mathbb{Z}$ . Suppose that the pair  $(G, H)$  is uniform spherical.*

*Then for any  $l$  there exists  $n$  such that for any  $n$ -close local fields  $F$  and  $E$ , the module  $\mathcal{S}(G(F)/H(F))^{K_\ell(F)}$  over the algebra  $\mathcal{H}_\ell(G(F))$  is isomorphic to the module  $\mathcal{S}(G(E)/H(E))^{K_\ell(E)}$  over the algebra  $\mathcal{H}_\ell(G(E))$ , where we identify  $\mathcal{H}_\ell(G(F))$  and  $\mathcal{H}_\ell(G(E))$  using Kazhdan's isomorphism.*

This implies the following corollary.

**Corollary C.** *Let  $(G, H)$  be a uniform spherical pair of reductive groups defined over  $\mathbb{Z}$ . Suppose that*

- *For any local field  $F$ , and any parabolic subgroup  $P \subset G$ , there is a finite number of double cosets  $P(F) \backslash G(F)/H(F)$ .*
- *For any local field  $F$  of characteristic zero the pair  $(G(F), H(F))$  is a Gelfand pair, i.e. for any irreducible smooth representation  $\rho$  of  $G(F)$  we have*

$$\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) \leq 1.$$

*Then for any local field  $F$  the pair  $(G(F), H(F))$  is a Gelfand pair.*

We prove stronger versions of the last two results, where we do not require the groups to be defined over  $\mathbb{Z}$ . See §3 for more details.

**Remark.** In a similar way one can deduce an analogous corollary for cuspidal representations. Namely, suppose that the first two conditions of the last corollary hold and the third condition holds for all cuspidal representations  $\rho$ . Then for any local field  $F$  the pair  $(G(F), H(F))$  is a cuspidal Gelfand pair: for any irreducible smooth cuspidal representation  $\rho$  of  $G(F)$  we have

$$\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) \leq 1.$$

**Remark.** Originally, we included in the formulation of Theorem B an extra condition: we demanded that the module  $\mathcal{S}(G(F)/H(F))^{K_\ell(F)}$  is finitely generated over the Hecke algebra  $\mathcal{H}_\ell(G(F))$  for any  $F$  and  $\ell$ . This was our original motivation for Theorem A. Later we realized that this condition just follows from the definition of uniform spherical pair. However, we think that Theorem A and the technique we use in its proof have importance of their own.

In the last part of the paper we apply our technique to show that  $(\operatorname{GL}_{n+1}, \operatorname{GL}_n)$  is a strong Gelfand pair over any local field and  $(\operatorname{GL}_{n+k}, \operatorname{GL}_n \times \operatorname{GL}_k)$  is a Gelfand pair over any local field of odd characteristic.

**Theorem D.** *Let  $F$  be any local field. Then  $(\operatorname{GL}_{n+1}(F), \operatorname{GL}_n(F))$  is a strong Gelfand pair, i.e. for any irreducible smooth representations  $\pi$  of  $\operatorname{GL}_{n+1}(F)$  and  $\tau$  of  $\operatorname{GL}_n(F)$  we have*

$$\dim \operatorname{Hom}_{\operatorname{GL}_n(F)}(\pi, \tau) \leq 1.$$

**Remark.** Recently, Henniart suggested a direct proof of this result, by generalizing the techniques of the proof in characteristic 0, rather than using it as a black box. This proof is not published yet.



**Theorem E.** *Let  $F$  be any local field. Suppose that  $\text{char } F \neq 2$ . Then*

$$(\text{GL}_{n+k}(F), \text{GL}_n(F) \times \text{GL}_k(F))$$

*is a Gelfand pair.*

We deduce these theorems from the zero characteristic case, which was proven in [AGRS] and [JR96] respectively. The proofs in [AGRS] and [JR96] cannot be directly adapted to the case of positive characteristic since they rely on Jordan decomposition which is problematic in positive characteristic, local fields of positive characteristic being non-perfect.

**Remark.** In [AGS08], a special case of Theorem D was proven for all local fields; namely the case when  $\tau$  is one-dimensional.

**Remark.** In [AG09a] and (independently) in [SZ], an analog of Theorem D was proven for Archimedean local fields. In [AG09b], an analog of Theorem E was proven for Archimedean local fields.

**0.1. Structure of the paper.** In Section 1 we introduce notation and give some general preliminaries.

In Section 2 we prove Theorem A.

In Subsection 2.1 we collect a few general facts for the proof. One is a criterion, due to Bernstein, for finite generation of the space of  $K$ -invariant vectors in a representation of a reductive group  $G$ ; the other facts concern homologies of  $l$ -groups. In Subsection 2.2 we prove the main inductive step in the proof of Theorem A, and in Subsection 2.3 we prove Theorem A. Subsection 2.4 is devoted to the proofs of some facts about the homologies of  $l$ -groups.

In Section 3 we prove Theorem B and derive Corollary C.

In Subsection 3.1 we introduce the notion of uniform spherical pair. In Subsection 3.2 we prove the theorem and the corollary.

We provide applications of our results in Section 4. In Subsection 4.1 we prove that the pair  $(\text{GL}_{n+k}, \text{GL}_n \times \text{GL}_k)$  satisfies the assumptions of Corollary C over fields of characteristic different from 2. In Subsections 4.3 and 4.2 we prove that the pair  $(\text{GL}_{n+1} \times \text{GL}_n, \Delta \text{GL}_n)$  satisfies the assumptions of Corollary C. These facts imply Theorems D and E.

**Acknowledgments.** We thank Joseph Bernstein for directing us to the paper [Kaz86]. We also thank Vladimir Berkovich, Joseph Bernstein, Pierre Deligne, Patrick Delorme, Jochen Heinloth, Anthony Joseph, David Kazhdan, Yiannis Sakellaridis, and Eitan Sayag for fruitful discussions and the referee for many useful remarks.

A.A. was supported by a BSF grant, a GIF grant, an ISF Center of excellency grant and ISF grant No. 583/09. N.A. was supported by NSF grant DMS-0901638. D.G.

was supported by NSF grant DMS-0635607. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

## 1. Preliminaries and notation

**Definition 1.1.** A local field is a locally compact complete non-discrete topological field. In this paper we will consider only non-Archimedean local fields. All such fields have discrete valuations.

**Remark 1.2.** Any local field of characteristic zero and residue characteristic  $p$  is a finite extension of the field  $\mathbb{Q}_p$  of  $p$ -adic numbers and any local field of characteristic  $p$  is a finite extension of the field  $\mathbb{F}_p((t))$  of formal Laurent series over the field with  $p$  elements.

**Notation 1.3.** For a local field  $F$  we denote by  $\text{val}_F$  its valuation, by  $O_F$  the ring of integers and by  $\mathcal{P}_F$  its unique maximal ideal. For an algebraic group  $G$  defined over  $O_F$  we denote by  $K_\ell(G, F)$  the kernel of the (surjective) morphism  $G(O_F) \rightarrow G(O_F/\mathcal{P}_F^\ell)$ . If  $\ell > 0$  then we call  $K_\ell(G, F)$  the  $\ell$ -th congruence subgroup.

We will use the terminology of  $l$ -spaces and  $l$ -groups introduced in [BZ76]. An  $l$ -space is a locally compact second countable totally disconnected topological space, an  $l$ -group is a  $l$ -space with a continuous group structure. For further background on  $l$ -spaces,  $l$ -groups and their representations we refer the reader to [BZ76].

**Notation 1.4.** Let  $G$  be an  $l$ -group. Denote by  $\mathcal{M}(G)$  the category of smooth complex representations of  $G$ .

Define the functor of coinvariants  $\text{CI}_G: \mathcal{M}(G) \rightarrow \text{Vect}$  by

$$\text{CI}_G(V) := V / (\text{Span}\{v - gv \mid v \in V, g \in G\}).$$

Sometimes we will also write  $V_G := \text{CI}_G(V)$ .

**Notation 1.5.** For an  $l$ -space  $X$  we denote by  $\mathcal{S}(X)$  the space of locally constant compactly supported complex valued functions on  $X$ . If  $X$  is an analytic variety over a non-Archimedean local field, we denote by  $\mathcal{M}(X)$  the space of locally constant compactly supported measures on  $X$ .

**Notation 1.6.** For an  $l$ -group  $G$  and an open compact subgroup  $K$  we denote by  $\mathcal{H}(G, K)$  or  $\mathcal{H}_K(G)$  the Hecke algebra of  $G$  w.r.t.  $K$ , i.e. the algebra of compactly supported measures on  $G$  that are invariant w.r.t. both left and right multiplication by  $K$ .

For a local field  $F$  and a reductive group  $G$  defined over  $O_F$  we will also write  $\mathcal{H}_\ell(G(F)) := \mathcal{H}_{K_\ell(G)}(G(F))$ .

**Notation 1.7.** By a reductive group over a ring  $R$ , we mean a smooth group scheme over  $\text{Spec}(R)$  all of whose geometric fibers are reductive and connected.

## 2. Finite generation of Hecke modules

The goal of this section is to prove Theorem A.

In this section  $F$  is a fixed (non-Archimedean) local field of arbitrary characteristic. All the algebraic groups and algebraic varieties that we consider in this section are defined over  $F$ . In particular, reductive means reductive over  $F$ .

For the reader's convenience, we now give an overview of the argument. In Lemma 2.10 we present a criterion, due to Bernstein, for the finite generation of spaces of  $K$ -invariants. The proof of the criterion uses the theory of Bernstein Center. This condition is given in terms of all parabolic subgroups of  $G$ . We directly prove this condition when the parabolic is  $G$  (this is Step 1 in the proof of Theorem A). The case of general parabolic is reduced to the case where the parabolic is  $G$ . For this, the main step is to show that the assumptions of the theorem imply similar assumptions for the Levi components of the parabolic subgroups of  $G$ . This is proved in Lemma 2.16 by stratifying the space  $G(F)/P(F)$  according to the  $H(F)$ -orbits inside it.

In the proof of this lemma we use two homological tools: Lemma 2.11 which gives a criterion for finite dimensionality of the first homology of a representation and Lemma 2.12 which connects the homologies of a representation and of its induction.

### 2.1. Preliminaries

**Notation 2.1.** For  $l$ -groups  $H < G$  we denote by  $\text{ind}_H^G: \mathcal{M}(H) \rightarrow \mathcal{M}(G)$  the compactly supported induction functor and by  $\text{Ind}_H^G: \mathcal{M}(H) \rightarrow \mathcal{M}(G)$  the full induction functor.

**Definition 2.2.** Let  $G$  be a reductive group, let  $P < G$  be a parabolic subgroup with unipotent radical  $U$ , and let  $M := P/U$ . Such  $M$  is called a Levi subquotient of  $G$ . Note that every representation of  $M(F)$  can be considered as a representation of  $P(F)$  using the quotient morphism  $P \twoheadrightarrow M$ . Define:

- (1) The Jacquet functor

$$r_{GM}: \mathcal{M}(G(F)) \rightarrow \mathcal{M}(M(F))$$

$$\text{by } r_{GM}(\pi) := (\pi|_{P(F)})_{U(F)}.$$

- (2) The parabolic induction functor

$$i_{GM}: \mathcal{M}(M(F)) \rightarrow \mathcal{M}(G(F))$$

$$\text{by } i_{GM}(\tau) := \text{ind}_{P(F)}^{G(F)}(\tau).$$

Note that  $i_{GM}$  is right adjoint to  $r_{GM}$ . A representation  $\pi$  of  $G(F)$  is called *cuspidal* if  $r_{GM}(\pi) = 0$  for any Levi subquotient  $M$  of  $G$ .

**Definition 2.3.** Let  $G$  be an  $l$ -group. A smooth representation  $V$  of  $G$  is called **compact** if for any  $v \in V$  and  $\xi \in \tilde{V}$  the matrix coefficient function defined by  $m_{v,\xi}(g) := \xi(gv)$  is a compactly supported function on  $G$ .

**Theorem 2.4** (Bernstein–Zelevinsky). *Let  $G$  be an  $l$ -group. Then any compact representation of  $G$  is a projective object in the category  $\mathcal{M}(G)$ .*

**Definition 2.5.** Let  $G$  be a reductive group.

(i) Denote by  $G^1$  the preimage in  $G(F)$  of the maximal compact subgroup of  $G(F)/[G, G](F)$ .

(ii) Let  $G_0 := G^1 Z(G(F))$ .

(iii) A complex character of  $G(F)$  is called *unramified* if it is trivial on  $G^1$ . We denote the set of all unramified characters by  $\Psi_G$ . Note that  $G(F)/G^1$  is a lattice and therefore we can identify  $\Psi_G$  with  $(\mathbb{C}^\times)^n$ . This defines a structure of algebraic variety on  $\Psi_G$ .

(iv) For any smooth representation  $\rho$  of  $G(F)$  we write  $\Psi(\rho) := \text{ind}_{G^1}^G(\rho|_{G^1})$ . Note that  $\Psi(\rho) \simeq \rho \otimes \mathcal{O}(\Psi_G)$ , where  $G(F)$  acts only on the first factor, but this action depends on the second factor. This identification gives a structure of  $\mathcal{O}(\Psi_G)$ -module on  $\Psi(\rho)$ .

**Remark 2.6.** The definition of unramified characters above is not the standard one, but it is more convenient for our purposes.

**Theorem 2.7** (Harish-Chandra). *Let  $G$  be a reductive group and  $V$  be a cuspidal representation of  $G(F)$ . Then  $V|_{G^1}$  is a compact representation of  $G^1$ .*

**Corollary 2.8.** *Let  $G$  be a reductive group and  $\rho$  be a cuspidal representation of  $G(F)$ . Then*

- (i)  $\rho|_{G^1}$  is a projective object in the category  $\mathcal{M}(G^1)$ .
- (ii)  $\Psi(\rho)$  is a projective object in the category  $\mathcal{M}(G(F))$ .

*Proof.* (i) is clear.

(ii) Note that

$$\text{Hom}_G(\Psi(\rho), \pi) \cong \text{Hom}_{G/G^1}(\mathcal{O}(\Psi_M), \text{Hom}_{G^1}(\rho, \pi)),$$

for any representation  $\pi$ . Therefore the functor  $\pi \mapsto \text{Hom}_G(\Psi(\rho), \pi)$  is a composition of two exact functors and hence is exact.  $\square$

**Definition 2.9.** Let  $G$  be a reductive group and  $K < G(F)$  be a compact open subgroup. Let

$$\mathcal{M}(G, K) := \{V \in \mathcal{M}(G(F)) \mid V \text{ is generated by } V^K\}$$

and

$$\mathcal{M}(G, K)^\perp := \{V \in \mathcal{M}(G(F)) \mid V^K = 0\}.$$

We call  $K$  a splitting subgroup if the category  $\mathcal{M}(G(F))$  is the direct sum of the categories  $\mathcal{M}(G, K)$  and  $\mathcal{M}(G, K)^\perp$ , and  $\mathcal{M}(G, K) \cong \mathcal{M}(\mathcal{H}_K(G))$ . Recall that an abelian category  $\mathcal{A}$  is a direct sum of two abelian subcategories  $\mathcal{B}$  and  $\mathcal{C}$ , if every object of  $\mathcal{A}$  is isomorphic to a direct sum of an object in  $\mathcal{B}$  and an object in  $\mathcal{C}$ , and, furthermore, that there are no non-trivial morphisms between objects of  $\mathcal{B}$  and  $\mathcal{C}$ .

We will use the following statements from Bernstein's theory on the center of the category  $\mathcal{M}(G)$ . Let  $P < G$  be a parabolic subgroup and  $M$  be the reductive quotient of  $P$ .

- (1) The set of splitting subgroups defines a basis at 1 for the topology of  $G(F)$ . If  $G$  splits over  $O_F$  then, for any large enough  $\ell$ , the congruence subgroup  $K_\ell(G, F)$  is splitting.
- (2) Let  $\bar{P}$  denote the parabolic subgroup of  $G$  opposite to  $P$ . Furthermore, let  $\bar{r}_{GM}: \mathcal{M}(G(F)) \rightarrow \mathcal{M}(M(F))$  denote the Jacquet functor defined using  $\bar{P}$ . Then  $\bar{r}_{GM}$  is right adjoint to  $i_{GM}$ . In particular,  $i_{GM}$  maps projective objects to projective ones and hence for any irreducible cuspidal representation  $\rho$  of  $M(F)$ ,  $i_{GM}(\Psi(\rho))$  is a projective object of  $\mathcal{M}(G(F))$ .
- (3) Denote by  $\mathcal{M}_\rho$  the subcategory of  $\mathcal{M}(G(F))$  generated by  $i_{GM}(\Psi(\rho))$ . Then

$$\mathcal{M}(G, K) = \bigoplus_{(M, \rho) \in B_K} \mathcal{M}_\rho,$$

where  $B_K$  is some finite set of pairs consisting of a Levi subquotient of  $G$  and its cuspidal representation. Moreover, for any Levi subquotient  $M < G$  and a cuspidal representation  $\rho$  of  $M(F)$  such that  $\mathcal{M}_\rho \subset \mathcal{M}(G, K)$  there exist  $(M', \rho') \in B_K$  such that  $\mathcal{M}_\rho = \mathcal{M}_{\rho'}$ .

- (4)  $\text{End}(i_{GM}(\Psi(\rho)))$  is finitely generated over  $\mathcal{O}(\Psi)$  which is finitely generated over the center of the ring  $\text{End}(i_{GM}(\Psi(\rho)))$ . The center of the ring  $\text{End}(i_{GM}(\Psi(\rho)))$  is equal to the center  $Z(\mathcal{M}_\rho)$  of the category  $\mathcal{M}_\rho$ .

For statement (1) see e.g. [Ber84], pp. 15–16, and §2 in [vD]. For statement (2) see [Ber87] or [Bus01], Theorem 3. For statements (3), (4) see [Ber84], Propositions 2.10, 2.11.

We now present a criterion, due to Bernstein, for finite generation of the space  $V^K$ , consisting of vectors in a representation  $V$  that are invariant with respect to a compact open subgroup  $K$ .

**Lemma 2.10.** *Let  $V$  be a smooth representation of  $G(F)$ . Suppose that for any parabolic  $P < G$  and any irreducible cuspidal representation  $\rho$  of  $M(F)$  (where  $M$  denotes the reductive quotient of  $P$ ),  $\text{Hom}_{G(F)}(i_{GM}(\Psi(\rho)), V)$  is a finitely generated module over  $\mathcal{O}(\Psi_M)$ . Then  $V^K$  is a finitely generated module over  $Z(\mathcal{H}_K(G(F)))$ , for any compact open subgroup  $K < G(F)$ .*

*Proof.* *Step 1.* Proof for the case when  $K$  is splitting and  $V = i_{GM}(\Psi(\rho))$  for some Levi subquotient  $M$  of  $G$  and an irreducible cuspidal representation  $\rho$  of  $M(F)$ . Let  $P$  denote the parabolic subgroup that defines  $M$  and  $U$  denote its unipotent radical. Let  $K_M := K/(U(F) \cap K) < M(F)$ . If  $V^K = 0$  there is nothing to prove. Otherwise  $\mathcal{M}_\rho$  is a direct summand of  $\mathcal{M}(G, K)$ . Now

$$V^K = \Psi(\rho)^{K_M} = \rho^{K_M} \otimes \mathcal{O}(\Psi).$$

Hence  $V^K$  is finitely generated over  $Z(\mathcal{M}_\rho)$ . Hence  $V^K$  is finitely generated over  $Z(\mathcal{M}(G, K)) = Z(\mathcal{H}_K(G))$ .

*Step 2.* Proof for the case when  $K$  is splitting and  $V \in \mathcal{M}_\rho$  for some Levi subquotient  $M < G$  and an irreducible cuspidal representation  $\rho$  of  $M(F)$ .

Let

$$\phi: i_{GM}(\Psi(\rho)) \otimes \text{Hom}(i_{GM}(\Psi(\rho)), V) \twoheadrightarrow V$$

be the natural epimorphism. We are given that  $\text{Hom}(i_{GM}(\Psi(\rho)), V)$  is finitely generated over  $\mathcal{O}(\Psi)$ . Hence it is finitely generated over  $Z(\mathcal{M}(\rho))$ . Choose some generators  $\alpha_1, \dots, \alpha_n \in \text{Hom}(i_{GM}(\Psi(\rho)))$ . Let

$$\psi: i_{GM}(\Psi(\rho))^n \hookrightarrow i_{GM}(\Psi(\rho)) \otimes \text{Hom}(i_{GM}(\Psi(\rho)), V)$$

be the corresponding morphism.  $\text{Im}(\phi \circ \psi)$  is  $Z(\mathcal{M}(\rho))$ -invariant and hence coincides with  $\text{Im}(\phi)$ . Hence  $\phi \circ \psi$  is onto. The statement now follows from the previous step.

*Step 3.* Proof for the case when  $K$  is splitting. Let  $W < V$  be the subrepresentation generated by  $V^K$ . By definition  $W \in \mathcal{M}(G, K)$  and hence  $W = \bigoplus_{i=1}^n W_i$  where  $W_i \in \mathcal{M}_{\rho_i}$  for some  $\rho_i$ . The lemma now follows from the previous step.

*Step 4.* General case. Let  $K'$  be a splitting subgroup s.t.  $K' < K$ . Let  $v_1, \dots, v_n \in V^{K'}$  be the generators of  $V^{K'}$  over  $Z(\mathcal{H}_{K'}(G(F)))$  given by the previous step. Define  $w_i := e_K v_i \in V^K$  where  $e_K \in \mathcal{H}_K(G(F))$  is the normalized Haar measure of  $K$ . Let us show that  $w_i$  generate  $V^K$  over  $Z(\mathcal{H}_K(G(F)))$ . Let  $x \in V^K$ . We can represent  $x$  as a sum  $\sum h_i v_i$ , where  $h_i \in Z(\mathcal{H}_{K'}(G(F)))$ . Now

$$\begin{aligned} x &= e_K x = \sum e_K h_i v_i = \sum e_K e_K h_i v_i = \sum e_K h_i e_K v_i \\ &= \sum e_K h_i e_K e_K v_i = \sum e_K h_i e_K w_i. \end{aligned} \quad \square$$

Finally, in this subsection, we state two facts about homologies of  $l$ -groups. The proofs and relevant definitions are in Subsection 2.4.

**Lemma 2.11.** *Let  $G$  be an algebraic group and  $U$  be its unipotent radical. Let  $\rho$  be an irreducible cuspidal representation of  $(G/U)(F)$ . We treat  $\rho$  as a representation of  $G(F)$  with trivial action of  $U(F)$ .*

*Let  $H < G$  be an algebraic subgroup. Suppose that the space of coinvariants  $\rho_{H(F)}$  is finite dimensional. Then  $\dim H_1(H(F), \rho) < \infty$ .*

We will also use the following version of Shapiro's lemma.

**Lemma 2.12.** *Let  $G$  be an  $l$ -group that acts transitively on an  $l$ -space  $X$ . Let  $\mathcal{F}$  be a  $G$ -equivariant sheaf over  $X$ . Choose a point  $x \in X$ , let  $\mathcal{F}_x$  denote the stalk of  $\mathcal{F}$  at  $x$  and let  $G_x$  denote the stabilizer of  $x$ . Then*

$$H_i(G, \mathcal{F}(X)) = H_i(G_x, \mathcal{F}_x).$$

## 2.2. Descent of finite multiplicity

**Definition 2.13.** We call a pair  $(G, H)$  consisting of a reductive group  $G$  and an algebraic subgroup  $H$  an  $F$ -spherical pair if for any parabolic subgroup  $P \subset G$ , there is a finite number of double cosets in  $P(F) \backslash G(F)/H(F)$ .

**Remark 2.14.** If  $\text{char } F = 0$  and  $G$  is quasi-split over  $F$  then  $(G, H)$  is an  $F$ -spherical pair if and only if it is a spherical pair of algebraic groups. However, we do not know whether this is true if  $\text{char } F > 0$ .

**Notation 2.15.** Let  $G$  be a reductive group and  $H$  be a subgroup. Let  $P < G$  be a parabolic subgroup and  $M$  be its Levi quotient. We denote by  $H_M$  the image of  $H \cap P$  under the projection  $P \twoheadrightarrow M$ .

The following lemma is the main step in the proof of Theorem A

**Lemma 2.16.** *Let  $(G, H)$  be an  $F$ -spherical pair. Let  $P < G$  be a parabolic subgroup and  $M$  be its Levi quotient. Then*

- (i)  $(M, H_M)$  is also an  $F$ -spherical pair.
- (ii) Suppose also that for any smooth irreducible representation  $\rho$  of  $G(F)$  we have

$$\dim \text{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) < \infty.$$

*Then for any irreducible cuspidal representation  $\sigma$  of  $M(F)$  we have*

$$\dim \text{Hom}_{H_M(F)}(\sigma|_{H_M(F)}, \mathbb{C}) < \infty.$$

**Remark 2.17.** One can show that the converse of (ii) is also true. Namely, if  $\dim \text{Hom}_{H_M(F)}(\sigma|_{H_M(F)}, \mathbb{C}) < \infty$  for any irreducible cuspidal representation  $\sigma$  of  $M(F)$  for any Levi subquotient  $M$  then  $\dim \text{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) < \infty$  for any smooth irreducible representation  $\rho$  of  $G(F)$ . We will not prove this since we will not use this.



We will need the following lemma.

**Lemma 2.18.** *Let  $M$  be an  $l$ -group and  $V$  be a smooth representation of  $M$ . Let  $0 = F^0 V \subset \cdots \subset F^{n-1} V \subset F^n V = V$  be a finite filtration of  $V$  by subrepresentations. Suppose that for any  $i$ , either*

$$\dim(F^i V / F^{i-1} V)_M = \infty$$

or

$$\text{both } \dim(F^i V / F^{i-1} V)_M < \infty \text{ and } \dim H_1(M, (F^i V / F^{i-1} V)) < \infty.$$

Suppose also that  $\dim V_M < \infty$ . Then  $\dim(F^i V / F^{i-1} V)_M < \infty$  for any  $i$ .

*Proof.* We prove by a decreasing induction on  $i$  that  $\dim(F^i V)_M < \infty$ , and, therefore,  $\dim(F^i V / F^{i-1} V)_M < \infty$ . Consider the short exact sequence

$$0 \rightarrow F^{i-1} V \rightarrow F^i V \rightarrow F^i V / F^{i-1} V \rightarrow 0,$$

and the corresponding long exact sequence

$$\begin{aligned} \cdots \rightarrow H_1(M, (F^i V / F^{i-1} V)) &\rightarrow (F^{i-1} V)_M \\ &\rightarrow (F^i V)_M \rightarrow (F^i V / F^{i-1} V)_M \rightarrow 0. \end{aligned}$$

In this sequence  $\dim H_1(M, (F^i V / F^{i-1} V)) < \infty$  and  $\dim(F^i V)_M < \infty$ , and hence  $\dim(F^{i-1} V)_M < \infty$ .  $\square$

Now we are ready to prove Lemma 2.16.

*Proof of Lemma 2.16.* (i) is trivial.

(ii) Let  $P < G$  be a parabolic subgroup,  $M$  be the Levi quotient of  $P$  and let  $\rho$  be a cuspidal representation of  $M(F)$ . We know that  $\dim(i_{GM}\rho)_{H(F)} < \infty$  and we have to show that  $\dim \rho_{H_M(F)} < \infty$ .

Let  $\mathcal{I}$  denote the natural  $G(F)$ -equivariant locally constant sheaf of complex vector spaces on  $G(F)/P(F)$  such that  $i_{GM}\rho \cong \mathcal{S}(G(F)/P(F), \mathcal{I})$ . Let  $Y_j$  denote the  $H(F)$  orbits on  $G(F)/P(F)$ . We know that there exists a natural filtration on  $\mathcal{S}(G(F)/P(F), \mathcal{I})|_{H(F)}$  with associated graded components isomorphic to  $\mathcal{S}(Y_j, \mathcal{I}_j)$ , where  $\mathcal{I}_j$  are  $H(F)$ -equivariant sheaves on  $Y_j$  corresponding to  $\mathcal{I}$ . For any  $j$  choose a representative  $y_j \in Y_j$ . Do it in such a way that there exists  $j_0$  such that  $y_{j_0} = [1]$ . Let  $P_j := G_{y_j}$  and  $M_j$  be its Levi quotient. Note that  $P_{j_0} = P$  and  $M_{j_0} = M$ . Let  $\rho_j$  be the stalk of  $\mathcal{I}_j$  at the point  $y_j$ . Clearly  $\rho_j$  is a cuspidal irreducible representation of  $M_j(F)$  and  $\rho_{j_0} = \rho$ . By Shapiro's lemma (Lemma 2.12)

$$H_i(H(F), \mathcal{S}(Y_j, \mathcal{I}_j)) \cong H_i((H \cap P_j)(F), \rho_j).$$

By Lemma 2.11 either  $\dim H_0((H \cap P_j)(F), \rho_j) = \infty$  or both  $\dim H_0((H \cap P_j)(F), \rho_j) < \infty$  and  $\dim H_1((H \cap P_j)(F), \rho_j) < \infty$ . Hence by Lemma 2.18  $\dim H_0((H \cap P_j)(F), \rho_j) < \infty$  and hence  $\dim \rho_{H_M(F)} < \infty$ .  $\square$



**2.3. Proof of Theorem A.** In this subsection we prove Theorem A. Let us remind its formulation.

**Theorem 2.19.** *Let  $(G, H)$  be an  $F$ -spherical pair. Suppose that for any irreducible smooth representation  $\rho$  of  $G(F)$  we have*

$$\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) < \infty. \quad (2)$$

*Then for any compact open subgroup  $K < G(F)$ ,  $\mathcal{S}(G(F)/H(F))^K$  is a finitely generated module over the Hecke algebra  $\mathcal{H}_K(G(F))$ .*

**Remark 2.20.** Conjecturally, any  $F$ -spherical pair satisfies the condition (2). In [Del] and in [SV] this is proven for wide classes of spherical pairs, which include all symmetric pairs over fields of characteristic different from 2.

We will need several lemmas and definitions.

**Lemma 2.21.** *Let  $(G, H)$  be an  $F$ -spherical pair, and let  $\tilde{H} = H(F)Z(G(F)) \cap G^1$ . Suppose that for any smooth (respectively cuspidal) irreducible representation  $\rho$  of  $G(F)$  we have  $\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) < \infty$ . Then for any smooth (respectively cuspidal) irreducible representation  $\rho$  of  $G(F)$  and for every character  $\tilde{\chi}$  of  $\tilde{H}$  whose restriction to  $H(F) \cap G^1$  is trivial, we have*

$$\dim \operatorname{Hom}_{\tilde{H}}(\rho|_{\tilde{H}}, \tilde{\chi}) < \infty.$$

*Proof.* Let  $\rho$  be a smooth (respectively cuspidal) irreducible representation of  $G(F)$ , and let  $\tilde{\chi}$  be a character of  $\tilde{H}$  whose restriction to  $H(F) \cap G^1$  is trivial.

$$\begin{aligned} & \operatorname{Hom}_{\tilde{H}}(\rho|_{\tilde{H}}, \tilde{\chi}) \\ &= \operatorname{Hom}_{(H(F)Z(G(F))) \cap G_0}(\rho|_{(H(F)Z(G(F))) \cap G_0}, \operatorname{Ind}_{\tilde{H}}^{(H(F)Z(G(F))) \cap G_0} \tilde{\chi}). \end{aligned}$$

Since

$$H(F)Z(G(F)) \cap G_0 = \tilde{H}Z(G(F)) \cap G_0 = \tilde{H}Z(G(F)),$$

the subspace of  $\operatorname{Ind}_{\tilde{H}}^{(H(F)Z(G(F))) \cap G_0} \tilde{\chi}$  that transforms under  $Z(G(F))$  according to the central character of  $\rho$  is at most one dimensional. If this subspace is trivial, then the lemma is clear. Otherwise, denote it by  $\tau$ . Since  $H(F) \cap G^1$  is normal in  $H(F)Z(G(F))$ , we get that the restriction of  $\operatorname{Ind}_{\tilde{H}}^{(H(F)Z(G(F))) \cap G_0} \tilde{\chi}$  to  $H(F) \cap G^1$  is trivial, and hence that  $\tau|_{H(F) \cap G^1}$  is trivial. Hence  $\operatorname{Hom}_{\tilde{H}}(\rho|_{\tilde{H}}, \tilde{\chi})$  is equal to

$$\begin{aligned} & \operatorname{Hom}_{(H(F)Z(G(F))) \cap G_0}(\rho|_{(H(F)Z(G(F))) \cap G_0}, \tau) \\ &= \operatorname{Hom}_{H(F) \cap G_0}(\rho|_{H(F) \cap G_0}, \tau|_{H(F) \cap G_0}) \\ &= \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \operatorname{Ind}_{H(F) \cap G_0}^{H(F)} \tau|_{H(F) \cap G_0}). \end{aligned}$$

As  $H(F)/H(F) \cap G_0$  is finite and abelian, the representation  $\text{Ind}_{H(F) \cap G_0}^{H(F)} \tau|_{H(F) \cap G_0}$  is a finite direct sum of characters of  $H(F)$ , the restrictions of all to  $H(F) \cap G^1$  are trivial. Any character  $\theta$  of  $H(F)$  whose restriction to  $H(F) \cap G^1$  is trivial can be extended to a character of  $G(F)$ , because  $H(F)/(H(F) \cap G^1)$  is a sub-lattice of  $G(F)/G^1$ . Denoting the extension by  $\Theta$ , we get that

$$\text{Hom}_{H(F)}(\rho|_{H(F)}, \theta) = \text{Hom}_{H(F)}((\rho \otimes \Theta^{-1})|_{H(F)}, \mathbb{C}),$$

but  $\rho \otimes \Theta^{-1}$  is again smooth (respectively cuspidal) irreducible representation of  $G(F)$ , so this last space is finite-dimensional.  $\square$

**Lemma 2.22.** *Let  $A$  be a commutative unital Noetherian algebra without zero divisors and let  $K$  be its field of fractions. Let  $K^{\mathbb{N}}$  be the space of all sequences of elements of  $K$ . Let  $V$  be a finite dimensional subspace of  $K^{\mathbb{N}}$  and let  $M := V \cap A^{\mathbb{N}}$ . Then  $M$  is finitely generated.*

*Proof.* Since  $A$  does not have zero divisors,  $M$  injects into  $K^{\mathbb{N}}$ . There is a number  $n$  such that the projection of  $V$  to  $K^{\{1, \dots, n\}}$  is injective. Therefore,  $M$  injects into  $A^{\{1, \dots, n\}}$ , and, since  $A$  is Noetherian,  $M$  is finitely generated.  $\square$

**Lemma 2.23.** *Let  $M$  be an  $l$ -group, let  $L \subset M$  be a closed subgroup, and let  $L' \subset L$  be an open normal subgroup of  $L$  such that  $L/L'$  is a lattice. Let  $\rho$  be a smooth representation of  $M$  of countable dimension. Suppose that for any character  $\chi$  of  $L$  whose restriction to  $L'$  is trivial we have*

$$\dim \text{Hom}_L(\rho|_L, \chi) < \infty.$$

*Consider  $\text{Hom}_{L'}(\rho, \mathcal{S}(L/L'))$  as a representation of  $L$ , where  $L$  acts by*

$$((hf)(x))([y]) = (f(x))([yh]).$$

*Then this representation is finitely generated.*

*Proof.* By assumption, the action of  $L$  on  $\text{Hom}_{L'}(\rho, \mathcal{S}(L/L'))$  factors through  $L/L'$ . Since  $L/L'$  is discrete,  $\mathcal{S}(L/L')$  is the group algebra  $\mathbb{C}[L/L']$ . We want to show that  $\text{Hom}_{L'}(\rho, \mathbb{C}[L/L'])$  is a finitely generated module over  $\mathbb{C}[L/L']$ .

Let  $\mathbb{C}(L/L')$  be the fraction field of  $\mathbb{C}[L/L']$ . Choosing a countable basis for the vector space of  $\rho$ , we can identify any  $\mathbb{C}$ -linear map from  $\rho$  to  $\mathbb{C}[L/L']$  with an element of  $\mathbb{C}[L/L']^{\mathbb{N}}$ . Moreover, the condition that the map intertwines the action of  $L/L'$  translates into a collection of linear equations that the tuple in  $\mathbb{C}[L/L']^{\mathbb{N}}$  should satisfy. Hence,  $\text{Hom}_{L'}(\rho, \mathbb{C}[L/L'])$  is the intersection of the  $\mathbb{C}(L/L')$ -vector space  $\text{Hom}_{L'}(\rho, \mathbb{C}(L/L'))$  and  $\mathbb{C}[L/L']^{\mathbb{N}}$ . By Lemma 2.22, it suffices to prove that  $\text{Hom}_{L'}(\rho, \mathbb{C}(L/L'))$  is finite dimensional over  $\mathbb{C}(L/L')$ .

Since  $L'$  is separable, and  $\rho$  is smooth and of countable dimension, there are only countably many linear equations defining  $\text{Hom}_{L'}(\rho, \mathbb{C}(L/L'))$ ; denote them by  $\phi_1, \phi_2, \dots \in (\mathbb{C}(L/L')^{\mathbb{N}})^*$ . Choose a countable subfield  $K \subset \mathbb{C}$  that contains all the coefficients of the elements of  $\mathbb{C}(L/L')$  that appear in any of the  $\phi_i$ 's. If we define  $W$  as the  $K(L/L')$ -linear subspace of  $K(L/L')^{\mathbb{N}}$  defined by the  $\phi_i$ 's, then  $\text{Hom}_{L'}(\rho, \mathbb{C}(L/L')) = W \otimes_{K(L/L')} \mathbb{C}(L/L')$ , so  $\dim_{\mathbb{C}(L/L')} \text{Hom}_{L'}(\rho, \mathbb{C}(L/L')) = \dim_{K(L/L')} W$ .

Since  $L/L'$  is a lattice generated by, say,  $g_1, \dots, g_n$ , we get that  $K(L/L') = K(t_1^{\pm 1}, \dots, t_n^{\pm 1}) = K(t_1, \dots, t_n)$ . Choosing elements  $\pi_1, \dots, \pi_n \in \mathbb{C}$  such that  $\text{tr.deg}_K(K(\pi_1, \dots, \pi_n)) = n$ , we get an injection  $\iota$  of  $K(L/L')$  into  $\mathbb{C}$ . As before, we get that if we denote the  $\mathbb{C}$ -vector subspace of  $\mathbb{C}^{\mathbb{N}}$  cut by the equations  $\iota(\phi_i)$  by  $U$ , then  $\dim_{K(L/L')} W = \dim_{\mathbb{C}} U$ . However,  $U$  is isomorphic to  $\text{Hom}_{L'}(\rho, \chi)$ , where  $\chi$  is the character of  $L/L'$  such that  $\chi(g_i) = \pi_i$ . By assumption, this last vector space is finite dimensional.  $\square$

Now we are ready to prove Theorem 2.19.

*Proof of Theorem 2.19.* By Lemma 2.10 it is enough to show that for any parabolic  $P < G$  and any irreducible cuspidal representation  $\rho$  of  $M(F)$  (where  $M$  denotes the Levi quotient of  $P$ ),  $\text{Hom}(i_{GM}(\Psi(\rho)), \mathcal{S}(G(F)/H(F)))$  is a finitely generated module over  $\mathcal{O}(\Psi_M)$ .

*Step 1.* Proof for the case  $P = G$ .

We have

$$\begin{aligned} \text{Hom}_{G(F)}(i_{GM}(\Psi(\rho)), \mathcal{S}(G(F)/H(F))) &= \text{Hom}_{G(F)}(\Psi(\rho), \mathcal{S}(G(F)/H(F))) \\ &= \text{Hom}_{G^1}(\rho, \mathcal{S}(G(F)/H(F))). \end{aligned}$$

Here we consider the space  $\text{Hom}_{G^1}(\rho, \mathcal{S}(G(F)/H(F)))$  with the natural action of  $G$ . Note that  $G^1$  acts trivially and hence this action gives rise to an action of  $G/G^1$ , which gives the  $\mathcal{O}(\Psi_G)$ -module structure.

Now consider the subspace

$$V := \text{Hom}_{G^1}(\rho, \mathcal{S}(G^1/(H(F) \cap G^1))) \subset \text{Hom}_{G^1}(\rho, \mathcal{S}(G(F)/H(F))).$$

It generates  $\text{Hom}_{G^1}(\rho, \mathcal{S}(G(F)/H(F)))$  as a representation of  $G(F)$ , and therefore also as an  $\mathcal{O}(\Psi_G)$ -module. Note that  $V$  is  $H(F)$  invariant. Therefore it is enough to show that  $V$  is finitely generated over  $H(F)$ . Write  $H' := H(F) \cap G^1$  and  $H'' := (H(F)Z(G(F))) \cap G^1$ . Note that

$$\mathcal{S}(G^1/H') \cong \text{ind}_{H''}^{G^1}(\mathcal{S}(H''/H')) \subset \text{Ind}_{H''}^{G^1}(\mathcal{S}(H''/H')).$$

Therefore  $V$  is canonically embedded into  $\text{Hom}_{H''}(\rho, \mathcal{S}(H''/H'))$ . The action of  $H$  on  $V$  is naturally extended to an action  $\Pi$  on  $\text{Hom}_{H''}(\rho, \mathcal{S}(H''/H'))$  by

$$((\Pi(h)(f))(v))([k]) = f(h^{-1}v)([h^{-1}kh]).$$

Let  $\Xi$  be the action of  $H''$  on  $\text{Hom}_{H''}(\rho, \mathcal{S}(H''/H'))$  as described in Lemma 2.23, i.e.

$$((\Xi(h)(f))(v))([k]) = f(v)([kh]).$$

By Lemmas 2.23 and 2.21 it is enough to show that for any  $h \in H''$  there exist an  $h' \in H$  and a scalar  $\alpha$  s.t.

$$\Xi(h) = \alpha \Pi(h').$$

In order to show this let us decompose  $h$  to a product  $h = zh'$  where  $h' \in H$  and  $z \in Z(G(F))$ . Now

$$\begin{aligned} ((\Xi(h)(f))(v))([k]) &= f(v)([kh]) = f(h^{-1}v)([h^{-1}kh]) \\ &= f(h'^{-1}z^{-1}v)([h'^{-1}kh']) = \alpha f(h'^{-1}v)([h'^{-1}kh']) \\ &= \alpha((\Pi(h')(f))(v))([k]), \end{aligned}$$

where  $\alpha$  is the scalar with which  $z^{-1}$  acts on  $\rho$ .

*Step 2. Proof in the general case.*

$$\begin{aligned} &\text{Hom}_{G(F)}(i_{GM}(\Psi(\rho)), \mathcal{S}(G(F)/H(F))) \\ &= \text{Hom}_{M(F)}(\Psi(\rho), \bar{r}_{MG}(\mathcal{S}(G(F)/H(F)))) \\ &= \text{Hom}_{M(F)}(\Psi(\rho), ((\mathcal{S}(G(F)/H(F)))|_{\bar{P}(F)})_{\bar{U}(F)}), \end{aligned}$$

where  $\bar{U}$  is the unipotent radical of  $\bar{P}$ , the parabolic opposite to  $P$ . Let  $\{Y_i\}_{i=1}^n$  be the orbits of  $\bar{P}(F)$  on  $G(F)/H(F)$ . We know that there exists a filtration on  $(\mathcal{S}(G(F)/H(F)))|_{\bar{P}(F)}$  such that the associated graded components are isomorphic to  $\mathcal{S}(Y_i)$ . Consider the corresponding filtration on  $((\mathcal{S}(G(F)/H(F)))|_{\bar{P}(F)})_{\bar{U}(F)}$ . Let  $V_i$  be the associated graded components of this filtration. We have a natural surjection  $\mathcal{S}(Y_i)_{\bar{U}} \twoheadrightarrow V_i$ . In order to prove that

$$\text{Hom}_{M(F)}(\Psi(\rho), ((\mathcal{S}(G(F)/H(F)))|_{\bar{P}(F)})_{\bar{U}(F)})$$

is finitely generated it is enough to prove that  $\text{Hom}_{M(F)}(\Psi(\rho), V_i)$  is finitely generated. Since  $\Psi(\rho)$  is a projective object of  $\mathcal{M}(M(F))$  (by Corollary 2.8), it is enough to show that  $\text{Hom}_{M(F)}(\Psi(\rho), \mathcal{S}(Y_i)_{\bar{U}(F)})$  is finitely generated. Let  $Z_i := \bar{U}(F) \setminus Y_i$ . It is easy to see that  $Z_i \cong M(F)/((H_i)_M(F))$ , where  $H_i$  is some conjugation of  $H$ .

Now the assertion follows from the previous step using Lemma 2.16.  $\square$

**2.4. Homologies of  $I$ -groups.** The goal of this subsection is to prove Lemma 2.11 and Lemma 2.12.

We start with some generalities on abelian categories.

**Definition 2.24.** Let  $\mathcal{C}$  be an abelian category. We call a family of objects  $\mathcal{A} \subset \text{Ob}(\mathcal{C})$  *generating family* if for any object  $X \in \text{Ob}(\mathcal{C})$  there exists an object  $Y \in \mathcal{A}$  and an epimorphism  $Y \twoheadrightarrow X$ .

**Definition 2.25.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be abelian categories and let  $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{D}$  be a right-exact additive functor. A family of objects  $\mathcal{A} \subset \text{Ob}(\mathcal{C})$  is called  $\mathcal{F}$ -adapted if it is generating, closed under direct sums and for any exact sequence  $0 \rightarrow A_1 \rightarrow A_2 \rightarrow \cdots$  with  $A_i \in \mathcal{A}$ , the sequence  $0 \rightarrow \mathcal{F}(A_1) \rightarrow \mathcal{F}(A_2) \rightarrow \cdots$  is also exact.

For example, a generating, closed under direct sums system consisting of projective objects is  $\mathcal{F}$ -adapted for any right-exact functor  $\mathcal{F}$ . For an  $l$ -group  $G$  the system of objects consisting of direct sums of copies of  $\mathcal{S}(G)$  is an example of such system.

The following results are well known.

**Theorem 2.26.** Let  $\mathcal{C}$  and  $\mathcal{D}$  be abelian categories and  $\mathcal{F} : \mathcal{C} \rightarrow \mathcal{D}$  be a right-exact additive functor. Suppose that there exists an  $\mathcal{F}$ -adapted family  $\mathcal{A} \subset \text{Ob}(\mathcal{C})$ . Then  $\mathcal{F}$  has derived functors.

**Lemma 2.27.** Let  $\mathcal{A}$ ,  $\mathcal{B}$  and  $\mathcal{C}$  be abelian categories. Let  $\mathcal{F} : \mathcal{A} \rightarrow \mathcal{B}$  and  $\mathcal{G} : \mathcal{B} \rightarrow \mathcal{C}$  be right-exact additive functors. Suppose that both  $\mathcal{F}$  and  $\mathcal{G}$  have derived functors.

(i) Suppose that  $\mathcal{F}$  is exact. Suppose also that there exists a class  $\mathcal{E} \subset \text{Ob}(\mathcal{A})$  which is  $\mathcal{G} \circ \mathcal{F}$ -adapted and such that  $\mathcal{F}(X)$  is  $\mathcal{G}$ -acyclic for any  $X \in \mathcal{E}$ . Then the functors  $L^i(\mathcal{G} \circ \mathcal{F})$  and  $L^i \mathcal{G} \circ \mathcal{F}$  are isomorphic.

(ii) Suppose that there exists a class  $\mathcal{E} \subset \text{Ob}(\mathcal{A})$  which is  $\mathcal{G} \circ \mathcal{F}$ -adapted and  $\mathcal{F}$ -adapted and such that  $\mathcal{F}(X)$  is  $\mathcal{G}$ -acyclic for any  $X \in \mathcal{E}$ . Let  $Y \in \mathcal{A}$  be an  $\mathcal{F}$ -acyclic object. Then  $L^i(\mathcal{G} \circ \mathcal{F})(Y)$  is (naturally) isomorphic to  $L^i \mathcal{G}(\mathcal{F}(Y))$ .

(iii) Suppose that  $\mathcal{G}$  is exact. Suppose that there exists a class  $\mathcal{E} \subset \text{Ob}(\mathcal{A})$  which is  $\mathcal{G} \circ \mathcal{F}$ -adapted and  $\mathcal{F}$ -adapted. Then the functors  $L^i(\mathcal{G} \circ \mathcal{F})$  and  $\mathcal{G} \circ L^i \mathcal{F}$  are isomorphic.

**Definition 2.28.** Let  $G$  be an  $l$ -group. For any smooth representation  $V$  of  $G$  let  $H_i(G, V) := L^i \text{CI}_G(V)$ . Recall that  $\text{CI}_G$  denotes the coinvariants functor.

*Proof of Lemma 2.12.* Note that  $\mathcal{F}(X) = \text{ind}_{G_x}^G \mathcal{F}_x$ . Note also that  $\text{ind}_{G_x}^G$  is an exact functor, and  $\text{CI}_{G_x} = \text{CI}_G \circ \text{ind}_{G_x}^G$ . The lemma follows now from Lemma 2.27 (i).  $\square$

**Lemma 2.29.** Let  $L$  be a lattice. Let  $V$  be a linear space. Let  $L$  act on  $V$  by a character. Then

$$H_1(L, V) = H_0(L, V) \otimes_{\mathbb{C}} (L \otimes_{\mathbb{Z}} \mathbb{C}).$$

The proof of this lemma is straightforward.

**Lemma 2.30.** Let  $L$  be an  $l$ -group and  $L' < L$  be a subgroup. Then:

(i) For any representation  $V$  of  $L$  we have

$$H_i(L', V) = L^i \mathcal{F}(V),$$

where  $\mathcal{F}: \mathcal{M}(L) \rightarrow \text{Vect}$  is the functor defined by  $\mathcal{F}(V) = V_{L'}$ .

(ii) Suppose that  $L'$  is normal. Let  $\mathcal{F}': \mathcal{M}(L) \rightarrow \mathcal{M}(L/L')$  be the functor defined by  $\mathcal{F}'(V) = V_{L'}$ . Then for any representation  $V$  of  $L$  we have  $H_i(L', V) = L^i \mathcal{F}'(V)$ .

*Proof.* (i) Consider the restriction functor  $\text{Res}_{L'}^L: \mathcal{M}(L) \rightarrow \mathcal{M}(L')$ . Note that it is exact. Consider also the functor  $\mathcal{G}: \mathcal{M}(L') \rightarrow \text{Vect}$  defined by  $\mathcal{G}(\rho) := \rho_{L'}$ . Note that  $\mathcal{F} = \mathcal{G} \circ \text{Res}_{L'}^L$ . The assertion follows now from Lemma 2.27 (i) using the fact that  $\mathcal{S}(L)$  is a projective object in  $\mathcal{M}(L')$ .

(ii) follows from (i) in a similar way, but using part (iii) of Lemma 2.27 instead part (i).  $\square$

**Lemma 2.31.** *Let  $G$  be a reductive group and  $H < G$  be a subgroup. Consider the functor*

$$\mathcal{F}: \mathcal{M}(G(F)) \rightarrow \mathcal{M}(H(F)/(H(F) \cap G^1)) \text{ defined by } \mathcal{F}(V) = V_{H(F) \cap G^1}.$$

*Then any finitely generated cuspidal representation of  $G(F)$  is an  $\mathcal{F}$ -acyclic object.*

*Proof.* Consider the restriction functors

$$\text{Res}_1^{H(F)/(H(F) \cap G^1)}: \mathcal{M}(H(F)/(H(F) \cap G^1)) \rightarrow \text{Vect}$$

and

$$\text{Res}_{G^1}^{G(F)}: \mathcal{M}(G(F)) \rightarrow \mathcal{M}(G^1).$$

Note that they are exact. Consider also the functor  $\mathcal{G}: \mathcal{M}(G^1) \rightarrow \text{Vect}$  defined by  $\mathcal{G}(\rho) := \rho_{G^1 \cap H(F)}$ . Let  $\mathcal{E} := \mathcal{G} \circ \text{Res}_{G^1}^{G(F)}$ . Note that  $\mathcal{E} = \text{Res}_1^{H(F)/(H(F) \cap G^1)} \circ \mathcal{F}$ .

$$\begin{array}{ccc} \mathcal{M}(G(F)) & \xrightarrow{\mathcal{F}} & \mathcal{M}(H(F)/(H(F) \cap G^1)) \\ \text{Res}_{G^1}^{G(F)} \downarrow & \searrow \mathcal{E} & \downarrow \text{Res}_1^{H(F) \cap G^1} \\ \mathcal{M}(G^1) & \xrightarrow{\mathcal{G}} & \text{Vect} \end{array}$$

Let  $\pi$  be a cuspidal finitely generated representation of  $G(F)$ . By Corollary 2.8,  $\text{Res}_{G^1}^{G(F)}(\pi)$  is projective and hence  $\mathcal{G}$ -acyclic. Hence by Lemma 2.27 (ii)  $\pi$  is  $\mathcal{E}$ -acyclic. Hence by Lemma 2.27 (iii)  $\pi$  is  $\mathcal{F}$ -acyclic.  $\square$

**Lemma 2.32.** *Let  $L$  be an  $l$ -group and  $L' < L$  be a normal subgroup. Suppose that  $H_i(L', \mathbb{C}) = 0$  for all  $i > 0$ . Let  $\rho$  be a representation of  $L/L'$ . Denote by  $\text{Ext}(\rho)$  the natural representation of  $L$  obtained from  $\rho$ . Then  $H_i(L/L', \rho) = H_i(L, \text{Ext}(\rho))$ .*

*Proof.* Consider the coinvariants functors  $\mathcal{E}: \mathcal{M}(L) \rightarrow \text{Vect}$  and  $\mathcal{F}: \mathcal{M}(L/L') \rightarrow \text{Vect}$  defined by  $\mathcal{E}(V) := V_L$  and  $\mathcal{F}(V) := V_{L/L'}$ . Note that  $\mathcal{F} = \mathcal{E} \circ \text{Ext}$  and  $\text{Ext}$  is exact. By Shapiro's lemma (Lemma 2.12),  $\mathcal{S}(L/L')$  is acyclic with respect to both  $\mathcal{E}$  and  $\mathcal{F}$ . The lemma follows now from Lemma 2.27 (ii).  $\square$

**Remark 2.33.** Recall that if  $L' = N(F)$  where  $N$  is a unipotent algebraic group, then  $H_i(L') = 0$  for all  $i > 0$ .

Now we are ready to prove Lemma 2.11

*Proof of Lemma 2.11.* By Lemma 2.32 we can assume that  $G$  is reductive.

Let  $\mathcal{F}: \mathcal{M}(G(F)) \rightarrow \text{Vect}$  be the functor defined by  $\mathcal{F}(V) := V_{H(F)}$ . Let

$$\mathcal{G}: \mathcal{M}(G(F)) \rightarrow \mathcal{M}(H(F)/(H(F) \cap G^1))$$

be defined by

$$\mathcal{G}(V) := V_{H(F) \cap G^1}.$$

Let

$$\mathcal{E}: \mathcal{M}(H(F)/(H(F) \cap G^1)) \rightarrow \text{Vect}$$

be defined by

$$\mathcal{E}(V) := V_{H(F)/(H(F) \cap G^1)}.$$

Clearly,  $\mathcal{F} = \mathcal{E} \circ \mathcal{G}$ . By Lemma 2.31,  $\rho$  is  $\mathcal{G}$ -acyclic. Hence by Lemma 2.27 (ii),  $L^i \mathcal{F}(\rho) = L^i \mathcal{E}(\mathcal{G}(\rho))$ .

Consider the coinvariants functors

$$\mathcal{K}: \mathcal{M}(H(F)/(H(F) \cap G^1)) \rightarrow \mathcal{M}(H(F)/(H(F) \cap G^0))$$

and

$$\mathcal{C}: \mathcal{M}(H(F)/(H(F) \cap G^0)) \rightarrow \text{Vect},$$

defined by  $\mathcal{K}(\rho) := \rho_{(H(F) \cap G^0)/(H(F) \cap G^1)}$  and  $\mathcal{C}(\rho) := \rho_{H(F)/(H(F) \cap G^1)}$ . Note that  $\mathcal{E} = \mathcal{C} \circ \mathcal{K}$ .

$$\begin{array}{ccccccc} & & \mathcal{F} & & & & \\ & \nearrow & & \searrow & & \nearrow & \\ & & \mathcal{E} & & & & \\ \mathcal{M}(G(F)) & \xrightarrow{\mathcal{G}} & \mathcal{M}(H(F)/(H(F) \cap G^1)) & \xrightarrow{\mathcal{K}} & \mathcal{M}(H(F)/(H(F) \cap G^0)) & \xrightarrow{\mathcal{C}} & \text{Vect} \end{array}$$

Note that  $\mathcal{C}$  is exact since the group  $H(F)/(H(F) \cap G^1)$  is finite. Hence by Lemma 2.27 (iii),  $L^i \mathcal{E} = \mathcal{C} \circ L^i \mathcal{K}$ .

Now, by Lemma 2.30,

$$\begin{aligned} H_i(H(F), \rho) &= L^i \mathcal{F}(\rho) = L^i \mathcal{E}(\mathcal{G}(\rho)) = \mathcal{C}(L^i \mathcal{K}(\mathcal{G}(\rho))) \\ &= \mathcal{C}(H_i((H(F) \cap G_0)/(H(F) \cap G^1), \mathcal{G}(\rho))). \end{aligned}$$

Hence, by Lemma 2.29, if  $H_0(H(F), \rho)$  is finite dimensional then  $H_1(H(F), \rho)$  is finite dimensional.  $\square$

### 3. Uniform spherical pairs

In this section we introduce the notion of uniform spherical pair and prove Theorem B.

We follow the main steps of [Kaz86], where the author constructs an isomorphism between the Hecke algebras of a reductive group over close enough local fields. First, he constructs a linear isomorphism between the Hecke algebras, using Cartan decomposition. Then, he shows that for two given elements of the Hecke algebra there exists  $m$  such that if the fields are  $m$ -close then the product of those elements will be mapped to the product of their images. Then he uses the fact that the Hecke algebras are finitely generated and finitely presented to deduce the theorem.

Roughly speaking, we call a pair  $H < G$  of reductive groups a uniform spherical pair if it possesses a relative analog of Cartan decomposition, i.e. a “nice” description of the set of double cosets  $K_0(G, F) \backslash G(F)/H(F)$  which in some sense does not depend on  $F$ . We give the precise definition in the first subsection and prove Theorem B in the second subsection.

**3.1. Definitions.** Let  $R$  be a complete and smooth local ring, let  $\mathfrak{m}$  denote its maximal ideal, and let  $\pi$  be an element in  $\mathfrak{m} \setminus \mathfrak{m}^2$ . A good example to keep in mind is the ring  $\mathbb{Z}_p[[\pi]]$ . An  $(R, \pi)$ -local field is a local field  $F$  together with an epimorphism of rings  $R \rightarrow \mathcal{O}_F$ , such that the image of  $\pi$  (which we will continue to denote by  $\pi$ ) is a uniformizer. Denote the collection of all  $(R, \pi)$ -local fields by  $\mathcal{F}_{R, \pi}$ .

Suppose that  $G$  is a reductive group defined and split over  $R$ . Let  $T$  be a fixed split torus, and let  $X_*(T)$  be the coweight lattice of  $T$ . For every  $\lambda \in X_*(T)$  and every  $(R, \pi)$ -local field  $F$ , we write  $\pi^\lambda = \lambda(\pi) \in T(F) \subset G(F)$ . We denote the subgroup  $G(\mathcal{O}_F)$  by  $K_0(F)$ , and denote its  $\ell$ 'th congruence subgroup by  $K_\ell(F)$ .

**Definition 3.1.** Let  $F$  be a local field. Let  $X \subset \mathbb{A}_{\mathcal{O}_F}^n$  be a closed subscheme. For any  $x, y \in X(F)$ , define the valuative distance between  $x$  and  $y$  to be  $\text{val}_F(x, y) := \min\{\text{val}_F(x_i - y_i)\}$ . Also, for any  $x \in X(F)$ , define  $\text{val}_F(x) := \min\{\text{val}_F(x_i)\}$ . The ball of valuative radius  $\ell$  around a point  $x$  in  $X(F)$  will be denoted by  $B(x, \ell)(F)$ .

**Definition 3.2.** Let  $G$  be a split reductive group defined over  $R$  and let  $H \subset G$  be a smooth reductive subgroup defined over  $R$ . We say that the pair  $(G, H)$  is uniform spherical if there are



- an  $R$ -split torus  $T \subset G$ ,
- an affine embedding  $G/H \hookrightarrow \mathbb{A}^n$ .
- a finite subset  $\mathfrak{X} \subset G(R)/H(R)$ .
- a subset  $\Upsilon \subset X_*(T)$ ,

such that

- (1) the map  $x \mapsto K_0(F)x$  from  $\pi^\Upsilon \mathfrak{X}$  to  $K_0(F) \backslash G(F)/H(F)$  is onto for every  $F \in \mathcal{F}_{R,\pi}$ ,
- (2) for every  $x, y \in \pi^\Upsilon \mathfrak{X} \subset (G/H)(R[\pi^{-1}])$ , the closure in  $G$  of the  $R[\pi^{-1}]$ -scheme

$$T_{x,y} := \{g \in G \times_{\text{Spec}(R)} \text{Spec}(R[\pi^{-1}]) \mid gx = y\}$$

is smooth over  $R$ ; we denote this closure by  $S_{x,y}$ ,

- (3) for every  $x \in \pi^\Upsilon \mathfrak{X}$ , the valuation  $\text{val}_F(x)$  does not depend on  $F \in \mathcal{F}_{R,\pi}$ ,
- (4) there exists  $l_0$  s.t. for any  $l > l_0$ , for any  $F \in \mathcal{F}_{R,\pi}$  and for every  $x \in \mathfrak{X}$  and  $\alpha \in \Upsilon$  we have  $K_l \pi^\alpha K_l x = K_l \pi^\alpha x$ .

If  $G, H$  are defined over  $\mathbb{Z}$ , we say that the pair  $(G, H)$  is uniform spherical if, for every  $R$  as above, the pair  $(G \times_{\text{Spec}(\mathbb{Z})} \text{Spec}(R), H \times_{\text{Spec}(\mathbb{Z})} \text{Spec}(R))$  is uniform spherical.

In Section 4 we give two examples of uniform spherical pairs. We will list now several basic properties of uniform spherical pairs. In light of the recent developments in the structure theory of symmetric and spherical pairs (e.g. [Del], [SV]), we believe that the majority of symmetric pairs and many spherical pairs defined over local fields are specializations of appropriate uniform spherical pairs.

From now and until the end of the section we fix a uniform spherical pair  $(G, H)$ . First note that, since  $H$  is smooth, the fibers of  $G \rightarrow G/H$  are smooth. Hence the map  $G \rightarrow G/H$  is smooth.

**Lemma 3.3.** *Let  $(G, H)$  be a uniform spherical pair. Let  $x, y \in \pi^\Upsilon \mathfrak{X}$ . Let  $F$  be an  $(R, \pi)$ -local field. Then*

$$S_{x,y}(O_F) = T_{x,y}(F) \cap G(O).$$

*Proof.* The inclusion  $S_{x,y}(O_F) \subset T_{x,y}(F) \cap G(O_F)$  is evident. In order to prove the other inclusion we have to show that any map  $\psi: \text{Spec}(O_F) \rightarrow G \times_{\text{Spec } R} \text{Spec } O_F$  such that  $\text{Im}(\psi|_{\text{Spec } F}) \subset T_{x,y} \times_{\text{Spec } R[\pi^{-1}]} \text{Spec } F$  satisfies  $\text{Im } \psi \subset S_{x,y} \times_{\text{Spec } R} \text{Spec } O_F$ .

This holds since  $S_{x,y} \times_{\text{Spec } R} \text{Spec } O_F$  lies in the closure of  $T_{x,y} \times_{\text{Spec } R[\pi^{-1}]} \text{Spec } F$  in  $G \times_{\text{Spec } R} \text{Spec } O_F$ .  $\square$

**Lemma 3.4.** *If  $(G, H)$  is uniform spherical, then there is a subset  $\Delta \subset \pi^\Upsilon \mathcal{X}$  such that, for every  $F \in \mathcal{F}_{R,\pi}$ , the map  $x \mapsto K_0(F)x$  is a bijection between  $\Delta$  and  $K_0(F) \backslash G(F)/H(F)$ .*

*Proof.* It is enough to show that for any  $F, F' \in \mathcal{F}_{R,\pi}$  and for any  $x, y \in \pi^\Upsilon \mathcal{X}$ , the equality  $K_0(F)x = K_0(F)y$  is equivalent to  $K_0(F')x = K_0(F')y$ .

The scheme  $S_{x,y} \otimes O_F$  is smooth over  $R$ , and hence it is smooth over  $O_F$ . Therefore, it is formally smooth. This implies that the map  $S_{x,y}(O_F) \rightarrow S_{x,y}(\mathbb{F}_q)$  is onto and hence  $\{g \in G(O_F) \mid gx = y\} \neq \emptyset$  if and only if  $S_{x,y}(\mathbb{F}_q) \neq \emptyset$ .

Hence, the two equalities  $K_0(F)x = K_0(F)y$  and  $K_0(F')x = K_0(F')y$  are equivalent to  $S_{x,y}(\mathbb{F}_q) \neq \emptyset$ .  $\square$

From now until the end of the section we fix  $\Delta$  as in the lemma.

**Proposition 3.5.** *If  $(G, H)$  is uniform spherical, then for every  $x \in \pi^\Upsilon \mathcal{X}$  and every  $\ell \in \mathbb{N}$ , there is  $M \in \mathbb{N}$  such that for every  $F \in \mathcal{F}_{R,\pi}$ , the set  $K_\ell(F)x$  contains a ball of radius  $M$  around  $x$ .*

*Proof.* Since, for every  $\delta \in X_*(T)$  and every  $\ell \in \mathbb{N}$ , there is  $n \in \mathbb{N}$  such that  $K_n(F) \subset \pi^\delta K_\ell(F) \pi^{-\delta}$  for every  $F$ , we can assume that  $x \in \mathcal{X}$ . The claim now follows from the following version of the implicit function theorem.

**Lemma 3.6.** *Let  $F$  be a local field. Let  $X$  and  $Y$  be affine schemes defined over  $O_F$ . Let  $\psi: X \rightarrow Y$  be a smooth morphism defined over  $O_F$ . Let  $x \in X(O_F)$  and  $y := \psi(x)$ . Then  $\psi(B(x, \ell)(F)) = B(y, \ell)(F)$  for any natural number  $\ell$ .*

*Proof.* The inclusion  $\psi(B(x, \ell)(F)) \subset B(y, \ell)(F)$  is clear. We prove the inclusion  $\psi(B(x, \ell)(F)) \supset B(y, \ell)(F)$ .

Case 1:  $X$  and  $Y$  are affine spaces and  $\psi$  is étale. The proof is standard.

Case 2:  $X = \mathbb{A}^m$ ,  $\psi$  is étale: We can assume that  $Y \subset \mathbb{A}^{m+n}$  is defined by  $f_1, \dots, f_n$  with independent differentials, and that  $\psi$  is the projection. The proof in this case follows from Case 1 by considering the map  $F: \mathbb{A}^{m+n} \rightarrow \mathbb{A}^{m+n}$  given by  $F(x_1, \dots, x_{m+n}) = (x_1, \dots, x_m, f_1, \dots, f_n)$ .

Case 3:  $\psi$  is étale: Follows from Case 2 by restriction from the ambient affine spaces.

Case 4: In general, a smooth morphism is a composition of an étale morphism and a projection, for which the claim is trivial.  $\square$

$\square$

**Lemma 3.7.** *For every  $\lambda \in X_*(T)$  and  $x \in \pi^\Upsilon \mathcal{X}$ , there is a finite subset  $B \subset \pi^\Upsilon \mathcal{X}$  such that  $\pi^\lambda K_0(F)x \subset \bigcup_{y \in B} K_0(F)y$  for all  $F \in \mathcal{F}_{R,\pi}$ .*

*Proof.* By Lemma 3.4, we can assume that the sets  $K_0(F)\pi^\lambda x_0$  for  $\lambda \in \Upsilon$  are disjoint. There is a constant  $C$  such that for every  $F$  and for every  $g \in \pi^\lambda K_0(F)\pi^\delta$ ,  $\text{val}_F(gx_0) \geq C$ . Fix  $F$  and assume that  $g \in K_0(F)\pi^\lambda K_0(F)\pi^\delta$ . From the proof of Proposition 3.5, it follows that  $K_0(F)gx_0$  contains a ball whose radius depends only on  $\lambda, \delta$ . Since  $F$  is locally compact, there are only finitely many disjoint such balls in the set  $\{x \in G(F)/H(F) \mid \text{val}_F(x) \geq C\}$ , so there are only finitely many  $\eta \in \Upsilon$  such that  $\text{val}_F(\pi^\eta x_0) \geq C$ . By definition, this finite set,  $S$ , does not depend on the field  $F$ . Therefore,  $\pi^\lambda K_0(F)\pi^\delta x_0 \subset \bigcup_{\eta \in S} K_0(F)\pi^\eta x_0$ .  $\square$

**Notation 3.8.** • Denote by  $\mathcal{M}_\ell(G(F)/H(F))$  the space of  $K_\ell(F)$ -invariant compactly supported measures on  $G(F)/H(F)$ .

• For a  $K_l$  invariant subset  $U \subset G(F)/H(F)$  we denote by  $1_U \in \mathcal{M}_\ell(G(F)/H(F))$  the Haar measure on  $G(F)/H(F)$  multiplied by the characteristic function of  $U$  and normalized s.t. its integral is 1. We define in a similar way  $1_V \in \mathcal{H}_\ell(G, F)$  for a  $K_l$ -double invariant subset  $V \subset G(F)$ .

**Proposition 3.9.** *If  $(G, H)$  is uniform spherical then  $\mathcal{M}_\ell(G(F)/H(F))$  is finitely generated over  $\mathcal{H}_\ell(G, F)$  for any  $\ell$ .*

*Proof.* As in step 4 of Lemma 2.10, it is enough to prove the assertion for large enough  $l$ . Thus we may assume that for every  $x \in \mathfrak{X}$  and  $\alpha \in \Upsilon$  we have  $K_l \pi^\alpha K_l x = K_l \pi^\alpha x$ . Therefore,  $1_{K_l \pi^\alpha K_l} 1_{K_l x} = 1_{K_l \pi^\alpha x}$ . Hence for any  $g \in K_0/K_l$  we have  $(g 1_{K_l \pi^\alpha K_l}) 1_{K_l x} = 1_{g K_l \pi^\alpha x}$ . Now, the elements  $1_{g K_l \pi^\alpha x}$  span  $\mathcal{M}_\ell(G(F)/H(F))$  by condition 1 in definition 3.2. This implies that the elements  $1_{K_l x}$  generate  $\mathcal{M}_\ell(G(F)/H(F))$ .  $\square$

### 3.2. Close local fields

**Definition 3.10.** Two  $(R, \pi)$ -local fields  $F, E \in \mathcal{F}_{R, \pi}$ , are  $n$ -close if there is an isomorphism  $\phi_{E, F}: O_F/\pi^n \rightarrow O_E/\pi^n$  such that the two maps  $R \rightarrow O_F \rightarrow O_F/\pi^n \rightarrow O_E/\pi^n$  and  $R \rightarrow O_E \rightarrow O_E/\pi^n$  coincide. In this case,  $\phi$  is unique.

**Theorem 3.11** ([Kaz86]). *Let  $F$  be an  $(R, \pi)$  local field. Then, for any  $\ell$ , there exists  $n$  such that, for any  $E \in \mathcal{F}_{R, \pi}$ , which is  $n$ -close to  $F$ , there exists a unique isomorphism  $\Phi_{\mathcal{H}, \ell}$  between the algebras  $\mathcal{H}_\ell(G, F)$  and  $\mathcal{H}_\ell(G, E)$  that maps the Haar measure on  $K_\ell(F)\pi^\lambda K_\ell(F)$  to the Haar measure on  $K_\ell(E)\pi^\lambda K_\ell(E)$ , for every  $\lambda \in X_*(T)$ , and intertwines the actions of the finite group  $K_0(F)/K_\ell(F) \xrightarrow{\phi_{F, E}} K_0(E)/K_\ell(E)$ .*

In this section we prove the following refinement of Theorem B from the Introduction:

**Theorem 3.12.** *Let  $(G, H)$  be a uniform spherical pair. Then, for any  $\ell \in \mathbb{N}$  and  $F \in \mathcal{F}_{R,\pi}$ , there exists  $n$  such that, for any  $E \in \mathcal{F}_{R,\pi}$  that is  $n$ -close to  $F$ , there exists a unique map*

$$\mathcal{M}_\ell(G(F)/H(F)) \rightarrow \mathcal{M}_\ell(G(E)/H(E))$$

*which is an isomorphism of modules over the Hecke algebra*

$$\mathcal{H}(G(F), K_\ell(F)) \stackrel{\Phi_{\mathcal{H},\ell}}{\cong} \mathcal{H}(G(E), K_\ell(E))$$

*that maps the Haar measure on  $K_\ell(F)x$  to the Haar measure on  $K_\ell(E)x$ , for every  $x \in \Delta \subset \pi^\lambda \Upsilon$ , and intertwines the actions of the finite group  $K_0(F)/K_\ell(F) \stackrel{\phi_{F,E}}{\cong} K_0(E)/K_\ell(E)$ .*

For the proof we will need notation and several lemmas.

**Notation 3.13.** For any valued field  $F$  with uniformizer  $\pi$  and any integer  $m \in \mathbb{Z}$ , we denote by  $\text{res}_m: F \rightarrow F/\pi^m O$  the projection. Note that the groups  $\pi^n O$  are naturally isomorphic for all  $n$ . Hence if two local fields  $F, E \in \mathcal{F}_{R,\pi}$  are  $n$ -close, then for any  $m$  we are given an isomorphism, which we also denote by  $\phi_{F,E}$  between  $\pi^{m-n} O_F/\pi^m O_F$  and  $\pi^{m-n} O_E/\pi^m O_E$ , which are subgroups of  $F/\pi^m O_F$  and  $E/\pi^m O_E$ .

**Lemma 3.14.** *Suppose that  $(G, H)$  is a uniform spherical pair, and suppose that  $F, E \in \mathcal{F}_{R,\pi}$  are  $\ell$ -close. Then for all  $\delta \in \Delta$ ,*

$$\phi_{F,E}(\text{Stab}_{K_0(F)/K_\ell(F)} K_\ell(F)\delta) = \text{Stab}_{K_0(E)/K_\ell(E)} K_\ell(E)\delta.$$

*Proof.* The stabilizer of  $K_\ell(F)\delta$  in  $K_0/K_\ell$  is the projection of the stabilizer of  $\delta$  in  $K_0$  to  $K_0/K_\ell$ . In other words, it is the image of  $S_{\delta,\delta}(O_F)$  in  $S_{\delta,\delta}(O_F/\pi^\ell)$ . Since  $S_{\delta,\delta}$  is smooth over  $R$ , it is smooth over  $O_F$ . Hence  $S_{\delta,\delta}$  is formally smooth, and so this map is onto. The same applies to the stabilizer of  $K_\ell(E)\delta$  in  $K_0(E)/K_\ell(E)$ , but  $\phi_{F,E}(S_{\delta,\delta}(O/\pi^\ell)) = S_{\delta,\delta}(O'/\pi'^\ell)$ .  $\square$

**Corollary 3.15.** *Let  $\ell \in \mathbb{N}$ . Then, for any  $F, E \in \mathcal{F}_{R,\pi}$  that are  $\ell$ -close, there exists a unique morphism of vector spaces*

$$\Phi_{\mathcal{M},\ell}: \mathcal{M}_\ell(G(F)/H(F)) \rightarrow \mathcal{M}_\ell(G(E)/H(E))$$

*that maps the Haar measure on  $K_\ell(F)x$  to the Haar measure on  $K_\ell(E)x$ , for every  $x \in \Delta$ , and intertwines the actions of the finite group  $K_0(F)/K_\ell(F) \stackrel{\phi_{F,E}}{\cong} K_0(E)/K_\ell(E)$ . Moreover, this morphism is an isomorphism.*

*Proof.* The uniqueness is evident. By Lemma 3.14 and Lemma 3.4, the map between  $K_\ell(F) \backslash G(F) / H(F)$  and  $K_\ell(E) \backslash G(E) / H(E)$  given by

$$K_\ell(F)g\delta \mapsto K_\ell(E)g'\delta,$$

where  $g \in K_0(F)$  and  $g' \in K_0(E)$  satisfy that  $\phi_{F,E}(\text{res}_\ell(g)) = \text{res}_\ell(g')$ , is a bijection. This bijection gives the required isomorphism.  $\square$

**Remark 3.16.** A similar construction can be applied to the pair  $(G \times G, \Delta G)$ . In this case, the main result of [Kaz86] is that the obtained linear map  $\Phi_{\mathcal{H},\ell}$  between the Hecke algebras  $\mathcal{H}(G(F), K_\ell(F))$  and  $\mathcal{H}(G(E), K_\ell(E))$  is an isomorphism of algebras if the fields  $F$  and  $E$  are close enough.

The following lemma is evident:

**Lemma 3.17.** *Let  $P(x) \in R[\pi^{-1}][x_1, \dots, x_d]$  be a polynomial. For any natural numbers  $M$  and  $k$ , there is  $N$  such that, if  $F, E \in \mathcal{F}_{R,\pi}$  are  $N$ -close, and  $x_0 \in \pi^{-k}O_F^d, y_0 \in \pi^{-k}O_E^d$  satisfy that  $P(x_0) \in \pi^{-k}O_F$  and  $\phi_{F,E}(\text{res}_N(x_0)) = \text{res}_N(y_0)$ , then  $P(y_0) \in \pi^{-k}O_E$  and  $\phi_{F,E}(\text{res}_M(P(x_0))) = \text{res}_M(P(y_0))$ .*

**Corollary 3.18.** *Suppose that  $(G, H)$  is a uniform spherical pair. Fix an embedding of  $G/H$  to an affine space  $\mathbb{A}^d$ . Let  $\lambda \in X_*(T)$ ,  $x \in \pi^\gamma \mathcal{X}$ ,  $F \in \mathcal{F}_{R,\pi}$ , and  $k \in G(O_F)$ . Choose  $m$  such that  $\pi^\lambda kx \in \pi^{-m}O_F^d$ . Then, for every  $M$ , there is  $N \geq M + m$  such that, for any  $E \in \mathcal{F}_{R,\pi}$  that is  $N$ -close to  $F$ , and for any  $k' \in G(O_E)$  such that  $\phi_{F,E}(\text{res}_N(k)) = \text{res}_N(k')$ ,*

$$\pi^\lambda k'x \in G(E)/H(E) \cap \pi^{-m}O_E^d \quad \text{and} \quad \phi_{F,E}(\text{res}_M(\pi^\lambda kx)) = \text{res}_M(\pi^\lambda k'x).$$

**Corollary 3.19.** *Suppose that  $(G, H)$  is a uniform spherical pair. Fix an embedding of  $G/H$  to an affine space  $\mathbb{A}^d$ . Let  $m$  be an integer. For every  $M$ , there is  $N$  such that, for any  $F, E \in \mathcal{F}_{R,\pi}$  that are  $N$ -close, any  $x \in G(F)/H(F) \cap \pi^{-m}O_F^d$  and any  $y \in G(E)/H(E) \cap \pi^{-m}O_E^d$ , such that  $\phi_{F,E}(\text{res}_{N-m}(x)) = \text{res}_{N-m}(y)$ , we have  $\Phi_{\mathcal{M}}(1_{K_M(F)}x) = 1_{K_M(E)}y$ .*

*Proof.* Let  $k_F \in G(O_F)$  and  $\delta \in \Delta$  such that  $x = k_F\delta$ . By Proposition 3.5, there is an  $l$  such that, for any  $L \in \mathcal{F}_{R,\pi}$  and any  $k_L \in G(O_L)$ , we have  $K_M(L)k_L\delta$  contains a ball of radius  $l$ .

Using the previous corollary, choose an integer  $N$  such that, for any  $F$  and  $E$  that are  $N$ -close and any  $k_E \in G(O_E)$ , such that  $\phi_{F,E}(\text{res}_N(k_F)) = \text{res}_N(k_E)$ , we have

$$k_E\delta \in (G(E)/H(E)) \cap \pi^{-m}O_E^d \quad \text{and} \quad \phi_{F,E}(\text{res}_l(x)) = \text{res}_l(k_E\delta).$$

Choose such  $k_E \in G(O_E)$  and let  $z = k_E \delta$ . Since  $\text{res}_l(z) = \phi_{F,E}(\text{res}_l(x)) = \text{res}_l(y)$ , we have that  $z \in B(y, l)$ , and hence  $z \in K_M(E)y$ . Hence

$$1_{K_M(E)}y = 1_{K_M(E)}z = \Phi_{\mathcal{M}}(1_{K_M(F)}x). \quad \square$$

From the last two corollaries we obtain the following one.

**Corollary 3.20.** *Given  $\ell \in \mathbb{N}$ ,  $\lambda \in X_*(T)$ , and  $\delta \in \Delta$ , there is  $n$  such that if  $F, E \in \mathcal{F}_{R,\pi}$  are  $n$ -close, and  $g_F \in G(O_F)$ ,  $g_E \in G(O_E)$  satisfy that  $\phi_{F,E}(\text{res}_n(g_F)) = \text{res}_n(g_E)$ , then  $\Phi_{\mathcal{M},\ell}(1_{K_\ell(F)}\pi^\lambda g_F \delta) = 1_{K_\ell(E)}\pi^\lambda g_E \delta$ .*

**Proposition 3.21.** *Let  $F \in \mathcal{F}_{R,\pi}$ . Then for every  $\ell$ , and every two elements  $f \in \mathcal{H}_\ell(F)$  and  $v \in \mathcal{M}_\ell(F)$ , there is  $n$  such that, if  $E \in \mathcal{F}_{R,\pi}$  is  $n$ -close to  $F$ , then  $\Phi_{\mathcal{M},\ell}(f \cdot v) = \Phi_{\mathcal{H},\ell}(f) \cdot \Phi_{\mathcal{M},\ell}(v)$ .*

*Proof.* By linearity, we can assume that  $f = 1_{K_\ell(F)}k_1\pi^\lambda k_2 1_{K_\ell(F)}$  and that  $v = 1_{K_\ell(F)}k_3\delta$ , where  $k_1, k_2, k_3 \in K_0(F)$ . Choose  $N \geq l$  big enough such that  $\pi^\lambda K_N(F)\pi^{-\lambda} \subset K_\ell(F)$ .

Choose  $k'_i \in G(O_E)$  such that  $\phi_{F,E}(\text{res}_N(k_i)) = \text{res}_N(k'_i)$ . Since  $\Phi_{\mathcal{M},\ell}$  and  $\Phi_{\mathcal{H},\ell}$  intertwine left multiplication by  $1_{K_\ell(F)}k_1 1_{K_\ell(F)}$  to left multiplication by  $1_{K_\ell(E)}k'_1 1_{K_\ell(E)}$ , we can assume that  $k_1 = 1 = k'_1$ . Also, since  $k_2$  normalizes  $K_\ell(F)$ , we can assume that  $k_2 = 1 = k'_2$ . Let  $K_\ell(F) = \bigcup_{i=1}^s K_N(F)g_i$  be a decomposition of  $K_\ell(F)$  into cosets. Choose  $g'_i \in K_\ell(E)$  such that  $\phi_{F,E}(\text{res}_N(g_i)) = \text{res}_N(g'_i)$ . Then

$$1_{K_\ell(F)} = c \sum_{i=1}^s 1_{K_N(F)}g_i \quad \text{and} \quad 1_{K_\ell(E)} = c \sum_{i=1}^s 1_{K_N(E)}g'_i$$

where  $c = |K_\ell(F)/K_N(F)| = |K_\ell(E)/K_N(E)|$ . Hence

$$\begin{aligned} f v &= 1_{K_\ell(F)}\pi^\lambda 1_{K_\ell(F)}k_3\delta = c \sum_{i=1}^s 1_{K_\ell(F)}\pi^\lambda 1_{K_N(F)}g_i k_3\delta \\ &= c \sum_{i=1}^s 1_{K_\ell(F)}\pi^\lambda g_i k_3\delta. \end{aligned}$$

and

$$\begin{aligned} \Phi_{\mathcal{H},\ell}(f)\Phi_{\mathcal{M},\ell}(v) &= 1_{K_\ell(E)}\pi^\lambda 1_{K_\ell(E)}k'_3\delta \\ &= c \sum_{i=1}^s 1_{K_\ell(E)}\pi^\lambda 1_{K_N(E)}g_i k'_3\delta \\ &= c \sum_{i=1}^s 1_{K_\ell(E)}\pi^\lambda g'_i k'_3\delta. \end{aligned}$$

The proposition follows now from Corollary 3.20.  $\square$

Now we are ready to prove Theorem 3.12.

*Proof of Theorem 3.12.* We have to show for any  $\ell$  there exists  $n$  such that if  $F, E \in \mathcal{F}_{R,\pi}$  are  $n$ -close then the map  $\Phi_{\mathcal{M},\ell}$  constructed in Corollary 3.15 is an isomorphism of modules over  $\mathcal{H}(G(F), K_\ell(F)) \xrightarrow{\Phi_{\mathcal{H},\ell}} \mathcal{H}(G(E), K_\ell(E))$ .

Since  $\mathcal{H}(G(F), K_\ell(F))$  is Noetherian,  $\mathcal{M}_\ell(G(F)/H(F))$  is generated by a finite set  $v_1, \dots, v_n$  satisfying a finite set of relations  $\sum_i f_{i,j} v_i = 0$ . Without loss of generality we may assume that for any  $x \in \mathfrak{X}$  the Haar measure on  $K_\ell(F)x$  is contained in the set  $\{v_i\}$ .

By Proposition 3.21, if  $E$  is close enough to  $F$ , then  $\Phi_{\mathcal{M},\ell}(v_i)$  satisfy the above relations.

Therefore there exists a homomorphism of Hecke modules

$$\Phi': \mathcal{M}_\ell(G(F)/H(F)) \rightarrow \mathcal{M}_\ell(G(E)/H(E))$$

given on the generators  $v_i$  by  $\Phi'(v_i) := \Phi_{\mathcal{M},\ell}(v_i)$ .

$\Phi'$  intertwines the actions of the finite group  $K_0(F)/K_\ell(F) \xrightarrow{\phi_{F,E}} K_0(E)/K_\ell(E)$ . Therefore, by Corollary 3.15, in order to show that  $\Phi'$  coincides with  $\Phi_{\mathcal{M},\ell}$  it is enough to check that  $\Phi'$  maps the normalized Haar measure on  $K_\ell(F)x$  to the normalized Haar measure on  $K_\ell(E)x$  for every  $x \in \Delta$ . In order to do this let us decompose  $x = \pi^\alpha x_0$  where  $x_0 \in \mathfrak{X}$  and  $\alpha \in \Upsilon$ . Now, since  $(G, H)$  is uniformly spherical we have

$$1_{K_n(F)x} = 1_{K_n(F)\pi^\alpha K_n(F)} 1_{K_n(F)x_0}$$

and

$$1_{K_n(E)x} = 1_{K_n(E)\pi^\alpha K_n(E)} 1_{K_n(E)x_0}.$$

Therefore, since  $\Phi'$  is a homomorphism, we have

$$\begin{aligned} \Phi'(1_{K_n(F)x}) &= \Phi'(1_{K_n(F)\pi^\alpha K_n(F)} 1_{K_n(F)x_0}) \\ &= 1_{K_n(E)\pi^\alpha K_n(E)} 1_{K_n(E)x_0} = 1_{K_n(E)x}. \end{aligned}$$

Hence the linear map  $\Phi_{\mathcal{M},\ell}: \mathcal{M}_\ell(G(F)/H(F)) \rightarrow \mathcal{M}_\ell(G(E)/H(E))$  is a homomorphism of Hecke modules. Since it is a linear isomorphism, it is an isomorphism of Hecke modules.  $\square$

Now we obtain the following generalization of Corollary C:

**Corollary 3.22.** *Let  $(G, H)$  be a uniform spherical pair. Suppose that*

- *For any  $F \in \mathcal{F}_{R,\pi}$ , the pair  $(G, H)$  is  $F$ -spherical.*
- *For any  $E \in \mathcal{F}_{R,\pi}$  and natural number  $n$ , there is a field  $F \in \mathcal{F}_{R,\pi}$  such that  $E$  and  $F$  are  $n$ -close and the pair  $(G(F), H(F))$  is a Gelfand pair, i.e. for any irreducible smooth representation  $\rho$  of  $G(F)$  we have*

$$\dim \operatorname{Hom}_{H(F)}(\rho|_{H(F)}, \mathbb{C}) \leq 1.$$

*Then  $(G(F), H(F))$  is a Gelfand pair for any  $F \in \mathcal{F}_{R,\pi}$ .*

**Remark 3.23.** Fix a prime power  $q = p^k$ . Let  $F$  be the unramified extension of  $\mathbb{Q}_p$  of degree  $k$ , let  $W$  be the ring of integers of  $F$ , and let  $R = W[[\pi]]$ . Then  $\mathcal{F}_{R,\pi}$  includes all local fields with residue field  $\mathbb{F}_q$ , and so Corollary 3.22 implies Corollary C.

Corollary 3.22 follows from Theorem 3.12 and the following lemma.

**Lemma 3.24.** *Let  $F$  be a local field and  $H < G$  be a pair of reductive groups defined over  $F$ . Suppose that  $G$  is split over  $F$ . Then  $(G(F), H(F))$  is a Gelfand pair if and only if for any large enough  $l \in \mathbb{Z}_{>0}$  and any simple module  $\rho$  over  $\mathcal{H}_l(G(F))$  we have*

$$\dim \operatorname{Hom}_{\mathcal{H}_l(G(F))}(\mathcal{M}_l(G(F)/H(F)), \rho) \leq 1.$$

This lemma follows from statement (1) formulated in Subsection 2.1.

## 4. Applications

In this section we will prove that the pair  $(\operatorname{GL}_{n+k}(F), \operatorname{GL}_n(F) \times \operatorname{GL}_k(F))$  is a Gelfand pair for any local field  $F$  of characteristic different from 2 and the pair  $(\operatorname{GL}_{n+1}(F), \operatorname{GL}_n(F))$  is a strong Gelfand pair for any local field  $F$ . We use Corollary 3.22 to deduce those results from the characteristic zero case which were proven in [JR96] and [AGRS] respectively. Let  $R = W[[\pi]]$ .

To verify condition (2) in Definition 3.2, we use the following straightforward lemma:

**Lemma 4.1.** *Let  $G = (\operatorname{GL}_{n_1})_R \times \cdots \times (\operatorname{GL}_{n_k})_R$  and let  $C < G \otimes_R R[\pi^{-1}]$  be a sub-group scheme defined over  $R[\pi^{-1}]$ . Suppose that  $C$  is defined by equations of the following type:*

$$\sum_{i=1}^l \epsilon_i a_{\mu_i} \pi^{\lambda_i} = \pi^v,$$

or

$$\sum_{i=1}^l \epsilon_i a_{\mu_i} \pi^{\lambda_i} = 0,$$

where  $\epsilon_i = \pm 1$ ,  $a_1, \dots, a_{n_1^2 + \dots + n_k^2}$  are entries of matrices,  $1 \leq \mu_i \leq n_1^2 + \dots + n_k^2$  are some indices, and  $v, \lambda_i$  are integers. Suppose also that the indices  $\mu_i$  are distinct for all the equations. Then the closure  $\bar{C}$  of  $C$  in  $G$  is smooth over  $R$ .

To verify condition (4) in Definition 3.2, we use the following straightforward lemma:



**Lemma 4.2.** *Suppose that there exists a natural number  $\ell_0$  such that, for any  $F \in \mathcal{F}_{R,\pi}$  and any  $\ell > \ell_0$ , there is a subgroup  $P_\ell < K_\ell(G, F)$  satisfying that for every  $x \in \mathcal{X}$*

- (1) *For any  $\alpha \in \Upsilon$  we have  $\pi^\alpha P_\ell \pi^{-\alpha} \subset K_\ell$ .*
- (2)  *$K_\ell x = P_\ell x$ .*

*Then condition (4) in Definition 3.2 is satisfied.*

In our applications, we use the following to show that the pairs we consider are  $F$ -spherical.

**Proposition 4.3.** *Let  $F$  be an infinite field, and consider  $G = \mathrm{GL}_{n_1} \times \cdots \times \mathrm{GL}_{n_k}$  embedded in the standard way in  $M = \mathrm{Mat}_{n_1} \times \cdots \times \mathrm{Mat}_{n_k}$ . Let  $A, B \subset G \otimes F$  be two  $F$ -subgroups whose closures in  $M$  are affine subspaces  $M_A, M_B$ .*

- (1) *For any  $x, y \in G(F)$ , if the variety  $\{(a, b) \in A \times B \mid axb = y\}$  is non-empty, then it has an  $F$ -rational point.*
- (2) *If  $(G, A)$  is a spherical pair, then it is also an  $F$ -spherical pair.*

*Proof.* (i) Denote the projections  $G \rightarrow \mathrm{GL}_{n_j}$  by  $\pi_j$ . Assume that  $x, y \in G(F)$ , and there is a pair  $(\bar{a}, \bar{b}) \in (A \times B)(\bar{F})$  such that  $\bar{a}x\bar{b} = y$ . Let  $L \subset M_A \times M_B$  be the affine subspace  $\{(\alpha, \beta) \mid \alpha x = y\beta\}$ , defined over  $F$ . By assumption, the functions  $(\alpha, \beta) \mapsto \det \pi_j(\alpha)$  and  $(\alpha, \beta) \mapsto \det \pi_j(\beta)$ , for  $j = 1, \dots, k$ , are non-zero on  $L(\bar{F})$ . Hence there is  $(a, b) \in L(F) \cap G$ , which means that  $axb^{-1} = y$ .

(ii) Applying (1) to  $A$  and any parabolic subgroup  $B \subset G$ , any  $(A \times B)(\bar{F})$ -orbit in  $G(\bar{F})$  contains at most one  $(A \times B)(F)$ -orbit. Since there are only finitely many  $(A \times B)(\bar{F})$ -orbits in  $G(\bar{F})$ , the pair  $(G, A)$  is  $F$ -spherical.  $\square$

**4.1. The pair  $(\mathrm{GL}_{n+k}, \mathrm{GL}_n \times \mathrm{GL}_k)$ .** In this subsection we assume  $p \neq 2$  and consider only local fields of characteristic different from 2.

Let  $G := (\mathrm{GL}_{n+k})_R$  and  $H := (\mathrm{GL}_n)_R \times (\mathrm{GL}_k)_R < G$  be the subgroup of block matrices. Note that  $H$  is a symmetric subgroup since it consists of fixed points of conjugation by  $\epsilon = \begin{pmatrix} \mathrm{Id}_k & 0 \\ 0 & -\mathrm{Id}_n \end{pmatrix}$ . We prove that  $(G, H)$  is a Gelfand pair using Corollary C. The pair  $(G, H)$  is a symmetric pair, hence it is a spherical pair and therefore by Proposition 4.3 it is  $F$ -spherical. The second condition of Corollary C is [JR96, Theorem 1.1]. It remains to prove that  $(G, H)$  is a uniform spherical pair.

**Proposition 4.4.** *The pair  $(G, H)$  is uniform spherical.*

*Proof.* Without loss of generality suppose that  $n \geq k$ . Let  $\mathcal{X} = \{x_0\}$ , where

$$x_0 := \begin{pmatrix} \mathrm{Id}_k & 0 & \mathrm{Id}_k \\ 0 & \mathrm{Id}_{n-k} & 0 \\ 0 & 0 & \mathrm{Id}_k \end{pmatrix} H$$

and

$$\Upsilon = \{(\mu_1, \dots, \mu_k, 0, \dots, 0) \in X_*(T) \mid \mu_1 \leq \dots \leq \mu_k \leq 0\}.$$

To show the first condition we show that every double coset in  $K_0 \backslash G/H$  includes an element of the form  $\begin{pmatrix} \text{Id}_k & 0 & \text{diag}(\pi^{\mu_1}, \dots, \pi^{\mu_k}) \\ 0 & \text{Id}_{n-k} & 0 \\ 0 & 0 & \text{Id}_k \end{pmatrix}$  s.t.  $\mu_1 \leq \dots \leq \mu_k \leq 0$ . Take any  $g \in G$ . By left multiplication by  $K_0$  we can bring it to upper triangular form. By right multiplication by  $H$  we can bring it to a form  $\begin{pmatrix} \text{Id}_n & A \\ 0 & \text{Id}_k \end{pmatrix}$ . Conjugating by a matrix  $\begin{pmatrix} k_1 & 0 \\ 0 & k_2 \end{pmatrix} \in K_0 \cap H$  we can replace it by  $\begin{pmatrix} \text{Id}_n & k_1 A k_2^{-1} \\ 0 & \text{Id}_k \end{pmatrix}$ . Hence we can bring  $A$  to be a  $k$ -by- $(n-k)$  block of zero, followed by the a diagonal matrix of the form  $\text{diag}(\pi^{\mu_1}, \dots, \pi^{\mu_k})$  s.t.  $\mu_1 \leq \dots \leq \mu_k$ . Multiplying by an element of  $K_0$  of the form  $\begin{pmatrix} \text{Id}_k & 0 & k \\ 0 & \text{Id}_{n-k} & 0 \\ 0 & 0 & \text{Id}_k \end{pmatrix}$  we can bring  $A$  to the desired form.

As for the second condition, we first compute the stabilizer  $G_{x_0}$  of  $x_0$  in  $G$ . Note that the coset  $x_0 \in G/H$  equals

$$\left\{ \begin{pmatrix} g_1 & g_2 & h \\ g_3 & g_4 & 0 \\ 0 & 0 & h \end{pmatrix} \mid \begin{pmatrix} g_1 & g_2 \\ g_3 & g_4 \end{pmatrix} \in (\text{GL}_n)_R, h \in (\text{GL}_k)_R \right\}$$

and

$$\begin{pmatrix} A & B & C \\ D & E & F \\ G & H & I \end{pmatrix} \begin{pmatrix} \text{Id}_k & 0 & \text{Id}_k \\ 0 & \text{Id}_{n-k} & 0 \\ 0 & 0 & \text{Id}_k \end{pmatrix} = \begin{pmatrix} A & B & A+C \\ D & E & D+F \\ G & H & G+I \end{pmatrix}.$$

Therefore

$$G_{x_0} = \left\{ \begin{pmatrix} g_1 & g_2 & h - g_1 \\ g_3 & g_4 & -g_3 \\ 0 & 0 & h \end{pmatrix} \mid \begin{pmatrix} g_1 & g_2 \\ g_3 & g_4 \end{pmatrix} \in (\text{GL}_n)_R, h \in (\text{GL}_k)_R \right\}.$$

Hence, for any  $\delta_1 = (\lambda_{1,1}, \dots, \lambda_{1,k}, 0, \dots, 0)$ ,  $\delta_2 = (\lambda_{2,1}, \dots, \lambda_{2,k}, 0, \dots, 0) \in \Upsilon$ ,

$$G(F)_{\pi^{\lambda_1} x_0, \pi^{\lambda_2} x_0}$$

$$\begin{aligned} &= \left\{ \begin{pmatrix} \pi^{\lambda_2} g_1 \pi^{-\lambda_1} & \pi^{\lambda_2} g_2 & \pi^{\lambda_2} (h - g_1) \\ g_3 \pi^{-\lambda_1} & g_4 & -g_3 \\ 0 & 0 & h \end{pmatrix} \mid \begin{pmatrix} g_1 & g_2 \\ g_3 & g_4 \end{pmatrix} \in (\text{GL}_n)_R, h \in (\text{GL}_k)_R \right\} \\ &= \left\{ \begin{pmatrix} A & B & C \\ D & E & F \\ 0 & 0 & I \end{pmatrix} \in \text{GL}_{n+k} \mid D = -F \pi^{-\lambda_1}, C = \pi^{\lambda_2} I - A \pi^{\lambda_1} \right\}. \end{aligned}$$

The second condition of Definition 3.2 follows now from Lemma 4.1.

As for the third condition, we use the embedding  $G/H \rightarrow G$  given by  $g \mapsto g\epsilon g^{-1}\epsilon$ . It is easy to see that  $\text{val}_F(\pi^\mu x_0) = \mu_1$ , which is independent of  $F$ .

Let us now prove the last condition using Lemma 4.2. Take  $l_0 = 1$  and

$$P := \left\{ \begin{pmatrix} \text{Id} & 0 & 0 \\ D & E & F \\ G & H & I \end{pmatrix} \in GL_{n+k} \right\}.$$

Let  $P_l := P(F) \cap K_l(GL_{n+k}, F)$ . The first condition of Lemma 4.2 obviously holds. To show the second condition, we have to show that for any  $F$ , any  $l \geq 1$  and any  $g \in K_l(GL_{n+k}, F)$  there exist  $p \in P_l$  and  $h \in H(F)$  such that  $gx_0 = px_0h$ . In other words, we have to solve the following equation:

$$\begin{pmatrix} \text{Id}_k + A & B & \text{Id}_k + A + C \\ D & \text{Id}_{n-k} + E & D + F \\ G & H & \text{Id}_k + G + I \end{pmatrix} = \begin{pmatrix} \text{Id}_k & 0 & \text{Id}_k \\ D' & \text{Id}_{n-k} + E' & D' + F' \\ G' & H' & \text{Id}_k + G' + I' \end{pmatrix} \begin{pmatrix} \text{Id}_k + x & y & 0 \\ z & \text{Id}_k + w & 0 \\ 0 & 0 & \text{Id}_k + h \end{pmatrix},$$

where all the capital letters denote matrices of appropriate sizes with entries in  $\pi^l \mathcal{O}_F$ , and the matrices in the left hand side are parameters and matrices in the right hand side are unknowns.

The solution is given by:

$$\begin{aligned} x &= A, \quad y = B, \quad z = D, \quad w = E, \quad h = A + C \\ D' &= 0, \quad E' = 0, \quad F' = (D + F)(\text{Id}_k + A + C)^{-1}, \\ H' &= (H - G(\text{Id}_k + A)^{-1}B)(-D(\text{Id}_k + A)^{-1}B + \text{Id}_{n-k} + E)^{-1} \\ G' &= (G - H'D)(\text{Id}_k + A)^{-1}, \quad I' = (G + I - A - C)(\text{Id}_k + A + C)^{-1} - G' \end{aligned}$$

□

**4.2. Structure of the spherical space  $(\text{GL}_{n+1} \times \text{GL}_n)/\Delta \text{GL}_n$ .** Consider the embedding  $\iota: \text{GL}_n \hookrightarrow \text{GL}_{n+1}$  given by

$$A \mapsto \begin{pmatrix} 1 & 0 \\ 0 & A \end{pmatrix}.$$

Let  $G = \text{GL}_{n+1}(F) \times \text{GL}_n(F)$  and  $H = \Delta \text{GL}_n(F)$ . The quotient space  $G/H$  is isomorphic to  $(\text{GL}_{n+1})_R$  via the map  $(g, h) \mapsto g\iota(h^{-1})$ . Under this isomorphism, the action of  $G$  on  $G/H$  becomes  $(g, h) \cdot X = gX\iota(h^{-1})$ .

The space  $G/H$  is spherical. Indeed, let  $B \subset G$  be the Borel subgroup consisting of pairs  $(b_1, b_2)$ , where  $b_1$  is lower triangular and  $b_2$  is upper triangular, and let  $x_0 \in G/H$  be the point represented by the matrix

$$x_0 = \begin{pmatrix} 1 & e \\ 0 & I \end{pmatrix},$$

where  $e$  is a row vector of 1's. We claim that  $Bx_0$  is open in  $G/H$ . Let  $\mathfrak{b}$  be the Lie algebra of  $B$ . It consists of pairs  $(X, Y)$  where  $X$  is lower triangular and  $Y$  is upper triangular. The infinitesimal action of  $\mathfrak{b}$  on  $X$  at  $x_0$  is given by  $(X, Y) \mapsto Xx_0 - x_0 d\iota(Y)$ . To show that the image is  $\text{Mat}_{n+1}$ , it is enough to show that the images of the maps  $X \mapsto Xx_0$  and  $Y \mapsto x_0 d\iota(Y)$  have trivial intersection. Suppose that  $Xx_0 = x_0 d\iota(Y)$ . Then  $X = x_0 d\iota(Y)x_0^{-1}$ , i.e.

$$X = \begin{pmatrix} 1 & e \\ 0 & I \end{pmatrix} \begin{pmatrix} 0 & 0 \\ 0 & Y \end{pmatrix} \begin{pmatrix} 1 & -e \\ 0 & I \end{pmatrix} = \begin{pmatrix} 0 & eY \\ 0 & Y \end{pmatrix}.$$

Since  $X$  is lower triangular and  $Y$  is upper triangular, both have to be diagonal. But  $eY = 0$  implies that  $Y = 0$ , and hence also  $X = 0$ . Proposition 4.3 implies that the pair  $(G, H)$  is  $F$ -spherical.

The following describes the quotient  $G(O_F) \backslash G(F)/H(F)$ .

**Lemma 4.5.** *For every matrix  $A \in \text{Mat}_{n+1}(F)$  there are  $k_1 \in \text{GL}_{n+1}(O)$  and  $k_2 \in \text{GL}_n(O)$  such that*

$$k_1 A \iota(k_2) = \begin{pmatrix} \pi^a & \pi^{b_1} & \pi^{b_2} & \dots & \pi^{b_n} \\ & \pi^{c_1} & & & \\ & & \pi^{c_2} & & \\ & & & \ddots & \\ & & & & \pi^{c_n} \end{pmatrix}, \quad (3)$$

where the numbers  $a, b_i, c_i$  satisfy that if  $i < j$  then  $c_i - c_j \leq b_i - b_j \leq 0$  and  $b_1 \leq c_1$ .

*Proof.* Let  $a$  be the minimal valuation of an element in the first column of  $A$ . There is an integral matrix  $w_1$  such that the first column of the matrix  $w_1 A$  is  $\pi^a, 0, 0, \dots, 0$ . Let  $C$  be the  $n \times n$  lower-right sub-matrix of  $w_1 A$ . By Cartan decomposition, there are integral matrices  $w_2, w_3$  such that  $w_2 C w_3^{-1}$  is diagonal, and its diagonal entries are  $\pi^{c_i}$  for a non-decreasing sequence  $c_i$ . Finally, there are integral and diagonal matrices  $d_1, d_2$  such that the matrix  $d_1 \iota(w_2) w_1 A \iota(w_3^{-1}) \iota(d_2^{-1})$  has the form (3).

Suppose that  $i < j$  and  $b_i > b_j$ . Then adding the  $j$ 'th column to the  $i$ 'th column and subtracting  $\pi^{c_j - c_i}$  times the  $i$ 'th row to the  $j$ 'th row, we can change the matrix (3) so that  $b_i = b_j$ . Similarly, if  $i < j$  and  $b_i - b_j < c_i - c_j$ , then adding  $\pi^{b_j - b_i - 1}$

times the  $i$ 'th column to the  $j$ 'th column, and subtracting  $\pi^{c_i+b_j-b_i-1-c_j}$  times the  $j$ 'th row to the  $i$ 'th row changes the matrix (3) so that  $b_i$  becomes smaller in 1. Finally, if  $c_1 < b_1$  then adding the second row to the first changes the matrix so that  $c_1 = b_1$ .  $\square$

Let  $T \subset G$  be the torus consisting of pairs  $(t_1, t_2)$  such that  $t_i$  are diagonal. The co-character group of  $T$  is the group  $\mathbb{Z}^{n+1} \times \mathbb{Z}^n$ . The positive Weyl chamber of  $T$  that is defined by  $B^1$  is the set  $\Delta \subset X_*(T)$  consisting of pairs  $(\mu, \nu)$  such that the  $\mu_i$ 's are non-decreasing and the  $\nu_i$ 's are non-increasing. Lemma 4.5 implies that the set  $\{\pi^\lambda x_0\}_{\lambda \in \Delta}$  is a complete set of orbit representatives for  $G(O) \backslash G(F)/H(F)$ .

We are ready to prove that  $(G, H)$  is uniform spherical.

**Proposition 4.6.** *The pair  $((\mathrm{GL}_{n+1})_R \times (\mathrm{GL}_n)_R, \Delta(\mathrm{GL}_n)_R)$  is uniform spherical.*

*Proof.* Let  $\Upsilon \subset X_*(T)$  be the positive Weyl chamber and let  $\mathfrak{X} := \{x_0\}$ . By the above, the first condition of Definition 3.2 holds. As for the second condition, an easy computation shows that if  $a, b_1, \dots, b_n, c_1, \dots, c_n \in \mathbb{Z}, a', b'_1, \dots, b'_n, c'_1, \dots, c'_n \in \mathbb{Z}$  satisfy the conclusion of Lemma 4.5, and  $(k_1, k_2) \in G(O)$  satisfy that

$$k_1 \begin{pmatrix} \pi^a & \pi^{b_1} & \pi^{b_2} & \dots & \pi^{b_n} \\ & \pi^{c_1} & & & \\ & & \pi^{c_2} & & \\ & & & \ddots & \\ & & & & \pi^{c_n} \end{pmatrix} \iota(k_1) = \begin{pmatrix} \pi^{a'} & \pi^{b'_1} & \pi^{b'_2} & \dots & \pi^{b'_n} \\ & \pi^{c'_1} & & & \\ & & \pi^{c'_2} & & \\ & & & \ddots & \\ & & & & \pi^{c'_n} \end{pmatrix},$$

then  $a = a', c_i = c'_i, k_1$  has the form  $\begin{pmatrix} 1 & B \\ 0 & D \end{pmatrix}$ , where  $B$  is a  $1 \times n$  matrix and  $D$  is an  $n \times n$  matrix that satisfy the equations  $D = \pi^c k_2 \pi^{-c}$  and  $B \pi^c = \pi^b - \pi^{b'} k_2$ , where  $\pi^c$  denotes the diagonal matrix with entries  $\pi^{c_1}, \dots, \pi^{c_n}$ ,  $\pi^b$  denotes the row vector with entries  $\pi^{b_i}$ , and  $\pi^{b'}$  denotes the row vector with entries  $\pi^{b'_i}$ . The second condition of Definition 3.2 holds by Lemma 4.1.

The third condition follows because, using the affine embedding as above,  $\pi^\lambda x_0$  has the form (3) and so  $\mathrm{val}_F(\pi^\lambda x_0)$  is independent of  $F$ .

Finally it is left to verify the last condition. In the following, we will distinguish between the  $\ell$ th congruence subgroup in  $\mathrm{GL}_{n+1}(F)$ , which we denote by  $K_\ell(\mathrm{GL}_{n+1}(F))$ , the  $\ell$ th congruence subgroup in  $\mathrm{GL}_n(F)$ , which we denote by  $K_\ell(\mathrm{GL}_n(F))$ , and the  $\ell$ th congruence subgroup in  $G = \mathrm{GL}_{n+1}(F) \times \mathrm{GL}_n(F)$ , which we denote by  $K_\ell$ . By Lemma 4.2 it is enough to show that  $(B \cap K_\ell)x_0 = K_\ell x_0$ . It is easy to see that  $K_\ell x_0 = x_0 + \pi^\ell \mathrm{Mat}_n(O_F)$ . Let  $y \in x_0 + \pi^\ell \mathrm{Mat}_n(O_F)$ . We have to show that  $y \in (B \cap K_\ell)x_0$ . In order to do this let us represent  $y$  as a block matrix

$$y = \begin{pmatrix} a & b \\ c & D \end{pmatrix},$$

<sup>1</sup>The positive Weyl chamber defined by the Borel  $B$  is the subset of co-weights  $\lambda$  such that  $\pi^\lambda B(O) \pi^{-\lambda} \subset B(O)$

where  $a$  is a scalar and  $D$  is  $n \times n$  matrix. Using left multiplication by lower triangular matrix from  $K_l(\mathrm{GL}_{n+1}(F))$  we may bring  $y$  to the form  $\begin{pmatrix} 1 & b' \\ 0 & D' \end{pmatrix}$ . We can decompose  $D' = LU$ , where  $L, U \in K_l(\mathrm{GL}_{n+1}(F))$  and  $L$  is lower triangular and  $U$  is upper triangular. Therefore by action of an element from  $B \cap K_l$  we may bring  $y$  to the form  $\begin{pmatrix} 1 & b'' \\ 0 & \mathrm{Id} \end{pmatrix}$ . Using right multiplication by diagonal matrix from  $K_l(\mathrm{GL}_{n+1}(F))$  (with first entry 1) we may bring  $y$  to the form  $\begin{pmatrix} 1 & e \\ 0 & D'' \end{pmatrix}$ , where  $e$  is a row vector of 1's and  $D''$  is a diagonal matrix. Finally, using left multiplication by diagonal matrix from  $K_l(\mathrm{GL}_{n+1}(F))$  we may bring  $y$  to be  $x_0$ .  $\square$

**4.3. The pair  $(\mathrm{GL}_{n+1} \times \mathrm{GL}_n, \Delta \mathrm{GL}_n)$ .** In this section we prove Theorem D which states that  $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$  is a strong Gelfand pair for any local field  $F$ , i.e. for any irreducible smooth representations  $\pi$  of  $\mathrm{GL}_{n+1}(F)$  and  $\tau$  of  $\mathrm{GL}_n(F)$  we have

$$\dim \mathrm{Hom}_{\mathrm{GL}_n(F)}(\pi, \tau) \leq 1.$$

It is well known (see e.g. [AGRS, section 1]) that this theorem is equivalent to the statement that  $(\mathrm{GL}_{n+1}(F) \times \mathrm{GL}_n(F), \Delta \mathrm{GL}_n(F))$ , where  $\Delta \mathrm{GL}_n$  is embedded in  $\mathrm{GL}_{n+1} \times \mathrm{GL}_n$  by the map  $\iota \times \mathrm{Id}$ , is a Gelfand pair.

By Corollary C this statement follows from Proposition 4.6, and the following theorem:

**Theorem 4.7** ([AGRS], Theorem 1). *Let  $F$  be a local field of characteristic 0. Then  $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$  is a strong Gelfand pair.*

## References

- [AG09a] A. Aizenbud and D. Gourevitch, Multiplicity one theorem for  $(\mathrm{GL}_{n+1}(\mathbb{R}), \mathrm{GL}_n(\mathbb{R}))$ . *Selecta Math.* **15** (2009), no. 2, 271–294. Zbl 1185.22006 MR 2529937
- [AG09b] A. Aizenbud and D. Gourevitch, Generalized Harish-Chandra descent, Gelfand pairs and an Archimedean analog of Jacquet-Rallis' Theorem. *Duke Math. J.* **149** (2009), no. 3, 509–567. Zbl 1221.22018 MR 2553879
- [AGRS] A. Aizenbud, D. Gourevitch, S. Rallis and G. Schiffmann, Multiplicity one theorems. *Ann. of Math. (2)* **172** (2010), no. 2, 1407–1434. Zbl 1202.22012 MR 2680495
- [AGS08] A. Aizenbud, D. Gourevitch and E. Sayag,  $(\mathrm{GL}_{n+1}(F), \mathrm{GL}_n(F))$  is a Gelfand pair for any local field  $F$ . *Compos. Math.* **144** (2008), no. 6, 1504–1524. Zbl 1157.22004 MR 2474319
- [Ber87] J. N. Bernstein, Second adjointness for representations of reductive  $p$ -adic groups. Unpublished, available at <http://www.math.uchicago.edu/~mitya/langlands.html>.
- [Ber84] J. N. Bernstein, Le “centre” de Bernstein. In *Représentations des groupes réductifs sur un corps local*, Travaux en Cours, Hermann, Paris 1984, 1–32. Zbl 0599.22016 MR 0771671

- [Bus01] C. J. Bushnell, Representations of reductive  $p$ -adic groups: localization of Hecke algebras and applications. *J. London Math. Soc.* **63** (2001), 364–386. Zbl 1017.22011 MR 1810135
- [BZ76] I. N. Bernšteĭn and A. V. Zelevinskiĭ, Representations of the group  $GL(n, F)$ , where  $F$  is a local non-Archimedean field. *Uspehi Mat. Nauk* **31** (1976), no. 3 (189), 5–70; English transl. *Russian Math. Surveys* **31** (1976), no. 3, 1–68. Zbl 0342.43017 MR 0425030
- [Del] P. Delorme, Constant term of smooth  $H_\psi$ -spherical functions on a reductive  $p$ -adic group. *Trans. Amer. Math. Soc.* **362** (2010), 933–955. Zbl 1193.22014 MR 2551511
- [vD] G. van Dijk, Some recent results of Harish-Chandra for  $p$ -adic groups. In *Colloque sur les fonctions sphériques et la théorie des groupes* (Univ. Nancy, Nancy, 1971), Exp. No. 2, Inst. Élie Cartan, Univ. Nancy, Nancy, 1971. Zbl 0352.22016 MR 0439998
- [JR96] H. Jacquet and S. Rallis, Uniqueness of linear periods. *Compos. Math.* **102** (1996), no. 1, 65–123. Zbl 0855.22018 MR 1394521
- [Kaz86] D. A. Kazhdan, Representations groups over close local fields. *J. Analyse Math.* **47** (1986), 175–179. Zbl 0634.22010 MR 0874049
- [Lag08] N. Lagier, Terme constant de fonctions sur un espace symétrique réductif  $p$ -adique. *J. Funct. Anal.* **254** (2008) 1088–1145. Zbl 1194.22010 MR 2381204
- [SV] Y. Sakellaridis and A. Venkatesh, Periods and harmonic analysis on spherical varieties. Preprint, arXiv:1203.0039.
- [SZ] B. Sun and C.-B. Zhu, Multiplicity one theorems: the archimedean case. *Ann. of Math.* **175** (2012), no. 1, 23–44.

Received November 3, 2009

Avraham Aizenbud, Faculty of Mathematics and Computer Science, The Weizmann Institute of Science, POB 26, Rehovot 76100, Israel

E-mail: aizner@yahoo.com

Nir Avni, Department of Mathematics, Harvard University, One Oxford Street, Cambridge, MA 02138, U.S.A.

E-mail: avni.nir@gmail.com

Dmitry Gourevitch, School of Mathematics, Institute for Advanced Study, 1 Einstein Drive, Princeton, NJ 08540, U.S.A.

E-mail: guredim@yahoo.com