

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 88 (2013)

Artikel: Knots with small rational genus
Autor: Calegari, Danny / Gordon, Cameron
DOI: <https://doi.org/10.5169/seals-515632>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Knots with small rational genus

Danny Calegari and Cameron Gordon

Abstract. If K is a rationally null-homologous knot in a 3-manifold M , the *rational genus* of K is the infimum of $-\chi(S)/2p$ over all embedded orientable surfaces S in the complement of K whose boundary wraps p times around K for some p (hereafter: S is a *p-Seifert surface* for K). Knots with very small rational genus can be constructed by “generic” Dehn filling, and are therefore extremely plentiful. In this paper we show that knots with rational genus less than $1/402$ are all *geometric* – i.e. they may be isotoped into a special form with respect to the geometric decomposition of M – and give a complete classification. Our arguments are a mixture of hyperbolic geometry, combinatorics, and a careful study of the interaction of small p -Seifert surfaces with essential subsurfaces in M of non-negative Euler characteristic.

Mathematics Subject Classification (2010). 57M27.

Keywords. Knots, rational genus, stable commutator length, Thurston norm, Berge conjecture.

1. Introduction

Let K be an oriented knot in a 3-manifold M . If K is null-homologous, it bounds an embedded oriented surface S , called a Seifert surface. The least genus of such a surface is called the *genus* of K , and is denoted by $g(K)$. More generally, define the *rational genus* of K , denoted by $\|K\|$, to be the infimum of $-\chi(S)/2p$ over all embedded orientable surfaces S in the complement of K without disk or sphere components, whose boundary wraps p times around K for some p (a precise definition is given in §2).

Largely because of an approach to the Berge conjecture via Knot Floer Homology, there has been recent interest in the question of finding knots in 3-manifolds with the property that they are the *unique* knot in their homology class with least rational genus. Since Knot Floer Homology detects the Thurston norm, and therefore the rational genus of a knot (see Ozsváth–Szabó [18]) such knots have the property that they are characterized by their Knot Floer Homology, and one can study such knots and surgeries on them using the surgery exact sequence. For example, the unknot in S^3 is the unique knot of genus 0, and various families of so-called “simple knots” in lens spaces are the unique knots of minimal rational genus in their respective

homology classes (see Baker [2], Hedden [15] or Rasmussen [19]). Other authors, e.g. [3], have studied *rational linking number*, and its relation to contact geometry.

One way to produce knots of small rational genus is by surgery. For example, let K' be a non-trivial knot in S^3 , and let M be the 3-manifold obtained by p/q surgery on K' . Let K in M be the core of the surgery solid torus. Then $[K]$ has order p in $H_1(M)$, and $\|K\| \leq g(K')/p - 1/2p$ where $g(K')$ denotes the (ordinary) genus of K' , since the boundary of a Seifert surface for K' wraps p times around K in the surgered manifold. For more detailed examples, see §2.4.

The purpose of this paper is firstly to initiate a systematic study of rational genus and some of its properties, and secondly to demonstrate that there is a universal positive constant such that knots in 3-manifolds with rational genus bounded above by this constant can be completely classified. The precise statement of this classification, and the best estimate for the relevant constant, falls into several cases depending on the geometric decomposition of M .

In the generic (i.e. hyperbolic) case, the strategy is to deduce information about K in two steps:

$$L^1\text{-homology} \longrightarrow \text{homotopy} \longrightarrow \text{isotopy}.$$

An estimate for the rational genus of K is really an estimate of the L^1 -norm of a certain relative homology class; such an estimate can be reinterpreted dually in terms of bounded cohomology. Low-dimensional bounded cohomology in hyperbolic manifolds is related to geometry (at the level of π_1) by the methods of Calegari [7]. Homotopy information is promoted to isotopy information by drilling and filling, using uniform geometric estimates due to Hodgson–Kerckhoff [16], and Gauss–Bonnet. One interesting technical aspect of the argument is that it involves finding a CAT(−1) representative of a surface in the complement of the cone locus of a hyperbolic cone manifold. Such a surface can be found either by the PL wrapping technique of Soma [25], or the shrinkwrapping technique of Calegari–Gabai [9].

The organization of the paper is as follows. §2 introduces definitions, proves some basic lemmas, and ends with several subsections enumerating examples. §3 introduces and develops the tools that power our combinatorial arguments (which apply especially to knots with non-hyperbolic complements). §4 treats knots with hyperbolic complements, and the arguments are more geometric and analytic. Finally, §5 assembles all this material, and contains the main theorems and their proofs.

1.1. Statement of results. The classification of knots with (sufficiently) small rational genus falls into several cases, depending on the prime and geometric decomposition of M . These theorems are proved in §5, and reproduced here for the convenience of the reader.

Reducible Theorem (5.1). *Let K be a knot in a reducible manifold M . Then either*

- (1) $\|K\| \geq 1/12$; or

- (2) *there is a decomposition $M = M' \# M''$, $K \subset M'$ and either*
- (a) *M' is irreducible, or*
 - (b) *$(M', K) = (\mathbb{RP}^3, \mathbb{RP}^1) \# (\mathbb{RP}^3, \mathbb{RP}^1)$.*

Lens Theorem (5.2). *Let K be a knot in a lens space M . Then either*

- (1) *$\|K\| \geq 1/24$; or*
- (2) *K lies on a Heegaard torus in M ; or*
- (3) *M is of the form $L(4k, 2k - 1)$ and K lies on a Klein bottle in M as a non-separating orientation-preserving curve.*

Hyperbolic Theorem (5.9). *Let K be a knot in a closed hyperbolic 3-manifold M . Then either*

- (1) *$\|K\| \geq 1/402$; or*
- (2) *K is trivial; or*
- (3) *K is isotopic to a cable of the core of a Margulis tube.*

Small SFS Theorem (5.10). *Let M be an atoroidal Seifert fiber space over S^2 with three exceptional fibers and let K be a knot in M . Then either*

- (1) *$\|K\| \geq 1/402$; or*
- (2) *K is trivial; or*
- (3) *K is a cable of an exceptional Seifert fiber of M ; or*
- (4) *M is a prism manifold and K is a fiber in the Seifert fiber structure of M over \mathbb{RP}^2 with at most one exceptional fiber.*

Toroidal Theorem (5.19). *Let M be a closed, irreducible, toroidal 3-manifold, and let K be a knot in M . Then either*

- (1) *$\|K\| \geq 1/402$; or*
- (2) *K is trivial; or*
- (3) *K is contained in a hyperbolic piece N of the JSJ decomposition of M and is isotopic either to a cable of a core of a Margulis tube or into a component of ∂N ; or*
- (4) *K is contained in a Seifert fiber piece N of the JSJ decomposition of M and either*
 - (A) *K is isotopic to an ordinary fiber or a cable of an exceptional fiber or into ∂N , or*
 - (B) *N contains a copy Q of the twisted S^1 bundle over the Möbius band and K is contained in Q as a fiber in this bundle structure;*

or

- (5) M is a T^2 -bundle over S^1 with Anosov monodromy and K is contained in a fiber.

The subsections §2.4 and §2.5 discuss constructions giving rise to eight families of examples of knots with arbitrarily small rational genus, illustrating that all the possibilities listed in the classification theorems really do occur.

1.2. Acknowledgements. The first author would like to thank Matthew Hedden and Jake Rasmussen for interesting and stimulating talks they gave at Caltech in 2007, which were the inspiration for this paper. He would also like to thank Marty Scharlemann and Yoav Rieck for useful conversations about thin position. The second author would like to thank Constance Leidy and Peter Ozsváth for useful comments. Danny Calegari was partially supported by NSF grant DMS 0707130.

2. p -Seifert surfaces

This section standardizes definitions, proves some basic facts about rational genus, and describes how to construct examples of knots with small rational genus, illustrating the significance of the cases we enumerate in our classification theorems.

2.1. Definitions. We formalize definitions in this section. Throughout, all 3-manifolds considered will be compact, connected and orientable. A *knot* K in a 3-manifold M is a tamely embedded S^1 . If K is null-homologous in M , a *Seifert surface* for K is a connected embedded two-sided surface S in M with $\partial S = K$. The *genus* of K is the least genus of a Seifert surface.

This can be generalized as follows. By analogy with the Thurston norm on $H_2(M)$, we adopt the following notation:

Notation 2.1. If S is a compact, orientable connected surface, define $\chi^-(S) := \min(0, \chi(S))$. If S is a compact, orientable surface with components S_i , define $\chi^-(S) = \sum_i \chi^-(S_i)$. Denote $\eta(S) = -\chi^-(S)/2$.

Remark 2.2. The normalizing factor of 2 in the denominator of η reflects the fact that Euler characteristic is “almost” -2 times genus for a surface with a bounded number of boundary components.

Definition 2.3. Let K be a knot in a 3-manifold M , with regular neighborhood $N(K)$. If p is a positive integer, a p -*Seifert surface* for K is a compact, oriented surface S embedded in $M - \text{int } N(K)$ such that $S \cap \partial N(K) = \partial S$, and $[\partial S] = p[K] \in$

$H_1(N(K))$ (for some choice of orientation on K). In this case we define the *norm*, or *rational genus* of K by

$$\|K\| = \inf_S \eta(S)/p$$

where the infimum is taken over all p and all p -Seifert surfaces S for K .

A p -Seifert surface S for K can be extended into $N(K)$ to give a map $S \rightarrow M$ which is an embedding on $\text{int } S$ and whose restriction $\partial S \rightarrow K$ is a (possibly disconnected) covering map of degree p . We will regard a p -Seifert surface for K as a singular surface in M in this way in §4.

Remark 2.4. A knot K has a p -Seifert surface for some p if and only if $[K]$ has finite order in $H_1(M)$.

Definition 2.5. A p -Seifert surface is *good* if it satisfies the following properties:

- (1) S is connected.
- (2) S is incompressible in $M - \text{int } N(K)$.
- (3) ∂S consists of q parallel, coherently oriented copies of an essential simple closed curve ρ on $\partial N(K)$ that represents r times a generator of $H_1(N(K))$, where $qr = p$.

Lemma 2.6. Let S be a p -Seifert surface for K . Then there is a good p' -Seifert surface S' for K satisfying

$$\eta(S)/p \geq \eta(S')/p'.$$

Proof. Boundary components of S that are inessential in $\partial N(K)$ may be capped off with disks, and closed components of S may be discarded; neither of these operations increases η or changes p . If some component of S is a disk D , then D is a good p' -Seifert surface for K for some p' , and $\eta(D)$ and $\|K\|$ are both zero. Hence we may assume that every component S_i of S satisfies $\chi(S_i) = \chi^-(S_i)$, and has non-empty boundary, each component of which is an essential curve in $\partial N(K)$.

Since a p -Seifert surface is embedded, the components of ∂S are all parallel in $\partial N(K)$, and are therefore all isotopic (though *a priori* they might have opposite orientations). Let S be a p -Seifert surface for K , and suppose there are a pair of adjacent components of ∂S on $\partial N(K)$ that are oppositely oriented. Tubing this pair of components does not affect $\eta(S)$ or p , so without loss of generality we may assume that all components are coherently oriented.

If S is compressible in $M - \text{int } N(K)$ then compressing it along a disk gives a surface S' with $\partial S' = \partial S$ and $\eta(S') \leq \eta(S)$. So we may assume that S is incompressible.

It remains to show that we can take S to be connected. Suppose S is the disjoint union of S_1 and S_2 , where ∂S_i consists of q_i copies of ρ , and each $q_i > 0$. Then we estimate

$$\frac{\eta(S)}{p} = \frac{\eta(S)}{qr} = \frac{\eta(S_1) + \eta(S_2)}{(q_1 + q_2)r} \geq \min \left\{ \frac{\eta(S_1)}{q_1 r}, \frac{\eta(S_2)}{q_2 r} \right\}. \quad \square$$

It will be important in the sequel to consider surfaces in manifolds M that meet ∂M .

Definition 2.7. A *relative p -Seifert surface* F for a knot K in M is an oriented surface, properly embedded in $M - \text{int } N(K)$ such that $[\partial F \cap \partial N(K)] = p[K] \in H_1(N(K))$.

The definition of *good* extends to relative p -Seifert surfaces, and Lemma 2.6 generalizes to such surfaces as well and with the same proof, so in the sequel we assume all our p -Seifert surfaces, relative or otherwise, are good.

Notation 2.8. In the sequel, we write $X := M - \text{int } N(K)$.

2.2. Thurston norm. A basic reference for this section is Thurston's paper [26].

If K is a knot in a closed 3-manifold M , then X is a compact 3-manifold with torus boundary, and as is well-known, the kernel of the inclusion map

$$H_1(\partial X; \mathbb{Q}) \rightarrow H_1(X; \mathbb{Q})$$

is 1-dimensional, and denoted L . Consequently, the kernel of

$$H_1(\partial X; \mathbb{Z}) \rightarrow H_1(X; \mathbb{Z})$$

is isomorphic to \mathbb{Z} , and is generated by a single element $m[\rho]$, where $[\rho]$ is primitive in $H_1(\partial X; \mathbb{Z})$, and $[\rho]$ is represented by a simple loop $\rho \subset \partial X$ which represents r times a generator of $H_1(N(K))$ (this is the same ρ as before).

Let $\partial^{-1}L$ denote the subspace of $H_2(X, \partial X; \mathbb{Q})$ that is the preimage of L under the connecting homomorphism in rational homology. If S is a p -Seifert surface for K , then $p = qr$ where $m|q$, and $[S] \in \partial^{-1}L$. Consider the affine rational subspace $\partial^{-1}[\rho]/r \subset \partial^{-1}L$. The multiple $[S]/p \in \partial^{-1}[\rho]/r$, and $\eta(S)/p = \|S\|_T/2p$ where $\|\cdot\|_T$ denotes the Thurston norm of a surface. Hence, by the definition of rational genus and of Thurston norm, we obtain the following formula:

Lemma 2.9. *There is an equality*

$$\|K\| = \inf_{A \in \partial^{-1}[\rho]/r} \|A\|_T/2$$

where $\|A\|_T$ denotes the Thurston norm of the (rational) homology class A .

Since $\|\cdot\|_T$ is a convex, non-negative, piecewise rational linear function on $H_2(X, \partial X; \mathbb{R})$, the infimum of $\|\cdot\|_T/2$ is *achieved* (on some rational subpolyhedron) on the rational affine subspace $\partial^{-1}[\rho]/r$. Consequently, we have:

Proposition 2.10. *Let K be a knot. The rational genus $\|K\|$ is equal to $\eta(S)/p$ for some p -Seifert surface S and some p . Therefore $\|K\|$ is rational. Moreover, there is an algorithm to find S and compute $\|K\|$.*

Proof. There is a (straightforward) algorithm to compute the Thurston norm, described in [26], and to find a norm-minimizing surface in any integral class (note that such a surface can be taken to be normal relative to a fixed triangulation, and therefore may be found by linear programming in normal surface space). \square

Although Proposition 2.10 is included for completeness, it is not essential for the remainder of the paper, and it is generally good enough in the sequel to work with a p -Seifert surface that comes close to realizing $\|K\|$.

The first thing one wants to know about an invariant is when it vanishes.

Theorem 2.11. *Let K be a knot in a 3-manifold M . Then $\|K\| = 0$ if and only if either*

- (1) K bounds a disk in M ; or
- (2) K is the core of a genus 1 Heegaard splitting of a lens space summand of M ; or
- (3) K is the fiber of multiplicity r in a Seifert fiber subspace of M whose base orbifold is a Möbius band with one orbifold point of order $r \geq 1$.

Proof. Suppose $\|K\| = 0$. Then by Lemma 2.6 and Proposition 2.10, K has a good p -Seifert surface with $\chi^-(S) = 0$, i.e. S is a disk or annulus.

First assume S is a disk. If $p = 1$ we have conclusion (1). If $p > 1$ then a regular neighborhood of $N(K) \cup S$ is a punctured lens space with fundamental group $\mathbb{Z}/p\mathbb{Z}$, with K as a core of a genus 1 Heegaard splitting.

If S is an annulus, note that both boundary components of S wrap with the same orientation r times around K . A regular neighborhood of $N(K) \cup S$ evidently has the desired structure. \square

Remark 2.12. Since M is orientable, the total space of the Seifert fiber subspace in bullet (3) of Theorem 2.11 is orientable; i.e. it is a twisted S^1 bundle over an orbifold Möbius band. There is no suggestion that it is essential in M . The situation described in (3) arises in case (2) (b) of Theorem 5.1, case (3) of Theorem 5.2, case (4) of Theorem 5.10, and case (4) (B) of Theorem 5.19.

Under suitable homological conditions on M and K , the analysis simplifies considerably:

Lemma 2.13. *Let K be a knot in a \mathbb{Q} -homology 3-sphere and let S be a connected p -Seifert surface for K . Then p is the order of $[K]$ in $H_1(M)$.*

Proof. Since M is a \mathbb{Q} -homology sphere, $H_2(X) = 0$ and hence the boundary map $H_2(X, \partial X) \rightarrow H_1(\partial X)$ is injective, with image $\ker(H_1(\partial X) \rightarrow H_1(X)) \cong \mathbb{Z}$. Let x be a generator of $H_2(X, \partial X)$. Then the image of x under the composition

$$H_2(X, \partial X) \rightarrow H_1(\partial X) \rightarrow H_1(N(K)) \cong \mathbb{Z}$$

is p_0 , say, the order of $[K]$ in $H_1(M)$. Let S be a connected p -Seifert surface for K . Then $[S] = kx$ and $p = kp_0$ for some $k > 0$. But since S is connected, $k = 1$ by Lemma 1 of [26]. \square

If K is a knot in a homology 3-sphere, the (ordinary) *genus of K* , denoted by $g(K)$, is the minimal genus of any Seifert surface for K . Lemma 2.13 reduces the study of rational genus to that of the usual genus in homology spheres:

Corollary 2.14. *If K is a knot in a homology 3-sphere then*

$$\|K\| = \begin{cases} 0, & \text{if } g(K) = 0; \\ g(K) - 1/2, & \text{if } g(K) > 0. \end{cases}$$

Proof. By Lemma 2.13, $\|K\| = \eta(S)$, where S is a minimal genus Seifert surface for K . If S is a disk, $\|K\| = g(K) = 0$. Otherwise, $\eta(S) = (2g(S) - 1)/2 = g(S) - 1/2$. \square

The following lemma will allow us to construct knots in 3-manifolds with arbitrarily small (non-zero) rational genus.

Lemma 2.15. *Let K' be a knot in a homology 3-sphere M' . Let M be the manifold obtained by m/n -Dehn surgery on K' , where $m > 0$, and let $K \subset M$ be the core of the surgery solid torus. Then $\|K\| = \|K'\|/m$.*

Proof. Note that $[K]$ has order m in $H_1(M) = \mathbb{Z}/m\mathbb{Z}$.

Let S be a good p -Seifert surface for K in M such that $\|K\| = \eta(S)/p$. The restriction of S to $M - K = M' - K'$ extends to a good p' -Seifert surface for K' (which by abuse of notation we call S) where $p' = p/m$. By the proof of Lemma 2.13, $p' = 1$ and S is a Seifert surface for K' in M' . Conversely a Seifert surface for K' can also be thought of as a good m -Seifert surface for K . Therefore $\|K'\| = \eta(S)$ and $p = m$. Hence $\|K\| = \|K'\|/m$. \square

2.3. Connect sums. We now examine the behavior of rational genus under connected sum. In this context it is convenient to say that a knot K in M is p -trivial if it has a p -Seifert surface that is a disk. If $p = 1$ then K bounds a disk, and is *trivial*. If $p > 1$ then K is the core of a genus 1 Heegaard splitting of a lens space summand of M with fundamental group $\mathbb{Z}/p\mathbb{Z}$.

Remark 2.16. If K is p -trivial for some p then clearly $\partial N(K)$ is compressible in X . Conversely, a compressing disk for $\partial N(K)$ in X is either a p -Seifert surface for K , for some $p \geq 1$, or has boundary a meridian of K , in which case K is isotopic to $S^1 \times \{\text{point}\}$ in some $S^1 \times S^2$ summand of M . So for rationally null-homologous knots, being p -trivial for some p is equivalent to $\partial N(K)$ being compressible in X .

Theorem 2.17. *Let K_1 and K_2 be knots in 3-manifolds M_1 and M_2 . Then*

- (1) *if K_1 is p_1 -trivial and K_2 is trivial then $K_1 \# K_2$ is p_1 -trivial;*
- (2)

$$\|K_1 \# K_2\| = \begin{cases} \|K_1\| + \|K_2\| + \frac{1}{2} & \text{if } K_1 \text{ and } K_2 \text{ are not } p\text{-trivial for any } p; \\ \|K_1\| + \frac{1}{2} - \frac{1}{2p_2} & \text{if } K_2 \text{ is } p_2\text{-trivial, } K_1 \text{ is not } p\text{-trivial for any } p; \\ \frac{1}{2} - \frac{(p_1 + p_2)}{2p_1 p_2} & \text{if } K_i \text{ is } p_i\text{-trivial, } p_i \geq 2, i = 1, 2. \end{cases}$$

Remark 2.18. The first case in bullet (2) says that for knots that are not p -trivial for any p , the quantity $\|K\| + \frac{1}{2}$ is additive under connected sum, and is the analog of the additivity of genus for knots in S^3 ; see Corollary 2.14.

Remark 2.19. Theorem 2.17 has an analog in the theory of stable commutator length (see Definition 4.1); compare with the Product formula (Theorem 2.93) from [8].

Remark 2.20. Note that K is 2-trivial if and only if K is contained in an \mathbb{RP}^3 summand as \mathbb{RP}^1 . Also, it follows from Theorem 2.17 that if $K = K_1 \# K_2$ with K_1 and K_2 non-trivial, then $\|K\| = 0$ if and only if K_1 and K_2 are 2-trivial, i.e. K is contained in an $\mathbb{RP}^3 \# \mathbb{RP}^3$ summand as $\mathbb{RP}^1 \# \mathbb{RP}^1$. This is a special case of Theorem 2.11 (3), with $r = 1$.

Proof of Theorem 2.17. Let $S_i \subset M_i$ be a good p_i -Seifert surface for K_i with $\|K_i\| = \eta(S_i)/p_i$, $i = 1, 2$. Then we may construct a $p_1 p_2$ -Seifert surface S for $K = K_1 \# K_2$ in $M_1 \# M_2$ by taking p_2 copies of S_1 and p_1 copies of S_2 and joining them along $p_1 p_2$ arcs. We have

$$\chi(S) = p_2 \chi(S_1) + p_1 \chi(S_2) - p_1 p_2. \quad (2.3.1)$$

Conversely, let S be a good p -Seifert surface for $K = K_1 \# K_2$ with $\|K\| = \eta(S)/p$. Suppose ∂S has q components, each having intersection number r with a meridian of K . Let A be the annulus in $X = M_1 \# M_2 - \text{int } N(K)$ that realizes the connected sum decomposition. By an isotopy of S we may assume that each component of ∂S meets each component of ∂A in r points, and that $S \cap A$ is a disjoint union of arcs and simple closed curves. Since the boundary components of S are oriented coherently on $\partial N(K)$, each arc must have one endpoint on each component of ∂A . It follows that any simple closed curve of intersection is inessential in A , and therefore in S , and so these can be removed by performing surgery on S and discarding the resulting 2-spheres. Hence we may assume that $S \cap A$ consists of $qr = p$ essential arcs in A . Cutting S along these arcs gives p -Seifert surfaces S_1, S_2 for K_1, K_2 in M_1, M_2 . Note that

$$\chi(S) = \chi(S_1) + \chi(S_2) - p. \quad (2.3.2)$$

To prove part (1), note that if S_1 and S_2 are disks and $p_2 = 1$ then the p_1 -Seifert surface for $K_1 \# K_2$ in (2.3.1) is a disk.

To prove (2), first suppose that K_1 and K_2 are not p -trivial (for any p). Then the surfaces S_1 and S_2 in (2.3.1) and (2.3.2) have no disk components, and from (2.3.1) we get

$$\eta(S) = p_2\eta(S_1) + p_1\eta(S_2) + p_1p_2/2,$$

and hence

$$\begin{aligned} \|K\| &\leq \eta(S)/p_1p_2 = \eta(S_1)/p_1 + \eta(S_2)/p_2 + \frac{1}{2} \\ &= \|K_1\| + \|K_2\| + \frac{1}{2}. \end{aligned} \quad (2.3.3)$$

Similarly, (2.3.2) gives

$$\eta(S) = \eta(S_1) + \eta(S_2) + p/2,$$

and hence

$$\begin{aligned} \|K\| &= \eta(S)/p = \eta(S_1)/p + \eta(S_2)/p + \frac{1}{2} \\ &\geq \|K_1\| + \|K_2\| + \frac{1}{2}. \end{aligned} \quad (2.3.4)$$

Together, (2.3.3) and (2.3.4) give the first assertion in part (2) of the theorem.

Second, suppose K_2 is p_2 -trivial and K_1 is not p -trivial for any p . Then in (2.3.1) S_2 is a disk while $\chi(S_1) \leq 0$. Hence $\chi(S) \leq 0$, and we get

$$\begin{aligned} \|K\| &\leq \eta(S)/p_1p_2 = \eta(S_1)/p_1 - \frac{1}{2p_2} + \frac{1}{2} \\ &= \|K_1\| - \frac{1}{2p_2} + \frac{1}{2}. \end{aligned} \quad (2.3.5)$$

In (2.3.2), since K_1 is not p -trivial for any p , no component of S_1 is a disk. Hence $\eta(S_1) = -\chi(S_1)/2$, $\chi(S_1) \leq 0$, and $\chi(S) \leq 0$. Let S_2 have d_2 disk components. Then $\chi(S_2) \leq d_2$ and

$$d_2 p_2 \leq p. \quad (2.3.6)$$

Now

$$\begin{aligned} \|K\| &= \eta(S)/p = -\chi(S)/2p \\ &\geq -\chi(S_1)/2p - \frac{d_2}{2p} + \frac{1}{2} \\ &\geq \|K_1\| - \frac{d_2}{2p} + \frac{1}{2}. \end{aligned} \quad (2.3.7)$$

Comparing (2.3.7) with (2.3.5) we get

$$\frac{d_2}{2p} \geq \frac{1}{2p_2}, \quad \text{i.e.} \quad d_2 p_2 \geq p.$$

By (2.3.6), this gives $d_2 p_2 = p$ and (2.3.5) is an equality.

Finally, suppose that K_i is p_i -trivial, $p_i \geq 2$, $i = 1, 2$. In (2.3.1), S_1 and S_2 are disks, and

$$\chi(S) = p_1 + p_2 - p_1 p_2 \leq 0.$$

Hence

$$\begin{aligned} \|K\| &\leq \eta(S)/p_1 p_2 = \frac{p_1 p_2 - (p_1 + p_2)}{2 p_1 p_2} \\ &= \frac{1}{2} - \frac{(p_1 + p_2)}{2 p_1 p_2}. \end{aligned} \quad (2.3.8)$$

Now consider (2.3.2), and let S_i have d_i disk components, $i = 1, 2$. Note that

$$d_i p_i \leq p, \quad i = 1, 2; \quad (2.3.9)$$

in particular, since $p_i \geq 2$, $d_i \leq p/2$, $i = 1, 2$. It follows that

$$\chi(S) \leq d_1 + d_2 - p \leq 0.$$

Therefore

$$\begin{aligned} \|K\| &= \eta(S)/p = -\chi(S)/2p \\ &\geq \frac{1}{2} - \frac{(d_1 + d_2)}{2p}. \end{aligned} \quad (2.3.10)$$

Comparing (2.3.10) with (2.3.8) gives

$$\frac{d_1}{p} + \frac{d_2}{p} \geq \frac{1}{p_1} + \frac{1}{p_2}.$$

On the other hand, by (2.3.9) we have $d_i/p \leq 1/p_i$, $i = 1, 2$, and hence (2.3.8) is an equality, as desired. \square

2.4. Examples of knots with small rational genus. We use Lemma 2.15 to construct examples of knots K in 3-manifolds M with arbitrarily small (but non-zero) rational genus. These examples illustrate cases from our main classification theorems, to be proved in §4 and §5, especially case (2) of Theorem 4.8, case (2) of Theorem 5.2, case (3) of Corollary 5.9, case (3) of Theorem 5.10, and cases (3) and (4) of Theorem 5.19. We recall the notation $\Delta(\cdot, \cdot)$ for the minimal number of points of intersection (i.e. the homological intersection number) of two unoriented isotopy classes of simple essential loops on a torus; in co-ordinates, $\Delta(a/b, c/d) = |ad - bc|$.

Case A. (M is hyperbolic and K is the core of a Margulis tube.) Let K' be a hyperbolic knot in S^3 . Let M be the result of m/n -Dehn surgery on K' , $m > 0$, and let K be the core of the surgery solid torus. By Lemma 2.15, $\|K\| = \|K'\|/m \rightarrow 0$ as $m \rightarrow \infty$. Also, for m sufficiently large M is hyperbolic and K is a geodesic in M whose length $\rightarrow 0$ as $m \rightarrow \infty$.

Case B. (M is a lens space and K lies on a genus 1 Heegaard surface for M .) Let (M, K) be as in Case A above, but with K' a (u, v) -torus knot in S^3 , where $u, v > 1$. If $d = \Delta(m/n, uv/1) = |m - nuv| = 1$ then M is a lens space, and one sees that K lies on a Heegaard torus in M . By choosing m large enough we can make $\|K\| = \|K'\|/m$ arbitrarily small.

Case C. (M is a Seifert fiber space and K is an ordinary or exceptional fiber.) By taking $d > 1$ in Case B above, M becomes a Seifert fiber space with base orbifold S^2 with three cone points of orders u, v and d , and K is the exceptional fiber of multiplicity d .

More generally, let M' be a Seifert fibered homology 3-sphere, with base S^2 and $k \geq 3$ exceptional fibers. Let K' be an ordinary fiber, and let a/b be the slope of the fiber on $\partial X'$, where $X' = M' - \text{int } N(K')$. Let M be m/n -Dehn surgery on K' , and K the core of the surgery solid torus. Let $d = \Delta(a/b, m/n) = |an - bm|$. If $d = 1$, then M is a Seifert fiber space over S^2 with k exceptional fibers, and K is an ordinary fiber. If $d > 1$, then M is a Seifert fiber space over S^2 with $(k + 1)$ exceptional fibers, and K is an exceptional fiber of multiplicity d . By Lemma 2.15, $\|K\| = \|K'\|/m$, and in both cases this can be made arbitrarily small by taking m sufficiently large.

Case D. (M is hyperbolic and K is a non-trivial cable of the core of a Margulis tube.) Let K_0 be a hyperbolic knot in S^3 . Fix coprime integers $p, q > 1$, and let k be any positive integer. Then kq^2 and $(1 + kpq)$ are coprime, so there exist integers a, b such that $akq^2 - b(1 + kpq) = 1$. Let M be the manifold obtained by $(-kq^2/b)$ -Dehn surgery on K_0 , and let $K' \subset M$ be the core of the surgery. Let X_0 be the exterior of K_0 ; then $M = X_0 \cup V$ where V is a tubular neighborhood of K' . Let μ_0, λ_0 (resp. μ, λ) be a canonical meridian-longitude pair of generators for $H_1(\partial X_0)$ (resp. $H_1(\partial V)$). Let $f: \partial V \rightarrow \partial X_0$ be the gluing homeomorphism. We can choose f so

that with respect to the above bases $f_*: H_1(\partial V) \rightarrow H_1(\partial X_0)$ is given by the matrix

$$A = \begin{pmatrix} kq^2 & -(1 + kpq) \\ -b & a \end{pmatrix}.$$

Then

$$A^{-1} = \begin{pmatrix} a & 1 + kpq \\ b & kq^2 \end{pmatrix}$$

in particular, under the gluing the longitude λ_0 on ∂X_0 is identified with a curve of slope $(1 + kpq)/kq^2$ on ∂V .

Let $K \subset \text{int } V$ be the (p, q) -cable of K' , the core of V . Then $X = M - \text{int } N(K) = X_0 \cup (V - \text{int } N(K)) = X_0 \cup C$, where C is a (p, q) -cable space (see [13], §3). There exists a planar surface $P \subset C$ with one boundary component on ∂V , with slope $(1 + kpq)/kq^2$, and q boundary components on $\partial N(K)$, with slope $(1 + kpq)/k$ (see Lemma 3.1 in [13]). Let S_0 be a Seifert surface for K_0 , and define $S = P \cup S_0$, glued along $\partial S_0 = P \cap \partial V$. Since each of the q components of $S \cap \partial N(K)$ has intersection number k with the meridian of K , it follows that S is a qk -Seifert surface for K . Therefore $\|K\| \leq \eta(S)/qk = (\eta(S_0) + (q - 1)/2)/qk$, which goes to 0 as k goes to infinity. Also, for k sufficiently large, M is hyperbolic and K' is the core of a Margulis tube in M .

Case E. (M is a Seifert fiber space and K is a cable of an exceptional fiber but not a fiber.) Repeat the construction in Case D above, but with K_0 a (u, v) -torus knot. Then M is a Seifert fiber space and K' is a fiber of multiplicity $\Delta(-kq^2/b, uv/1) = |kq^2 + buv|$, which we can arrange to be > 1 . Then K is the (p, q) -cable of an exceptional fiber. On the other hand it is easy to see that K can be a fiber in the Seifert fibration of M for at most one value of k .

Case F. (M is toroidal and K lies in a torus in the JSJ decomposition of M .) Let K_1, K_2 be non-trivial knots in S^3 , and let $K' = K_1 \# K_2$. Let M be the manifold obtained by m -surgery on K' for some $m > 0$, and let K be the core of the surgery solid torus. By Lemma 2.15 $\|K\| = \|K'\|/m$; thus $\|K\|$ is non-zero but can be made arbitrarily small by taking m sufficiently large.

Let $X_i = S^3 - \text{int } N(K_i)$ be the exterior of K_i , $i = 1, 2$; then $X = S^3 - \text{int } N(K') = X_1 \cup_A X_2$ where A is a meridional annulus in ∂X_i , $i = 1, 2$. Let V be the surgery solid torus. Note that the boundary slope of A on ∂X (the meridian of K') intersects the meridian of V once. It follows that $X_1 \cup V \cong X_1$, and so $M = (X_1 \cup X_2) \cup V \cong X_1 \cup_T X_2$, where $T = \partial X_1 = \partial X_2$. Also K , the core of V , is isotopic into T .

If, for example, we take K_1 to be hyperbolic and K_2 to be either hyperbolic or a torus knot, then T is the unique torus in the JSJ decomposition of M . Note also that in the second case K is not a Seifert fiber in X_2 . If we take K_i to be the (p_i, q_i) torus

knot, $i = 1, 2$, and $m \neq p_1q_1 + p_2q_2$, then again T is the unique JSJ torus in M , and K is not a Seifert fiber of either X_1 or X_2 .

Further examples are given in the next subsection.

2.5. Torus bundles. In this section we analyze the case where M is a T^2 -bundle over S^1 and K is an essential simple closed curve in a fiber. It turns out that this gives further examples of knots with arbitrarily small rational genus, which need to be taken account of in the statement of Theorem 5.19.

Let $f: T^2 \rightarrow T^2$ be an orientation preserving diffeomorphism, and let M_f be the mapping torus of f , obtained from $T^2 \times I$ by identifying $(x, 0)$ with $(f(x), 1)$ for all $x \in T^2$. If we choose a basis for $H_1(T^2)$, the automorphism f_* of $H_1(T^2)$ induced by f is represented by a matrix $A_f \in \text{SL}(2, \mathbb{Z})$; the diffeomorphism type of M_f depends only on the conjugacy class of A_f in $\text{GL}(2, \mathbb{Z})$. Let K in M_f be an essential simple closed curve in a fiber. If $\text{trace } f_* \neq 2$ then $\det(A_f - I) \neq 0$, and so every such K has finite order in $H_1(M_f)$. If $\text{trace } f_* = 2$ then A_f is conjugate in $\text{GL}(2, \mathbb{Z})$ to $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$ for some $p \geq 0$. If $p = 0$ then $M_f = T^3$ and no K has finite order in $H_1(M_f)$, so in the sequel we shall always assume that $f_* \neq \text{Id}$. If $p \geq 1$ then there is a unique K with finite order ($= p$) in $H_1(M_f)$.

Let a and b be oriented simple closed curves in T^2 meeting transversely in a single point, such that K is the image in M_f of the curve $a \times \{1/2\} \subset T^2 \times I$. Thinking of $T^2 \times I$ as $(a \times b) \times I = a \times (b \times I)$ shows that $(T^2 \times I, K) = (S^1 \times A^2, S^1 \times \{\text{point}\})$, and therefore $T^2 \times I - \text{int } N(K) \cong S^1 \times P^2$, where P^2 is a pair of pants. Let the boundary components of P^2 be B_0, B_1 and C , where $S^1 \times B_i = T_i = T^2 \times \{i\}$, $i = 0, 1$, and $S^1 \times C = \partial N(K)$. Let b_0, b_1, c be the homology classes of B_0, B_1 and C , respectively, oriented so that $[\partial P^2] = b_0 - b_1 + c$, and so that b_0 and b_1 map to the class b above in $H_1(T^2 \times I)$ (we will abuse notation by not distinguishing between a, b and their classes in $H_1(T^2)$).

We now wish to describe certain horizontal surfaces in $S^1 \times P^2$. Consider the homomorphism $H_1(P^2) \rightarrow \mathbb{Z}$ defined by $b_0 \mapsto \ell_0, b_1 \mapsto \ell_1$ (so $c \mapsto \ell_1 - \ell_0$), where ℓ_0 and ℓ_1 are arbitrary integers. Composing with the Hurewicz map we get a homomorphism $\pi_1(P^2) \rightarrow \mathbb{Z}$, which is induced by a map $\pi: P^2 \rightarrow S^1$. Let m be an integer ≥ 1 and let $\sigma: S^1 \rightarrow S^1$ be the connected covering of degree m . Let $F \rightarrow P^2$ be the $\mathbb{Z}/m\mathbb{Z}$ covering corresponding to the composition of π_* with the quotient map $\mathbb{Z} \rightarrow \mathbb{Z}/m\mathbb{Z}$. This covering is the pull-back of σ under the map π , in other words we have a commutative diagram

$$\begin{array}{ccc} F & \xrightarrow{p_1} & S^1 \\ p_2 \downarrow & & \downarrow \sigma \\ P^2 & \xrightarrow{\pi} & S^1 \end{array}$$

where F is identified with the subset $\{(x, y) : \sigma(x) = \pi(y)\}$ of $S^1 \times P^2$, and p_1 and p_2 are the restrictions of the projections of $S^1 \times P^2$ onto the factors.

Consider the ordered bases (a, b_0) , (a, b_1) and (a, c) for $H_1(T_0)$, $H_1(T_1)$ and $H_1(\partial N(K))$, respectively. Let \tilde{B}_0 , \tilde{B}_1 , \tilde{C} be the respective inverse images in ∂F of B_0 , B_1 and C . Since the coverings $\tilde{B}_i \rightarrow B_i$, $i = 0, 1$, and $\tilde{C} \rightarrow C$ have degree m , we have

$$\begin{aligned} [\tilde{B}_i] &= \ell_i a + m b_i, \quad i = 0, 1, \\ [\tilde{C}] &= \ell a + m c \quad \text{where } \ell = \ell_1 - \ell_0. \end{aligned} \quad (2.5.1)$$

In particular, if $\ell \neq 0$ then F is a relative $|\ell|$ -Seifert surface for K in $T^2 \times I$.

Let f_* be represented with respect to the basis (a, b) by $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{SL}(2, \mathbb{Z})$. Assume for the moment that we are not in the case $\gamma = 0$, $\alpha = \delta = -1$. Then we can choose ℓ_0 and $m \geq 1$ such that

$$\gamma \ell_0 + (\delta - 1)m = 0, \quad (2.5.2)$$

and define

$$\ell_1 = \alpha \ell_0 + \beta m; \quad (2.5.3)$$

so

$$\ell = (\alpha - 1)\ell_0 + \beta m. \quad (2.5.4)$$

Then $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} \ell_0 \\ m \end{pmatrix} = \begin{pmatrix} \ell_1 \\ m \end{pmatrix}$, which implies that we may isotope f so that $f(\tilde{B}_0) = \tilde{B}_1$, and hence F becomes an orientable surface S in M_f . If $\text{trace } f_* \neq 2$ then (2.5.2) and (2.5.4) imply that $\ell \neq 0$, and so S is an $|\ell|$ -Seifert surface for K . Since $\chi(S) = \chi(F) = m\chi(P^2) = -m$, we get

$$\|K\| \leq m/2|\ell|. \quad (2.5.5)$$

We note that if $\text{trace } f_* \neq 2$ and $\gamma \neq 0$, it follows easily from (2.5.2) and (2.5.4) that

$$m/\ell = \gamma/(\text{trace } f_* - 2). \quad (2.5.6)$$

In the case $\text{trace } f_* = 2$, A_f is conjugate to $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$, $p \geq 1$, where the first element of the corresponding ordered basis for $H_1(T^2)$ is represented by the unique K that has finite order in $H_1(M_f)$. Thus $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$, and in (2.5.2) we choose $\ell_0 = 0$, $m = 1$, giving $\ell = \ell_1 = p$.

The above discussion shows that unless $\gamma = 0$ and $\text{trace } f_* = -2$, any K in a fiber of M_f having finite order in $H_1(M_f)$ has an $|\ell|$ -Seifert surface, $\ell \neq 0$, such that the corresponding surface F in $T^2 \times I - \text{int } N(K) = S^1 \times P^2$ is horizontal (i.e. transverse to the S^1 fibers), and (2.5.5) holds.

It remains to discuss the case $\gamma = 0$, $\text{trace } f_* = -2$, i.e. where our matrix is $\begin{pmatrix} -1 & \beta \\ 0 & -1 \end{pmatrix}$. Let A_i be a vertical annulus in $S^1 \times P^2$ with one boundary component on

each of T_i and $\partial N(K)$, $i = 0, 1$. Since $f_*(a) = -a$, A_0 and A_1 glue up to give an annulus A in $M_f - \text{int } N(K)$ whose boundary components are coherently oriented on $\partial N(K)$. Hence $\|K\| = 0$.

Finally, we show that the inequality (2.5.5) is an equality. Let S be a good p -Seifert surface for K in M_f such that $\|K\| = \eta(S)/p$. Let $T \subset M_f$ be the fiber that is the image of $T^2 \times \partial I$; note that $K \cap T = \emptyset$. Isotoping S to minimize the number of components of $S \cap T$ we get a relative p -Seifert surface F for K in $T^2 \times I$ that is essential in $T^2 \times I - \text{int } N(K) = S^1 \times P^2$. Therefore F is either horizontal or vertical.

First we dispose of the vertical case. Here F must consist of either a single annulus with both boundary components on $\partial N(K)$, or two annuli, one running from $\partial N(K)$ to T_0 and the other from $\partial N(K)$ to T_1 . In the first case, $[\partial S] = [\partial F] = 0 \in H_1(\partial N(K))$, a contradiction. In the second case, we also get $[\partial S] = 0 \in H_1(\partial N(K))$ unless $f_*(a) = -a$, in which case A_0 and A_1 glue up to give an annulus A in $M_f - \text{int } N(K)$ that is a 2-Seifert surface for K , implying that $\|K\| = 0$. This is precisely the case $\gamma = 0$, trace $f_* = -2$ discussed above.

Now suppose F is horizontal. Then the restriction to F of $p_2: S^1 \times P^2 \rightarrow P^2$ is a covering projection, of degree $m \geq 1$, say. Then, with the same notation as used earlier, we see that (2.5.1) must hold, and the subsequent discussion shows that (2.5.2) holds for some ℓ_0 , that $p = |\ell|$ where $\ell = \ell_1 - \ell_0$, and thence that $\|K\| = \eta(S)/|\ell| = m/2|\ell|$.

The following theorem summarizes our conclusions.

Theorem 2.21. *Let M_f be a T^2 -bundle over S^1 with monodromy f not isotopic to the identity, and let K be an essential simple closed curve in a fiber.*

- (1) *If trace $f_* = 2$ then there is a unique K that has finite order in $H_1(M_f)$, and $\|K\| = 1/2p$ where f_* is represented by the matrix $\begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}$, $p \geq 1$.*
- (2) *If trace $f_* \neq 2$ then every K has finite order in $H_1(M_f)$, and*

$$\|K\| = |\gamma/2(\text{trace } f_* - 2)|$$

where f_ is represented by the matrix $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ with respect to an ordered basis of $H_1(T^2)$ whose first member is $[K]$.*

We see immediately from Theorem 2.21 that knots in fibers of torus bundles provide additional examples of knots with arbitrarily small rational genus. These examples are relevant to Theorem 5.19, cases (4) and (5). We now describe this in more detail, continuing the list in §2.4.

Case G. (M is a Seifert fiber space with no exceptional fibers and K is a fiber.) Let M be M_f where trace $f_* = 2$, as in part (1) of Theorem 2.21. Then M can also be described as an S^1 -bundle over T^2 with Euler number p , and K is a fiber. We remark

that M has a Nil geometric structure (see [24] pp. 467–470). By taking p large we can make $\|K\| = 1/2p$ arbitrarily small.

Case H. (M is a T^2 -bundle over S^1 with Anosov monodromy and K lies in a fiber.) Let M be M_f with $|\text{trace } f_*| > 2$, so the monodromy f is Anosov. Note that M has a Sol geometric structure (see [24], pp. 470–472). It is clear from the formula in part (2) of Theorem 2.21 that we can choose M and K so that $\|K\|$ is arbitrarily small. For example, choose any matrix $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \in \text{SL}(2, \mathbb{Z})$ with $\gamma \neq 0$ and let $f_n: T^2 \rightarrow T^2$ be given by the matrix $\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} \alpha+n\gamma & \beta+n\delta \\ \gamma & \delta \end{pmatrix}$ with respect to some basis (a, b) , say. Let M_n be the corresponding T^2 -bundle and let K_n be the knot in a fiber such that $[K_n] = a$. Then $\|K_n\| = |\gamma/2(\alpha + \delta + n\gamma - 2)| \rightarrow 0$ as $n \rightarrow \infty$.

Remark 2.22. It follows from Theorem 2.21 that if we are not in Case G or Case H, i.e. if $\text{trace } f_* = -2$ or $|\text{trace } f_*| \leq 1$, then either $\|K\| = 0$ or $\|K\| \geq 1/8$.

2.6. Knots in vertical tori in Seifert fiber spaces. In this section we analyze relative p -Seifert surfaces for knots that lie in essential vertical tori in Seifert fiber spaces. First we have the following lemma.

Lemma 2.23. *Let M be a Seifert fiber space with non-empty boundary and let $\pi: M \rightarrow \mathcal{B}$ be the projection of M onto its base orbifold \mathcal{B} . Let F be a horizontal surface in M and let k be the degree of the induced branched covering $\pi|_F: F \rightarrow \mathcal{B}$. If $\chi(F) < 0$ then $\chi(F) \leq -k/6$.*

Proof. Let q_1, \dots, q_n be the multiplicities of the exceptional fibers of M . Then (see for example [14], §2.1)

$$\chi(F) = k \left(\chi(\mathcal{B}) - \sum_{i=1}^n \left(1 - \frac{1}{q_i} \right) \right).$$

Since $\chi(F) < 0$, the maximal value of the expression in parentheses is attained when $\chi(\mathcal{B}) = 1$, $n = 2$, $q_1 = 2$, $q_2 = 3$, which gives the value $-1/6$. \square

Proposition 2.24. *Let M be a Seifert fiber space, T a vertical essential torus in M , K an essential simple closed curve in T , and F a relative p -Seifert surface for K . Then either*

- (1) $-\chi^-(F) \geq p/6$; or
- (2) K is an ordinary fiber in the Seifert fibration of M ; or
- (3) M contains a submanifold Q that is a twisted S^1 -bundle over the Möbius band and K is a fiber in Q .

Note that if (3) holds but (2) does not then the Seifert fibration of Q induced from M is the one with base orbifold a disk with two cone points of order 2.

Proof. Let $T^2 \times I$ be a regular neighborhood of $T = T^2 \times \{1/2\}$. First suppose that T separates M . Then $M = X_0 \cup T^2 \times I \cup X_1$, where $T_i = T^2 \times \{i\}$ is a component of ∂X_i , $i = 0, 1$. Let $N(K)$ be a regular neighborhood of K in $T^2 \times I$ and let $Y = T^2 \times I - \text{int } N(K)$. Then $X = M - \text{int } N(K) = X_0 \cup Y \cup X_1$. Let F be a good relative p -Seifert surface for K in M . Let $F_i = F \cap X_i$, $i = 0, 1$, and $G = F \cap Y$. We may assume that F_0 , F_1 and G are essential in X_0 , X_1 and Y respectively. The Seifert fibration of M induces Seifert fiber structures on X_0 and X_1 , with base orbifolds \mathcal{B}_0 and \mathcal{B}_1 , say. Recall from §2.5 that $Y \cong S^1 \times P^2$ where P^2 is a pair of pants.

Note that F_i is horizontal or vertical in X_i , $i = 0, 1$, and G is horizontal or vertical in $Y \cong S^1 \times P^2$. Write $\partial_i F_i = \partial F_i \cap T_i = \partial G \cap T_i = \partial_i G$, $i = 0, 1$.

Case I. G is vertical. Since a vertical annulus in Y that has both its boundary components on $\partial N(K)$ has these boundary components oriented oppositely on $\partial N(K)$, and since F is good, it follows that G consists of p parallel copies of an annulus with one boundary component on $\partial N(K)$ and the other on (say) T_0 , $F_1 = \emptyset$, and F_0 is connected.

Subcase (a). $\chi(F_0) < 0$. Then F_0 is horizontal in X_0 . Since $\partial_0 F_0$ has p components the index of the covering $F_0 \rightarrow \mathcal{B}_0$ is at least p . Therefore, by Lemma 2.23, $\chi(F) = \chi(F_0) \leq -p/6$.

Subcase (b). $\chi(F_0) = 0$. Then F_0 is an annulus. First suppose that $\partial_0 F_0$ has a single component. If F_0 is horizontal then $X_0 \cong F_0 \times S^1 \cong T_0 \times I$, contradicting the assumption that T is essential in M . If F_0 is vertical then K is an ordinary fiber in the Seifert fibration of M .

If $\partial_0 F_0$ has two components then F is an annulus with both boundary components on $\partial N(K)$ and by Theorem 2.11 K is contained in a submanifold N of M where N is a Seifert fiber space over the Möbius band with at most one orbifold point of order $r \geq 1$ and K is a fiber of multiplicity r . If $r > 1$ then, since the Seifert fibration of N is unique, K is an exceptional fiber in M . But this contradicts the fact that K is contained in a vertical torus. Hence $r = 1$ and we have conclusion (3).

This completes the proof in Case I.

Case II. G is horizontal. Here we will adopt the notation of §2.5. Let m be the index of the covering $G \rightarrow P^2$; so $\chi(G) = -m$. Also we have

$$\begin{aligned} [\partial_i F_i] &= [\partial_i G] = \ell_i a + m b_i, \quad i = 0, 1, \\ [\partial G \cap N(K)] &= (\ell_1 - \ell_0) a + m c, \end{aligned}$$

where $p = |\ell_1 - \ell_0|$.

Let φ_i be the Seifert fiber of M on T_i , $i = 0, 1$. Then $[\varphi_i] = \alpha a + \beta b_i$, say, $i = 0, 1$. If $\beta = 0$ then $[K] = [\varphi_i]$ so K is an ordinary fiber in the Seifert fibration of M . We will therefore assume that $\beta \neq 0$.

If F_i is horizontal in X_i , let k_i denote the index of the associated covering $F_i \rightarrow \mathcal{B}_i$, $i = 0, 1$. Then $k_i = |\partial_i F_i \cdot \varphi_i| = |(\ell_i a + m b_i) \cdot (\alpha a + \beta b_i)| = |\beta \ell_i - \alpha m|$, where \cdot denotes algebraic intersection number.

Sublemma 2.25. (1) *If both F_0 and F_1 are horizontal then $k_0 + k_1 \geq p$.*

(2) *If F_i is horizontal and $\chi(F_i) = 0$ then $k_i \leq m$.*

(3) *If F_0 is horizontal and F_1 is vertical then $k_0 \geq p$.*

Proof. (1) $k_0 + k_1 = |\beta \ell_0 - \alpha m| + |\beta \ell_1 - \alpha m| \geq |\beta| |\ell_1 - \ell_0| = |\beta| p \geq p$.

(2) Here F_i consists of parallel copies of a horizontal annulus A in X_i . If A has one boundary component on T_i and one on ∂M then $X_i \cong A \times S^1 \cong T_i \times I$, contradicting the assumption that T is essential in M . Also, since the components of $\partial_i G$ are coherently oriented on T_i the same holds for $\partial_i F_i$. It follows that A is non-separating and \mathcal{B}_i is a disk with two cone points of order 2. In particular each boundary component of A has intersection number 1 with the Seifert fiber φ_i . Therefore $k_i = |\partial_i F_i \cdot \varphi_i|$ is the number of components of $\partial_i F_i = \partial_i G$, which is $\leq m$ since G is an m -fold covering of P^2 .

(3) If F_1 is vertical then $[\partial_1 F_1] = \ell_1 a + m b_1 = s[\varphi_1] = s(\alpha a + \beta b_1)$ where s is the number of components of $\partial_1 F_1 = \partial_1 G$. Hence $s \leq m$. Now

$$\begin{aligned} s k_0 &= |\partial_0 F_0 \cdot s \varphi_0| \\ &= (\ell_0 a + m b_0) \cdot (\ell_1 a + m b_0) \\ &= m |\ell_1 - \ell_0| = m p. \end{aligned}$$

Therefore $k_0 = m p / s \geq p$. □

We now complete the proof of Proposition 2.24 in Case II.

First note that F_0 and F_1 cannot both be vertical, for then we would have $\ell_0 = \ell_1$ and hence $p = 0$.

Subcase (a). F_0 and F_1 horizontal. If $\chi(F_0), \chi(F_1) < 0$ then

$$\begin{aligned} |\chi(F)| &= |\chi(F_0)| + |\chi(F_1)| + |\chi(G)| \\ &\geq (k_0 + k_1)/6 + m, \text{ by Lemma 2.23} \\ &\geq p/6 + m, \text{ by Sublemma 2.25 (1)} \\ &> p/6. \end{aligned}$$

If $\chi(F_0) = 0$ and $\chi(F_1) < 0$ then $k_0 \leq m$ by Sublemma 2.25 (2), and hence $k_1 \geq p - m$ by Sublemma 2.25 (1). Therefore

$$\begin{aligned} |\chi(F)| &= |\chi(F_1)| + |\chi(G)| \\ &\geq k_1/6 + m \\ &\geq (p - m)/6 + m > p/6. \end{aligned}$$

Finally, if $\chi(F_0) = \chi(F_1) = 0$, then by Sublemma 2.25 parts (1) and (2), we have $p \leq k_0 + k_1 \leq 2m$. Hence

$$|\chi(F)| = |\chi(G)| = m \geq p/2.$$

Subcase (b). F_0 horizontal, F_1 vertical. If $\chi(F_0) < 0$ then

$$\begin{aligned} |\chi(F)| &= |\chi(F_0)| + |\chi(G)| \\ &\geq k_0/6 + m \\ &\geq p/6 + m, \text{ by Sublemma 2.25 (3)} \\ &> p/6. \end{aligned}$$

If $\chi(F_0) = 0$ then by Sublemma 2.25 (2) $k_0 \leq m$, while by Sublemma 2.25 (3) $k_0 \geq p$. Hence $|\chi(F)| = |\chi(G)| = m \geq p$.

This completes the proof of Proposition 2.24 when T separates M .

If T is non-separating, let $\pi: M \rightarrow \mathcal{B}$ be the projection from M to its base orbifold \mathcal{B} . Let N be a regular neighborhood of either the union of the exceptional fibers of M or, if M is closed and has no exceptional fibers, an ordinary fiber. Let $M_0 = M - \text{int } N$, with corresponding base orbifold \mathcal{B}_0 . Then M_0 is an S^1 -bundle over \mathcal{B}_0 and $T = \pi^{-1}(C)$ for some non-separating orientation-preserving simple closed curve C in \mathcal{B}_0 . Now $H_1(T)$ has basis φ, γ , where φ is the class of the S^1 -fiber of M_0 and $\pi_*(\gamma) = [C] \in H_1(\mathcal{B}_0)$. Therefore $[K] = r\varphi + s\gamma$ for some pair of relatively prime integers r, s .

Isotoping F to be transverse to the core of the components of N , let $F_0 = F \cap M_0$. Then F_0 defines a homology of pK into ∂M_0 . Therefore, considering the map $\pi_*: H_1(M_0, \partial M_0) \rightarrow H_1(\mathcal{B}_0, \partial \mathcal{B}_0)$, we have $0 = \pi_*(p[K]) = ps[C]$. Since C is orientation-preserving and non-separating, $[C]$ has infinite order in $H_1(\mathcal{B}_0, \partial \mathcal{B}_0)$, and so we conclude that $s = 0$. Therefore K is an ordinary fiber in the Seifert fibration of M . \square

3. Graphs

Several of our arguments concern the interaction between a relative p -Seifert surface F for a knot K in M , and another surface \hat{G} properly embedded in M . Such

arguments are handled in a uniform manner, by making the surfaces intersect as simply as possible, and then by analyzing cases depending on the combinatorics of this intersection. The arguments in this section are mostly combinatorial.

3.1. Graphs on surfaces. Fix the following notation. Let K denote a knot in M , let F be a good relative p -Seifert surface for K , and let \hat{G} be a properly embedded surface in M (usually a Heegaard surface, or an essential surface; usually of low complexity). Under such a circumstance, we perform the following procedure.

Isotop $N(K)$ so that it meets \hat{G} in n meridian disks, and let $G = \hat{G} \cap X$, so that F and G are both proper surfaces in X . After an isotopy, we may assume that F and G meet transversely in a finite disjoint union of circles and properly embedded arcs and that each of the q components of $\partial F \cap \partial N(K)$ meets each of the n components of $\partial G \cap \partial N(K)$ in r points, with notation as in Definition 2.5.

Formally cap off the components of $\partial F \cap \partial N(K)$ with disks to obtain a surface \hat{F} (note: if F is an honest p -Seifert surface, then \hat{F} is closed. Otherwise, $\partial \hat{F} = \partial F \cap \partial M$). The intersection $F \cap G$ determines graphs Γ_F and Γ_G in \hat{F} and \hat{G} respectively, where the vertices of Γ_F (resp. the vertices of Γ_G) correspond to the disks of $\hat{F} - F$ (resp. the disks of $\hat{G} - G$) and the edges correspond to the arc components of $F \cap G$ with at least one endpoint on $\partial N(K)$. We distinguish between two kinds of edges of Γ_F and Γ_G : *interior edges*, which have both endpoints on $\partial N(K)$ (i.e. at the vertices), and *boundary edges*, which have one endpoint on $\partial N(K)$, and the other on $\partial \hat{F} = F \cap \partial M$ or $\partial \hat{G} = G \cap \partial M$.

Choose orientations on F , G and X . This induces orientations on ∂F , ∂G and ∂X , and an arc of $F \cap G$ joins points of intersection of ∂F with ∂G of opposite sign.

Number the components of $\partial G \cap \partial N(K)$ (equivalently, the vertices of Γ_G) with the integers $1, 2, \dots, n$ in the (cyclic) order they occur along $\partial N(K)$. Hereafter an *index* means an element of this index set; i.e. an element $i \in \{1, \dots, n\}$. We imagine that the vertices of Γ_F and Γ_G are thickened, so that distinct edges end at distinct “edge-endpoints” on a (thickened) vertex. With this convention, an edge-endpoint at a vertex of Γ_F is a point of intersection of the corresponding component of $\partial F \cap \partial N(K)$ with a component of $\partial G \cap \partial N(K)$, and we label the edge-endpoint with the index corresponding to the label on the component of $\partial G \cap \partial N(K)$. Notice that it is the *vertices* of Γ_G and the *edge-endpoints* of Γ_F that are labeled with indices. Since F is good (by hypothesis), all components of $\partial F \cap \partial N(K)$ are coherently oriented on $\partial N(K)$, and therefore at each vertex of Γ_F we see the index labels $1, 2, \dots, n, 1, 2, \dots, n, \dots$ repeated r times in (say) anticlockwise order on the edge-endpoints around the vertex. Notice that if Γ_F and Γ_G have e_i interior edges and e_∂ boundary edges, then $2e_i + e_\partial = pn$.

In applications, the surface \hat{G} will always be either essential, or a Heegaard surface. In the former case we will choose $n = |K \cap \hat{G}|$ to be minimal; and in the latter case we will put K in thin position.

Remark 3.1. Thin position for knots in S^3 was introduced by Gabai [12], and for knots in arbitrary 3-manifolds by Rieck [20], [21], and we refer to these references for details. Technically, a knot is in thin position with respect to a sweepout of a 3-manifold (associated to a Heegaard splitting). The Heegaard surface \hat{G} is one of the nonsingular level sets of this sweepout, chosen depending on F .

Lemma 3.2. *With notation and conventions as above, we can arrange that no arc component of $F \cap G$ with both endpoints in $\partial N(K)$ is boundary parallel in either F or G . Equivalently, the graphs Γ_F and Γ_G have no monogon (disk) faces.*

Proof. The arguments are standard. Since components of ∂F are oriented coherently on $\partial N(K)$, every point of intersection of a given component of ∂G with ∂F has the same sign. Hence in particular, every interior edge of Γ_G has endpoints on distinct vertices of Γ_G , and there are never any complementary monogons.

If there is a monogon complementary to Γ_F then \hat{G} can be pushed over such a monogon by an isotopy, thereby reducing the number of intersections with K ; this is ruled out by hypothesis when \hat{G} is essential.

It remains to rule out monogon regions for Γ_F when \hat{G} is a Heegaard surface (note that such monogons *may* contain interior loops of $F \cap \hat{G}$ that bound compressing disks for \hat{G} ; see footnote 12 on page 635 of [21]). Such a monogon region is either a *high* disk or a *low* disk for \hat{G} , in the terminology of [21]. The existence of disjoint high and low disks at some level violates thinness; Gabai's argument in [12] (also see Theorem 6.2 in [21]) shows that for a knot in thin position, some level set of the sweepout admits neither. Choosing \hat{G} to be such a level set, Γ_F has no monogons. \square

Remark 3.3. If $F \cap G$ has a simple closed curve component that bounds a disk in G , let γ be an innermost such, i.e. γ bounds a disk D in G such that $F \cap (\text{int } D) = \emptyset$. Since F is incompressible in X , the loop γ bounds a disk E in F . Surgering F along D produces a 2-sphere $\Sigma = E \cup D$ together with a surface F' that is essential in X , has $\partial F' = \partial F$, and is homeomorphic to F , and which may be isotoped so that $|F' \cap G| < |F \cap G|$. If M is irreducible then Σ bounds a 3-ball and F' is isotopic to F . So in that case we may assume that no simple closed curve component of $F \cap G$ bounds a disk in G .

If \hat{G} is essential we will always assume that $n = |K \cap \hat{G}|$ is minimal over all essential surfaces \hat{G} in M of the given homeomorphism type. This implies that G is essential in X . Hence by the remarks above, interchanging the roles of F and G , we may assume that no simple closed curve component of $F \cap G$ bounds a disk in F .

Let Γ_F as above be a graph on \hat{F} without monogons. If every complementary region to Γ_F is a bigon, then either \hat{F} is a sphere, Γ_F has exactly two vertices, with parallel interior edges running between them, or \hat{F} is a disk, Γ_F has exactly one

vertex, with parallel boundary edges running from the vertex to the boundary. We call such a Γ_F a *beachball* (Figure 1 indicates why), of the *first kind* and *second kind* respectively.

Remark 3.4. If Γ_F is a beachball of the first kind then $\|K\| = 0$ and K satisfies conclusion (3) of Theorem 2.11.




Figure 1. A beachball of the first kind, with 18 complementary bigon regions.

Suppose Γ_F is not a beachball. The *reduced graph* $\bar{\Gamma}_F$ associated to Γ_F is obtained from Γ_F by collapsing all bigon regions. More generally, a reduced graph $\bar{\Gamma}$ in a surface \hat{F} is any graph with no complementary monogon or bigon regions.

Lemma 3.5. Let $\bar{\Gamma}$ be a reduced graph in \hat{F} with \bar{e} edges. Then $-\chi(F) \geq \bar{e}/3$.

Proof. Let \bar{v} be the number of vertices of $\bar{\Gamma}$, and \bar{f} the number of complementary disk faces. Non-disk faces contribute non-positively to Euler characteristic, so $\chi(\hat{F}) \leq \bar{v} - \bar{e} + \bar{f}$. Hence $\chi(F) = \chi(\hat{F}) - \bar{v} \leq \bar{f} - \bar{e}$. Since $\bar{\Gamma}$ is reduced, it has no monogon or bigon faces, so $2\bar{e} \geq 3\bar{f}$. Hence $3\chi(F) \leq 3\bar{f} - 3\bar{e} \leq -\bar{e}$. \square

Edges in Γ_F that cobound a bigon are said to be *parallel*. If Γ_F is complicated, either $-\chi(F)$ is large by Lemma 3.5, or else there are many parallel edges. The next lemma discusses the latter possibility. But first we introduce some terminology.

Notation 3.6. An interior edge of Γ_G that joins vertices with index labels i and j will be called an (i, j) -edge.

Lemma 3.7. If Γ_F contains $(mn + 1)$ parallel interior edges where $m \geq 1$, then there exists an index k such that

- (1) for each index i , the graph Γ_G has $2m$ edges which are $(i, k - i)$ -edges; and
- (2) for some index i_0 , the graph Γ_G has $(2m + 1)$ edges which are $(i_0, k - i_0)$ -edges.

Proof. Since all boundary components of F are oriented coherently on K (one says the vertices of Γ_F have the same *sign*), there is some (odd) index k such that the index labels of any edge in the family are i and $k - i$ (taken mod n). Since the family contains $(mn + 1)$ edges, there is an index label i_0 that appears $(m + 1)$ times at one end of the family, and m times at the other end; this proves the second claim. Moreover, any index label i appears at least m times at each end, proving the first claim. \square

Every edge of Γ_G is an arc of intersection of F with G , and therefore corresponds to an edge of Γ_F , and conversely. The next two lemmas control what happens when there are pairs of edges that are parallel on both graphs simultaneously.

Lemma 3.8. *Suppose there are interior edges that are parallel on both Γ_F and Γ_G . Then $(M, K) = (M', K') \# (\mathbb{RP}^3, \mathbb{RP}^1)$.*

Proof. Since all vertices of Γ_F have the same sign, this follows from the argument in [13], proof of Proposition 1.3. We observe that this argument is still valid if $F \cap G$ has simple closed curve components. \square

Lemma 3.9. *Suppose there are boundary edges that are parallel on both Γ_F and Γ_G . Then K is isotopic into ∂M .*

Proof. By [11], Lemma 2.5.4, such a pair of boundary edges gives an essential annulus A in X with one boundary component on ∂M and one on $\partial N(K)$, the latter having intersection number 1 with the meridian of K . Again this argument is valid in the presence of simple closed curve components of $F \cap G$. This annulus can be used to define an isotopy of K into ∂M . \square

3.2. Cables, satellites and tori. In the sequel, many arguments will depend on relativizing to a knot in a simple 3-manifold with boundary (i.e. a submanifold of M). In this section, we analyze the most important special cases.

Definition 3.10. Let K be a knot in M , with regular neighborhood $N(K)$. Let K' be a simple closed curve on $\partial N(K)$ that is essential in $N(K)$. Then we call K' a *cable* of K .

A knot K' contained in $N(K)$ is called a *satellite* of K if it is not contained in a 3-ball in $N(K)$. If $[K'] = k[K] \in H_1(N(K))$ then k is called the *winding number* of the satellite.

Remark 3.11. Note that our definitions of a satellite and of a cable include the trivial cases where K' is isotopic to K in $N(K)$.

Proposition 3.12. *Let K_0 be a knot in a 3-manifold M whose exterior has incompressible boundary, and let K be a satellite of K_0 with winding number $k > 0$. Then $\|K\| \geq k\|K_0\|$.*

Proof. By the definition of satellite, K is contained in a solid torus V in M whose core is K_0 . Let $X_0 = M - \text{int } V$. Let S be a good p -Seifert surface for K , $F = S \cap (V - \text{int } N(K))$, and $S_0 = S \cap X_0$.

If K_0 is p -trivial for some p the result is obvious. So we may assume that ∂V is incompressible in X_0 (see Remark 2.16). Thus ∂V is incompressible in $M - K$, and we may therefore assume that no component of S_0 or F is a disk. Hence $\eta(S) = \eta(F) + \eta(S_0)$.

In $H_1(V)$, there is equality $[\partial S_0] = [\partial S] = p[K] = pk[K_0]$. Therefore

$$\|K_0\| \leq \eta(S_0)/pk \leq \eta(S)/pk.$$

Since S can be chosen so that $\eta(S)/p$ is arbitrarily close to $\|K\|$, the result follows (or one can just apply Proposition 2.10). \square

Proposition 3.13. *Let M be a 3-manifold whose boundary contains a compressible torus T , and let K be a knot in M such that T is incompressible in $M - K$. Let F be a relative p -Seifert surface for K . Then either*

- (1) $-\chi^-(F) > p/6$; or
- (2) K is isotopic into T .

Proof. It is enough to prove the proposition under the assumption that F is good.

Let D be a compressing disk for ∂M in M , such that $D \cap N(K)$ consists of n meridian disks of $N(K)$, with n minimal. By hypothesis, $n > 0$. Let $X = M - \text{int } N(K)$ as usual.

Let $P = D \cap X$, a planar surface. By the minimality of n , the surface P is incompressible in X . Let Γ_F and Γ_P be the intersection graphs in \hat{F} and D respectively. There are two cases to consider.

Case A. ($n = 1$) In this case, P is an annulus, Γ_F and Γ_P have p edges, and all edges are boundary edges. In particular, all edges of Γ_P are parallel. If Γ_F has a pair of parallel edges then K is isotopic into T by the proof of Lemma 3.9. If not, then either Γ_F is a beachball of the second kind with a single edge, or the reduced graph $\bar{\Gamma}_F$ is defined and is equal to Γ_F . In the first case the annulus F defines an isotopy of K into T . In the second case $-\chi(F) \geq p/3$ by Lemma 3.5.

Case B. ($n > 1$) This case depends on two sublemmas.

Sublemma 3.14. *The graph Γ_F contains no family of $(\lfloor n/2 \rfloor + 1)$ parallel interior edges.*

Proof. Since all the vertices of Γ_F have the same sign, such a family would contain a (length 2) *Scharlemann cycle* (hereafter referred to as an *S-cycle*) i.e. a configuration of the form depicted in Figure 2. As in [22], Proposition 4.7, this *S-cycle* can be




Figure 2. An *S-cycle*.

used to tube and compress D , giving a new compressing disk D' with $\partial D' = \partial D$ and $|D \cap K| = n - 2$, contradicting minimality of n . \square

Sublemma 3.15. *If Γ_F contains $(2n - 1)$ parallel boundary edges then K is isotopic into T .*

Proof. In a family of $(2n - 1)$ parallel boundary edges, the labels (on the vertex at one end of the family) cycle through a full set of labels twice, with exactly one exception. Hence in Γ_P we get a pair of boundary edges at each vertex except (at most) one. Since D is a disk, a pair of boundary edges together with a common vertex separates D , so there is an outermost pair with the property that one of the complementary regions contains no other vertex of Γ_P . But this means that the outermost pair of boundary edges are in fact parallel, so we obtain a pair of boundary edges that are parallel in both Γ_F and Γ_P . The desired result now follows from the proof of Lemma 3.9. \square

We now complete the proof in Case B. First note that Sublemma 3.14 implies that Γ_F is not a beachball of the first kind. If Γ_F is a beachball of the second kind then by Sublemma 3.15 we may assume that $p = 1$. But then the boundary component of F that lies on $\partial N(K)$ intersects the meridian of K exactly once, and so F defines an isotopy of K into T . We may therefore suppose that the reduced graph $\bar{\Gamma}_F$ of Γ_F on \hat{F} exists, with, say, \bar{e}_i interior edges and \bar{e}_∂ boundary edges. By Sublemma 3.14 each interior edge of $\bar{\Gamma}_F$ corresponds to at most $\lfloor n/2 \rfloor$ edges of Γ_F , and by Sublemma 3.15 we may assume that each boundary edge of $\bar{\Gamma}_F$ corresponds to at most $(2n - 2)$ edges of Γ_F . Thus edges of $\bar{\Gamma}_F$ contribute at most n and $(2n - 2)$ to the sum of valences at all vertices of Γ_F respectively. Since this total sum is pn , and since $n \leq 2n - 2$, we get $\bar{e} = \bar{e}_i + \bar{e}_\partial \geq pn/(2n - 2) > p/2$. The conclusion now follows from Lemma 3.5. \square

Corollary 3.16. *Let K_0 be a knot in a 3-manifold M that is not m -trivial for any m , and let K be a non-trivial cable of a non-trivial cable of K_0 . Then $\|K\| > 1/12$.*

Proof. Let V be a regular neighborhood of K_0 containing K , and let $X_0 = M - \text{int } V$. By the hypothesis on K_0 , ∂V is incompressible in X_0 . Let S be a good p -Seifert

surface for K in M , such that $\|K\| = \eta(S)/p$, and let $F = S \cap V$ and $S_0 = S \cap X_0$. Then F is a relative p -Seifert surface for K in V . Since ∂V is incompressible in $M - K$, we may assume that no component of F or S_0 is a disk, and hence $\eta(S) = \eta(F) + \eta(S_0)$, giving $\eta(F) \leq \eta(S)$.

Since K is not a cable of K_0 , Proposition 3.13, applied to $K \subset V$, implies that $-\chi(F) > p/6$. Therefore $\|K\| = \eta(S)/p \geq \eta(F)/p > 1/12$. \square

The next proposition considers knots contained in a neighborhood of a torus.

Proposition 3.17. *Let K be a knot in $T^2 \times I$, and let F be a relative p -Seifert surface for K . Then either*

- (1) $-\chi^-(F) \geq p/3$; or
- (2) K is isotopic into $T^2 \times \{1/2\}$.

Proof. We assume F is good.

Put K in thin position with respect to the torus $\hat{T} := T^2 \times \{1/2\}$. Let $n = |K \cap \hat{T}|$, an even integer ≥ 0 . If $n = 0$ the second conclusion holds, so assume $n \geq 2$. Construct Γ_F, Γ_T without monogons, as in Lemma 3.2. Note that all edges are interior edges. We require a sublemma.

Sublemma 3.18. Γ_F does not contain a family of $(n/2 + 1)$ parallel edges.

Proof. Any such family contains an S -cycle. This gives rise to a Möbius band properly embedded in (say) $T^2 \times [1/2, 1]$, which is absurd. \square

By Sublemma 3.18, we obtain an estimate $\bar{e} \geq e/(n/2) = p$, where \bar{e} is the number of edges in the reduced graph $\bar{\Gamma}_F$. (Note that $\bar{\Gamma}_F$ exists, since otherwise Γ_F is a beachball of the first kind and K satisfies part (3) of Theorem 2.11. But this would give a Klein bottle embedded in $T^2 \times I$, which is absurd.) Now apply Lemma 3.5. \square

As a corollary, we deduce the following:

Corollary 3.19. *Let T be an incompressible torus in a 3-manifold M , and let K be a knot in M that lies in a regular neighborhood of T . Let F be a relative p -Seifert surface for K in M . Then either*

- (1) $-\chi^-(F) \geq p/3$; or
- (2) K is isotopic into T .

Proof. Let N be a regular neighborhood of T , and define $F' = F \cap N$, $F'' = F \cap (\overline{M - N})$. Since T is incompressible in M we may assume that F'' has no disk (or sphere) components. Therefore $-\chi^-(F) \geq -\chi^-(F')$, and the result follows from Proposition 3.17. \square

4. Hyperbolic knots

In this section we consider the case that $M - K$ is hyperbolic; i.e. that K is a *hyperbolic knot*. The arguments in this section use more geometry and analysis.

4.1. Stable commutator length

Definition 4.1. Let G be a group, and a an element in $[G, G]$. The *commutator length* of a , denoted by $\text{cl}(a)$, is the minimal number of commutators in G whose product is a , and the *stable commutator length*, denoted by $\text{scl}(a)$, is the limit

$$\text{scl}(a) = \liminf_{n \rightarrow \infty} \frac{\text{cl}(a^n)}{n}.$$

From the definition one sees that cl (and therefore also scl) is a characteristic function, and therefore in particular it is constant on conjugacy classes. The function scl can be extended to conjugacy classes which represent torsion elements in $H_1(G; \mathbb{Z})$ by the formula $\text{scl}(a) = \text{scl}(a^n)/n$ for any positive integer n .

For an introduction to stable commutator length and its properties, see [4] or [8].

There is a straightforward relationship between (stable) commutator length and norm, as follows.

Lemma 4.2. *Let M be a 3-manifold, and $K \subset M$ a knot. Let $a \in \pi_1(M)$ be an element in the conjugacy class determined by the free homotopy class of K . Then $\text{scl}(a) \leq \|K\|$.*

Proof. Proposition 2.10 from [8] says that $\text{scl}(a) = \inf_S -\chi^-(S)/2n$ where the infimum is taken over all oriented surfaces S mapping to M with boundary ∂S mapping to K with total degree n . If S is a p -Seifert surface for K , collapsing $N(K)$ to K wraps ∂S around K with total degree p . The result follows. \square

In general, $\text{scl}(a)$ can be smaller than $\|K\|$, since the infimum in the geometric definition of $\text{scl}(a)$ (in the proof of Lemma 4.2) is taken over *all* surfaces S in M which bound K , and not just *embedded* surfaces whose interior is disjoint from K . In other words, $\text{scl}(a)$ is the best lower bound on $\|K\|$ which can be estimated from the *homotopy* class of K . If $\|K\|$ is small, then $\text{scl}(a)$ is small and we will deduce information about the homotopy class of K from this.

4.2. Stable commutator length in hyperbolic 3-manifolds. There are strong interactions between geometry and scl , especially in dimension 3. The most interesting case is that of hyperbolic geometry, summarized in the following theorem:

Theorem 4.3 ([7], Theorem C). *For every $\epsilon > 0$ there is a constant $\delta(\epsilon) > 0$ such that, if M is a complete hyperbolic 3-manifold and a nontrivial $a \in \pi_1(M)$ has*

$\text{scl}(a) \leq \delta$, then either a is parabolic, or otherwise if γ is the unique geodesic in the free homotopy class associated to the conjugacy class of a ,

$$\text{length}(\gamma) \leq \epsilon.$$

The dependence of δ on ϵ is not proper: in every finite volume hyperbolic 3-manifold, conjugacy classes a with $\text{scl}(a) = 1/2$ correspond to arbitrarily long geodesics. However, if K is a knot with *sufficiently* small norm in a closed hyperbolic 3-manifold, Theorem 4.3 implies that K is homotopic to a power of the core geodesic of a Margulis tube, and the length of the geodesic can be bounded from above by a constant depending on $\|K\|$.

In more detail, recall that Margulis showed that in each dimension n , there is a positive constant $\epsilon(n)$ so that every geodesic in a closed hyperbolic n -manifold of length at most $\epsilon(n)$ is simple, and is contained in an embedded solid tube whose diameter can be estimated from below by a function of length. The exact value of the biggest constant $\epsilon(n)$ with this property is not known when $n > 2$, so for the sake of precision, we make the following definition.

Definition 4.4. A *Margulis tube* in a hyperbolic 3-manifold is an embedded solid tube of radius at least 0.531 around a simple geodesic (the *core* of the Margulis tube) of length at most 0.162286.

The precise choice of constants are somewhat arbitrary, but are chosen to be compatible with the estimates obtained by Hodgson–Kerckhoff [16]; see the discussion in the next subsection.

4.3. Deformation of cone-manifold structure. We have seen in the previous subsection that if K is a knot of sufficiently small norm, K is homotopic into the core of a Margulis tube.

We would like to conclude in fact that K is *isotopic* to (a cable of) the core of the tube. To do this we must use the fact that $\|K\|$ is small, not just scl of the corresponding conjugacy class in $\pi_1(M)$. To make use of this fact, we must study the geometry of $M - K$. In what follows we make use of some well-known results from the theory of hyperbolic cone manifolds (with non-singular cone locus). For a reference see e.g. [6] or [16].

We assume throughout this section that $M - K$ is hyperbolic. Then as Thurston already showed (see e.g. [27]), for all sufficiently small positive real numbers θ , there exists a hyperbolic cone manifold M_θ , unique up to isometry, whose underlying manifold is homeomorphic to M , and whose cone locus is a geodesic in the isotopy class of K with cone angle equal to θ .

The manifolds M_θ can be deformed (by increasing θ) until one of the following happens:

- (1) The cone angle can be increased all the way to 2π , and one obtains a complete hyperbolic structure on M for which K is isotopic to a geodesic.
- (2) The volume goes to 0 (and either converges after rescaling to a Euclidean cone manifold, or the injectivity radius goes to 0 everywhere after rescaling to have a fixed diameter).
- (3) The cone locus bumps into itself (this can only happen for $\theta > \pi$).

In the second or third case we say that the cone manifold structure becomes singular. Under deformation, the length of the cone geodesic strictly increases. For each θ , let $l(\theta)$ denote the length of the cone geodesic isotopic to K in M_θ , and let $R(\theta)$ be the radius of a maximal open embedded tube around the cone geodesic.

Following Hodgson–Kerckhoff [16] we define

$$h(R) := 3.3957 \frac{\tanh(R)}{\cosh(2R)}$$

which is non-negative for positive R , is asymptotic to 0 as R goes to 0 or to ∞ , and which has a single maximum value ≈ 1.019675 , achieved at $r \approx 0.531$.

The following is proved in §5 of [16]:

Theorem 4.5 (Hodgson–Kerckhoff). *Let $h(R)$ be as above. Let $l(\theta)$ be the length of the singular geodesic in M_θ . Then M_θ can be deformed (by increasing θ) either until $\theta = 2\pi$, or until $\theta \cdot l$ is equal to the maximum of $h(R)$ (which occurs at approximately $R = 0.531$ and is equal to approximately $h(0.531) = h_{\max} = 1.019675$) and for all smaller values of θ , the radius of a maximal embedded tube about the cone geodesic is at least 0.531.*

It follows that we can deform M_θ either all the way to $\theta = 2\pi$ with $l \leq 0.162286$ and $R \geq 0.531$, in which case K is isotopic to the core of a Margulis tube, or else we can deform M_θ until $\theta \cdot l = h(0.531) \approx 1.019675$ for some $\theta < 2\pi$. In the second case we can estimate $l(\theta) \geq 1.019675/2\pi \approx 0.162286$ and $R(\theta) \geq 0.531$. In the next few sections we will obtain a positive lower bound on the rational genus of K in the second case.

4.4. 1-forms from tubes. Let M_θ be a cone manifold, with a cone geodesic γ with length l and tube radius R . Let T denote an embedded tube around γ whose radius is R , and let $p: T \rightarrow \gamma$ denote radial projection. Let $\phi: \gamma \rightarrow \mathbb{R}/l \cdot \mathbb{Z}$ be a parameterization of γ so that $d\phi$ is the length form on γ . Pulling back the 1-form $d\phi$ by p^* defines a 1-form on all of T which, by abuse of notation, we denote by $d\phi$. Let $r: T \rightarrow [0, R]$ be the function on T which is equal to the radial distance to γ . Define a 1-form α on M_θ by

$$\alpha = d\phi \cdot (\sinh(R) - \sinh(r))$$

on T , and extend it by 0 outside T . Let β be a C^∞ function on $[0, R]$ taking the value 1 in a neighborhood of 0 and the value 0 in a neighborhood of R , and satisfying $|\beta'| < 1/(R - \epsilon)$ throughout $[0, R]$, for some small fixed ϵ . Finally define $\alpha_\epsilon = \beta(r)\alpha$. Then the form α_ϵ is C^∞ on $M - \gamma$, and satisfies $\|d\alpha_\epsilon\| \leq 1 + 1/(R - \epsilon)$ pointwise. Moreover, the integral of α_ϵ along γ is $l \cdot \sinh(R)$. For a proof of these estimates, see Lemma 4.3 from [7].

4.5. Wrapping. Suppose K is a knot of sufficiently small rational genus such that $M - K$ is hyperbolic. If M is hyperbolic, and K is isotopic to an embedded geodesic, then K is isotopic to the core of a Margulis tube (whose length may be estimated from above in terms of $\|K\|$, by Theorem 4.3). Otherwise, we can find a hyperbolic cone manifold structure M_θ on the underlying topological manifold M , with cone angle $\theta < 2\pi$ along a single cone geodesic in the isotopy class of K , whose length is at least 0.162286, and is contained in an embedded tube whose radius is at least 0.531. For each $\epsilon > 0$, let α_ϵ be a 1-form constructed as in §4.4. Let S be a good p -Seifert surface for K in M_θ realizing $\|K\|$, and let S'' be another (possibly immersed) surface, homotopic to S rel. boundary, with interior disjoint from K . For each ϵ ,

$$p \cdot l \cdot \sinh(R) = \int_{\partial S''} \alpha_\epsilon = \int_{S''} d\alpha_\epsilon \leq \text{area}(S'') \cdot (1 + 1/(R - \epsilon)).$$

Taking $\epsilon \rightarrow 0$ we obtain an estimate

$$l \cdot \sinh(R) \cdot R/(R + 1) \leq \text{area}(S'')/p. \quad (4.5.1)$$

By the discussion above, we can estimate

$$0.03131 \approx 0.162286 \cdot \sinh(0.531) \cdot 0.531/1.531 \leq l \cdot \sinh(R) \cdot R/(R + 1).$$

We claim that one can find a representative surface S'' homotopic to S rel. boundary in $M - K$, of area at most $\epsilon - 2\pi\chi(S)$ for any $\epsilon > 0$. This will imply that $\|K\| \geq 0.03131/4\pi \approx 2.491 \times 10^{-3}$.

The representative surface S'' is obtained by *wrapping*; there are two (essentially equivalent) methods to construct a “wrapped” surface: the shrinkwrapping method from [9], and the PL wrapping method from [25]. For technical ease, we use the PL wrapping method. Roughly speaking, given a surface S in a hyperbolic 3-manifold and a prescribed family Γ of geodesics, the PL wrapping technique finds a $\text{CAT}(-1)$ representative in the homotopy class of S , which can be approximated by surfaces homotopic to S in the complement of Γ .

Our situation is analogous to, but not strictly equivalent to, the situation in [25] or [9]. In our context Γ will be a singular cone geodesic, in the isotopy class of K , in a cone manifold structure on M ; and the surface S will have boundary wrapping some number of times around Γ but interior disjoint from Γ . In fact, this extra complication

does not add any difficulty to the argument, and we will obtain the same conclusion – namely, the existence of a $\text{CAT}(-1)$ surface which can be approximated by surfaces homotopic to S (rel. boundary) in the complement of Γ . For the sake of completeness, we explain the construction in detail.

The key point of the construction is that the universal cover of $M - \Gamma$ has a metric completion which is intrinsically $\text{CAT}(-1)$; this is Lemma 1.2 from [25]. Let N denote the universal cover of $M - \Gamma$, and \bar{N} its metric completion. Notice that \bar{N} is obtained from N by adding geodesics which project to components of Γ .

Let T be a triangulation of S with all vertices on ∂S . After an isotopy, we can assume that the map $\partial S \rightarrow \Gamma$ takes all vertices of the triangulation to distinct points in Γ .

The interior of each edge e of T not on ∂S lifts to an open interval in N whose closure is a closed interval \tilde{e} in \bar{N} . Since \bar{N} is $\text{CAT}(-1)$, there is a unique geodesic \tilde{e}' in \bar{N} with the same endpoints as \tilde{e} . This projects to a piecewise geodesic segment e' in M with vertices on Γ (note that the projection e' does not depend on the choice of the lift \tilde{e}). For each triangle Δ of T , we can choose lifts \tilde{e}_i of the edges e_i which together span a triangle $\tilde{\Delta}$ in \bar{N} . After replacing each \tilde{e}_i with a geodesic \tilde{e}'_i , we straighten $\tilde{\Delta}$ to a piecewise linear surface by coning one vertex to the points on the opposite edge; i.e. if v is the vertex opposite \tilde{e}'_1 (say), for each point p on \tilde{e}'_1 there is a unique geodesic in \bar{N} from v to p , and the union of these geodesics is a piecewise totally geodesic disk $\tilde{\Delta}'$ spanning the union of the \tilde{e}'_i . Let Δ' denote the projection of $\tilde{\Delta}'$. Then the union of the Δ' is a $\text{CAT}(-1)$ surface S' which can be approximated by surfaces whose interior is disjoint from Γ , and which are homotopic to S through surfaces with interior disjoint from Γ . The approximating surfaces S'' can be chosen to have area as close to the area of S' as desired; since S' is $\text{CAT}(-1)$, by Gauss–Bonnet, we have $\text{area}(S'') \leq \epsilon - 2\pi\chi(S)$, as claimed.

Remark 4.6. The argument in [25] uses extra hypotheses on S , namely that it is incompressible and 2-incompressible rel. Γ . In fact, these hypotheses are superfluous in the case that S has boundary. In fact, even when S is a closed surface, one really only needs to know that S contains *some* embedded loop which is covered by a nondegenerate infinite geodesic in \bar{N} .

Remark 4.7. The reader familiar with the construction of pleated surfaces in hyperbolic or $\text{CAT}(-1)$ spaces will recognize the similarity with PL wrapping.

In fact, for our applications, it is important to construct surfaces S'' as above when M has boundary consisting of tori, and S is a relative p -Seifert surface for K . There are no extra difficulties in this case. One can proceed by either of two methods. The easiest is to construct the PL wrapped surface directly. The triangulation of the relative p -Seifert surface can include vertices on a “cusp” boundary component; one just needs to observe that semi-infinite rays in \bar{N} have geodesic representatives

(constructed e.g. by taking limits of sequences of geodesic arcs with one endpoint going to infinity along the ray). Triangles with two vertices on a single “cusp” boundary component degenerate to a geodesic ray under straightening. The resulting surface, while non-compact, is complete, and satisfies $\text{area} \leq -2\pi\chi$.

Alternately, one can deform the metric in a neighborhood of infinity to make it $\text{CAT}(0)$, in such a way that each end is foliated as a metric product by Euclidean totally geodesic tori; such a metric is described explicitly in the proof of Lemma 7.12 in [9]. One obtains a compact PL wrapped surface as above with some boundary components on Γ , and some on a fixed family of Euclidean tori, one for each cusp component of M . The PL wrapped surface so obtained is $\text{CAT}(0)$, and its restriction to any prescribed compact region of M can be taken to be $\text{CAT}(-1)$; in particular, we can assume that the surface is $\text{CAT}(-1)$ in the support of the 2-form $d\alpha_\epsilon$, so that the area of the part of the surface in the support of $d\alpha_\epsilon$ is at most $-2\pi\chi(S)$, and we obtain the desired bound on $-\chi(S)$.

We have therefore obtained a proof of the following theorem:

Theorem 4.8. *Let K be a knot in a 3-manifold M , possibly with boundary consisting of a union of tori. Suppose $M - K$ is hyperbolic. Then either*

- (1) $\|K\| \geq 1/402$; or
- (2) M is hyperbolic and K is isotopic to the core of a Margulis tube.

Moreover, if F is a relative p -Seifert surface for K in M , then either

- (1) $\eta(F)/p \geq 1/402$; or
- (2) M is hyperbolic and K is isotopic to the core of a Margulis tube.

4.6. Better estimates. In fact, though the wrapping technique explains in a direct geometric way the relationship between $\|K\|$ and the geometry of K in M , one can obtain better estimates at the cost of appealing to some more refined technology of Agol and Cao–Meyerhoff (which we treat as a black box). For the benefit of the reader we include a sketch of a proof of the following:

Proposition 4.9. *Let K be a knot in a 3-manifold M . Suppose $M - K$ is hyperbolic. Then either*

- (1) $\|K\| \geq 1/50$; or
- (2) M is hyperbolic and K is isotopic to the core of a Margulis tube.

Proof. Let S be a p -Seifert surface for K . Let T be a maximal horospherical cusp torus in $M - K$; we may regard S as satisfying $\partial S \subset T$. Since S is essential, Theorem 5.1 from [1] gives $|\chi(S)| \geq \ell(\partial S)/6$ where ℓ denotes Euclidean length measured on T . Note that $\ell(\partial S) = q\ell(\sigma)$, where q is the number of boundary

components of S , and σ is the boundary slope. Also, $p = q\Delta(\sigma, \mu)$, where $\Delta(\sigma, \mu)$ is the geometric intersection number of σ with the meridian slope μ . Hence

$$|\chi(S)|/2p \geq \ell(\partial S)/12p = q\ell(\sigma)/12q\Delta(\sigma, \mu) = \ell(\sigma)/12\Delta(\sigma, \mu).$$

Let A be the area of T . Then by the proof of Theorem 8.1 from [1],

$$A \leq \ell(\sigma)\ell(\mu)/\Delta(\sigma, \mu).$$

Hence

$$|\chi(S)|/2p \geq A/12\ell(\mu) \geq \sqrt{A}/12\ell_N(\mu)$$

where $\ell_N(\mu) := \ell(\mu)/\sqrt{A}$ is the *normalized length* of μ (see [10]). By [10] one knows without hypothesis that $A \geq 3.35$. If we define

$$C := \sqrt{3.35}/12(7.515) \sim 0.0203 > 1/50$$

then we conclude

$$|\chi(S)|/2p < C \text{ implies that } \ell_N(\mu) \geq 7.515$$

and therefore by [10] p. 410 we deduce that M is hyperbolic and K is isotopic to the core of a Margulis tube (i.e. a geodesic of length ≤ 0.162 with tube radius ≥ 0.531). \square

It is therefore probably safe to replace $1/402$ by $1/50$ throughout the sequel by appealing to Proposition 4.9 in place of Theorem 4.8 (it is, however, unlikely that $1/50$ is sharp).

5. General knots

We are now in a position to discuss the most general case of a knot K in a closed, orientable 3-manifold M such that $[K]$ has finite order in $H_1(M)$. The discussion is case-by-case, and depends on the (well-known) prime and JSJ decomposition theorems.

The first step is to consider the interaction of (M, K) with the essential 2-spheres in M . Such spheres are treated by the following theorem.

Theorem 5.1. *Let K be a knot in a reducible manifold M . Then either*

- (1) $\|K\| \geq 1/12$; or
- (2) *there is a decomposition $M = M' \# M''$, $K \subset M'$ and either*
 - (a) *M' is irreducible, or*

$$(b) (M', K) = (\mathbb{RP}^3, \mathbb{RP}^1) \# (\mathbb{RP}^3, \mathbb{RP}^1).$$

Note that in case (2) (b), $\|K\| = 0$.

Proof. Let S be a good p -Seifert surface for K .

First assume that $M - K$ is irreducible. Let Σ be an essential 2-sphere in M , chosen so that $n = |\Sigma \cap K|$ is minimal. Since $[K]$ has finite order in $H_1(M)$, the algebraic intersection number of K and Σ is zero; thus n is even and > 0 .

Let P be the planar surface $\Sigma - \text{int } N(K)$. Let Γ_S, Γ_P be the intersection graphs in \hat{S} and Σ respectively, defined by the arc components of $S \cap P$.

Case A. ($n \geq 4$) An *extended S-cycle* is a configuration of the form depicted in Figure 3; i.e. a series of four parallel edges, whose middle pair form an ordinary S -cycle.




Figure 3. An extended S -cycle.

In this context, and with the assumption that $n \geq 4$, Lemma 2.3 from [28] says precisely that Γ_S does not contain an extended S -cycle. Hence Γ_S does not contain a family of $(n/2 + 3)$ parallel edges. Let $\bar{\Gamma}_S$ denote the reduced graph of Γ_S . Since $e = pn/2$, we can estimate $\bar{e} \geq p(n/2)/(n/2 + 2) \geq p/2$ (because $n \geq 4$), where \bar{e} denotes the number of edges of $\bar{\Gamma}_S$. Hence by Lemma 3.5, we have $-\chi(S) \geq p/6$ and therefore $\|K\| \geq 1/12$.

Case B. ($n = 2$) Here P is an annulus, so all edges of Γ_P are parallel (i.e. Γ_P is a beachball).

If Γ_S has a pair of parallel edges, consider an innermost such pair e_1, e_2 , i.e. e_1 and e_2 cobound a bigon face of Γ_S . The corresponding edges of Γ_P are necessarily parallel, and it follows that their endpoints, x_1 and x_2 , say, at a given vertex of Γ_P are diametrically opposite, in the sense that there are the same number of edge-endpoints in each of the two intervals around the vertex on either side of the pair $\{x_1, x_2\}$; see Figure 4 in [13]. Therefore Γ_S cannot have three mutually parallel edges.

If Γ_S is a beachball with exactly two edges, the argument in the proof of Proposition 1.3, case (1) from [13] shows that Σ decomposes (M, K) as a connected sum $(\mathbb{RP}^3, \mathbb{RP}^1) \# (\mathbb{RP}^3, \mathbb{RP}^1)$, which we think of as a degenerate case of case (2) (b) in the statement of the theorem.

We may therefore assume that Γ_S is not a beachball, and that no family of parallel edges in Γ_S has more than two edges. Consequently the reduced graph $\bar{\Gamma}_S$ exists and its number of edges $\bar{e} \geq p/2$, so Lemma 3.5 gives $\chi(S) \leq -p/6$, and hence $\|K\| \geq 1/12$. This completes the analysis when $M - K$ is irreducible.

If $M - K$ is reducible, we can write $M = M' \# M''$ where $K \subset M'$ and $M' - K$ is irreducible. Let Σ' be a 2-sphere in $M - K$ realizing the connected sum decomposition $M - K = (M' - K) \# M''$. By surgering S along the curves of intersection $S \cap \Sigma'$ we get a p -Seifert surface S' for K in M' , with $-\chi^-(S') \leq -\chi^-(S)$. The theorem now follows from the argument in the case that $M - K$ is irreducible. \square

After Theorem 5.1 we may assume that M is irreducible. Moreover, if $M - K$ is reducible then K lies in a 3-ball in M and so either $\|K\| \geq 1/2$ or K bounds a disk (in which case $\|K\| = 0$). We therefore assume in the sequel that both M and $M - K$ are irreducible.

A closed, irreducible 3-manifold M is either S^3 , a lens space, an atoroidal Seifert fiber space over S^2 with three exceptional fibers, hyperbolic, or toroidal. The next theorem treats the case that M is a lens space.

Theorem 5.2. *Let K be a knot in a lens space M . Then either*

- (1) $\|K\| \geq 1/24$; or
- (2) K lies on a Heegaard torus in M ; or
- (3) M is of the form $L(4k, 2k - 1)$ and K lies on a Klein bottle in M as a non-separating orientation-preserving curve.

Note that in case (3) we have $\|K\| = 0$; this is a special case of Theorem 2.11 (3).

Proof. Let \hat{T} be a Heegaard torus in M . Then either (2) holds, or we can put K in thin position with respect to \hat{T} . Assuming the latter, let $n = |K \cap \hat{T}|$; so n is even and ≥ 2 . Let S be a good p -Seifert surface for K in M . We then get intersection graphs Γ_S, Γ_T . Thin position of K and incompressibility of S guarantee by Lemma 3.2 that Γ_S and Γ_T have no monogon disk faces.

We need a sublemma.

Sublemma 5.3. *Suppose there is a pair of edges that are parallel on both Γ_S and Γ_T . Then $M = \mathbb{RP}^3$, and either*

- (1) $\|K\| \geq 3/4$; or
- (2) $(M, K) = (\mathbb{RP}^3, \mathbb{RP}^1)$.

Proof. By Lemma 3.8 we have $(M, K) = (M', K') \# (\mathbb{RP}^3, \mathbb{RP}^1)$. Since M is a lens space we must have $M' = S^3$. If K' is trivial we have (2). Suppose K' is non-trivial. Then $\|K'\| \geq 1/2$ and, since \mathbb{RP}^1 in \mathbb{RP}^3 is 2-trivial, Theorem 2.17 implies that $\|K\| = \|K'\| - \frac{1}{4} + \frac{1}{2} \geq \frac{3}{4}$. \square

Note that in case (2) (i.e. $(M, K) = (\mathbb{RP}^3, \mathbb{RP}^1)$) we have $\|K\| = 0$.

If Γ_S has $(2n + 1)$ parallel edges then by Lemma 3.7 bullet (2) there are indices i_0 and k so that Γ_T has 5 edges labeled $(i_0, k - i_0)$. Remember that this means that there are vertices i_0 and $k - i_0$ in Γ_T that are joined to each other by at least 5 edges. On a torus, one can find at most four embedded pairwise non-parallel arcs joining two points, that are disjoint except at their endpoints. To see this, “engulf” one of the edges by a disk, and observe that it is equivalent to show that there are at most three pairwise non-parallel essential embedded loops that intersect each other in one point; this latter fact can be shown using intersection number. Consequently we can deduce that two of the $(i_0, k - i_0)$ -edges must be parallel in \hat{T} .

If they are parallel in Γ_T (i.e. if they cobound an embedded complementary bigon) then there is a pair of edges that are parallel in both graphs, and Sublemma 5.3 applies. If not, then the disk D in \hat{T} realizing the parallelism of the two edges must contain vertices of Γ_T in its interior. By Lemma 3.7 bullet (1), these vertices come in pairs i , $k - i$, and each pair are joined by 4 edges. An easy innermost argument in the disk D shows that, for some i , some pair of $(i, k - i)$ edges are parallel in Γ_T . Hence again we get edges that are parallel in both graphs.

By Sublemma 5.3 we may therefore suppose that there is no family of $(2n + 1)$ parallel edges in Γ_S . If Γ_S is not a beachball then the reduced graph $\bar{\Gamma}_S$ exists and satisfies $\bar{e} \geq e/2n = pn/4n = p/4$. Hence $-\chi^-(S) \geq p/12$ by Lemma 3.5, giving $\|K\| \geq 1/24$.

If Γ_S is a beachball then as observed in Remark 3.4, K satisfies conclusion (3) of Theorem 2.11, i.e. K is the fiber of multiplicity r in a Seifert fiber subspace N of M whose base orbifold is a Möbius band with one orbifold point, of order $r \geq 1$. Since M is a lens space, $\overline{M - N}$ is a solid torus, and the meridian of this solid torus is not the Seifert fiber of N . Hence M is a Seifert fiber space with orbifold \mathbb{RP}^2 and one orbifold point of order $k \geq 1$, where $k = r$ if $r > 1$. Thus M is the lens space $L(4k, 2k - 1)$. If $r > 1$ then K lies on a Heegaard torus in M . If $r = 1$ then K satisfies conclusion (3). \square

Remark 5.4. The examples in Case B of §2.4 show that for any $\epsilon > 0$ there exists a lens space M and a knot K lying on a Heegaard torus in M with $0 < \|K\| < \epsilon$.

Remark 5.5. The bound $1/24$ in Theorem 5.2 is almost certainly not best possible. The smallest value of $\|K\|$ that we know of for a knot K in a lens space not satisfying (2) or (3) of Theorem 5.2, comes from the following example.

Let K' be the $(-2, 3, 7)$ -pretzel knot in S^3 . Then 19-surgery on K' gives a knot K in the lens space $L(19, 7)$. By Lemma 2.15, $\|K\| = (g(K') - 1/2)/19 = (5 - 1/2)/19 = 9/38$. Since $M - K \cong S^3 - K'$ is hyperbolic, K does not lie on a Heegaard torus in M . In fact K' is a Berge knot, and so K is 1-bridge in M .

Remark 5.6. Baker has shown in [2] that if K is a knot in a lens space M such that $\|K\|$ is realized by a p -Seifert surface S with a single boundary component then

either K is 1-bridge in M or $\|K\| > 1/4$. By the proof of Lemma 2.15, the hypothesis holds for pairs (M, K) that come from surgery on a knot in S^3 (or any homology sphere).

Remark 5.7. In our proof of Theorem 5.2, the argument shows that if $n > 2$ then Γ_S cannot have $(n + 1)$ parallel edges. Hence (1) can be improved to $\|K\| \geq 1/12$ if (2) is weakened to say that K is 0- or 1-bridge in M .

The next proposition considers knots in hyperbolic 3-manifolds, possibly with boundary. The case that the complement of the knot is hyperbolic was already treated in Theorem 4.8. Here we treat the general case.

Proposition 5.8. *Let M be a hyperbolic 3-manifold, possibly with boundary consisting of a union of tori. Let K be a knot in M such that $M - K$ is irreducible, and let F be a relative p -Seifert surface for K . Then either*

- (1) $\eta(F)/p \geq 1/402$; or
- (2) K is isotopic to a cable of a core of a Margulis tube; or
- (3) K is isotopic into ∂M .

Corollary 5.9. *Let K be a knot in a closed hyperbolic 3-manifold M . Then either*

- (1) $\|K\| \geq 1/402$; or
- (2) K is trivial; or
- (3) K is isotopic to a cable of the core of a Margulis tube.

Proof of Proposition 5.8. If $M - K$ is hyperbolic then the result follows from Theorem 4.8. So we may assume that $M - K$ is toroidal.

Since M is hyperbolic, every essential torus in $M - K$ is separating, so there exists an *extremal* such torus, i.e. an essential torus T in $M - K$ such that the component X_0 of M cut along T that does not contain K is atoroidal. Let $N = \overline{M - X_0}$; thus K is contained in N .

Let F be a good relative p -Seifert surface for K in M , and let $F' = F \cap N$, $F_0 = F \cap X_0$. Since T is incompressible in $M - K$, and $M - K$ is irreducible, we may assume that no component of F' or F_0 is a disk. Hence $\eta(F) = \eta(F') + \eta(F_0)$.

Since M is atoroidal, either T compresses in N or N is a product $T^2 \times I$ where $T^2 \times \{0\}$ is a component of ∂M and $T^2 \times \{1\} = T$. In the latter case, Proposition 3.17 implies that either $\eta(F) \geq \eta(F') \geq p/6$ or K is isotopic into ∂M .

We may therefore assume that T compresses in N . By Proposition 3.13 either $\eta(F')/p > 1/12$ or K is isotopic into T . So we may suppose the latter holds.

First suppose that N is a solid torus. Let K_0 be the core of N . Then K is a non-trivial cable of K_0 . The exterior of K_0 in M is X_0 , which is atoroidal. Since

M is hyperbolic X_0 cannot be Seifert fibered, and therefore X_0 is hyperbolic. By Proposition 3.12 $\|K\| \geq 2\|K_0\|$. Hence either we have conclusion (1) or $\|K_0\| < 1/804$, in which case K_0 is isotopic to the core of a Margulis tube by Theorem 4.8.

If N is not a solid torus then X_0 lies in a 3-ball in M . Since K is isotopic into $T = \partial X_0$, K also lies in a 3-ball, contradicting our assumption that $M - K$ is irreducible. \square

It remains to consider toroidal manifolds, and small Seifert fiber spaces. The small Seifert fiber spaces are treated by the following theorem. Recall that a *prism manifold* is a Seifert fiber space M with base S^2 and three exceptional fibers of multiplicities 2, 2, n . Then M has another Seifert fiber structure with base \mathbb{RP}^2 and at most one exceptional fiber.

Theorem 5.10. *Let M be an atoroidal Seifert fiber space over S^2 with three exceptional fibers and let K be a knot in M . Then either*

- (1) $\|K\| \geq 1/402$; or
- (2) K is trivial; or
- (3) K is a cable of an exceptional Seifert fiber of M ; or
- (4) M is a prism manifold and K is a fiber in the Seifert fiber structure of M over \mathbb{RP}^2 with at most one exceptional fiber.

Note that in case (4) we have $\|K\| = 0$. This is a special case of assertion (3) in Theorem 2.11.

Proof. We may assume that X (i.e. $M - \text{int } N(K)$) is irreducible.

If X is hyperbolic then (1) holds by Theorem 4.8.

If X is Seifert fibered then (since M is irreducible) the Seifert fibration of X extends to M . Thus K is a Seifert fiber of this fibration. If M is not a prism manifold then the Seifert fiber structure on M is unique and (since an ordinary fiber is a cable of an exceptional fiber) we have conclusion (3). If M is a prism manifold then the two possible Seifert fiber structures on M give either (3) or (4).

We may therefore assume that X is toroidal. Since M is irreducible and atoroidal, every torus in X is separating. As in the proof of Proposition 5.8, let T be an extremal essential torus in X , so $M = X_0 \cup_T N$ where X_0 is atoroidal. Again as in the proof of Proposition 5.8, we may assume that K is isotopic into T , and that N is a solid torus, with core K_0 . By Proposition 3.12, $\|K\| \geq 2\|K_0\|$. If $M - K_0$ is hyperbolic, and (1) does not hold, then $\|K_0\| < 1/804$ and so M is hyperbolic by Theorem 4.8, a contradiction.

Hence we may assume that X_0 is Seifert fibered. The Seifert fibration of X_0 extends to M , so K_0 is a fiber in the fibration of M , and K is a non-trivial cable of K_0 . Since an ordinary fiber is a non-trivial cable of an exceptional fiber, by Corollary 3.16 either $\|K\| > 1/12$ or K_0 is an exceptional fiber. \square

The toroidal case involves several subcases, which are treated in the next few propositions, culminating in Theorem 5.19.

Proposition 5.11. *Let M be an irreducible, toroidal 3-manifold, and let K be a knot in M such that $M - K$ is irreducible and atoroidal. Let F be a relative p -Seifert surface for K . Then $-\chi^-(F) \geq p/12$.*

Proof. We may suppose that F is good.

Let \hat{T} be an essential torus in M such that $n = |K \cap \hat{T}|$ is minimal (over all essential tori in M). Note that $n > 0$ since $M - K$ is atoroidal. Also, the existence of F shows that $p[K] = 0 \in H_1(M, \partial M)$, which implies that n is even.

Let T be the punctured torus $\hat{T} \cap (M - \text{int } N(K))$ and let Γ_F, Γ_T be the intersection graphs in \hat{F} and \hat{T} respectively. Note that all edges are interior edges.

We need some sublemmas.

Sublemma 5.12. *If $n \geq 4$ then Γ_F contains no family of more than $(n/2 + 2)$ parallel edges.*

Proof. Since all vertices of Γ_F are of the same sign, a family of more than $(n/2 + 2)$ parallel edges would contain an extended S -cycle. But this is impossible, again by Lemma 2.3 from [28] (compare with Case A in the proof of Theorem 5.1). \square

Sublemma 5.13. *If $n = 2$ then Γ_F contains no family of 5 parallel edges.*

Proof. The condition $n = 2$ implies that Γ_T has 2 vertices. Recall from the proof of Theorem 5.2 that on a torus one can find at most four embedded pairwise non-parallel arcs joining two points, that are disjoint except at their endpoints. Consequently if $n = 2$ and Γ_F contains a family of 5 parallel edges, at least two of these edges are also parallel in Γ_T , so Lemma 3.8 implies that $M = \mathbb{RP}^3$, contrary to the assumption that M is toroidal. \square

We now complete the proof of Proposition 5.11. If F is not an annulus, let \bar{e} be the number of edges in the reduced graph $\bar{\Gamma}_F$. If $n \geq 4$ then Sublemma 5.12 shows that $\bar{e} \geq pn/(n+4) \geq p/2$. If $n = 2$ Sublemma 5.13 shows that $\bar{e} \geq 2p/8 = p/4$. Hence $-\chi^-(F) \geq p/12$ by Lemma 3.5.

If F is an annulus then by Theorem 2.11 (3) K is contained as a fiber of multiplicity r in a Seifert fiber submanifold N of M , where N has base orbifold a Möbius band with one orbifold point of order $r \geq 1$. In particular $N - K$ is toroidal. Since $M - K$ is atoroidal by hypothesis, ∂N must be compressible in $W = \overline{M - N}$. Since $M - K$ is irreducible it follows that W is a solid torus, and that the meridian of W is not identified with a fiber in ∂N . Hence $M - \text{int } N(K)$ is a Seifert fiber space over the Möbius band with at most one exceptional fiber (the core of W). Since this manifold is atoroidal by hypothesis, the core of W is in fact an ordinary fiber. Then M is a

Seifert fiber space over \mathbb{RP}^2 with at most one exceptional fiber, namely K . But this contradicts our assumption that M is toroidal. \square

Proposition 5.14. *Let M be an irreducible 3-manifold containing an essential annulus A . Let K be a knot in M and let F be a good relative p -Seifert surface for K . Then either*

- (1) $-\chi^-(F) \geq p/6$; or
- (2) K can be isotoped to be either disjoint from A or to lie in ∂M ; or
- (3) F is an annulus.

Proof. We may assume that F is good. Let $X = M - \text{int } N(K)$.

Isotop K so that $n = |K \cap A|$ is minimal. If $n = 0$ we are done, so assume in what follows that $n > 0$. Let $P = A \cap X$, and let Γ_F, Γ_P be the intersection graphs in \hat{F} and A respectively.

Sublemma 5.15. Γ_F contains no family of $(n + 1)$ parallel interior edges.

Proof. By Lemma 3.7 bullet (2), Γ_P has 3 $(i_0, k - i_0)$ -edges for some i_0, k . Two of these must be parallel in A (since the union of any two is an embedded loop, and there is only one isotopy class of embedded essential loops in an annulus). If they are parallel in Γ_P then we have interior edges parallel in both graphs and Lemma 3.8 readily gives a contradiction. If not, then the disk D in A realizing the parallelism must have vertices of Γ_P in its interior. By Lemma 3.7 bullet (1) these vertices come in pairs $i, k - i$, each with 2 edges joining them. By taking an innermost such pair in D we again get a pair of edges that are parallel in both graphs, a contradiction as before. \square

Sublemma 5.16. *If Γ_F contains a family of $(2n + 1)$ parallel boundary edges then K is isotopic into ∂M .*

Proof. At the vertex end of such a family every label in $\{1, 2, \dots, n\}$ appears twice, and one label appears three times. The three corresponding boundary edges in Γ_P all share a common vertex; at least two must be parallel in A . If they are parallel in Γ_P , we get boundary edges parallel in both graphs, so the result follows from Lemma 3.9. If not, then the disk D in A realizing the parallelism must have vertices of Γ_P in its interior. Each such vertex has two boundary edges coming from the given family; an innermost pair is parallel in both graphs, so we are done by Lemma 3.9. \square

We now complete the proof of Proposition 5.14. If F is an annulus we are done. If not, then the reduced graph $\bar{\Gamma}_F$ exists. Let $\bar{e}_i, \bar{e}_\partial$ be the number of interior and boundary edges of $\bar{\Gamma}_F$. By Sublemma 5.15 each interior edge of $\bar{\Gamma}_F$ contributes at most $2n$ to the sum of the valences of the vertices on Γ_F , and by Sublemma 5.16 we

may assume that each boundary edge of $\bar{\Gamma}_F$ also contributes at most $2n$ to this sum. Hence $\bar{e} = \bar{e}_i + \bar{e}_\partial \geq pn/2n = p/2$, and the estimate in bullet (1) follows from Lemma 3.5. \square

In the next proposition we consider the orientable Seifert fiber spaces $D^2(2)$, $M^2(0)$, $A^2(1)$ and $P^2(0)$, where $F^2(n)$ denotes a Seifert fiber space with base surface F^2 and n exceptional fibers, and D^2 , M^2 , A^2 , P^2 are the disk, Möbius band, annulus, and pair of pants, respectively. These, together with $S^1 \times D^2$ and $T^2 \times I$, are precisely the atoroidal Seifert fiber spaces with non-empty boundary.

Remark 5.17. This notation does not uniquely determine the space up to homeomorphism, since we do not specify the type of singular fiber.

Proposition 5.18. *Let M be an atoroidal Seifert fiber space of the kind $D^2(2)$, $M^2(0)$, $A^2(1)$ or $P^2(0)$, and let K be a knot in M . Let F be a relative p -Seifert surface for K . Then either*

- (1) $-\chi^-(F) \geq p/6$; or
- (2) K is isotopic into ∂M ; or
- (3) K is isotopic to a cable of an exceptional Seifert fiber.

Proof. We may assume that F is good and $M - K$ is irreducible.

First assume that F is not an annulus. Let A be an essential annulus in M such that M cut along A is two fibered solid tori if $M = D^2(2)$, one fibered solid torus if $M = M^2(0)$ or $A^2(1)$, and $T^2 \times I$ if $M = P^2(0)$. By Proposition 5.14 either bullet (1) or (2) holds, or K can be isotoped to be disjoint from A .

If $M = P^2(0)$ and K can be isotoped to be disjoint from A , then $K \subset T \times I$ where T is a boundary component of M . Therefore bullet (1) or (2) holds, by Corollary 3.19.

If M is one of $D^2(2)$, $M^2(0)$ or $A^2(1)$ and K can be isotoped to be disjoint from A , then K is contained in a fibered solid torus $V \subset M$. Let $F' = F \cap V$ and $F'' = F \cap (M - \text{int } V)$. By hypothesis $M - K$ is irreducible, so ∂V is incompressible in $M - \text{int } N(K)$. Therefore we can assume that F'' has no disk (or sphere) components. Hence $-\chi^-(F) \geq -\chi^-(F')$. Therefore by Proposition 3.13 either $-\chi^-(F) > p/6$ or K is a cable of K_0 , the core of V . If K_0 is an ordinary fiber then it lies in a vertical incompressible torus T , so by Corollary 3.19 either bullet (1) holds or K is isotopic to K_0 , in which case bullet (2) holds. If K_0 is an exceptional fiber we get conclusion (3).

Finally we consider the case where F is an annulus, i.e. Γ_F is a beachball.

If Γ_F is a beachball of the first kind then (see Theorem 2.11 (3)) K is contained in a submanifold N of M of the form $M^2(0)$ or $M^2(1)$, as an ordinary or exceptional fiber respectively. Since M is irreducible, atoroidal, and has non-empty boundary, the torus ∂N is boundary parallel in M , and so $M \cong N$. Therefore N is of the form $M^2(0)$, and K is isotopic into ∂M .

If Γ_F is a beachball of the second kind, then $M = V \cup_B W$, where V is a solid torus, B is an annulus with winding number $r \geq 1$ in V , and K is a core of V . If $r = 1$ then K is isotopic into ∂M . If $r > 1$ then the form of M implies that the (separating) annulus B is essential in M , and hence is vertical. Therefore K is an exceptional fiber in M . \square

We are finally ready to treat the case that M is toroidal.

Theorem 5.19. *Let M be a closed, irreducible, toroidal 3-manifold, and let K be a knot in M . Then either*

- (1) $\|K\| \geq 1/402$; or
 - (2) K is trivial; or
 - (3) K is contained in a hyperbolic piece N of the JSJ decomposition of M and is isotopic either to a cable of a core of a Margulis tube or into a component of ∂N ; or
 - (4) K is contained in a Seifert fiber piece N of the JSJ decomposition of M and either
 - (A) K is isotopic to an ordinary fiber or a cable of an exceptional fiber or into ∂N , or
 - (B) N contains a copy Q of the twisted S^1 bundle over the Möbius band and K is contained in Q as a fiber in this bundle structure;
- or
- (5) M is a T^2 -bundle over S^1 with Anosov monodromy and K is contained in a fiber.

Remark 5.20. If K is disjoint from the hyperbolic pieces in the JSJ decomposition of M , the constant $1/402$ can be improved to $1/24$.

Remark 5.21. In Case (4) (B), $\|K\| = 0$ (see Theorem 2.11 (3)). Also, the Seifert fibration of Q induced from N may be the one with base orbifold a disk with two cone points of order 2, in which case K is not a Seifert fiber in N .

Remark 5.22. Strictly speaking, (5) is a special case of (4) where the Seifert fiber piece N is $T^2 \times I$, but for clarity we list it separately.

Proof. We may assume that $M - K$ is irreducible. By Proposition 5.11, either $\|K\| \geq 1/24$ or $M - K$ is toroidal. So we may assume that $M - K$ is irreducible and toroidal.

Let \mathcal{T} be a maximal disjoint union of non-parallel essential tori in $M - K$ (note that \mathcal{T} is nonempty). Let N be the component of M cut along \mathcal{T} that contains K .

Then N is irreducible, has boundary a non-empty disjoint union of tori, and $N - K$ is irreducible and atoroidal.

Let S be a good p -Seifert surface for K in M . Then $F = S \cap N$ is a relative p -Seifert surface for K in N . Let $S_0 = \overline{S - F}$. Since ∂N is incompressible in $M - K$ we may assume that no component of S_0 or F is a disk (or sphere). Hence $\eta(S) = \eta(F) + \eta(S_0)$ and so $\eta(S) \geq \eta(F)$.

If N is toroidal then we are done by Proposition 5.11. If N is atoroidal then N is either hyperbolic or Seifert fibered. In the former case, N is a piece of the JSJ decomposition of M , and by Proposition 5.8, either (1) or (3) holds.

Suppose N is a Seifert fiber space. Since N is atoroidal and has non-empty boundary it is either homeomorphic to $S^1 \times D^2$ or $T^2 \times I$, or has Seifert fiber structure $D^2(2)$, $M^2(0)$, $A^2(1)$ or $P^2(0)$. In the last four cases the result follows from Proposition 5.18 (to get conclusion (4) of the theorem we replace the N considered here with the Seifert fiber piece of the JSJ decomposition of M that contains it).

If $N = T^2 \times I$ then by Proposition 3.17 either (1) holds or K is isotopic onto $T = T^2 \times \{1/2\}$. Since T is an incompressible torus in M there are four possibilities:

- (i) T is a torus in the JSJ decomposition of M ;
- (ii) T is a vertical essential torus in a Seifert fiber piece of the JSJ decomposition of M ;
- (iii) M is a Seifert fiber space and T is horizontal;
- (iv) M is a T^2 -bundle over S^1 and T is a fiber.

If (i) holds we are done. If (ii) holds we are done by Proposition 2.24.

Suppose (iii) holds but not (iv). Then T separates M into two twisted I -bundles over the Klein bottle. The Seifert fibering of M must have Euler number 0, and M is either a twisted S^1 -bundle over the Klein bottle or has base orbifold \mathbb{RP}^2 with two orbifold points of order 2. In both cases M has an isomorphic (but non-isotopic) Seifert fibering in which T is vertical.

If (iv) holds then by Remark 2.22 either $\|K\| = 0$, or $\|K\| \geq 1/8$, or we are in Case G or Case H of §2.5. In Case G, M is Seifert fibered and K is a fiber. Case H is conclusion (5). So suppose $\|K\| = 0$. Then by part (2) of Theorem 2.21 $\begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} = \begin{pmatrix} -1 & n \\ 0 & -1 \end{pmatrix}$, with K representing the first element of the corresponding basis. Thus M also has the structure of an S^1 -bundle over the Klein bottle with Euler number n , where K is a fiber.

Finally, suppose $N = S^1 \times D^2$. Then by Proposition 3.13, and the fact that \mathcal{T} is essential in $M - K$, we may assume that K is a non-trivial cable of K_0 , the core of N . By Proposition 3.12, $\|K_0\| < \|K\|$.

Now repeat the whole argument with K_0 in place of K . We conclude that either (1) holds or

- (a) K_0 is contained in a hyperbolic piece N_0 of the JSJ decomposition of M and is isotopic either to a cable of a core of a Margulis tube or into ∂N_0 ; or

- (b) K_0 is contained in a Seifert fiber piece N_0 of the JSJ decomposition of M and is isotopic either to an ordinary fiber or a cable of an exceptional fiber of N_0 or into ∂N_0 , or N_0 contains a copy Q_0 of the twisted S^1 bundle over the Möbius band and K is contained in Q_0 as an S^1 fiber; or
- (c) M is a T^2 -bundle over S^1 and K_0 lies in a fiber; or
- (d) K_0 is a non-trivial cable of a knot K_1 .

By Proposition 3.13, K cannot be a non-trivial cable of a non-trivial cable. Similarly, by Proposition 3.17, K cannot be a non-trivial cable of an essential curve in an incompressible torus in M . This completes the proof of the theorem. \square

References

- [1] I. Agol, Bounds on exceptional Dehn filling. *Geom. Topol.* **4** (2000), 431–449. [Zbl 0959.57009](#) [MR 1799796](#)
- [2] K. Baker, Small genus knots in lens spaces have small bridge number. *Algebr. Geom. Topol.* **6** (2006), 1519–1621. [Zbl 1130.57004](#) [MR 2253458](#)
- [3] K. Baker and J. Etnyre, Rational linking and contact geometry. In *Perspectives in analysis, geometry, and topology* (in honor of Oleg Viro), Progr. Math. 296, Birkhäuser/Springer, New York 2012, 19–37. [Zbl 06082160](#) [MR 2884030](#)
- [4] C. Bavard, Longueur stable des commutateurs. *L'Enseign. Math.* **37** (1991), 109–150. [Zbl 0810.20026](#) [MR 1115747](#)
- [5] R. Benedetti and C. Petronio, *Lectures on hyperbolic geometry*. Universitext, Springer-Verlag, New York 1992. [Zbl 0768.51018](#) [MR 1219310](#)
- [6] M. Boileau and J. Porti, Geometrization of 3-orbifolds of cyclic type. *Astérisque* **272**, 2001. [Zbl 0971.57004](#) [MR 1844891](#)
- [7] D. Calegari, Length and stable length. *Geom. Funct. Anal.* **18** (2008), no. 1, 50–76. [Zbl 1155.57015](#) [MR 2399095](#)
- [8] D. Calegari, *scl*. MSJ Memoirs 20, Mathematical Society of Japan, Tokyo 2009. [Zbl 1187.20035](#) [MR 2527432](#)
- [9] D. Calegari and D. Gabai, Shrinkwrapping and the taming of hyperbolic 3-manifolds. *J. Amer. Math. Soc.* **19** (2006), no. 2, 385–446. [Zbl 1090.57010](#) [MR 2188131](#)
- [10] C. Cao and R. Meyerhoff, The orientable cusped hyperbolic 3-manifolds of minimum volume. *Invent. Math.* **146** (2001), 451–478. [Zbl 1028.57010](#) [MR 1869847](#)
- [11] M. Culler, C. Gordon, J. Luecke and P. Shalen, Dehn surgery on knots. *Ann. of Math.* **125** (1987), 237–300. [Zbl 0633.57006](#) [MR 0881270](#)
- [12] D. Gabai, Foliations and the topology of 3-manifolds III. *J. Differential Geom.* **26** (1987), no. 3, 479–536. [Zbl 0639.57008](#) [MR 0910018](#)
- [13] C. Gordon and R. Litherland, Incompressible planar surfaces in 3-manifolds. *Topology Appl.* **18** (1984), 121–144. [Zbl 0554.57010](#) [MR 0769286](#)
- [14] A. Hatcher, *Notes on basic 3-manifold topology*. Available from the author's webpage <http://www.math.cornell.edu/~hatcher/3M/3Mdownloads.html>

- [15] M. Hedden, On Floer Homology and the Berge Conjecture on knots admitting lens space surgeries. *Trans. Amer. Math. Soc.* **363** (2011), no. 2, 949–968. [Zbl 05863899](#) [MR 2728591](#)
- [16] C. Hodgson and S. Kerckhoff, Universal bounds for hyperbolic Dehn surgery. *Ann. of Math.* **162** (2005), 367–421. [Zbl 1087.57011](#) [MR 2178964](#)
- [17] W. Meeks, L. Simon and S.-T. Yau, Embedded minimal surfaces, exotic spheres, and manifolds with positive Ricci curvature. *Ann. of Math. (2)* **116**, no. 3, 621–659. [Zbl 0521.53007](#) [MR 0678484](#)
- [18] P. Ozsváth and Z. Szabó, Holomorphic disks and genus bounds. *Geom. Topol.* **8** (2004), 311–334. [Zbl 1056.57020](#) [MR 2023281](#)
- [19] J. Rasmussen, Lens space surgeries and L -space homology spheres. Preprint, [arXiv:0710.2531](#) [math.GT].
- [20] Y. Rieck, Heegaard surfaces and Dehn fillings: $g(M) + 1 \leq t(X) \leq g(M)$. PhD thesis, UT Austin 1997.
- [21] Y. Rieck, Heegaard structures of manifolds in the Dehn filling space. *Topology* **39** (2000), 619–641. [Zbl 0944.57013](#) [MR 1746912](#)
- [22] M. Scharlemann, Smooth spheres in \mathbb{R}^4 with four critical points are standard. *Invent. Math.* **79** (1985), 125–141. [Zbl 0559.57019](#) [MR 0774532](#)
- [23] R. Schoen and S.-T. Yau, Existence of incompressible minimal surfaces and the topology of three-dimensional manifolds with non-negative scalar curvature. *Ann. of Math. (2)* **110**, no. 1, 127–142. [Zbl 0431.53051](#) [MR 0541332](#)
- [24] P. Scott, The geometries of 3-manifolds. *Bull. London Math. Soc.* **15** (1983), no. 5, 401–487. [Zbl 0561.57001](#) [MR 0705527](#)
- [25] T. Soma, Existence of ruled wrappings in hyperbolic 3-manifolds. *Geom. Topol.* **10** (2006), 1173–1184. [Zbl 1130.57025](#) [MR 2255495](#)
- [26] W. Thurston, A norm for the homology of 3-manifolds. *Mem. Amer. Math. Soc.* **59** (1986), no. 339, i–vi and 99–130. [Zbl 0585.57006](#) [MR 0823443](#)
- [27] W. Thurston, *Geometry and topology of 3-manifolds*. Available at <http://library.msri.org/books/gt3m/>.
- [28] Y.-Q. Wu, The reducibility of surgered 3-manifolds. *Topology Appl.* **43** (1992), no. 3, 213–218. [Zbl 0817.57019](#) [MR 1158868](#)

Received December 10, 2009

Danny Calegari, Department of Mathematics, Caltech, Pasadena, CA 91125, U.S.A.

E-mail: dannyc@its.caltech.edu

Cameron Gordon, Department of Mathematics, University of Texas, Austin TX, 78712, U.S.A.

E-mail: gordon@math.utexas.edu