**Zeitschrift:** Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

**Band:** 88 (2013)

**Artikel:** Moduli spaces of hyperbolic 3-manifolds and dynamics on character

varieties

Autor: Canary, Richard D. / Storm, Peter A.

DOI: https://doi.org/10.5169/seals-515637

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF: 29.11.2025** 

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

# Moduli spaces of hyperbolic 3-manifolds and dynamics on character varieties

Richard D. Canary and Peter A. Storm\*

**Abstract.** The space AH(M) of marked hyperbolic 3-manifold homotopy equivalent to a compact 3-manifold with boundary M sits inside the  $PSL_2(\mathbb{C})$ -character variety X(M) of  $\pi_1(M)$ . We study the dynamics of the action of  $Out(\pi_1(M))$  on both AH(M) and X(M). The nature of the dynamics reflects the topology of M.

The quotient  $AI(M) = AH(M)/Out(\pi_1(M))$  may naturally be thought of as the moduli space of unmarked hyperbolic 3-manifolds homotopy equivalent to M and its topology reflects the dynamics of the action.

Mathematics Subject Classification (2010). 57M50.

**Keywords.** Hyperbolic 3-manifolds, outer automorphism group, character variety, moduli spaces.

## 1. Introduction

For a compact, orientable, hyperbolizable 3-manifold M with boundary, the deformation space AH(M) of marked hyperbolic 3-manifolds homotopy equivalent to M is a familiar object of study. This deformation space sits naturally inside the  $PSL_2(\mathbb{C})$ -character variety X(M) and the outer automorphism group  $Out(\pi_1(M))$  acts by homeomorphisms on both AH(M) and X(M). The action of  $Out(\pi_1(M))$  on AH(M) and X(M) has largely been studied in the case when M is an interval bundle over a closed surface (see, for example, [8], [22], [49], [18]) or a handlebody (see, for example, [43], [54]). In this paper, we initiate a study of this action for general hyperbolizable 3-manifolds.

We also study the topological quotient

$$AI(M) = AH(M)/Out(\pi_1(M))$$

which we may think of as the moduli space of unmarked hyperbolic 3-manifolds homotopy equivalent to M. The space AH(M) is a rather pathological topological

<sup>\*</sup>The first author was partially supported by National Science Foundation grants DMS-0504791 and DMS-0554239. The second author was partially supported by National Science Foundation grant DMS-0904355 and the Roberta and Stanley Bogen Visiting Professorship at Hebrew University.

object itself, often failing to even be locally connected (see Bromberg [12] and Magid [35]). However, since AH(M) is a closed subset of an open submanifold of the character variety, it does retain many nice topological properties. We will see that the topology of AI(M) can be significantly more pathological.

The first hint that the dynamics of  $Out(\pi_1(M))$  on AH(M) are complicated was Thurston's [51] proof that if M is homeomorphic to  $S \times I$ , then there are infinite order elements of  $Out(\pi_1(M))$  which have fixed points in AH(M). (These elements are pseudo-Anosov mapping classes.) One may further show that  $AI(S \times I)$  is not even  $T_1$ , see [18] for a closely related result. Recall that a topological space is  $T_1$  if all its points are closed. On the other hand, we show that in all other cases AI(M) is  $T_1$ .

**Theorem 1.1.** Let M be a compact hyperbolizable 3-manifold with non-abelian fundamental group. Then the moduli space AI(M) is  $T_1$  if and only if M is not an untwisted interval bundle.

We next show that  $Out(\pi_1(M))$  does not act properly discontinuously on AH(M) if M contains a primitive essential annulus. A properly embedded annulus in M is a primitive essential annulus if it cannot be properly isotoped into the boundary of M and its core curve generates a maximal abelian subgroup of  $\pi_1(M)$ . In particular, if M has compressible boundary and no toroidal boundary components, then M contains a primitive essential annulus (see Corollary 7.5).

**Theorem 1.2.** Let M be a compact hyperbolizable 3-manifold with non-abelian fundamental group. If M contains a primitive essential annulus then  $Out(\pi_1(M))$  does not act properly discontinuously on AH(M). Moreover, if M contains a primitive essential annulus, then AI(M) is not Hausdorff.

On the other hand, if M is acylindrical, i.e. has incompressible boundary and contains no essential annuli, then  $Out(\pi_1(M))$  is finite (see Johannson [29], Proposition 27.1), so  $Out(\pi_1(M))$  acts properly discontinuously on AH(M) and X(M). It is easy to see that  $Out(\pi_1(M))$  fails to act properly discontinuously on X(M) if M is not acylindrical, since it will contain infinite order elements with fixed points in X(M).

If M is a compact hyperbolizable 3-manifold which is not acylindrical, but does not contain any primitive essential annuli, then  $\operatorname{Out}(\pi_1(M))$  is infinite. However, if, in addition, M has no toroidal boundary components, we show that  $\operatorname{Out}(\pi_1(M))$  acts properly discontinuously on an open neighborhood of AH(M) in X(M). In particular, we see that AI(M) is Hausdorff in this case.

**Theorem 1.3.** If M is a compact hyperbolizable 3-manifold with no primitive essential annuli whose boundary has no toroidal boundary components, then there exists

an open  $Out(\pi_1(M))$ -invariant neighborhood W(M) of AH(M) in X(M) such that  $Out(\pi_1(M))$  acts properly discontinuously on W(M). In particular, AI(M) is Hausdorff.

If M is a compact hyperbolizable 3-manifold with no primitive essential annuli whose boundary has no toroidal boundary components, then  $\operatorname{Out}(\pi_1(M))$  is virtually abelian (see the discussion in Sections 5 and 9). However, we note that the conclusion of Theorem 1.3 relies crucially on the topology of M, not just the group theory of  $\operatorname{Out}(\pi_1(M))$ . In particular, if M is a compact hyperbolizable 3-manifold M with incompressible boundary, such that every component of its characteristic submanifold is a solid torus, then  $\operatorname{Out}(\pi_1(M))$  is always virtually abelian, but M may contain primitive essential annuli, in which case  $\operatorname{Out}(\pi_1(M))$  does not act properly discontinuously on  $\operatorname{AH}(M)$ .

One may combine Theorems 1.2 and 1.3 to completely characterize when  $Out(\pi_1(M))$  acts properly discontinuously on AH(M) in the case that M has no toroidal boundary components.

**Corollary 1.4.** Let M be a compact hyperbolizable 3-manifold with no toroidal boundary components and non-abelian fundamental group. The group  $Out(\pi_1(M))$  acts properly discontinuously on AH(M) if and only if M contains no primitive essential annuli. Moreover, AI(M) is Hausdorff if and only if M contains no primitive essential annuli.

It is a consequence of the classical deformation theory of Kleinian groups (see Bers [5] or Canary and McCullough, Chapter 7 in [17], for a survey of this theory) that  $Out(\pi_1(M))$  acts properly discontinuously on the interior int(AH(M)) of AH(M). If  $H_n$  is the handlebody of genus  $n \geq 2$ , Minsky [43] exhibited an explicit  $Out(\pi_1(H_n))$ -invariant open subset  $PS(H_n)$  of  $X(H_n)$  such that  $int(AH(H_n))$  is a proper subset of  $PS(H_n)$  and  $Out(\pi_1(H_n))$  acts properly discontinuously on  $PS(H_n)$ .

If M is a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components, which is not an interval bundle, then we find an open set W(M) strictly bigger than  $\operatorname{int}(AH(M))$  which  $\operatorname{Out}(\pi_1(M))$  acts properly discontinuously on. See Theorem 9.1 and its proof for a more precise description of W(M). We further observe, see Lemma 8.1, that  $W(M) \cap \partial AH(M)$  is a dense open subset of  $\partial AH(M)$  in this setting.

**Theorem 1.5.** Let M be a compact hyperbolizable 3-manifold with nonempty incompressible boundary and no toroidal boundary components, which is not an interval bundle. Then there exists an open  $Out(\pi_1(M))$ -invariant subset W(M) of X(M) such that  $Out(\pi_1(M))$  acts properly discontinuously on W(M) and int(AH(M)) is a proper subset of W(M).

It is conjectured that if M is an untwisted interval bundle over a closed surface S, then int(AH(M)) is the maximal open  $Out(\pi_1(M))$ -invariant subset of X(M) on which  $Out(\pi_1(M))$  acts properly discontinuously. One may show that no open domain of discontinuity can intersect  $\partial AH(S \times I)$  (see Lee [34]). Further evidence for this conjecture is provided by results of Bowditch [8], Goldman [21], Souto-Storm [49], Tan-Wong-Zhang [54] and Cantat [19].

Michelle Lee [34] has recently shown that if M is an twisted interval bundle over a closed surface, then there exists an open  $\operatorname{Out}(\pi_1(M))$ -invariant subset W of X(M) such that  $\operatorname{Out}(\pi_1(M))$  acts properly discontinuously on W and  $\operatorname{int}(AH(M))$  is a proper subset of W. Moreover, W contains points in  $\partial AH(M)$ . As a corollary, she proves that if M has incompressible boundary and no toroidal boundary components, then there is open  $\operatorname{Out}(\pi_1(M))$ -invariant subset W of X(M) such that  $\operatorname{Out}(\pi_1(M))$  acts properly discontinuously on W,  $\operatorname{int}(AH(M))$  is a proper subset of W, and  $W \cap \partial AH(M) \neq \emptyset$  if and only if M is not an untwisted interval bundle.

**Outline of paper.** In Section 2, we recall background material from topology and hyperbolic geometry which will be used in the paper.

In Section 3, we prove Theorem 1.1. The proof that  $AI(S \times I)$  is not  $T_1$  follows the arguments in [18], Proposition 3.1, closely. We now sketch the proof that AI(M) is  $T_1$  otherwise. In this case, let  $N \in AI(M)$  and let R be a compact core for N. We show that N is a closed point, by showing that any convergent sequence  $\{\rho_n\}$  in the pre-image of N is eventually constant. For all n, there exists a homotopy equivalence  $h_n \colon M \to N$  such that  $(h_n)_* = \rho_n$ . If G is a graph in M carrying  $\pi_1(M)$ , then, since  $\{\rho_n\}$  is convergent, we can assume that the length of  $h_n(G)$  is at most K, for all n and some K. But, we observe that  $h_n(G)$  cannot lie entirely in the complement of R, if R is not a compression body. In this case, each  $h_n(G)$  lies in the compact neighborhood of radius K of R, so there are only finitely many possible homotopy classes of maps of G. Thus, there are only finitely many possibilities for  $\rho_n$ , so  $\{\rho_n\}$  is eventually constant. The proof in the case that R is a compression body is somewhat more complicated and uses the Covering Theorem.

In Section 4, we prove Theorem 1.2. Let A be a primitive essential annulus in M. If  $\alpha$  is a core curve of A, then the complement  $\widehat{M}$  of a regular neighborhood of  $\alpha$  in M is hyperbolizable. We consider a geometrically finite hyperbolic manifold  $\widehat{N}$  homeomorphic to the interior of  $\widehat{M}$  and use the Hyperbolic Dehn Filling Theorem to produce a convergent sequence  $\{\rho_n\}$  in AH(M) and a sequence  $\{\varphi_n\}$  of distinct elements of  $\mathrm{Out}(\pi_1(M))$  such that  $\{\rho_n \circ \varphi_n\}$  also converges. Therefore,  $\mathrm{Out}(\pi_1(M))$  does not act properly discontinuously on AH(M). Moreover, we show that  $\{\rho_n\}$  projects to a sequence in AI(M) with two distinct limits, so AI(M) is not Hausdorff.

In Section 5 we recall basic facts about the characteristic submanifold and the mapping class group of compact hyperbolizable 3-manifolds with incompressible boundary and no toroidal boundary components. We identify a finite index subgroup J(M) of  $Out(\pi_1(M))$  and a projection of J(M) onto the direct product of mapping

class groups of the base surfaces whose kernel K(M) is the free abelian subgroup generated by Dehn twists in frontier annuli of the characteristic submanifold.

In Section 6, we organize the frontier annuli of the characteristic submanifold into characteristic collections of annuli and describe free subgroups of  $\pi_1(M)$  which register the action of the subgroup of  $\mathrm{Out}(\pi_1(M))$  generated by Dehn twists in the annuli in such a collection.

In Section 7, we show that compact hyperbolizable 3-manifolds with compressible boundary and no toroidal boundary components contain primitive essential annuli.

In Section 8, we introduce a subset  $AH_n(M)$  of AH(M) which contains all purely hyperbolic representations. We see that int(AH(M)) is a proper subset of  $AH_n(M)$  and that  $AH_n(M) = AH(M)$  if M does not contain any primitive essential annuli.

In Section 9, we prove that if M has incompressible boundary and no toroidal boundary components, but is not an interval bundle, there is an open neighborhood W(M) of  $AH_n(M)$  in X(M) such that  $Out(\pi_1(M))$  preserves and acts properly discontinuously on W(M). Theorems 1.3 and 1.5 are immediate corollaries. We finish the outline by sketching the proof in a special case.

Let X be an acylindrical, compact hyperbolizable 3-manifold and let A be an incompressible annulus in its boundary. Let V be a solid torus and let  $\{B_1, \ldots, B_n\}$  be a collection of disjoint parallel annuli in  $\partial V$  whose core curves are homotopic to the  $n^{th}$  power of the core curve of V where  $|n| \geq 2$ . Let  $\{M_1, \ldots, M_n\}$  be copies of X and let  $\{A_1, \ldots, A_n\}$  be copies of A in  $M_i$ . We form M by attaching each  $M_i$  to V by identifying  $A_i$  and  $B_i$ . Then M contains no primitive essential annuli, is hyperbolizable, and  $\operatorname{Out}(\pi_1(M))$  has a finite index subgroup J(M) generated by Dehn twists about  $\{A_1, \ldots, A_n\}$ . In particular,  $J(M) \cong \mathbb{Z}^{n-1}$ .

In this case,  $\{A_1, \ldots, A_n\}$  is the only characteristic collection of annuli. We say that a group H registers J(M) if it is freely generated by the core curve of V and, for each i, a curve contained in  $V \cup M_i$  which is not homotopic into V. So  $H \cong F_{n+1}$ . There is a natural map  $r_H: X(M) \to X(H)$  where X(H) is the  $\mathrm{PSL}_2(\mathbb{C})$ -character variety of the group H. Notice that J(M) preserves H and injects into  $\mathrm{Out}(H)$ . Let

$$S_{n+1} = \operatorname{int}(AH(H)) \subset X(H)$$

denote the space of Schottky representations (i.e. representations which are purely hyperbolic and geometrically finite.) Since Out(H) acts properly discontinuously on  $S_{n+1}$ , we see that J(M) acts properly discontinuously on

$$W_H = r_H^{-1}(S_{n+1})$$

Let  $W(M) = \bigcup W_H$  where the union is taken over all subgroups which register J(M). Notice that W(M) is open and J(M) acts properly discontinuously on W(M). One may use a ping pong argument to show that  $AH(M) \subset W(M)$ , see Lemma 8.3. Johannson's Classification Theorem is used to show that W(M) is invariant under Out $(\pi_1(M))$ , see Lemma 9.3. (Actually, we define a somewhat larger set, in general, by using the space of primitive-stable representations in place of Schottky space.)

**Acknowledgements.** Both authors would like to thank Yair Minsky for helpful and interesting conversations. The second author thanks the organizers of the workshop "Dynamics of  $Aut(F_n)$ -actions on representation varieties," held in Midreshet Sde Boker, Israel in January, 2009. We thank Michelle Lee, Darryl McCullough and the referee for helpful comments on a preliminary version of this paper.

#### 2. Preliminaries

As a convention, the letter M will denote a compact connected oriented hyperbolizable 3-manifold with boundary. We recall that M is said to be hyperbolizable if the interior of M admits a complete hyperbolic metric. We will use N to denote a hyperbolic 3-manifold. All hyperbolic 3-manifolds are assumed to be oriented, complete, and connected.

**2.1. The deformation spaces.** Recall that  $PSL_2(\mathbb{C})$  is the group of orientation-preserving isometries of  $\mathbb{H}^3$ . Given a 3-manifold M, a discrete, faithful representation  $\rho \colon \pi_1(M) \to PSL_2(\mathbb{C})$  determines a hyperbolic 3-manifold  $N_\rho = \mathbb{H}^3/\rho(\pi_1(M))$  and a homotopy equivalence  $m_\rho \colon M \to N_\rho$ , called the marking of  $N_\rho$ .

We let D(M) denote the set of discrete, faithful representations of  $\pi_1(M)$  into  $PSL_2(\mathbb{C})$ . The group  $PSL_2(\mathbb{C})$  acts by conjugation on D(M) and we let

$$AH(M) = D(M)/PSL_2(\mathbb{C}).$$

Elements of AH(M) are hyperbolic 3-manifolds homotopy equivalent to M equipped with (homotopy classes of) markings.

The space AH(M) is a closed subset of the character variety

$$X(M) = \operatorname{Hom}_T(\pi_1(M), \operatorname{PSL}_2(\mathbb{C})) /\!\!/ \operatorname{PSL}_2(\mathbb{C}),$$

which is the Mumford quotient of the space  $\operatorname{Hom}_T(\pi_1(M),\operatorname{PSL}_2(\mathbb{C}))$  of representations  $\rho\colon \pi_1(M)\to\operatorname{PSL}_2(\mathbb{C})$  such that  $\rho(g)$  is parabolic if  $g\neq \operatorname{id}$  lies in a rank two free abelian subgroup of  $\pi_1(M)$ . If M has no toroidal boundary components, then  $\operatorname{Hom}_T(\pi_1(M),\operatorname{PSL}_2(\mathbb{C}))$  is simply  $\operatorname{Hom}(\pi_1(M),\operatorname{PSL}_2(\mathbb{C}))$ . Moreover, each point in AH(M) is a smooth point of X(M) (see Kapovich [30], Sections 4.3 and 8.8, and Heusener-Porti [24] for more details on this construction).

The group  $\operatorname{Aut}(\pi_1(M))$  acts naturally on  $\operatorname{Hom}_T(\pi_1(M),\operatorname{PSL}_2(\mathbb{C}))$  via

$$(\varphi \cdot \rho)(\gamma) := \rho(\varphi^{-1}(\gamma)).$$

This descends to an action of  $Out(\pi_1(M))$  on AH(M) and X(M). This action is not free, and it often has complex dynamics. Nonetheless, we can define the topological quotient space

$$AI(M) = AH(M)/Out(\pi_1(M)).$$

Elements of AI(M) are naturally oriented hyperbolic 3-manifolds homotopy equivalent to M without a specified marking.

**2.2. Topological background.** A compact 3-manifold M is said to have *incompressible boundary* if whenever S is a component of  $\partial M$ , the inclusion map induces an injection of  $\pi_1(S)$  into  $\pi_1(M)$ . In our setting, this is equivalent to  $\pi_1(M)$  being freely indecomposable. A properly embedded annulus A in M is said to be *essential* if the inclusion map induces an injection of  $\pi_1(A)$  into  $\pi_1(M)$  and A cannot be properly homotoped into  $\partial M$  (i.e. there does not exist a homotopy of pairs of the inclusion  $(A, \partial A) \to (M, \partial M)$  to a map with image in  $\partial M$ ). An essential annulus A is said to be *primitive* if the image of  $\pi_1(A)$  in  $\pi_1(M)$  is a maximal abelian subgroup.

If M does not have incompressible boundary, it is said to have *compressible boundary*. The fundamental examples of 3-manifolds with compressible boundary are compression bodies. A *compression body* is either a handlebody or is formed by attaching 1-handles to disjoint disks on the boundary surface  $R \times \{1\}$  of a 3-manifold  $R \times [0,1]$  where R is a closed, but not necessarily connected, surface (see, for example, Bonahon [6]). The resulting 3-manifold C (assumed to be connected) will have a single boundary component  $\partial_+ C$  intersecting  $R \times \{1\}$ , called the positive (or external) boundary of C. If C is not an untwisted interval bundle over a closed surface, then  $\partial_+ C$  is the unique compressible boundary component of C. Notice that the induced homomorphism  $\pi_1(\partial_+ C) \to \pi_1(C)$  is surjective. In fact, a compact irreducible 3-manifold M is a compression body if and only if there exists a component S of  $\partial M$  such that  $\pi_1(S) \to \pi_1(M)$  is surjective.

Every compact hyperbolizable 3-manifold can be constructed from compression bodies and manifolds with incompressible boundary. Bonahon [6] and McCullough–Miller [40] showed that there exists a neighborhood  $C_M$  of  $\partial M$ , called the *characteristic compression body*, such that each component of  $C_M$  is a compression body and each component of  $\partial C_M - \partial M$  is incompressible in M.

Dehn filling will play a key role in the proof of Theorem 1.2. Let F be a toroidal boundary component of compact 3-manifold M and let (m, l) be a choice of meridian and longitude for F. Given a pair (p, q) of relatively prime integers, we may form a new manifold M(p, q) by attaching a solid torus V to M by an orientation-reversing homeomorphism  $g \colon \partial V \to F$  so that, if c is the meridian of V, then g(c) is a (p, q) curve on F with respect to the chosen meridian-longitude system. We say that M(p, q) is obtained from M by (p, q)-Dehn filling along F.

**2.3. Hyperbolic background.** If  $N = \mathbb{H}^3/\Gamma$  is a hyperbolic 3-manifold, then  $\Gamma \subset \mathrm{PSL}_2(\mathbb{C})$  acts on  $\widehat{\mathbb{C}}$  as a group of conformal automorphisms. The *domain* of discontinuity  $\Omega(\Gamma)$  is the largest open  $\Gamma$ -invariant subset of  $\widehat{\mathbb{C}}$  on which  $\Gamma$  acts properly discontinuously. Note that  $\Omega(\Gamma)$  may be empty. Its complement  $\Lambda(\Gamma) = \widehat{\mathbb{C}} - \Omega(\Gamma)$  is called the *limit set*. The quotient  $\partial_c N = \Omega(\Gamma)/\Gamma$  is naturally a Riemann surface called the *conformal boundary*.

Thurston's Hyperbolization theorem, see Morgan [44], Theorem B', guarantees that if M is compact and hyperbolizable, then there exists a hyperbolic 3-manifold N and a homeomorphism

$$\psi: M - \partial_T M \to N \cup \partial_c N$$

where  $\partial_T M$  denotes the collection of toroidal boundary components of M.

The convex core C(N) of N is the smallest convex submanifold whose inclusion into N is a homotopy equivalence. More concretely, it is obtained as the quotient, by  $\Gamma$ , of the convex hull, in  $\mathbb{H}^3$ , of the limit set  $\Lambda(\Gamma)$ . There is a well-defined retraction  $r: N \to C(N)$  obtained by taking x to the (unique) point in C(N) closest to x. The nearest point retraction r is a homotopy equivalence and is  $\frac{1}{\cosh s}$ -Lipschitz on the complement of the neighborhood of radius s of C(N).

There exists a universal constant  $\mu$ , called the Margulis constant, such that if  $\epsilon < \mu$ , then each component of the  $\epsilon$ -thin part

$$N_{\text{thin}(\epsilon)} = \{ x \in N \mid \text{inj}_N(x) < \epsilon \}$$

(where  $\operatorname{inj}_N(x)$  denotes the injectivity radius of N at x) is either a metric regular neighborhood of a geodesic or is homeomorphic to  $T \times (0, \infty)$  where T is either a torus or an open annulus (see Benedetti–Petronio [4] for example). The  $\epsilon$ -thick part of N is defined simply to be the complement of the  $\epsilon$ -thin part

$$N_{\operatorname{thick}(\epsilon)} = N - N_{\operatorname{thin}(\epsilon)}.$$

It is also useful to consider the manifold  $N_{\epsilon}^{0}$  obtained from N by removing the non-compact components of  $N_{\text{thin}(\epsilon)}$ .

If N is a hyperbolic 3-manifold with finitely generated fundamental group, then it admits a compact core, i.e. a compact submanifold whose inclusion into M is a homotopy equivalence (see Scott [48]). More generally, if  $\epsilon < \mu$ , then there exists a relative compact core R for  $N_{\epsilon}^0$ , i.e. a compact core which intersects each component of  $\partial N_{\epsilon}^0$  in a compact core for that component (see Kulkarni–Shalen [33] or McCullough [38]). Let  $P = \partial R - \partial N_{\epsilon}^0$  and let  $P^0$  denote the interior of P. The Tameness Theorem of Agol [1] and Calegari–Gabai [14] assures us that we may choose R so that  $N_{\epsilon}^0 - R$  is homeomorphic to  $(\partial R - P^0) \times (0, \infty)$ . In particular, the ends of  $N_{\epsilon}^0$  are in one-to-one correspondence with the components of  $\partial R - P^0$ . (We will blur this distinction and simply regard an end as a component of  $N_{\epsilon}^0 - R$  once we have chosen  $\epsilon$  and a relative compact core R for  $N_{\epsilon}^0$ .) We say that an end U of  $N_{\epsilon}^0$  is geometrically finite if the intersection of C(N) with U is bounded (i.e. admits a compact closure). N is said to be geometrically finite if all the ends of  $N_{\epsilon}^0$  are geometrically finite.

Thurston [53] showed that if M is a compact hyperbolizable 3-manifold whose boundary is a torus F, then all but finitely many Dehn fillings of M are hyperbolizable. Moreover, as the Dehn surgery coefficients approach  $\infty$ , the resulting hyperbolic

manifolds "converge" to the hyperbolic 3-manifold homeomorphic to  $\operatorname{int}(M)$ . If M has other boundary components, then there is a version of this theorem where one begins with a geometrically finite hyperbolic 3-manifold homeomorphic to  $\operatorname{int}(M)$  and one is allowed to perform the Dehn filling while fixing the conformal structure on the non-toroidal boundary components of M. The proof uses the cone-manifold deformation theory developed by Hodgson–Kerckhoff [25] in the finite volume case and extended to the infinite volume case by Bromberg [11] and Brock–Bromberg [9]. (The first statement of a Hyperbolic Dehn Filling Theorem in the infinite volume setting was given by Bonahon–Otal [7], see also Comar [20].) For a general statement of the Filling Theorem, and a discussion of its derivation from the previously mentioned work, see Bromberg [12] or Magid [35].

**Hyperbolic Dehn Filling Theorem.** Let M be a compact, hyperbolizable 3-manifold and let F be a toroidal boundary component of M. Let  $N = \mathbb{H}^3/\Gamma$  be a hyperbolic 3-manifold admitting an orientation-preserving homeomorphism  $\psi: M - \partial_T M \to N \cup \partial_c N$ . Let  $\{(p_n, q_n)\}$  be an infinite sequence of distinct pairs of relatively prime integers.

Then, for all sufficiently large n, there exists a (non-faithful) representation  $\beta_n \colon \Gamma \to \mathrm{PSL}_2(\mathbb{C})$  with discrete image such that

- (1)  $\{\beta_n\}$  converges to the identity representation of  $\Gamma$ , and
- (2) if  $i_n: M \to M(p_n, q_n)$  denotes the inclusion map, then for each n, there exists an orientation-preserving homeomorphism

$$\psi_n: M(p_n, q_n) - \partial_T M(p_n, q_n) \to N_{\beta_n} \cup \partial_c N_{\beta_n}$$

such that  $\beta_n \circ \psi_*$  is conjugate to  $(\psi_n)_* \circ (i_n)_*$ , and the restriction of  $\psi_n \circ i_n \circ \psi^{-1}$  to  $\partial_c N$  is conformal.

## 3. Points are usually closed

If S is a closed orientable surface, we showed in [18] that

$$AI(S) = AH(S \times I)/Mod_{+}(S)$$

is not  $T_1$  where  $\operatorname{Mod}_+(S)$  is the group of (isotopy classes of) orientation-preserving homeomorphisms of S. We recall that a topological space is  $T_1$  if all points are closed sets. Since  $\operatorname{Mod}_+(S)$  is identified with an index two subgroup of  $\operatorname{Out}(\pi_1(S))$ , one also expects that

$$AI(S \times I) = AH(S \times I)/Out(\pi_1(S))$$

is not  $T_1$ .

In this section, we show that if M is an untwisted interval bundle, which also includes the case that M is a handlebody, then AI(M) is not  $T_1$ , but that AI(M) is  $T_1$  for all other compact, hyperbolizable 3-manifolds.

**Theorem 1.1.** Let M be a compact hyperbolizable 3-manifold with non-abelian fundamental group. Then the moduli space AI(M) is  $T_1$  if and only if M is not an untwisted interval bundle.

*Proof.* We first show that AI(M) is  $T_1$  if M is not an untwisted interval bundle. Let  $p: AH(M) \to AI(M)$  be the quotient map and let N be a hyperbolic manifold in AI(M). We must show that  $p^{-1}(N)$  is a closed subset of AH(M). Since AH(M) is Hausdorff and second countable, it suffices to show that if  $\{\rho_n\}$  is a convergent sequence in  $p^{-1}(N)$ , then  $\lim \rho_n \in p^{-1}(N)$ .

An element  $\rho \in p^{-1}(N)$  is a representation such that  $N_{\rho}$  is isometric to N. Let  $\{\rho_n\}$  be a convergent sequence of representations in  $p^{-1}(N)$ . Let  $G \subset M$  be a finite graph such that the inclusion map induces a surjection of  $\pi_1(G)$  onto  $\pi_1(M)$ . Each  $\rho_n$  gives rise to a homotopy equivalence  $h_n \colon M \to N$ , and hence to a map  $j_n = h_n|_G \colon G \to N$ , both of which are only well-defined up to homotopy. Since  $\{\rho_n\}$  is convergent, there exists K such that  $j_n(G)$  has length at most K for all n, after possibly altering  $h_n$  by a homotopy.

Let R be a compact core for N. Assume first that R is not a compression body. In this case, if S is any component of  $\partial R$ , then the inclusion map does not induce a surjection of  $\pi_1(S)$  to  $\pi_1(R)$  (see the discussion in Section 2). Since  $j_n(G)$  carries the fundamental group it cannot lie entirely outside of R. It follows that  $j_n(G)$  lies in the closed neighborhood  $\mathcal{N}_K(R)$  of radius K about R. By compactness, there are only finitely many homotopy classes of maps of G into  $\mathcal{N}_K(R)$  with total length at most K. Hence, there are only finitely many different representations among the  $\rho_n$ , up to conjugacy. The deformation space AH(M) is Hausdorff, and the sequence  $\{\rho_n\}$  converges, implying that  $\{\rho_n\}$  is eventually constant. Therefore  $\lim \rho_n$  lies in the preimage of N, implying that the fiber  $p^{-1}(N)$  is closed and that N is a closed point of AI(M).

Next we assume that R is a compression body. If R were an untwisted interval bundle, then M would also have to be a untwisted interval bundle (by Theorems 5.2 and 10.6 in Hempel [23]) which we have disallowed. So R must have at least one incompressible boundary component and only one compressible boundary component  $\partial_+ R$ . We are free to assume that M is homeomorphic to R, since the definition of AI(M) depends only on the homotopy type of M. Let D denote the union of R and the component of N-R bounded by  $\partial_+ R$ . Since the fundamental group of a component of N-D never surjects onto  $\pi_1(N)$ , with respect to the map induced by inclusion, we see as above that each  $j_n(G)$  must intersect D, so is contained in the neighborhood of radius K of D.

Recall that there exists  $\epsilon_K > 0$  so that the distance from the  $\epsilon_K$ -thin part of N to the  $\mu$ -thick part of N is greater than K (where  $\mu$  is the Margulis constant). It follows

that  $j_n(G)$  must be contained in the  $\epsilon_K$ -thick part of N.

Let F be an incompressible boundary component of M. Then  $h_n(F)$  is homotopic to an incompressible boundary component of R (see, for example, the proof of Proposition 9.2.1 in [17]). As there are finitely many possibilities, we may pass to a subsequence so that  $h_n(F)$  is homotopic to a fixed boundary component F'. We may choose G so that there is a proper subgraph  $G_F \subset G$  such that the image of  $\pi_1(G_F)$  in  $\pi_1(M)$  (under the inclusion map) is conjugate to  $\pi_1(F)$ . Let  $p_F \colon N_F \to N$  be the covering map associated to  $\pi_1(F') \subset \pi_1(N)$ . Then  $j_n|_{G_F}$  lifts to a map  $k_n$  of  $G_F$  into  $N_F$ .

Assume first that F is a torus. Then  $k_n(G_F)$  must lie in the portion X of  $N_F$  with injectivity radius between  $\epsilon_K$  and K/2, which is compact. It follows that  $j_n(G)$  must lie in the closed neighborhood of radius K of  $p_F(X)$ . Since  $p_F(X)$  is compact, we may conclude, as in the general case, that  $\{\rho_n\}$  is eventually constant and hence that  $p^{-1}(N)$  is closed.

We now suppose that F has genus at least 2. We first establish that there exists L such that  $k_n(G_F)$  must be contained in a neighborhood of radius L of the convex core  $C(N_F)$ . It is a consequence of the thick-thin decomposition, that if G' is a graph in  $N_F$  which carries the fundamental group then G' must have length at least  $\mu$ . We also recall that the nearest point retraction  $r_F: N_F \to C(N_F)$  is a homotopy equivalence which is  $\frac{1}{\cosh s}$ -Lipschitz on the complement of the neighborhood of radius s of C(N). Therefore, if  $k_n(G_F)$  lies outside of  $\mathcal{N}_s(C(N_F))$ , then  $r_F(k_n(G_F))$  has length at most  $\frac{K}{\cosh s}$ . It follows that  $k_n(G_F)$  must intersect the neighborhood of radius  $\cosh^{-1}(\frac{K}{\mu})$  of  $C(N_F)$ , so we may choose  $L = K + \cosh^{-1}(\frac{K}{\mu})$ .

If  $N_F$  is geometrically finite, then  $X = C(N_F) \cap N_{\text{thick}}(\epsilon_K)$  is compact and  $j_n(G)$  must be contained in the neighborhood of radius L + K of  $p_F(X)$  which allows us to complete the proof as before.

If  $N_F$  is not geometrically finite, we will need to invoke the Covering Theorem to complete the proof. Let  $\tilde{F}$  denote the lift of F' to  $N_F$ . Then  $\tilde{F}$  divides  $N_F$  into two components, one of which, say  $A_-$ , is mapped homeomorphically to the component of N-R bounded by F'. Let  $A_+=N_F-A_-$ . We may choose a relative compact core  $R_F$  for  $(N_F)^0_\epsilon$  (for some  $\epsilon<\epsilon_K$ ) so that  $\tilde{F}$  is contained in the interior of  $R_F$ . Since  $p_F$  is infinite-to-one on each end of  $(N_F)^0_\epsilon$  which is contained in  $A_+$ , the Covering Theorem (see [15] or [53]) implies that all such ends are geometrically finite. Therefore,

$$Y = A_{+} \cap C(N_{F}) \cap (N_{F})_{\text{thick}(\epsilon_{K})}$$

is compact. If we let  $Z = A_- \cup Y$ , then we see that  $k_n(G_F)$  is contained in the closed neighborhood of radius L about Z (since  $C(N_F) \cap N_{\text{thick}(\epsilon_K)} \subset Z$ ). Therefore,  $j_n(G)$  is contained in the closed (L + K)-neighborhood of

$$D \cap p_F(Z) = D \cap p_F(Y).$$

Since  $D \cap p_F(Y)$  is compact, we conclude, exactly as in the previous cases, that

 $p^{-1}(N)$  is closed. This case completes the proof that AI(M) is  $T_1$  if M is not an untwisted interval bundle.

We now deal with the case where  $M = S \times I$  is an untwisted interval bundle over a compact surface S. (In the special case that M is a handlebody of genus 2, we choose S to be the punctured torus.) In our previous paper [18], we consider the space AH(S) of (conjugacy classes of) discrete faithful representations  $\rho \colon \pi_1(S) \to \mathrm{PSL}_2(\mathbb{C})$  such that if  $g \in \pi_1(S)$  is peripheral, then  $\rho(g)$  is parabolic. In Proposition 3.1, we use work of Thurston [51] and McMullen [41] to exhibit a sequence  $\{\rho_n\}$  in AH(S) which converges to  $\rho \in AH(S)$  such that  $\Lambda(\rho) = \widehat{\mathbb{C}}$ ,  $\Lambda(\rho_1) \neq \widehat{\mathbb{C}}$  and for all n there exists  $\varphi_n \in \mathrm{Mod}_+(S)$  such that  $\rho_n = \rho_1 \circ \varphi_n$ . Since  $AH(S) \subset AH(S \times I)$  and  $\mathrm{Mod}_+(S)$  is identified with a subgroup of  $\mathrm{Out}(\pi_1(S))$ , we see that  $\{\rho_n\}$  is a sequence in  $p^{-1}(N_{\rho_1})$  which converges to a point outside of  $p^{-1}(N_{\rho_1})$ . Therefore,  $N_{\rho_1}$  is a point in  $AI(S \times I)$  which is not closed.

**Remark.** One may further show, as in the remark after Proposition 3.1 in [18], that if  $N \in AI(S \times I)$  is a degenerate hyperbolic 3-manifold with a lower bound on its injectivity radius, then N is not a closed point in  $AI(S \times I)$ . We recall that  $N = \mathbb{H}^3/\Gamma$  is degenerate if  $\Omega(\Gamma)$  is connected and simply connected and  $\Gamma$  is finitely generated.

## 4. Primitive essential annuli and the failure of proper discontinuity

In this section, we show that if M contains a primitive essential annulus, then  $\operatorname{Out}(\pi_1(M))$  does not act properly discontinuously on AH(M). We do so by using the Hyperbolic Dehn Filling Theorem to produce a convergent sequence  $\{\rho_n\}$  in AH(M) and a sequence  $\{\varphi_n\}$  of distinct element of  $\operatorname{Out}(\pi_1(M))$  such that  $\{\rho_n \circ \varphi_n\}$  is also convergent. The construction is a generalization of a construction of Kerckhoff–Thurston [31]. One may also think of the argument as a simple version of the "wrapping" construction (see Anderson–Canary [2]) which was also used to show that components of  $\operatorname{int}(AH(M))$  self-bump whenever M contains a primitive essential annulus (see McMullen [42] and Bromberg–Holt [13]).

**Theorem 1.2.** Let M be a compact hyperbolizable 3-manifold with non-abelian fundamental group. If M contains a primitive essential annulus then  $Out(\pi_1(M))$  does not act properly discontinuously on AH(M). Moreover, if M contains a primitive essential annulus, then AI(M) is not Hausdorff.

*Proof.* Let A be a primitive essential annulus in M with core curve  $\alpha$ . Let  $\widehat{M} = M - \mathcal{N}(\alpha)$  where  $\mathcal{N}(\alpha)$  is an open regular neighborhood of  $\alpha$ . Lemma 10.2 in [3] observes that  $\widehat{M}$  is hyperbolizable. Since  $\widehat{M}$  is hyperbolizable, Thurston's Hyperbolization Theorem implies that there exists a hyperbolic manifold  $\widehat{N}$  and a homeomorphism

 $\psi: \widehat{M} - \partial_T \widehat{M} \to \widehat{N} \cup \partial_c \widehat{N}$ . The classical deformation theory of Kleinian groups (see Bers [5] or [17]) implies that we may choose any conformal structure on  $\partial_c \widehat{N}$ .

Let  $A_0$  and  $A_1$  denote the components of  $A \cap \widehat{M}$ . Let  $M_i$  be the complement in  $\widehat{M}$  of a regular neighborhood of  $A_i$ . Let  $h_i : M \to \widehat{M}$  be an embedding with image  $M_i$  which agrees with the identity map off of a (somewhat larger) regular neighborhood of A.

Let F be the toroidal boundary component of  $\widehat{M}$  which is the boundary of  $\mathcal{N}(\alpha)$  in M. Choose a meridian-longitude system for F so that the meridian for F bounds a disk in M and the longitude is isotopic to  $A_1 \cap F$ . Lemma 10.3 in [3] implies that if  $i_n : \widehat{M} \to \widehat{M}(1,n)$  is the inclusion map, then  $i_n \circ h_i : M \to \widehat{M}(1,n)$  is homotopic to a homeomorphism for each i=0,1 and all  $n\in\mathbb{Z}$ . Moreover, we may similarly check that  $i_n \circ h_1$  is homotopic to  $i_n \circ h_0 \circ D_A^n$  for all n, where  $D_A$  denotes a Dehn twist along A. Notice first that  $j_n = D_{A_0}^n$  takes a (1,0)-curve on F to a (1,n)-curve on F, so extends to a homeomorphism  $j_n : M = \widehat{M}(1,0) \to \widehat{M}(1,n)$ . Therefore, since  $i_0 \circ h_0$  and  $i_0 \circ h_1$  are homotopic, so are  $j_n \circ i_0 \circ h_0$  and  $j_n \circ i_0 \circ h_1$ . But,  $j_n \circ i_0 \circ h_0$  is homotopic to  $i_n \circ h_0 \circ D_A^n$  and  $j_n \circ i_0 \circ h_1$ , which completes the proof that  $i_n \circ h_1$  is homotopic to  $i_n \circ h_0 \circ D_A^n$  for all n.

Let  $\rho_0 = (\psi \circ h_0)_*$  and  $\rho_1 = (\psi \circ h_1)_*$ . Since  $(h_i)_*$  induces an injection of  $\pi_1(M)$  into  $\pi_1(\hat{M})$ ,  $\rho_i \in AH(M)$ . We next observe that one can choose  $\hat{N}$  so that  $N_{\rho_0}$  and  $N_{\rho_1}$  are not isometric. Let  $a_i = A_i \cap (\partial M - \partial_T \hat{M})$  and let  $a_i^*$  denote the geodesic representative of  $\psi(a_i)$  in  $\partial_c \hat{N}$ . Notice that for each i = 0, 1 there is a conformal embedding of  $\partial_c \hat{N} - a_i^*$  into  $\partial_c N_{\rho_i}$  such that each component of the complement of the image of  $\partial_c \hat{N} - a_i^*$  is a neighborhood of a cusp. One may therefore choose the conformal structure on  $\partial_c \hat{N}$  so that there is not a conformal homeomorphism from  $\partial_c N_{\rho_0}$  to  $\partial_c N_{\rho_1}$ . Therefore,  $N_{\rho_0}$  and  $N_{\rho_1}$  are not isometric.

Let  $\{N_n = N_{\beta_n}\}$  be the sequence of hyperbolic 3-manifolds provided by the Hyperbolic Dehn Filling Theorem applied to the sequence  $\{(1,n)\}_{n\in\mathbb{Z}_+}$  and let  $\{\psi_n : \widehat{M}(1,n) - \partial_T \widehat{M}(1,n) \to N_n \cup \partial_c N_n\}$  be the homeomorphisms such that  $\psi_n \circ i_n \circ \psi^{-1}$  is conformal on  $\partial_c N$ . Let

$$\rho_{n,i} = \beta_n \circ \rho_i$$

for all n large enough that  $N_n$  and  $\psi_n$  exist. Since  $\beta_n \circ \psi_*$  is conjugate to  $(\psi_n \circ i_n)_*$  (by applying part (2) of the Hyperbolic Dehn Filling Theorem) and  $i_n \circ h_i$  is homotopic to a homeomorphism, we see that  $\rho_{n,i} = (\psi_n \circ i_n \circ h_i)_*$  lies in AH(M) for all n and each i. It follows from part (1) of the Hyperbolic Dehn Filling Theorem that  $\{\rho_{n,i}\}$  converges to  $\rho_i$  for each i. Moreover,  $\rho_{n,1} = \rho_{n,0} \circ (D_A)_*^n$  for all n, since  $i_n \circ h_1$  is homotopic to  $i_n \circ h_0 \circ D_A^n$  for all n. Therefore,  $\operatorname{Out}(\pi_1(M))$  does not act properly discontinuously on AH(M).

Moreover,  $\{\rho_{n,0}\}$  and  $\{\rho_{n,1}\}$  project to the same sequence in AI(M) and both  $N_{\rho_0}$  and  $N_{\rho_1}$  are limits of this sequence. Since  $N_{\rho_0}$  and  $N_{\rho_1}$  are distinct manifolds in AI(M), it follows that AI(M) is not Hausdorff.

**Remark.** One can also establish Theorem 1.2 using deformation theory of Kleinian groups and convergence results of Thurston [52]. This version of the argument follows the same outline as the proof of Proposition 3.3 in [18].

We provide a brief sketch of this argument. The classical deformation theory of Kleinian groups (in combination with Thurston's Hyperbolization Theorem) guarantees that there exists a component B of  $\operatorname{int}(AH(M))$  such that if  $\rho \in B$ , then there exists a homeomorphism  $\bar{h}_{\rho} \colon M - \partial_T M \to N_{\rho} \cup \partial_c N_{\rho}$  and the point  $\rho$  is determined by the induced conformal structure on  $\partial M - \partial_T M$ . Moreover, every possible conformal structure on  $\partial M - \partial_T M$  arises in this manner.

Let  $a_0$  and  $a_1$  denote the components of  $\partial A$  and let  $t_{a_0}$  and  $t_{a_1}$  denote Dehn twists about  $a_0$  and  $a_1$  respectively. We choose orientations so that  $D_A$  induces  $t_{a_0} \circ t_{a_1}$  on  $\partial M$ . We then let  $\rho_{n,0} \in B$  have associated conformal structure  $t_{a_0}^n(X)$  and let  $\rho_{n,1}$  have associated conformal structure  $t_{a_0}^{-n}(X)$  for some conformal structure X on  $\partial M$ . Thurston's convergence results [51], [52] can be used to show that there exists a subsequence  $\{n_j\}$  of  $\mathbb Z$  such that  $\{\rho_{n_j,0}\}$  and  $\{\rho_{n_j,1}\}$  both converge. One can guarantee, roughly as above, that the limiting hyperbolic manifolds are not isometric. Moreover,  $\rho_{n,1} = \rho_{n,0} \circ (D_A)_*^n$  for all n, so we are the same situation as in the proof above.

# 5. The characteristic submanifold and mapping class groups

In order to further analyze the case where *M* has incompressible boundary we will make use of the characteristic submanifold (developed by Jaco–Shalen [27] and Johannson [29]) and the theory of mapping class groups of 3-manifolds developed by Johannson [29] and extended by McCullough and his co-authors [39], [26], [17].

We begin by recalling the definition of the characteristic submanifold, specialized to the hyperbolic setting. In the general setting, the components of the characteristic submanifold are interval bundles and Seifert fibred spaces. In the hyperbolic setting, the only Seifert fibred spaces which occur are the solid torus and the thickened torus (see Morgan [44], Section 11, or Canary–McCullough [17], Chapter 5).

**Theorem 5.1.** Let M be a compact oriented hyperbolizable 3-manifold with incompressible boundary. There exists a codimension zero submanifold  $\Sigma(M) \subseteq M$  with frontier  $\operatorname{Fr}(\Sigma(M)) = \overline{\partial \Sigma(M)} - \overline{\partial M}$  satisfying the following properties:

- (1) Each component  $\Sigma_i$  of  $\Sigma(M)$  is either
  - (i) an interval bundle over a compact surface with negative Euler characteristic which intersects  $\partial M$  in its associated  $\partial I$ -bundle,
  - (ii) a thickened torus such that  $\partial M \cap \Sigma_i$  contains a torus, or
  - (iii) a solid torus.

- (2) The frontier  $Fr(\Sigma(M))$  is a collection of essential annuli.
- (3) Any essential annulus or incompressible torus in M is properly isotopic into  $\Sigma(M)$ .
- (4) If X is a component of  $M \Sigma(M)$ , then either  $\pi_1(X)$  is non-abelian or  $(\overline{X}, \operatorname{Fr}(X)) \cong (S^1 \times [0,1] \times [0,1], S^1 \times [0,1] \times \{0,1\})$  and X lies between an interval bundle component of  $\Sigma(M)$  and a thickened or solid torus component of  $\Sigma(M)$ .

Moreover, such a  $\Sigma(M)$  is unique up to isotopy, and is called the **characteristic** submanifold of M.

The existence and the uniqueness of the characteristic submanifold in general follows from The Characteristic Pair Theorem in [27] or Proposition 9.4 and Corollary 10.9 in [29]. Theorem 5.1(1) follows from Theorem 5.3.4 of [17], part (2) follows from (1) and the definition of the characteristic submanifold, part (3) follows from Theorem 12.5 of [29], and part (4) follows from Theorem 2.9.3 of [17].

Johannson's Classification Theorem [29] asserts that every homotopy equivalence between compact, irreducible 3-manifolds with incompressible boundary may be homotoped so that it preserves the characteristic submanifold and is a homeomorphism on its complement. Therefore, the study of  $\operatorname{Out}(\pi_1(M))$  often reduces to the study of mapping class groups of interval bundles and Seifert-fibered spaces.

**Johannson's Classification Theorem** ([29], Theorem 24.2). Let M and Q be irreducible 3-manifolds with incompressible boundary and let  $h: M \to Q$  be a homotopy equivalence. Then h is homotopic to a map  $g: M \to Q$  such that

- $(1) g^{-1}(\Sigma(Q)) = \Sigma(M),$
- (2)  $g|_{\Sigma(M)}: \Sigma(M) \to \Sigma(Q)$  is a homotopy equivalence, and
- (3)  $g|_{\overline{M-\Sigma(M)}}: \overline{M-\Sigma(M)} \to \overline{Q-\Sigma(Q)}$  is a homeomorphism.

Moreover, if h is a homeomorphism, then g is a homeomorphism.

We let the mapping class group  $\operatorname{Mod}(M)$  denote the group of isotopy classes of self-homeomorphisms of M. Since M is irreducible and has (non-empty) incompressible boundary, any two homotopic homeomorphisms are isotopic (see Waldhausen [55], Theorem 7.1), so  $\operatorname{Mod}(M)$  is naturally a subgroup of  $\operatorname{Out}(\pi_1(M))$ . For simplicity, we will assume that M is a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components. Notice that this implies that  $\Sigma(M)$  contains no thickened torus components. Let  $\Sigma$  be the characteristic submanifold of M and denote its components by  $\{\Sigma_1, \ldots, \Sigma_k\}$ .

Following McCullough [39], we let  $Mod(\Sigma_i, Fr(\Sigma_i))$  denote the group of homotopy classes of homeomorphisms  $h: \Sigma_i \to \Sigma_i$  such that h(F) = F for each component F of  $Fr(\Sigma_i)$ . We let  $G(\Sigma_i, Fr(\Sigma_i))$  denote the subgroup consisting of

(homotopy classes of) homeomorphisms which have representatives which are the identity on  $Fr(\Sigma_i)$ . Define

$$G(\Sigma, \operatorname{Fr}(\Sigma)) = \bigoplus_{i=1}^k G(\Sigma_i, \operatorname{Fr}(\Sigma_i)).$$

Notice that using these definitions, the restriction of a Dehn twist along a component of  $Fr(\Sigma)$  is trivial in  $G(\Sigma, Fr(\Sigma))$ .

In our case, each  $\Sigma_i$  is either an interval bundle over a compact surface  $F_i$  with negative Euler characteristic or a solid torus. If  $\Sigma_i$  is a solid torus, then  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is finite (see Lemma 10.3.2 in [17]). If  $\Sigma_i$  is an interval bundle over a compact surface  $F_i$ , then  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is isomorphic to the group  $G(F_i, \partial F_i)$  of proper isotopy classes of self-homeomorphisms of F which are the identity on  $\partial F$  (see Proposition 3.2.1 in [39] and Lemma 6.1 in [26]). Moreover,  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  injects into  $\operatorname{Out}(\pi_1(\Sigma_i))$  (see Proposition 5.2.3 in [17] for example). We say that  $\Sigma_i$  is tiny if its base surface  $F_i$  is either a thrice-punctured sphere or a twice-punctured projective plane. If  $\Sigma_i$  is not tiny, then  $F_i$  contains a 2-sided, non-peripheral homotopically non-trivial simple closed curve, so  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is infinite. If  $\Sigma_i$  is tiny, then  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is finite (see Korkmaz [32] for the case when  $F_i$  is a twice-punctured projective plane).

Let J(M) be the subgroup of  $\operatorname{Mod}(M)$  consisting of classes represented by homeomorphisms fixing  $M-\Sigma$  pointwise. Lemma 4.2.1 of McCullough [39] implies that J(M) has finite index in  $\operatorname{Mod}(M)$ . (Instead of J(M), McCullough writes  $\mathcal{K}(M, \Sigma_1, \Sigma_2, \ldots, \Sigma_k)$ .) Lemma 4.2.2 of McCullough [39] implies that the kernel K(M) of the natural surjective homomorphism

$$p_{\Sigma}: J(M) \to G(\Sigma, \operatorname{Fr}(\Sigma))$$

is abelian and is generated by Dehn twists about the annuli in  $Fr(\Sigma)$ .

We summarize the discussion above in the following statement.

**Theorem 5.2.** Let M be a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components. Then there is a finite index subgroup J(M) of Mod(M) and an exact sequence

$$1 \longrightarrow K(M) \longrightarrow J(M) \xrightarrow{p_{\Sigma}} G(\Sigma, \operatorname{Fr}(\Sigma)) \longrightarrow 1$$

such that K(M) is an abelian group generated by Dehn twists about essential annuli in  $Fr(\Sigma)$ .

Suppose that  $\Sigma_i$  is a component of  $\Sigma(M)$ . If  $\Sigma_i$  is a solid torus or a tiny interval bundle, then  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is finite. Otherwise,  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is infinite and injects into  $\operatorname{Out}(\pi_1(\Sigma_i))$ .

#### 6. Characteristic collections of annuli

We continue to assume that M has incompressible boundary and no toroidal boundary components and that  $\Sigma(M)$  is its characteristic submanifold. In this section, we organize K(M) into subgroups generated by collections of annuli with homotopic core curves, called characteristic collection of annuli, and define a class of free subgroups of  $\pi_1(M)$  which "register" these subgroups of K(M).

A characteristic collection of annuli for M is either a) the collection of all frontier annuli in a solid torus component of  $\Sigma(M)$ , or b) an annulus in the frontier of an interval bundle component of  $\Sigma(M)$  which is not properly isotopic to a frontier annulus of a solid torus component of  $\Sigma(M)$ .

If  $C_j$  is a characteristic collection of annuli for M, let  $K_j$  be the subgroup of K(M) generated by Dehn twists about the annuli in  $C_j$ . Notice that  $K_i \cap K_j = \{id\}$  for  $i \neq j$ , since each element of  $K_j$  fixes any curve disjoint from  $C_j$ . Then  $K(M) = \bigoplus_{j=1}^m K_j$ , since every frontier annulus of  $\Sigma(M)$  is properly isotopic to a component of some characteristic collection of annuli. Let  $q_j : K(M) \to K_j$  be the projection map.

We next introduce free subgroups of  $\pi_1(M)$ , called  $C_j$ -registering subgroups, which are preserved by  $K_i$  and such that  $K_j$  acts effectively on the subgroup.

We first suppose that  $C_j = \operatorname{Fr}(T_j)$  where  $T_j$  is a solid torus component of  $\Sigma(M)$ . Let  $\{A_1, \ldots, A_l\}$  denote the components of  $\operatorname{Fr}(T_j)$ . For each  $i=1,\ldots,l$ , let  $X_i$  be the component of  $M-(T_j\cup C_1\cup C_2\cup\ldots\cup C_m)$  abutting  $A_i$ . (Notice that each  $X_i$  is either a component of  $M-\Sigma(M)$  or properly isotopic to the interior of an interval bundle component of  $\Sigma(M)$ .) Let a be a core curve for  $T_j$  and let  $x_0$  be a point on a. We say that a subgroup H of  $\pi_1(M,x_0)$  is  $C_j$ -registering if it is freely (and minimally) generated by a and, for each  $i=1,\ldots,l$ , a loop  $g_i$  in  $T_j\cup X_i$  based at  $x_0$  intersecting  $A_i$  exactly twice. In particular, every  $C_j$ -registering subgroup of  $\pi_1(M,x_0)$  is isomorphic to  $F_{l+1}$ .

Notice that a Dehn twist  $D_{A_i}$  along any  $A_i$  preserves H in  $\pi_1(M, x_0)$ . It acts on H by the map  $t_i$  which fixes a and  $g_m$  for  $m \neq i$ , and conjugates  $g_i$  by  $a^n$  (where the core curve of  $A_i$  is homotopic to  $a^n$ ). Let  $s_H: K_j \to \operatorname{Out}(H)$  be the homomorphism which takes each  $D_{A_i}$  to  $t_i$ . Simultaneously twisting along all l annuli induces conjugation by  $a^n$ , which is an inner automorphism of H. Moreover, it is easily checked that  $s_H(K_j)$  is isomorphic to  $\mathbb{Z}^{l-1}$  and is generated by  $\{t_1, \ldots, t_{l-1}\}$ . The set  $\{a, g_1, \ldots, g_l\}$  may be extended to a generating set for  $\pi_1(M, x_0)$  by appending curves which intersect  $\operatorname{Fr}(T_j)$  exactly twice, so  $D_{A_1} \circ \cdots \circ D_{A_l}$  acts as conjugation by  $a^n$  on all of  $\pi_1(M, x_0)$ . Therefore,  $K_j$  itself is isomorphic to  $\mathbb{Z}^{l-1}$  and  $s_H$  is injective. (In particular, if  $C_j$  is a single annulus in the boundary of a solid torus component of  $\Sigma(M)$ , then  $K_j$  is trivial and we could have omitted  $C_j$ .)

Now suppose that  $C_j = \{A\}$  is a frontier annulus of an interval bundle component  $\Sigma_i$  of  $\Sigma$  which is not properly isotopic into a solid torus component of  $\Sigma$ . Let a be a core curve for A and let  $x_0$  be a point on a. We say that a subgroup H of  $\pi_1(M, x_0)$ 

is  $C_j$ -registering if it is freely (and minimally) generated by a and two loops  $g_1$  and  $g_2$  based at  $x_0$  each of whose interiors misses A, and which lie in the two distinct components of  $M - (C_1 \cup C_2 \cup \ldots \cup C_m)$  abutting A. In this case, H is isomorphic to  $F_3$ . Arguing as above, it follows that  $K_j$  is an infinite cyclic subgroup of  $Out(\pi_1(M))$  and there is an injection  $s_H : K_j \to Out(H)$ .

In either situation, if H is a  $C_j$ -registering group for a characteristic collection of annuli  $C_j$ , then we may consider the map

$$r_H: X(M) \to X(H)$$

simply obtained by taking  $\rho$  to  $\rho|_H$ . (Here, X(H) is the  $PSL_2(\mathbb{C})$ -character variety of the abstract group H.) One easily checks from the description above that if  $\alpha \in K_j$ , then  $r_H(\rho \circ \alpha) = r_H(\rho) \circ s_H(\alpha)$  for all  $\rho \in X(M)$ . Notice that if  $\varphi \in K_l$  and  $j \neq l$ , then  $K_l$  acts trivially on H, since each generating curve of H is disjoint from  $C_l$ . Therefore,

$$r_H(\rho \circ \alpha) = r_H(\rho) \circ s_H(q_i(\alpha))$$

for all  $\rho \in X(M)$  and  $\alpha \in K(M)$ .

We summarize the key points of this discussion for use later:

**Lemma 6.1.** Let M be a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components. If  $C_j$  is a characteristic collection of annuli for M and H is a  $C_j$ -registering subgroup of  $\pi_1(M)$ , then H is preserved by each element of  $K_j$  and there is a natural injective homomorphism  $s_H: K_j \to \operatorname{Out}(H)$ . Moreover, if  $\alpha \in K(M)$ , then  $r_H(\rho \circ \alpha) = r_H(\rho) \circ s_H(q_j(\alpha))$  for all  $\rho \in X(M)$ .

# 7. Primitive essential annuli and manifolds with compressible boundary

In this section we use a result of Johannson [29] to show that every compact hyperbolizable 3-manifolds with compressible boundary and no toroidal boundary components contains a primitive essential annulus. It then follows from Theorem 1.2 that if M has compressible boundary and no toroidal boundary components, then  $Out(\pi_1(M))$  fails to act properly discontinuously on AH(M) and AI(M) is not Hausdorff.

We first find indivisible curves in the boundary of compact hyperbolizable 3-manifolds with incompressible boundary and no toroidal boundary components. We call a curve a in M indivisible if it generates a maximal cyclic subgroup of  $\pi_1(M)$ .

**Lemma 7.1.** Let M be a compact hyperbolizable 3-manifold with (non-empty) incompressible boundary. Then, if F is a component of  $\partial M$ , there exists an indivisible simple closed curve in F.

*Proof.* We use a special case of a result of Johannson [29] (see also Jaco–Shalen [28]) which characterizes divisible simple closed curves in  $\partial M$ .

**Lemma 7.2** ([29], Lemma 32.1). Let M be a compact hyperbolizable 3-manifold with incompressible boundary. An essential simple closed curve  $\alpha$  in  $\partial M$  which is not indivisible is either isotopic into a solid torus component of  $\Sigma(M)$  or is isotopic to a boundary component of an essential Möbius band in an interval bundle component of  $\Sigma(M)$ .

Therefore, if  $\Sigma(M)$  is not all of M, then any simple closed curve in F which cannot be isotoped into a solid torus or interval bundle component of  $\Sigma(M)$  is indivisible.

If  $\Sigma(M) = M$ , then M is an interval bundle over a closed surface with negative Euler characteristic and the proof is completed by the following lemma, whose full statement will be used later in the paper.

**Lemma 7.3.** Let M be a compact hyperbolizable 3-manifold with no toroidal boundary components. Let  $\Sigma_i$  be an interval bundle component of  $\Sigma(M)$  which is not tiny, then there is a primitive essential annulus (for M) contained in  $\Sigma_i$ .

<i>Proof.</i> Let $F_i$ be the base surface of $\Sigma_i$ .	Since $\Sigma_i$ is not tiny, $F_i$ contains a non-
peripheral simple closed curve a which is	two-sided and homotopically non-trivial.
Then $a$ is an indivisible curve in $F_i$ and he	nce in $M$ . The sub-interval bundle $A$ over
a is thus a primitive essential annulus.	

We are now prepared to prove the main result of the section.

**Proposition 7.4.** If M is a compact hyperbolizable 3-manifold with compressible boundary and no toroidal boundary components, then M contains a primitive essential annulus.

*Proof.* We first observe that under our assumptions every maximal abelian subgroup of  $\pi_1(M)$  is cyclic (since every non-cyclic abelian subgroup of the fundamental group of a compact hyperbolizable 3-manifold is conjugate into the fundamental group of a toroidal component of  $\partial M$ , see [44], Corollary 6.10). Therefore, in our case an essential annulus is primitive if and only if its core curve is indivisible.

We first suppose that M is a compression body. If M is a handlebody, then it is an interval bundle, so contains a primitive essential annulus by Lemma 7.3. Otherwise, M is formed from  $R \times I$  by appending 1-handles to  $R \times \{1\}$ , where R is a closed, but not necessarily connected, orientable surface. Let  $\alpha$  be an essential simple closed curve in  $R \times \{1\}$  which lies in  $\partial M$ . Let D be a disk in  $R \times \{1\} - \partial M$ . We may assume that  $\alpha$  intersects  $\partial D$  in exactly one point. Let  $\beta \subset (\partial M \cap R \times \{1\})$  be a simple

closed curve homotopic to  $\alpha * \partial D$  (in  $\partial M$ ) and disjoint from  $\alpha$ . Then  $\alpha$  and  $\beta$  bound an embedded annulus in  $R \times \{1\}$ , which may be homotoped to a primitive essential annulus in M (by pushing the interior of the annulus into the interior of  $R \times I$ ).

If M is not a compression body, let  $C_M$  be a characteristic compression body neighborhood of  $\partial M$  (as discussed in Section 2). Let C be a component of  $C_M$  which has a compressible boundary component  $\partial_+ C$  and an incompressible boundary component F. Let X be the component of  $\overline{M-C_M}$  which contains F in its boundary and let  $\alpha$  be an essential simple closed curve in F which is indivisible in X (which exists by Lemma 7.1). Let  $\alpha'$  be a curve in  $\partial_+ C \subset \partial M$  which is homotopic to  $\alpha$ . One may then construct as above a primitive essential annulus A in C with  $\alpha'$  as one boundary component. It is clear that A remains essential in M. Since  $\pi_1(M) = \pi_1(X) * H$  for some group H, the core curve of A, which is homotopic to  $\alpha$ , is indivisible in  $\pi_1(M)$ . Therefore, A is our desired primitive essential annulus in M.

**Remark.** The above argument is easily extended to the case where M is allowed to have toroidal boundary components (but is still hyperbolizable), unless M is a compression body all of whose boundary components are tori. In fact, the only counterexamples in this situation occur when M is obtained from one or two untwisted interval bundles over tori by attaching exactly one 1-handle.

We have thus already established Corollary 1.4 in the case that M has compressible boundary.

**Corollary 7.5.** If M is a compact hyperbolizable 3-manifold with compressible boundary, no toroidal boundary components, and non-abelian fundamental group, then  $Out(\pi_1(M))$  does not act properly discontinuously on AH(M). Moreover, the moduli space AI(M) is not Hausdorff.

# 8. The space $AH_n(M)$

In this section, we assume that M has incompressible boundary and no toroidal boundary components. We identify a subset  $AH_n(M)$  of AH(M) which contains all purely hyperbolic representations in AH(M). We will see later that  $Out(\pi_1(M))$  acts properly discontinuously on an open neighborhood of  $AH_n(M)$  in X(M) if M is not an interval bundle.

We define  $AH_n(M)$  to be the set of (conjugacy classes of) representations  $\rho \in AH(M)$  such that

(1) If  $\Sigma_i$  is a component of the characteristic submanifold which is not a tiny interval bundle, then  $\rho(\pi_1(\Sigma_i))$  is purely hyperbolic (i.e. if g is a non-trivial element of  $\pi_1(M)$  which is conjugate into  $\pi_1(\Sigma_i)$ , then  $\rho(g)$  is hyperbolic), and

(2) if  $\Sigma_i$  is a tiny interval bundle, then  $\rho(\pi_1(\operatorname{Fr}(\Sigma_i)))$  is purely hyperbolic.

We observe that int(AH(M)) is a proper subset of  $AH_n(M)$  and that  $AH(M) = AH_n(M)$  if and only if M contains no primitive essential annuli.

**Lemma 8.1.** Let M be a compact hyperbolizable 3-manifold with non-empty incompressible boundary and no toroidal boundary components. Then

- (1) the interior of AH(M) is a proper subset of  $AH_n(M)$ ,
- (2)  $AH_n(M)$  contains a dense subset of  $\partial AH(M)$ , and
- (3)  $AH_n(M) = AH(M)$  if and only if M contains no primitive essential annuli.

*Proof.* Sullivan [50] proved that all representations in int(AH(M)) are purely hyperbolic (if M has no toroidal boundary components), so clearly int(AH(M)) is contained in  $AH_n(M)$ . On the other hand,  $\partial AH(M)$  is non-empty (see Lemma 4.1 in Canary–Hersonsky [16]) and purely hyperbolic representations are dense in  $\partial AH(M)$  (which follows from Lemma 4.2 in [16] and the Density Theorem [9], [10], [45], [47]). This establishes claims (1) and (2).

If M contains a primitive essential annulus A, then there exist  $\rho \in AH(M)$  such that  $\rho(\alpha)$  is parabolic (where  $\alpha$  is the core curve of A), so  $AH_n(M)$  is not all of AH(M) in this case (see Ohshika [46]).

Now suppose that M contains no primitive essential annuli. We first note that every component of  $\Sigma(M)$  is a solid torus or tiny interval bundle (by Lemma 7.3). Moreover, if  $\Sigma_i$  is a tiny interval bundle component of  $\Sigma(M)$ , then any component A of its frontier must be isotopic to a component of the frontier of a solid torus component of  $\Sigma(M)$ . Otherwise, A would be a primitive essential annulus (by Lemma 7.2). Therefore, it suffices to prove that  $\rho(\Sigma_i)$  is purely hyperbolic whenever  $\Sigma_i$  is a solid torus component of  $\Sigma(M)$ .

Let T be a solid torus component of  $\Sigma(M)$ . A frontier annulus A of T is an essential annulus in M, so it must not be primitive. It follows that the core curve a of T is not peripheral in M (see [29], Theorem 32.1).

Let  $\rho \in AH(M)$  and let R be a relative compact core for  $(N_{\rho})^{0}_{\epsilon}$  (for some  $\epsilon < \mu$ ). Let  $h: M \to R$  be a homotopy equivalence in the homotopy class determined by  $\rho$ . By Johannson's Classification Theorem ([29], Theorem 24.2) h may be homotoped so that h(T) is a component T' of  $\Sigma(R)$ ,  $h|_{Fr(T)}$  is an embedding with image Fr(T') and  $h|_{T}: (T, Fr(T)) \to (T', Fr(T'))$  is a homotopy equivalence of pairs. It follows that h(a) is homotopic to the core curve of T' which is not peripheral in R.

If  $\rho(a)$  were parabolic, then h(a) would be homotopic into a non-compact component of  $(N_{\rho})_{\text{thin}(\epsilon)}$  and hence into  $P = R \cap \partial(N_{\rho})^0_{\epsilon} \subset \partial R$ , so h(a) would be peripheral in R. It follows that  $\rho(a)$  is hyperbolic. Since a generates  $\pi_1(T)$ , we see that  $\rho(\pi_1(T))$  is purely hyperbolic. Since T is an arbitrary solid torus component of  $\Sigma(M)$ , we see that  $\rho \in AH_n(M)$ .

We next check that the restriction of  $\rho \in AH_n(M)$  to the fundamental group of an interval bundle component of  $\Sigma(M)$  (which is not tiny) is Schottky. By definition, a Schottky group is a free, geometrically finite, purely hyperbolic subgroup of  $PSL_2(\mathbb{C})$  (see Maskit [36] for a discussion of the equivalence of this definition with more classical definitions).

**Lemma 8.2.** Let M be a compact hyperbolizable 3-manifold with incompressible boundary with no toroidal boundary components which is not an interval bundle. If  $\Sigma_i$  is an interval bundle component of  $\Sigma(M)$  which is not tiny and  $\rho \in AH_n(M)$ , then  $\rho(\pi_1(\Sigma_i))$  is a Schottky group.

*Proof.* By definition  $\rho(\pi_1(\Sigma_i))$  is purely hyperbolic, so it suffices to prove it is free and geometrically finite. Since  $\Sigma_i$  is an interval bundle whose base surface  $F_i$  has non-empty boundary,  $\pi_1(\Sigma_i) \cong \pi_1(F_i)$  is free. Let  $\pi_i : N_i \to N_\rho$  be the cover of  $N_\rho$  associated to  $\rho(\pi_1(\Sigma_i))$ . Since  $\pi_1(\Sigma_i)$  has infinite index in  $\pi_1(M)$ ,  $\pi_i : N_i \to N$  is a covering with infinite degree. Let  $R_i$  be a compact core for  $N_i$ . Since  $\pi_1(R_i)$  is free and  $R_i$  is irreducible,  $R_i$  is a handlebody ([23], Theorem 5.2). Therefore,  $N_i = (N_i)_{\epsilon}^0$  has one end and  $\pi_i$  is infinite-to-one on this end, so the Covering Theorem (see [15]) implies that this end is geometrically finite, and hence that  $N_i$  is geometrically finite. Therefore,  $\rho(\pi_1(\Sigma_i))$  is geometrically finite, completing the proof that it is a Schottky group.

Finally, we check that if  $\rho \in AH_n(M)$  and  $C_j$  is a characteristic collection of annuli, then there exists a  $C_j$ -registering subgroup whose image under  $\rho$  is Schottky.

**Lemma 8.3.** Suppose that M is a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components and  $C_j$  is a characteristic collection of frontier annuli for M. If  $\rho \in AH_n(M)$ , then there exists a  $C_j$ -registering subgroup H of  $\pi_1(M)$  such that  $\rho(H)$  is a Schottky group.

*Proof.* We first suppose that  $C_j = \{A\}$  is a frontier annulus of an interval bundle component of  $\Sigma(M)$  (and that A is not properly isotopic to a frontier annulus of a solid torus component of  $\Sigma(M)$ ) and let  $x_0 \in A$ . We identify  $\pi_1(M)$  with  $\pi_1(M, x_0)$ . Let  $X_1$  and  $X_2$  be the (distinct) components of M-Fr( $\Sigma$ ) abutting A. Notice that each  $X_i$  must have non-abelian fundamental group, since it either contains (the interior of) an interval bundle component of  $\Sigma(M)$  or (the interior of) a component of M- $\Sigma(M)$  which is not a solid torus lying between an interval bundle component of  $\Sigma(M)$  and a solid torus component of  $\Sigma(M)$ .

Let a be the core curve of A (based at  $x_0$ ). By assumption,  $\rho(a)$  is a hyperbolic element. Let F be a fundamental domain for the action of  $\langle \rho(a) \rangle$  on  $\Omega(\langle \rho(a) \rangle)$  which is an annulus in  $\widehat{\mathbb{C}}$ . Since each  $\rho(\pi_1(\overline{X_i}, x_0))$  is discrete, torsion-free and non-abelian, hence non-elementary, we may choose hyperbolic elements  $\gamma_i \in \rho(\pi_1(\overline{X_i}, x_0))$  whose fixed points lie in the interior of F. There exists s > 0 such that one

may choose (round) disks  $D_i^{\pm} \subset \operatorname{int}(F)$  about the fixed points of  $\gamma_i$ , such that  $\gamma_i^s(\operatorname{int}(D_i^-)) = \widehat{\mathbb{C}} - D_i^+$ , and  $D_1^+$ ,  $D_1^-$ ,  $D_2^+$  and  $D_2^-$  are disjoint. Then, the Klein Combination Theorem (commonly referred to as the ping pong lemma), guarantees that  $\rho(a)$ ,  $\gamma_1^s$  and  $\gamma_2^s$  freely generate a Schottky group, see, for example, Theorem C.2 in Maskit [37]. Then each  $\rho^{-1}(\gamma_i^s)$  is represented by a curve  $g_i$  in  $\overline{X}_i$  based at  $x_0$  and a,  $g_1$  and  $g_2$  generate a  $C_j$ -registering subgroup H such that  $\rho(H)$  is Schottky.

Now suppose that  $C_j = \{A_1, \ldots, A_l\}$  is the collection of frontier annuli of a solid torus component  $T_j$  of  $\Sigma(M)$ . Let  $X_i$  be the component of  $M - (T_j \cup C_1 \cup \cdots \cup C_m)$  abutting  $A_i$ . Pick  $x_0$  in  $T_j$  and let a be a core curve of  $T_j$  passing through  $x_0$ . Again each  $X_i$  must have non-abelian fundamental group.

Let F be an annular fundamental domain for the action of  $\langle \rho(a) \rangle$  on the complement in  $\widehat{\mathbb{C}}$  of the fixed points of  $\rho(a)$ . For each i, let  $Y_i = X_i \cup A_i \cup \operatorname{int}(T'_j)$  and pick a hyperbolic element  $\gamma_i$  in  $\rho(\pi_1(Y_i, x_0))$  both of whose fixed points lie in the interior of F. (Notice that even though it could be the case that  $X_i = X_k$  for  $i \neq k$ , we still have that  $\pi_1(Y_i, x_0)$  intersects  $\pi_1(Y_k, x_0)$  only in the subgroup generated by a, so these hyperbolic elements are all distinct.) Then, just as in the previous case, there exists s > 0 such that the elements  $\{\rho(a), \gamma_1^s, \ldots, \gamma_l^s\}$  freely generate a Schottky group. Each  $\rho^{-1}(\gamma_i^s)$  can be represented by a loop  $g_i$  based at  $x_0$  which lies in  $Y_i$  and intersects  $A_i$  exactly twice. Therefore, the group H generated by  $\{a, g_1, \ldots, g_l\}$  is  $C_i$ -registering and  $\rho(H)$  is Schottky.

# 9. Proper discontinuity on $AH_n(M)$

We are finally prepared to prove that  $Out(\pi_1(M))$  acts properly discontinuously on an open neighborhood of  $AH_n(M)$  if M is a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components which is not an interval bundle.

**Theorem 9.1.** Let M be a compact hyperbolizable 3-manifold with nonempty incompressible boundary and no toroidal boundary components which is not an interval bundle. Then there exists an open  $Out(\pi_1(M))$ -invariant neighborhood W(M) of  $AH_n(M)$  in X(M) such that  $Out(\pi_1(M))$  acts properly discontinuously on W(M).

Notice that Theorem 1.3 is an immediate consequence of Proposition 7.4, Lemma 8.1 and Theorem 9.1. Moreover, Theorem 1.5 is an immediate corollary of Lemma 8.1 and Theorem 9.1.

We now provide a brief outline of the section. In Section 9.1 we recall Minsky's work which shows that  $Out(\pi_1(H_n))$  acts properly discontinuously on the open set  $PS(H_n)$  of primitive-stable representations in  $X(H_n)$  where  $H_n$  is the handlebody of genus g. In Section 9.2, we consider the set  $Z(M) \subset X(M)$  such that if  $\rho \in Z(M)$ 

and  $C_j$  is a characteristic collection of annuli, then there exists a  $C_j$ -registering subgroup H of  $\pi_1(M)$  such that  $\rho|_H$  is primitive stable. We use Minsky's work to show that K(M) acts properly discontinuously on Z(M). In Section 9.3, we consider the set V(M) of all representation such that  $\rho|_{\pi_1(\Sigma_i)}$  is primitive-stable whenever  $\Sigma_i$  is an interval bundle component of  $\Sigma(M)$  which is not tiny. We show that if  $\{\alpha_n\}$  is a sequence in J(M) such that  $\{\rho_{\Sigma}(\alpha_n)\}$  is a sequence of distinct elements and K is compact subset of V(M), then  $\{\alpha_n(K)\}$  leaves every compact set. In Section 9.4, we let  $W(M) = Z(M) \cap V(M)$  and combine the work in the previous sections to show that J(M) acts properly discontinuously on W(M). Since J(M) has finite index in  $Out(\pi_1(M))$  (see [17]), this immediately implies Theorem 9.1. Johannson's Classification Theorem is used to show that J(M) is invariant under  $Out(\pi_1(M))$ .

**9.1. Schottky groups and primitive-stable groups.** In this section, we recall Minsky's work [43] on primitive-stable representations of the free group  $F_n$ , where  $n \ge 2$ . An element of  $F_n$  is called *primitive* if it is an element of a minimal free generating set for  $F_n$ . Let X be a bouquet of n circles with base point b and fix a specific identification of  $\pi_1(X,b)$  with  $F_n$ . To a conjugacy class [w] in  $F_n$  one can associated an infinite geodesic in X which is obtained by concatenating infinitely many copies of a cyclically reduced representative of w (here the cyclic reduction is in the generating set associated to the natural generators of  $\pi_1(X,b)$ ). Let  $\mathcal{P}$  denote the set of infinite geodesics in the universal cover  $\widetilde{X}$  of X which project to geodesics associated to primitive words of  $F_n$ .

Given a representation  $\rho \colon F_n \to \mathrm{PSL}_2(\mathbb{C}), x \in \mathbb{H}^3$  and a lift  $\tilde{b}$  of b, one obtains a unique  $\rho$ -equivariant map  $\tau_{\rho,x} \colon \tilde{X} \to \mathbb{H}^3$  which takes  $\tilde{b}$  to x and maps each edge of  $\tilde{X}$  to a geodesic. A representation  $\rho \colon F_n \to \mathrm{PSL}_2(\mathbb{C})$  is *primitive-stable* if there are constants  $K, \delta > 0$  such that  $\tau_{\rho,x}$  takes all the geodesics in  $\mathcal{P}$  to  $(K, \delta)$ -quasi-geodesics in  $\mathbb{H}^3$ . We let  $PS(H_n)$  denote the set of (conjugacy classes) of primitive-stable representations in  $X(H_n)$  where  $H_n$  is the handlebody of genus n.

We summarize the key points of Minsky's work which we use in the remainder of the section. We recall that Schottky space  $S_n \subset X(H_n)$  is the space of discrete faithful representations whose image is a Schottky group and that  $S_n$  is the interior of  $AH(H_n)$ .

# **Theorem 9.2** (Minsky [43]). *If* $n \ge 2$ , *then*

- (1) Out $(F_n)$  acts properly discontinuously on  $PS(H_n)$ ,
- (2)  $PS(H_n)$  is an open subset of  $X(H_n)$ , and
- (3) Schottky space  $S_n$  is a proper subset of  $PS(H_n)$ .

Moreover, if K is any compact subset of  $PS(H_n)$ , and  $\{\alpha_n\}$  is a sequence of distinct elements of  $Out(F_n)$ , then  $\{\alpha_n(K)\}$  exits every compact subset of  $X(H_n)$  (i.e. for any compact subset C of  $X(H_n)$  there exists N such that if  $n \geq N$ , then  $\alpha_n(K) \cap C = \emptyset$ ).

**Remark.** In order to prove our main theorem it would suffice to use Schottky space  $S_n$  in place of  $PS(H_n)$ . However, the subset W(M) we obtain using  $PS(H_n)$  is larger than the one we would obtain using simply  $S_n$ .

**9.2.** Characteristic collection of annuli. We will assume for the remainder of the section that M is a compact hyperbolizable 3-manifold with incompressible boundary and no toroidal boundary components which is not an interval bundle. Main Topological Theorem 2 in Canary and McCullough [17] (which is itself an exercise in applying Johannson's theory) implies that if M has incompressible boundary and no toroidal boundary components, then Mod(M) has finite index in  $Out(\pi_1(M))$ . Therefore, applying Theorem 5.2, we see that J(M) has finite index in  $Out(\pi_1(M))$ . In particular, if M is acylindrical, then J(M) is trivial and  $Out(\pi_1(M))$  acts properly discontinuously on X(M).

Let  $C_j$  be a characteristic collection of annuli in M. If H is a  $C_j$ -registering subgroup of  $\pi_1(M)$ , then the inclusion of H in  $\pi_1(M)$  induces a natural injection  $s_H: K_j \to \operatorname{Out}(H)$  such that if  $\alpha \in K(M)$ , then

$$r_H(\rho \circ \alpha) = r_H(\rho) \circ s_H(q_i(\alpha))$$

where  $r_H(\rho) = \rho|_H$  (see Lemma 6.1). Let

$$Z_H = r_H^{-1}(PS(H))$$

where  $PS(H) \subset X(H)$  is the set of (conjugacy classes of) primitive-stable representations of H. Let

$$Z(C_j) = \bigcup Z_H$$

where the union is taken over all  $C_i$ -registering subgroups H of  $\pi_1(M)$ .

If  $\{C_1, \ldots, C_m\}$  is the set of all characteristic collections of annuli for M, then we define

$$Z(M) = \bigcap_{i=1}^{m} Z(C_i).$$

If there are no characteristic collection of annuli, then M is acylindrical and we set Z(M) = X(M).

We use Lemma 8.3, Theorem 9.2, and Johannson's Classification Theorem to prove:

**Lemma 9.3.** Let M be a compact hyperbolizable 3-manifold with nonempty incompressible boundary and no toroidal boundary components. Then

- (1) Z(M) is an  $Out(\pi_1(M))$ -invariant open neighborhood of  $AH_n(M)$  in X(M), and
- (2) if  $K \subset Z(M)$  is compact and  $\{\alpha_n\}$  is a sequence of distinct elements of K(M), then  $\{\alpha_n(K)\}$  exits every compact set of X(M).

*Proof.* Lemma 8.3 implies that  $AH_n(M) \subset Z(C_j)$  for each j, so  $AH_n(M) \subset Z(H)$ . Moreover, since  $r_H$  is continuous for all H, each  $Z(C_j)$  is open, and hence Z(M) is open.

Johannson's Classification Theorem implies that if  $C_j$  is a characteristic collection of annuli for M and  $\varphi \in \operatorname{Out}(\pi_1(M))$ , then there exists a homotopy equivalence  $h \colon M \to M$  such that  $h_* = \varphi$  and  $h(C_j)$  is also a characteristic collection of annuli for M. Moreover, if H is a  $C_j$ -registering subgroup of  $\pi_1(M)$ , then  $\varphi(H)$  is a  $h(C_j)$ -registering subgroup of  $\pi_1(M)$ . Therefore, Z(M) is  $\operatorname{Out}(\pi_1(M))$ -invariant, completing the proof of claim (1).

If (2) fails to hold, then there is a compact subset K of Z(M), a compact subset C of X(M) and a sequence  $\{\alpha_n\}$  of distinct elements of K(M) such that  $\alpha_n(K) \cap C$  is non-empty for all n. We may pass to a subsequence, still called  $\{\alpha_n\}$ , so that there exists j such that  $\{q_j(\alpha_n)\}$  is a sequence of distinct elements. Since X(M) is locally compact, for each  $x \in K$ , there exists an open neighborhood  $U_x$  of x and a  $C_j$ -registering subgroup  $H_x$  such that the closure  $\overline{U}_x$  is a compact subset of  $Z_{H_x}$ . Since K is compact, there exists a finite collection of points  $\{x_1, \ldots, x_r\}$  such that  $K \subset U_{x_1} \cup \cdots \cup U_{x_r}$ . Therefore, again passing to subsequence if necessary, there must exists  $x_i$  such that  $\alpha_n(U_{x_i}) \cap C$  is non-empty for all n. Let  $U' = U_{x_i}$  and  $H' = H_{x_i}$ . Lemma 6.1 implies that  $\{s_{H'}(q_j(\alpha_n))\}$  is a sequence of distinct elements of  $\mathrm{Out}(H')$  and that  $s_{H'}(q_j(\alpha_n))(r_{H'}(\overline{U}')) = r_{H'}(\alpha_n(\overline{U}'))$ . Theorem 9.2 then implies that  $\{s_{H'}(q_j(\alpha_n))(r_{H'}(\overline{U}'))\} = \{r_{H'}(\alpha_n(\overline{U}'))\}$  exits every compact subset of X(M) which is a contradiction. We have thus established (2).

**9.3. Interval bundle components of**  $\Sigma(M)$ **.** Let  $\Sigma_i$  be an interval bundle component of  $\Sigma(M)$  with base surface  $F_i$  and let  $X(\Sigma_i)$  be its associated character variety. There exists a natural restriction map  $r_i: X(M) \to X(\Sigma_i)$  taking  $\rho$  to  $\rho|_{\pi_1(\Sigma_i)}$ . Recall that  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  injects into  $\operatorname{Out}(\pi_1(\Sigma_i))$  (by Lemma 5.2), so acts effectively on  $X(\Sigma_i)$ . Moreover, if  $\alpha \in J(M)$ , then  $r_i(\rho \circ \alpha) = r_i(\rho) \circ p_i(\alpha)$  where  $p_i$  is the projection of J(M) onto  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$ . If  $\Sigma_i$  is not tiny, we define

$$V(\Sigma_i) = r_i^{-1}(PS(\Sigma_i)).$$

If  $\{\Sigma_1, \ldots, \Sigma_n\}$  denotes the collection of all interval bundle components of  $\Sigma(M)$  which are not tiny, then we let

$$V(M) = \bigcap_{i=1}^{n} V(\Sigma_i).$$

If every interval bundle component of  $\Sigma(M)$  is tiny, then we let V(M) = X(M).

We use Lemma 8.2, Theorem 9.2, and Johannson's Classification Theorem to prove:

**Lemma 9.4.** Let M be a compact hyperbolizable 3-manifold with nonempty incompressible boundary and no toroidal boundary components which is not an interval bundle. Then

- (1) V(M) is an  $Out(\pi_1(M))$ -invariant open neighborhood of  $AH_n(M)$  in X(M), and
- (2) if K is a compact subset of V(M) and  $\{\alpha_n\}$  is a sequence in J(M) such that  $\{p_{\Sigma}(\alpha_n)\}$  is a sequence of distinct elements of  $G(\Sigma, Fr(\Sigma))$ , then  $\{\alpha_n(K)\}$  exits every compact subset of X(M).

*Proof.* Lemma 8.2 implies that  $AH_n(M) \subset V(\Sigma_i)$ , for each i, and each  $V(\Sigma_i)$  is open since  $r_i$  is continuous. Therefore, V(M) is an open neighborhood of  $AH_n(M)$ .

Johannson's Classification Theorem implies that if  $\varphi \in \operatorname{Out}(\pi_1(M))$ , then there exists a homotopy equivalence  $h \colon M \to M$  such that  $h(\Sigma(M)) \subset \Sigma(M), h|_{\operatorname{Fr}(\Sigma)}$  is a self-homeomorphism of  $\operatorname{Fr}(\Sigma)$  and h induces  $\varphi$ . Therefore, if  $\Sigma_i$  is an interval bundle component of  $\Sigma(M)$ , then  $\varphi(\pi_1(\Sigma_i))$  is conjugate to  $\pi_1(\Sigma_j)$  where  $\Sigma_j$  is also an interval bundle component of  $\Sigma(M)$ . Moreover, if  $\Sigma_i$  is not tiny, then  $\pi_1(\Sigma_j)$  is also not tiny (since  $h|_{\Sigma_i} \colon \Sigma_i \to \Sigma_j$  is a homotopy equivalence which is a homeomorphism on the frontier). It follows that V(M) is invariant under  $\operatorname{Out}(\pi_1(M))$ , completing the proof of claim (1).

If (2) fails to hold, then there is a compact subset K of Z(M), a compact subset C of X(M) and a sequence  $\{\alpha_n\}$  of elements of J(M) such that  $\{p_{\Sigma}(\alpha_n)\}$  is a sequence of distinct elements of  $G(\Sigma, \operatorname{Fr}(\Sigma))$  and  $\alpha_n(K) \cap C$  is non-empty for all n. If a component  $\Sigma_i$  of  $\Sigma(M)$  is a tiny interval bundle or a solid torus, then  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$  is finite, by Lemma 5.2. So, we may pass to a subsequence, so that there exists an interval bundle  $\Sigma_i$  which is not tiny such that  $\{p_i(\alpha_n)\}$  is a sequence of distinct elements of  $G(\Sigma_i, \operatorname{Fr}(\Sigma_i))$ . Theorem 9.2 then implies that  $\{p_i(\alpha_n)(r_i(K))\}$  leaves every compact subset of  $X(\Sigma_i)$ . Therefore, since  $r_i(\alpha_n(K)) = p_i(\alpha_n)(r_i(K))$  for all n,  $\{\alpha_n(K)\}$  leaves every compact subset of X(M). This contradiction establishes claim (2).

**9.4.** Assembly. Let  $W(M) = V(M) \cap Z(M)$ . Since V(M) and Z(M) are open  $\operatorname{Out}(\pi_1(M))$ -invariant neighborhoods of  $AH_n(M)$ , so is W(M). It remains to prove that  $\operatorname{Out}(\pi_1(M))$  acts properly discontinuously on W(M). Since J(M) is a finite index subgroup of  $\operatorname{Out}(\pi_1(M))$ , it suffices to prove that J(M) acts properly discontinuously on W(M). We will actually establish the following stronger fact, which will complete the proof of Theorem 9.1.

**Lemma 9.5.** If K is a compact subset of W(M) and  $\{\alpha_n\}$  is a sequence of distinct elements of J(M), then  $\{\alpha_n(K)\}$  leaves every compact subset of X(M).

*Proof.* If our claim fails, then there exists a compact subset K of W(M), a compact subset C of X(M) and a sequence  $\{\alpha_n\}$  of distinct elements of J(M) such that

 $\alpha_n(K) \cap C$  is non-empty. We may pass to an infinite subsequence, still called  $\{\alpha_n\}$ , such that either  $\{p_{\Sigma}(\alpha_n)\}$  is a sequence of distinct elements or  $\{p_{\Sigma}(\alpha_n)\}$  is constant.

If  $\{p_{\Sigma}(\alpha_n)\}$  is a sequence of distinct elements, Lemma 9.4 immediately implies that  $\{\alpha_n(K)\}$  leaves every compact subset of X(M) and we obtain the desired contradiction.

If  $\{\rho_{\Sigma}(\alpha_n)\}$  is constant, then, by Theorem 5.2, there exists a sequence  $\{\beta_n\}$  of distinct elements of K(M) such that  $\alpha_n = \alpha_1 \circ \beta_n$  for all n. Lemma 9.3 implies that  $\{\beta_n(K)\}$  exits every compact subset of X(M). Since  $\alpha_1$  induces a homeomorphism of X(M), it follows that  $\{\alpha_n(K) = \alpha_1(\beta_n(K))\}$  also leaves every compact subset of X(M). This contradiction completes the proof.

#### References

- [1] I. Agol, Tameness of hyperbolic 3-manifolds. Preprint, arXiv:math/0405568 [math.GT].
- [2] J. W. Anderson and R. D. Canary, Algebraic limits of Kleinian groups which rearrange the pages of a book. *Invent. Math.* **126** (1996), 205–214. Zbl 0874.57012 MR 1411128
- [3] J. W. Anderson, R. D. Canary and D. McCullough, On the topology of deformation spaces of Kleinian groups. *Ann. of Math.* **152** (2000), 693–741. Zbl 0976.57016 MR 1815699
- [4] R. Benedetti and C. Petronio, Lectures on hyperbolic geometry. Springer-Verlag Universitext, New York 1992. Zbl 0768.51018 MR 1219310
- [5] L. Bers, On moduli of Kleinian groups. Russian Math. Surveys 29 (1974), 88–102.Zbl 0304.30013 MR 0422691
- [6] F. Bonahon, Cobordism of automorphisms of surfaces. *Ann. Sci. École Norm. Sup.* **16** (1983), 237–270. Zbl 0535.57016 MR 0732345
- [7] F. Bonahon and J. P. Otal, Variétés hyperboliques à géodésiques arbitrairement courtes. Bull. London Math. Soc. 20 (1988), 255–261. Zbl 0648.53027 MR 0931188
- [8] B. Bowditch, Markoff triples and quasi-Fuchsian groups. Proc. London Math. Soc. 77 (1998), 697–736. Zbl 0928.11030 MR 1643429
- [9] J. Brock and K. Bromberg, On the density of geometrically finite Kleinian groups. *Acta Math.* 192 (2004), 33–93. Zbl 1055.57020 MR 2079598
- [10] J. Brock, R. Canary and Y. Minsky, The classification of Kleinian surface groups, II: The Ending Lamination Conjecture. Ann. of Math. (2) 176 (2012), no. 1, 1–149. Zbl 06074012 MR 2925381
- [11] K. Bromberg, Rigidity of geometrically finite hyperbolic cone-manifolds. *Geom. Dedicata* **105** (2004), 143–170. Zbl 1057.53029 MR 2057249
- [12] K. Bromberg, The space of Kleinian punctured torus groups is not locally connected. *Duke Math. J.* **156** (2011), no. 3, 387–427. Zbl 1213.30078 MR 2772066
- [13] K. Bromberg and J. Holt, Self-bumping of deformation spaces of hyperbolic 3-manifolds. *J. Differential Geom.* **57** (2001), 47–65. Zbl 1030.57028 MR 1871491
- [14] D. Calegari and D. Gabai, Shrinkwrapping and the taming of hyperbolic 3-manifolds. *J. Amer. Math. Soc.* **19** (2006), 385–446. Zbl 1090.57010 MR 2188131

- [15] R. D. Canary, A covering theorem for hyperbolic 3-manifolds and its applications. *Topology* **35** (1996), 751–778. Zbl 0863.57010 MR 1396777
- [16] R. D. Canary and S. Hersonsky, Ubiquity of geometric finiteness in boundaries of deformation spaces of hyperbolic 3-manifolds. Amer. J. Math. 126 (2004), 1193–1220. Zbl 1062.57021 MR 2102392
- [17] R. D. Canary and D. McCullough, Homotopy equivalences of 3-manifolds and deformation theory of Kleinian groups. *Mem. Amer. Math. Soc.* 172 (2004), no. 812. Zbl 1062.57027 MR 2096234
- [18] R. D. Canary and P. A. Storm, The curious moduli spaces of unmarked Kleinian surface groups. *Amer. J. Math.* **134** (2012), no. 1, 71–85. Zbl 06019021 MR 2876139
- [19] S. Cantat, Bers and Hénon, Painlevé and Schrödinger. Duke Math. J. 149 (2009), 411–460.
  Zbl 1181.37068 MR 2553877
- [20] T. D. Comar, Hyperbolic Dehn surgery and convergence of Kleinian groups. Ph.D. thesis, University of Michigan, 1996.
- [21] W. Goldman, The complex-symplectic geometry of  $SL_2(\mathbb{C})$ -characters over surfaces. In *Algebraic groups and arithmetic*, Tata Inst. Fund. Res., Mumbai 2004, 375–407. Zbl 1089.53060 MR 2094117
- [22] W. Goldman, Mapping class group dynamics on surface group representations. In *Problems on mapping class groups and related topics*, Proc. Sympos. Pure Math. 74, Amer. Math. Soc., Providence, RI, 2006, 189–214. Zbl 05124684 MR 2264541
- [23] J. Hempel, 3-manifolds. Ann. of Math. Stud. 86, Princeton University Press, Princeton, NJ, 1976. Zbl 0345.57001 MR 0415619
- [24] M. Heusener and J. Porti, The variety of characters in  $PSL_2(\mathbb{C})$ . Bol. Soc. Mat. Mex. 10 (2004), 221–238. Zbl 1100.57014 MR 2199350
- [25] C. Hodgson and S. P. Kerckhoff, Universal bounds for hyperbolic Dehn surgery. Ann. of Math. 162 (2005), 367–421. Zbl 1087.57011 MR 2178964
- [26] S. Hong and D. McCullough, Ubiquity of geometric finiteness in mapping class groups of Haken 3-manifolds. *Pacific J. Math.* **188** (1999), 275–301. Zbl 0926.57015 MR 1684204
- [27] W. Jaco and P. Shalen, Seifert fibered spaces in 3-manifolds. Mem. Amer. Math. Soc. 21 (1979), no. 220. Zbl 0415.57005 MR 0539411
- [28] W. Jaco and P. Shalen, Peripheral structure of 3-manifolds. *Invent. Math.* 38 (1976), 55–87.
  Zbl 0395.57004 MR 0428332
- [29] K. Johannson, *Homotopy equivalences of 3-manifolds with boundary*. Lecture Notes in Math. 761, Springer-Verlag, Berlin 1979. Zbl 0412.57007 MR 0551744
- [30] M. Kapovich, *Hyperbolic manifolds and discrete groups*. Progr. Math. 183, Birkhäuser, Boston 2001. Zbl 0958.57001 MR 1792613
- [31] S.P. Kerckhoff and W.P. Thurston, Noncontinuity of the action of the modular group at Bers' boundary of Teichmüller space. *Invent. Math.* 100(1990), 25–47. Zbl 0698.32014 MR 1037141
- [32] M. Korkmaz, Mapping class groups of non-orientable surfaces. *Geom. Dedicata* **89** (2002), 109–133. Zbl 1016.57013 MR 1890954
- [33] R. Kulkarni and P. Shalen, On Ahlfors' finiteness theorem. Adv. in Math. 76 (1989), 155– 169. Zbl 0684.57019 MR 1013665

- [34] M. Lee, Dynamics on the  $PSL_2(\mathbb{C})$ -character variety of a twisted *I*-bundle. Preprint arXiv:1103.3479 [math.GT].
- [35] A. Magid, Deformation spaces of Kleinian surface groups are not locally connected. *Geom. Topol.* **16** (2012), no. 3, 1247–1320. Zbl 06068625
- [36] B. Maskit, On free Kleinian groups. Duke Math. J. 48 (1981), 755–765. Zbl 0475.30039 MR 0782575
- [37] B. Maskit, Kleinian groups. Grundlehren Math. Wiss. 287, Springer-Verlag, Berlin 1988. Zbl 0627.30039 MR 0959135
- [38] D. McCullough, Compact submanifolds of 3-manifolds with boundary. *Quart. J. Math. Oxford* **37** (1986), 299–307. Zbl 0628.57008 MR 0854628
- [39] D. McCullough, Virtually geometrically finite mapping class groups of 3-manifolds. J. Differential Geom. 33 (1991), 1–65. Zbl 0721.57008 MR 1085134
- [40] D. McCullough and A. Miller, Homeomorphisms of 3-manifolds with compressible boundary. *Mem. Amer. Math. Soc.* **61** (1986), no. 344. Zbl 0602.57011 MR 0840832
- [41] C. McMullen, *Renormalization and 3-manifolds which fiber over the circle*. Ann. of Math. Stud. 142, Princeton University Press, Princeton, NJ, 1996. Zbl 0860.58002 MR 1401347
- [42] C. T. McMullen, Complex earthquakes and Teichmüller theory. J. Amer. Math. Soc. 11 (1998), 283–320. Zbl 0890.30031 MR 1478844
- [43] Y. Minsky, On dynamics of  $Out(F_n)$  on  $PSL_2(C)$ -characters. *Israel J. Math.*, to appear; Doi 10.1007/s11856-012-0086-0.
- [44] J. Morgan, On Thurston's uniformization theorem for three-dimensional manifolds. In The Smith conjecture, Pure Appl. Math. 112, Academic Press, Orlando, FL, 1984, 37–125. Zbl 0599.57002 MR 0758464
- [45] H. Namazi and J. Souto, Non-realizability and ending lamination Proof of the Density Conjecture. *Acta. Math.*, to appear; preprint, <a href="http://www.ma.utexas.edu/users/hossein/contents/ending6.6.pdf">http://www.ma.utexas.edu/users/hossein/contents/ending6.6.pdf</a>.
- [46] K. Ohshika, Geometrically finite Kleinian groups and parabolic elements. *Proc. Edin. Math. Soc.* 41 (1998), 141–159. Zbl 0903.30032 MR 1604337
- [47] K. Ohshika, Realising end invariants by limits of minimally parabolic, geometrically finite groups. *Geom. Topol.* **15** (2011), no. 2, 827–890.
- [48] P. Scott, Compact submanifolds of 3-manifolds. J. London Math. Soc. 7 (1974), 246–250.
  Zbl 0266.57001 MR 0326737
- [49] J. Souto and P. Storm, Dynamics of the mapping class group action on the variety of  $PSL_2(\mathbb{C})$ -characters. *Geom. Topol.* 10 (2006), 715–736. Zbl 1129.57022 MR 2240903
- [50] D. Sullivan, Quasiconformal homeomorphisms and dynamics. II. Structural stability implies hyperbolicity for Kleinian groups. *Acta Math.* 155 (1985), 243–260. Zbl 0606.30044 MR 0806415
- [51] W. P. Thurston, Hyperbolic structures on 3-manifolds, II: Surface groups and 3-manifolds which fiber over the circle. Preprint.
- [52] W. P. Thurston, Hyperbolic structures on 3-manifolds, III: Deformations of 3-manifolds with incompressible boundary. Preprint.
- [53] W. P. Thurston, *The geometry and topology of three-manifolds*. Princeton University course notes, available at http://www.msri.org/publications/books/gt3m/.

- [54] S. P. Tan, Y. L. Wong, and Y. Zhang, Generalized Markoff maps and McShane's identity. Adv. Math. 217 (2008), 761–813. Zbl 1158.57023 MR 2370281
- [55] F. Waldhausen, On irreducible 3-manifolds which are sufficiently large. *Ann. of Math.* **87** (1968), 56–88. Zbl 0157.30603 MR 0224099

Received February 2, 2010

Richard D. Canary, Department of Mathematics, University of Michigan, Ann Arbor, MI 48109 U.S.A.

E-mail: canary@umich.edu

Peter A. Storm, 303 W 21st St, Apt 9B, New York, NY 10011, U.S.A.

E-mail: peterastorm@gmail.com