

**Zeitschrift:** Commentarii Mathematici Helvetici  
**Herausgeber:** Schweizerische Mathematische Gesellschaft  
**Band:** 88 (2013)

**Artikel:** Entropy on Riemann surfaces and the Jacobians of finite covers  
**Autor:** McMullen, Curtis T.  
**DOI:** <https://doi.org/10.5169/seals-515661>

#### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

#### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 23.01.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

## Entropy on Riemann surfaces and the Jacobians of finite covers

Curtis T. McMullen\*

**Abstract.** This paper characterizes those pseudo-Anosov mappings whose entropy can be detected homologically by taking a limit over finite covers. The proof is via complex-analytic methods. The same methods show the natural map  $\mathcal{M}_g \rightarrow \prod \mathcal{A}_h$ , which sends a Riemann surface to the Jacobians of all of its finite covers, is a contraction in most directions.

**Mathematics Subject Classification (2010).** 32G15, 37E30.

**Keywords.** Entropy, pseudo–Anosov maps, Kobayashi metric, Siegel space, Teichmüller space, homology.

### 1. Introduction

Let  $f: S \rightarrow S$  be a pseudo-Anosov mapping on a surface of genus  $g$  with  $n$  punctures. It is well-known that the topological entropy  $h(f)$  is bounded below in terms of the spectral radius of  $f^*: H^1(S, \mathbb{C}) \rightarrow H^1(S, \mathbb{C})$ ; we have

$$\log \rho(f^*) \leq h(f).$$

If we lift  $f$  to a map  $\tilde{f}: \tilde{S} \rightarrow \tilde{S}$  on a finite cover of  $S$ , then its entropy stays the same but the spectral radius of the action on homology can increase. We say the entropy of  $f$  can be *detected homologically* if

$$h(f) = \sup \log \rho(\tilde{f}^*: H^1(\tilde{S}) \rightarrow H^1(\tilde{S})),$$

where the supremum is taken over all finite covers to which  $f$  lifts.

In this paper we will show:

**Theorem 1.1.** *The entropy of a pseudo-Anosov mapping  $f$  can be detected homologically if and only if the invariant foliations of  $f$  have no odd-order singularities in the interior of  $S$ .*

The proof is via complex analysis. Hodge theory provides a natural embedding  $\mathcal{M}_g \rightarrow \mathcal{A}_g$  from the moduli space of Riemann surfaces into the moduli space of

---

\*Research supported in part by the NSF.

Abelian varieties, sending  $X$  to its Jacobian. Any characteristic covering map from a surface of genus  $h$  to a surface of genus  $g$ , branched over  $n$  points, provides a similar map

$$\mathcal{M}_{g,n} \rightarrow \mathcal{M}_h \rightarrow \mathcal{A}_h. \quad (1.1)$$

It is known that the hyperbolic metric on a Riemann surface  $X$  can be reconstructed using the metrics induced from the Jacobians of its finite covers ([Kaz]; see the Appendix). Similarly, it is natural to ask if the Teichmüller metric on  $\mathcal{M}_{g,n}$  can be recovered from the Kobayashi metric on  $\mathcal{A}_h$ , by taking the limit over all characteristic covers  $\mathcal{C}_{g,n}$ . We will show such a construction is impossible.

**Theorem 1.2.** *The natural map  $\mathcal{M}_{g,n} \rightarrow \prod_{\mathcal{C}_{g,n}} \mathcal{A}_h$  is not an isometry for the Kobayashi metric, unless  $\dim \mathcal{M}_{g,n} = 1$ .*

It is an open problem to determine if the Kobayashi and Carathéodory metrics on moduli space coincide when  $\dim \mathcal{M}_{g,n} > 1$  (see e.g. [FM], Problem 5.1). An equivalent problem is to determine if Teichmüller space embeds holomorphically and isometrically into a (possibly infinite) product of bounded symmetric domains. Theorem 1.2 provides some support for a negative answer to this question.

Here is a more precise version of Theorem 1.2, stated in terms of the lifted map

$$\mathcal{T}_{g,n} \rightarrow \mathcal{T}_h \xrightarrow{J} \mathcal{S}_h$$

from Teichmüller space to Siegel space determined by a finite cover.

**Theorem 1.3.** *Suppose the Teichmüller mapping between a pair of distinct points  $X, Y \in \mathcal{T}_{g,n}$  comes from a quadratic differential with an odd order zero. Then*

$$\sup d(J(\tilde{X}), J(\tilde{Y})) < d(X, Y),$$

where the supremum is taken over all compatible finite covers of  $X$  and  $Y$ .

Conversely, if the Teichmüller map from  $X$  to  $Y$  has only even order singularities, then there is a double cover such that  $d(J(\tilde{X}), J(\tilde{Y})) = d(X, Y)$  (cf. [Kra]). In particular, the complex geodesics generated by squares of holomorphic 1-forms map isometrically into  $\mathcal{A}_g$ . The only directions contracted by the map  $\mathcal{M}_g \rightarrow \prod \mathcal{A}_h$  are those identified by Theorem 1.3.

Theorem 1.1 follows from Theorem 1.3 by taking  $X$  and  $Y$  to be points on the Teichmüller geodesic stabilized by the mapping-class  $f$ . It would be interesting to find a direct topological proof of Theorem 1.1.

As a sample application, let  $\beta \in B_n$  be a pseudo-Anosov braid whose monodromy map  $f : S \rightarrow S$  (on the  $n$ -times punctured plane) has an odd order singularity. Then Theorem 1.1 implies the image of  $\beta$  under the Burau representation satisfies

$$\log \sup_{|q|=1} \rho(B(q)) < h(f).$$

Indeed,  $\rho(B(q))$  at any  $d$ -th root of unit is bounded by  $\rho(\tilde{f}^*)$  on a  $\mathbb{Z}/d$  cover  $S$  [Mc2]. This improves a result in [BB]. Similar statements hold for other homological representations of the mapping-class group.

**Notes and references.** For  $C^\infty$  diffeomorphisms of a compact smooth manifold, one has  $h(f) \geq \log \sup_i \rho(f^*|H^i(X))$  [Ym], and equality holds for holomorphic maps on Kähler manifolds [Gr]. The lower bound  $h(f) \geq \log \rho(f^*|H^1(X))$  also holds for homeomorphisms [Mn]. For more on pseudo-Anosov mappings, see e.g. [FLP], [Bers] and [Th].

A proof that the inclusion of  $\mathcal{T}_{g,n}$  into universal Teichmüller space is a contraction, based on related ideas, appears in [Mc1].

## 2. Odd order zeros

We begin with an analytic result, which describes how well a monomial  $z^k$  of odd order can be approximated by the square of an analytic function.

**Theorem 2.1.** *Let  $k \geq 1$  be odd, and let  $f(z)$  be a holomorphic function on the unit disk  $\Delta$  such that  $\int |\mathcal{f}(z)|^2 = 1$ . Then*

$$\left| \int_{\Delta} f(z)^2 \left( \frac{\bar{z}}{|z|} \right)^k \right| \leq C_k = \frac{\sqrt{k+1} \sqrt{k+3}}{k+2} < 1.$$

Here the integral is taken with respect to Lebesgue measure on the unit disk.

*Proof.* Consider the orthonormal basis  $e_n(z) = a_n z^n$ ,  $n \geq 0$ ,  $a_n = \sqrt{n+1}/\sqrt{\pi}$ , for the Bergman space  $L^2_\alpha(\Delta)$  of analytic functions on the disk with  $\|f\|_2^2 = \int |\mathcal{f}(z)|^2 < \infty$ . With respect to this basis, the nonzero entries in the matrix of the symmetric bilinear form  $Z(f, g) = \int f(z)g(z)\bar{z}^k/|z|^k$  are given by

$$Z(e_n, e_{k-n}) = a_n a_{k-n} \int_{\Delta} |z|^k = \frac{2\sqrt{n+1} \sqrt{k-n+1}}{k+2}.$$

In particular,  $Z(e_i, e_i) = 0$  for all  $i$  (since  $k$  is odd), and  $Z(e_i, e_j) = 0$  for all  $i, j > k$ .

Note that the ratio above is less than one, by the inequality between the arithmetic and geometric means, and it is maximized when  $n < k/2 < n+1$ . Thus the maximum of  $|Z(f, f)|/\|f\|^2$  over  $L^2_\alpha(\Delta)$  is achieved when  $f = e_n + e_{n+1}$ ,  $n = (k-1)/2$ , at which point it is given by  $C_k$ .  $\square$

### 3. Siegel space

In this section we describe the Siegel space of Hodge structures on a surface  $S$ , and its Kobayashi metric.

**Hodge structures.** Let  $S$  be a closed, smooth, oriented surface of genus  $g$ . Then  $H^1(S) = H^1(S, \mathbb{C})$  carries a natural involution  $C(\alpha) = \bar{\alpha}$  fixing  $H^1(S, \mathbb{R})$ , and a natural Hermitian form

$$\langle \alpha, \beta \rangle = \frac{\sqrt{-1}}{2} \int_S \alpha \wedge \bar{\beta}$$

of signature  $(g, g)$ . A *Hodge structure* on  $H^1(S)$  is given by an orthogonal splitting

$$H^1(S) = V^{1,0} \oplus V^{0,1}$$

such that  $V^{1,0}$  is positive-definite and  $V^{0,1} = C(V^{1,0})$ . We have a natural norm on  $V^{1,0}$  given by  $\|\alpha\|^2 = \langle \alpha, \alpha \rangle$ .

The set of all possible Hodge structures forms the *Siegel space*  $\mathfrak{H}(S)$ . To describe this complex symmetric space in more detail, fix a splitting  $H^1(S) = W^{1,0} \oplus W^{0,1}$ . Then for any other Hodge structure  $V^{1,0} \oplus V^{0,1}$ , there is a unique operator

$$Z: W^{1,0} \rightarrow W^{0,1}$$

such that  $V^{1,0} = (I + Z)(W^{1,0})$ . This means  $V^{1,0}$  coincides with the graph of  $Z$  in  $W^{1,0} \oplus W^{0,1}$ .

The operator  $Z$  is determined uniquely by the associated bilinear form

$$Z(\alpha, \beta) = \langle \alpha, C Z(\beta) \rangle$$

on  $W^{1,0}$ , and the condition that  $V^{1,0} \oplus V^{0,1}$  is a Hodge structure translates into the conditions

$$Z(\alpha, \beta) = Z(\beta, \alpha) \quad \text{and} \quad |Z(\alpha, \alpha)| < 1 \text{ if } \|\alpha\| = 1. \quad (3.1)$$

Since the second inequality above is an open condition, the tangent space at the base point  $p \sim W^{1,0} \oplus W^{0,1}$  is given by

$$T_p \mathfrak{H}(S) = \{\text{symmetric bilinear maps } Z: W^{1,0} \times W^{1,0} \rightarrow \mathbb{C}\}.$$

**Comparison maps.** Any Hodge structure on  $H^1(S)$  determines an isomorphism

$$V^{1,0} \cong H^1(S, \mathbb{R}) \quad (3.2)$$

sending  $\alpha$  to  $\Re(\alpha) = (\alpha + C(\alpha))/2$ . Thus  $H^1(S, \mathbb{R})$  inherits a norm and a complex structure from  $V^{1,0}$ .

Put differently, (3.2) gives a *marking* of  $V^{1,0}$  by  $H^1(S, \mathbb{R})$ . By composing one marking with the inverse of another, we obtain the real-linear *comparison map*

$$T = (I + Z)(I + CZ)^{-1} : W^{1,0} \rightarrow V^{1,0} \quad (3.3)$$

between any pair of Hodge structures. It is characterized by  $\Re(\alpha) = \Re(T(\alpha))$ .

**Symmetric matrices.** The classical Siegel domain is given by

$$\mathfrak{H}_g = \{Z \in M_g(\mathbb{C}) : Z_{ij} = Z_{ji} \text{ and } I - Z\bar{Z} \gg 0\}.$$

(cf. [Sat], Chapter II.7). It is a convex, bounded symmetric domain in  $\mathbb{C}^N$ ,  $N = g(g+1)/2$ . The choice of an orthonormal basis for  $W^{1,0}$  gives an isomorphism  $Z \mapsto Z(\omega_i, \omega_j)$  between  $\mathfrak{H}(S)$  and  $\mathfrak{H}_g$ , sending the basepoint  $p$  to zero.

**The Kobayashi metric.** Let  $\Delta \subset \mathbb{C}$  denote the unit disk, equipped with the metric  $|dz|/(1-|z|^2)$  of constant curvature  $-4$ . The *Kobayashi metric* on  $\mathfrak{H}(S)$  is the largest metric such that every holomorphic map  $f : \Delta \rightarrow \mathfrak{H}(S)$  satisfies  $\|Df(0)\| \leq 1$ . It determines both a norm on the tangent bundle and a distance function on pairs of points [Ko].

**Proposition 3.1.** *The Kobayashi norm on  $T_p \mathfrak{H}(S)$  is given by*

$$\|Z\|_K = \sup\{Z(\alpha, \alpha) : \|\alpha\| = 1\},$$

and the Kobayashi distance is given in terms of the comparison map (3.3) by

$$d(V^{1,0}, W^{1,0}) = \log \|T\|.$$

*Proof.* Choosing a suitable orthonormal basis for  $W^{1,0}$ , we can assume that

$$Z(\omega_i, \omega_j) = \lambda_i \delta_{ij}$$

with  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_g \geq 0$ . Since  $\mathfrak{H}_g$  is a convex symmetric domain, the Kobayashi norm at the origin and the Kobayashi distance satisfy

$$\|Z\|_K = r \quad \text{and} \quad d(0, Z) = \frac{1}{2} \log \frac{1+r}{1-r},$$

where  $r = \inf\{s > 0 : Z \in s\mathfrak{H}_g\}$  (see [Ku]). Clearly  $r = \lambda_1 = \sup |Z(\alpha, \alpha)|/\|\alpha\|^2$ , and by (3.3), we have

$$\|T\|^2 = \|T(\sqrt{-1}\omega_1)\|^2 = \left\| \frac{\omega_1}{1-\lambda_1} + \frac{\lambda_1 \bar{\omega}_1}{1-\lambda_1} \right\|^2 = \frac{1+\lambda_1}{1-\lambda_1},$$

which gives the expressions above. □

#### 4. Teichmüller space

This section gives a functorial description of the derivative of the map from Teichmüller space to Siegel space.

**Markings.** Let  $\bar{S}$  be a compact oriented surface of genus  $g$ , and let  $S \subset \bar{S}$  be a subsurface obtained by removing  $n$  points.

Let  $\text{Teich}(S) \cong \mathcal{T}_{g,n}$  denote the Teichmüller space of Riemann surfaces marked by  $S$ . A point in  $\text{Teich}(S)$  is specified by a homeomorphism  $f: S \rightarrow X$  to a Riemann surface of finite type. This means there is a compact Riemann surface  $\bar{X} \supset X$  and an extension of  $f$  to a homeomorphism  $\bar{f}: \bar{S} \rightarrow \bar{X}$ .

**Metrics.** Let  $Q(X)$  denote the space of holomorphic quadratic differentials on  $X$  such that

$$\|q\|_X = \int_X |q| < \infty.$$

There is a natural pairing  $(q, \mu) \mapsto \int_X q\mu$  between the space  $Q(X)$  and the space  $M(X)$  of  $L^\infty$ -measurable Beltrami differentials  $\mu$ . The tangent and cotangent spaces to Teichmüller space at  $X$  are isomorphic to  $M(X)/Q(X)^\perp$  and  $Q(X)$  respectively.

The Teichmüller and Kobayashi metrics on  $\text{Teich}(S)$  coincide [Roy1], [Hub], Chapter 6. They are given by the norm

$$\|\mu\|_T = \sup \left\{ \left| \int q\mu \right| : \|q\|_X = 1 \right\}$$

on the tangent space at  $X$ ; the corresponding distance function

$$d(X, Y) = \inf \frac{1}{2} \log K(\phi)$$

measures the minimal dilatation  $K(\phi)$  of a quasiconformal map  $\phi: X \rightarrow Y$  respecting their markings.

**Hodge structure.** The periods of holomorphic 1-forms on  $X$  serve as classical moduli for  $X$ . From a modern perspective, these periods give a map

$$J: \text{Teich}(S) \rightarrow \mathfrak{H}(\bar{S}) \cong \mathfrak{H}_g,$$

sending  $X$  to the Hodge structure

$$H^1(\bar{S}) \cong H^1(\bar{X}) \cong H^{1,0}(\bar{X}) \oplus H^{0,1}(\bar{X}).$$

Here the first isomorphism is provided by the marking  $\bar{f}: \bar{S} \rightarrow \bar{X}$ . We also have a natural isomorphism between  $H^{1,0}(\bar{X})$  and the space of holomorphic 1-forms  $\Omega(\bar{X})$ . The image  $J(X)$  encodes the complex analytic structure of the Jacobian variety  $\text{Jac}(\bar{X}) = \Omega(\bar{X})^*/H_1(\bar{X}, \mathbb{Z})$ . (It does not depend on the location of the punctures of  $X$ .)

**Proposition 4.1.** *The derivative of the period map sends  $\mu \in M(X)$  to the quadratic form  $Z = DJ(\mu)$  on  $\Omega(\bar{X})$  given by*

$$Z(\alpha, \beta) = \int_{\bar{X}} \alpha \beta \mu.$$

This is a basis-free reformulation of Ahlfors' variational formula [Ah], §5; see also [Ra], [Roy2] and Proposition 1 of [Kra]. Note that  $\alpha \beta \in Q(X)$ .

## 5. Contraction

This section brings finite covers into play, and establishes a uniform estimate for contraction of the mapping  $\mathcal{T}_{g,n} \rightarrow \mathcal{T}_h \rightarrow \mathfrak{H}_h$ .

**Jacobians of finite covers.** A finite connected covering space  $S_1 \rightarrow S_0$  determines a natural map

$$P: \text{Teich}(S_0) \rightarrow \text{Teich}(S_1)$$

sending each Riemann surface to the corresponding covering space  $X_1 \rightarrow X_0$ . By taking the Jacobian of  $X_1$ , we obtain a map  $J \circ P: \text{Teich}(S_0) \rightarrow \mathfrak{H}(\bar{S}_1)$ .

Let  $q_0 \in Q(X_0)$  be a holomorphic quadratic differential with a zero of odd order  $k$ , say at  $p \in X_0$ . Let  $\mu = \bar{q}_0/|q_0| \in M(X_0)$ ; then  $\|\mu\|_T = 1$ . Let  $\pi: X_1 \rightarrow X_0$  denote the natural covering map, and let  $q_1 = \pi^*(q_0)$ .

We will show that  $J(X_1)$  cannot change too rapidly under the unit deformation  $\mu$  of  $X_0$ . Indeed, if  $J(X_1)$  were to move at nearly unit speed, then  $\pi^*(\mu) = \bar{q}_1/|q_1|$  would pair efficiently with  $\alpha^2$  for some unit-norm  $\alpha \in \Omega(\bar{X}_1)$ , which is impossible because of the many odd-order zeros of  $q_1$ .

To make a quantitative estimate, choose a holomorphic chart  $\phi: (\Delta, 0) \rightarrow (X_0, p)$  such that  $\phi^*(\mu) = z^k/|z|^k d\bar{z}/dz$ . Let  $U = \phi(\Delta)$ , and let

$$m(U) = \inf\{\|q\|_U : q \in Q(X_0), \|q\|_X = 1\}.$$

(Here  $\|q\|_U = \int_U |q|$ .) Since  $Q(X_0)$  is finite-dimensional, we have  $m(U) > 0$ .

**Theorem 5.1.** *The image  $Z$  of the vector  $[\mu]$  under the derivative of  $J \circ P$  satisfies*

$$\|Z\|_K \leq \delta < 1 = \|\mu\|_T,$$

where  $\delta = \max(1/2, 1 - (1 - C_k)m(U)/2)$  does not depend on the finite cover  $S_1 \rightarrow S_0$ .

*Proof.* The derivative of  $P$  sends  $\mu$  to  $\pi^*(\mu)$ . By Proposition 3.1, to show  $\|Z\|_K \leq \delta$  it suffices to show that

$$|Z(\alpha, \alpha)| = \left| \int_{X_1} \alpha^2 \pi^* \mu \right| \leq \delta$$

for all  $\alpha \in \Omega(\bar{X}_1)$  with  $\|\alpha^2\|_{X_1} = 1$ . Setting  $q = \pi_*(\alpha^2)$ , we also have

$$|Z(\alpha, \alpha)| = \left| \int_{X_0} q \mu \right| \leq \|q\|_{X_0},$$

so the proof is complete if  $\|q\|_{X_0} \leq 1/2$ . Thus we may assume that

$$\|\alpha^2\|_V \geq \|q\|_U \geq m(U)\|q\|_{X_0} \geq m(U)/2,$$

where  $V = \pi^{-1}(U) = \bigcup_1^d V_i$  is a finite union of disjoint disks. Using the coordinate charts  $V_i \cong U \cong \Delta$  and Theorem 2.1, we find that on each of these disks we have

$$\left| \int_{V_i} \alpha^2 \pi^*(\mu) \right| = \left| \int_{\Delta} \alpha(z)^2 \left( \frac{z}{|z|} \right)^k \right| \leq C_k \|\alpha^2\|_{V_i}.$$

Summing these bounds and using the fact that  $\|\alpha^2\|_{(X_1-V)} + \|\alpha^2\|_V = 1$ , we obtain

$$\left| \int_{X_1} \alpha^2 \pi^*(\mu) \right| \leq \|\alpha^2\|_{(X_1-V)} + C_k \|\alpha^2\|_V \leq 1 - \frac{(1-C_k)m(U)}{2} \leq \delta. \quad \square$$

## 6. Conclusion

It is now straightforward to establish the results stated in the Introduction.

*Proof of Theorem 1.3.* Assume the Beltrami coefficient of the Teichmüller mapping between  $X, Y \in \mathcal{T}_{g,n}$  has the form  $\mu = k\bar{q}/q$ , where  $q \in Q(X)$  has an odd order zero. Then the same is true for the tangent vectors to the Teichmüller geodesic  $\gamma$  joining  $X$  to  $Y$ . Theorem 5.1 then implies that  $D(J \circ P)|_{\gamma}$  is contracting by a factor  $\delta < 1$  independent of  $P$ , and therefore

$$d(J \circ P(X), J \circ P(Y)) = d(J(\tilde{X}), J(\tilde{Y})) < \delta \cdot d(X, Y). \quad \square$$

*Proof of Theorem 1.2.* The contraction of  $\mathcal{M}_{g,n} \rightarrow \prod_{\mathcal{C}_{g,n}} \mathcal{A}_h$  in some directions is immediate from the uniformity of the bound in Theorem 1.3, using the fact that the Kobayashi metric on a product is the sup of the Kobayashi metrics on each term, and that there exist  $q \in Q(X)$  with simple zeros whenever  $X \in \mathcal{M}_{g,n}$  and  $\dim \mathcal{M}_{g,n} > 1$ .  $\square$

*Proof of Theorem 1.1.* Let  $f : S_0 \rightarrow S_0$  be a pseudo-Anosov mapping. If  $f$  has only even order singularities, then its expanding foliation is locally orientable, and hence there is a double cover  $\tilde{S} \rightarrow S$  such that  $\log \rho(\tilde{f}^*) = h(f)$ .

Now suppose  $f$  has an odd-order singularity. Let  $X_0 \in \text{Teich}(S_0)$  be a point on the Teichmüller geodesic stabilized by the action of  $f$  on  $\text{Teich}(S_0)$ . Then  $h(f) = d(f \cdot X_0, X_0) > 0$  (see e.g. [FLP] and [Bers]).

Let  $\tilde{f}: S_1 \rightarrow S_1$  be a lift of  $f$  to a finite covering of  $S_0$ , and let  $X_1 = P(X_0) \in \text{Teich}(S_1)$ . Using the marking of  $X_1$  and the isomorphism  $H^1(X_1, \mathbb{R}) \cong H^{1,0}(X_1)$ , we obtain a commutative diagram

$$\begin{array}{ccc} H^1(S_1, \mathbb{R}) & \xrightarrow{\tilde{f}^*} & H^1(S_1, \mathbb{R}) \\ \downarrow & & \downarrow \\ H^{1,0}(\bar{X}_1) & \xrightarrow{T} & H^{1,0}(\bar{X}_1) \end{array}$$

where  $T$  is the comparison map between  $J(X_1)$  and  $J(\tilde{f} \cdot X_1)$  (see equation (3.3)). Then Theorem 1.3 and Proposition 3.1 yield the bound

$$\log \rho(\tilde{f}^*) \leq \log \|T\| = d(J(X_1), \tilde{f} \cdot J(X_1)) \leq \delta d(X_0, f \cdot X_0) = \delta h(f),$$

where  $\delta < 1$  does not depend on the finite covering  $S_1 \rightarrow S_0$ . Consequently,  $\sup \log \rho(\tilde{f}^*) < h(f)$ .  $\square$

## Appendix. The hyperbolic metric via Jacobians of finite covers

Let  $X = \Delta/\Gamma$  be a compact Riemann surface, presented as a quotient of the unit disk by a Fuchsian group  $\Gamma$ . Let  $Y_n \rightarrow X$  be an ascending sequence of finite Galois covers which converge to the universal cover, in the sense that

$$Y_n = \Delta/\Gamma_n, \quad \Gamma \supset \Gamma_1 \supset \Gamma_2 \supset \Gamma_3 \dots, \quad \text{and} \quad \bigcap \Gamma_i = \{e\}. \quad (\text{A.1})$$

The Bergman metric on  $Y_n$  (defined below) is invariant under automorphisms, so it descends to a metric  $\beta_n$  on  $X$ . This appendix gives a short proof of:

**Theorem A.1** (Kazhdan). *The Bergman metrics inherited from the finite Galois covers  $Y_n \rightarrow X$  converge to a multiple of the hyperbolic metric; more precisely, we have*

$$\beta_n \rightarrow \frac{\lambda_X}{2\sqrt{\pi}}$$

uniformly on  $X$ .

The argument below is based on [Kaz], §3; for another, somewhat more technical approach, see [Rh].

**Metrics.** We begin with some definitions. Let  $\Omega(X)$  denote the Hilbert space of holomorphic 1-forms on a Riemann surface  $X$  such that

$$\|\omega\|_X^2 = \int_X |\omega|^2 < \infty.$$

The area form of the *Bergman metric* on  $X$  is given by

$$\beta_X^2 = \sum |\omega_i|^2, \quad (\text{A.2})$$

where  $(\omega_i)$  is any orthonormal basis of  $\Omega(X)$ . Equivalently, the Bergman length of a tangent vector  $v \in TX$  is given by

$$\langle \beta_X, v \rangle = \sup_{\omega \neq 0} \frac{|\omega(v)|}{\|\omega\|_X}. \quad (\text{A.3})$$

This formula shows that inclusions are contracting: if  $Y$  is a subdomain of  $X$ , then  $\beta_Y \geq \beta_X$ .

Now suppose  $X$  is a compact surface of genus  $g > 0$ . Then (A.2) shows its Bergman area is given by

$$\int_X \beta_X^2 = \dim \Omega(X) = g. \quad (\text{A.4})$$

In this case  $\beta_X$  is also the pullback, via the Abel–Jacobi map, of the natural Kähler metric on the Jacobian of  $X$ .

Finally suppose  $X = \Delta/\Gamma$ . Then the hyperbolic metric of constant curvature  $-1$ ,

$$\lambda_\Delta = \frac{2|dz|}{1 - |z|^2},$$

descends to give the *hyperbolic metric*  $\lambda_X$  on  $X$ . Using the fact that  $\|dz\|_\Delta = \pi$ , it is easy to check that  $4\pi\beta_\Delta^2 = \lambda_\Delta^2$ .

*Proof of Theorem A.1.* We will regard the Bergman metric  $\beta_n$  on  $Y_n$  as a  $\Gamma_n$ -invariant metric on  $\Delta$ . It suffices to show that  $\beta_n/\beta_\Delta \rightarrow 1$  uniformly on  $\Delta$ .

Let  $g$  and  $g_n$  denote the genus of  $X$  and  $Y_n$  respectively, and let  $d_n$  denote the degree of  $Y_n/X$ ; then  $g_n - 1 = d_n(g - 1)$ . By (A.1), the injectivity radius of  $Y_n$  tends to infinity. In particular, there is a sequence  $r_n \rightarrow 1$  such that  $\gamma(r_n\Delta)$  injects into  $Y_n$  for any  $\gamma \in \Gamma$ . Since inclusions are contracting, this shows

$$\beta_n \leq (1 + \epsilon_n)\beta_\Delta \quad (\text{A.5})$$

where  $\epsilon_n \rightarrow 0$ .

Next, note that both  $\beta_n$  and  $\beta_\Delta$  are  $\Gamma$ -invariant, so they determine metrics on  $X$ . By (A.4), we have

$$\int_X \beta_n^2 = \frac{1}{d_n} \int_{Y_n} \beta_n^2 = \frac{g_n}{d_n} \rightarrow (g - 1) = \int_X \beta_\Delta^2$$

(since  $\int_X \lambda_X^2 = 2\pi(2g - 2)$  by Gauss–Bonnet). Together with (A.5), this implies

$$\int_X |\beta_n - \beta_\Delta|^2 \rightarrow 0. \quad (\text{A.6})$$

To show  $\beta_n \rightarrow \beta_\Delta$  uniformly, consider any sequence  $p_n \in \Delta$  and let  $x \in [0, 1]$  be a limit point of  $(\beta_n/\beta_\Delta)(p_n)$ . It suffices to show  $x = 1$ .

Passing to a subsequence and using compactness of  $X$ , we can assume that  $p_n \rightarrow p \in \Delta$  and that  $\beta_n(p_n) \rightarrow x\beta_\Delta(p)$ . By changing coordinates on  $\Delta$ , we can also assume  $p = 0$ . By (A.6) we can find  $q_n \rightarrow 0$  such that  $\beta_n(q_n) \rightarrow \beta_\Delta(0)$ . Then by (A.3), there exist  $\Gamma_n$ -invariant holomorphic 1-forms  $\omega_n(z) dz$  on  $\Delta$  such that  $\int_{Y_n} |\omega_n|^2 = 1$  and

$$|\omega_n(q_n)| = \beta_n(q_n) \rightarrow \beta_\Delta(0) = \frac{|dz|}{\pi}.$$

Since  $\omega_n$  is holomorphic and  $\int_{r_n \Delta} |\omega_n|^2 < 1$ , the equation above easily implies that  $|\omega_n| \rightarrow |dz|/\pi$  uniformly on compact subsets of  $\Delta$ . But we also have

$$\beta_n(p_n) \geq |\omega_n(p_n)| \rightarrow \beta_\Delta(0),$$

and thus  $\beta_n(p_n) \rightarrow \beta_\Delta(0)$  and hence  $x = 1$ .  $\square$

## References

- [Ah] L. Ahlfors, The complex analytic structure of the space of closed Riemann surfaces. In *Analytic functions*, Princeton University Press, Princeton, NJ, 1960, 45–66. [Zbl 0100.28903](#) [MR 0124486](#)
- [BB] G. Band and P. Boyland, The Burau estimate for the entropy of a braid. *Algebr. Geom. Topol.* **7** (2007), 1345–1378. [Zbl 1128.37028](#) [MR 2350285](#)
- [Bers] L. Bers, An extremal problem for quasiconformal maps and a theorem by Thurston. *Acta Math.* **141** (1978), 73–98. [Zbl 0389.30018](#) [MR 0477161](#)
- [FLP] A. Fathi, F. Laudenbach, and V. Poénaru, *Travaux de Thurston sur les surfaces*. Astérisque, vol. 66–67, Soc. Math. France, Paris 1979. [Zbl 0406.00016](#) [MR 0568308](#)
- [FM] A. Fletcher and V. Markovic, Infinite dimensional Teichmüller spaces. In *Handbook of Teichmüller theory*, ed. by A. Papadopoulos, Volume II, Eur. Math. Soc., Zürich 2009, 65–92. [Zbl 1210.30015](#) [MR 2497790](#)
- [Gr] M. Gromov, On the entropy of holomorphic maps. *Enseign. Math.* **49** (2003), 217–235. [Zbl 1080.37051](#) [MR 2026895](#)
- [Hub] J. H. Hubbard, *Teichmüller theory and applications to geometry, topology, and dynamics. Vol. I*. Matrix Editions, Ithaca, NY, 2006. [Zbl 1102.30001](#) [MR 2245223](#)
- [Kaz] D. A. Kazhdan, On arithmetic varieties. In *Lie groups and their representations* (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971), Halsted, New York 1975, 151–217. [Zbl 0308.14007](#) [MR 0486316](#)
- [Ko] S. Kobayashi, *Hyperbolic manifolds and holomorphic mappings*. Marcel Dekker, Inc., New York 1970. [Zbl 0207.37902](#) [MR 0277770](#)
- [Kra] I. Kra, The Carathéodory metric on abelian Teichmüller disks. *J. Analyse Math.* **40** (1981), 129–143. [Zbl 0487.32017](#) [MR 0659787](#)

- [Ku] Y. Kubota, On the Kobayashi and Carathéodory distances of bounded symmetric domains. *Kodai Math. J.* **12** (1989), 41–48. [Zbl 0672.32010](#) [MR 0987140](#)
- [Mn] A. Manning, Topological entropy and the first homology group. In *Dynamical systems—Warwick 1974*, Lecture Notes in Math. 468, Springer, Berlin 1975, 185–190. [Zbl 0307.54042](#) [MR 0650661](#)
- [Mc1] C. McMullen, Amenability, Poincaré series and quasiconformal maps. *Invent. Math.* **97** (1989), 95–127. [Zbl 0672.30017](#) [MR 0999314](#)
- [Mc2] C. McMullen, Braid groups and Hodge theory. *Math. Ann.* **355** (2013), no. 3, 893–946. [Zbl 06149478](#) [MR 3020148](#)
- [Ra] H. E. Rauch, A transcendental view of the space of algebraic Riemann surfaces. *Bull. Amer. Math. Soc.* **71** (1965), 1–39. [Zbl 0154.33002](#) [MR 0213543](#)
- [Rh] J. A. Rhodes, Sequences of metrics on compact Riemann surfaces. *Duke Math. J.* **72** (1993), 725–738. [Zbl 0798.11018](#) [MR 1253622](#)
- [Roy1] H. L. Royden, Automorphisms and isometries of Teichmüller space. In *Advances in the theory of Riemann surfaces*, Ann. of Math. Studies 66, Princeton University Press, Princeton, NJ, 1971, 369–384. [Zbl 0222.32011](#) [MR 0288254](#)
- [Roy2] H. L. Royden, Invariant metrics on Teichmüller space. In *Contributions to analysis*, Academic Press, New York 1974, 393–399. [Zbl 0302.32022](#) [MR 0377116](#)
- [Sat] I. Satake, *Algebraic structures of symmetric domains*. Kanô Memorial Lectures 4, Princeton University Press, Princeton, NJ, 1980. [Zbl 0483.32017](#) [MR 0591460](#)
- [Th] W. P. Thurston, On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc.* **19** (1988), 417–432. [Zbl 0674.57008](#) [MR 0956596](#)
- [Ym] Y. Yomdin, Volume growth and entropy. *Israel J. Math.* **57** (1987), 285–300. [Zbl 0641.54036](#) [MR 0889979](#)

Received March 29, 2011

Curtis T. McMullen, Mathematics Department, Harvard University, 1 Oxford St, Cambridge, MA 02138-2901, U.S.A

E-mail: [ctm@math.harvard.edu](mailto:ctm@math.harvard.edu)