

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 89 (2014)

Artikel: Harmonic diffeomorphisms between domains in the Euclidean 2-sphere
Autor: Alarcón, Antonio / Souam, Rabah
DOI: <https://doi.org/10.5169/seals-515671>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Harmonic diffeomorphisms between domains in the Euclidean 2-sphere

Antonio Alarcón and Rabah Souam*

Abstract. We study the existence or non-existence of harmonic diffeomorphisms between certain domains in the Euclidean 2-sphere. In particular, we show the existence of harmonic diffeomorphisms from circular domains in the complex plane onto finitely punctured spheres, with at least two punctures. This result follows from a general existence theorem for maximal graphs in the Lorentzian product $\mathbb{M} \times \mathbb{R}_1$, where \mathbb{M} is an arbitrary n -dimensional compact Riemannian manifold, $n \geq 2$. In contrast, we show that there is no harmonic diffeomorphism from the unit complex disc onto the once-punctured sphere and no harmonic diffeomorphisms from finitely punctured spheres onto circular domains in the Euclidean 2-sphere.

Mathematics Subject Classification (2010). 53C43, 53C42, 53C50.

Keywords. Harmonic diffeomorphisms, maximal graphs.

1. Introduction

In 1952, Heinz [He] proved there is no harmonic diffeomorphism from the unit complex disk \mathbb{D} onto the complex plane \mathbb{C} , with the Euclidean metric. Later, Schoen and Yau [SY] asked whether Riemannian surfaces which are related by a harmonic diffeomorphism are quasiconformally related, and proposed to investigate whether there is a harmonic diffeomorphism from \mathbb{C} onto the hyperbolic plane \mathbb{H}^2 . Markovic [Ma] answered the first question in the negative, by showing an example consisting of a pair of Riemann surfaces of infinite topological type. He also gave conditions under which the question has a positive answer in the case of surfaces of finite topology. Finally, Collin and Rosenberg [CR] gave an example of a harmonic diffeomorphism from \mathbb{C} onto \mathbb{H}^2 , disproving the conjecture by Schoen and Yau [SY]. To do that, they constructed an entire minimal graph Σ over \mathbb{H}^2 in the Riemannian product $\mathbb{H}^2 \times \mathbb{R}$, with the conformal type of \mathbb{C} . Then the vertical projection $\Sigma \rightarrow \mathbb{H}^2$ is a surjective harmonic diffeomorphism. See [GR] for further generalizations.

*A. Alarcón is supported by Vicerrectorado de Política Científica e Investigación de la Universidad de Granada, and is partially supported by MCYT-FEDER grants MTM2007-61775 and MTM2011-22547, Junta de Andalucía Grant P09-FQM-5088, and the grant PYR-2012-3 CEI BioTIC GENIL (CEB09-0010) of the MICINN CEI Program.

Let \mathbb{S}^2 and $\bar{\mathbb{C}}$ denote the 2-dimensional Euclidean unit sphere and the Riemann sphere, respectively. A domain in $\bar{\mathbb{C}}$ is said to be a circular domain if every connected component of its boundary is a circle.

In this paper we study the existence or non-existence of harmonic diffeomorphisms between certain domains in \mathbb{S}^2 . Our main result asserts:

Theorem I. (i) For any $m \in \mathbb{N}$, $m \geq 2$, and any subset $\{p_1, \dots, p_m\} \subset \mathbb{S}^2$ there exist a circular domain $U \subset \bar{\mathbb{C}}$ and a harmonic diffeomorphism $\phi: U \rightarrow \mathbb{S}^2 - \{p_1, \dots, p_m\}$.

(ii) There exists no harmonic diffeomorphism $\varphi: \mathbb{D} \rightarrow \mathbb{S}^2 - \{p\}$.

(iii) For any $m \in \mathbb{N}$, any subset $\{z_1, \dots, z_m\} \subset \bar{\mathbb{C}}$ and any pairwise disjoint closed discs D_1, \dots, D_m in \mathbb{S}^2 there exists no harmonic diffeomorphism $\psi: \bar{\mathbb{C}} - \{z_1, \dots, z_m\} \rightarrow \mathbb{S}^2 - \bigcup_{j=1}^m D_j$.

Notice that Theorem I is related to Schoen and Yau's questions [SY], since circular domains are of hyperbolic conformal type whereas $\bar{\mathbb{C}}$ with a finite set removed is parabolic. Moreover, it is worth mentioning that Items (i) and (iii) actually follow from much more general results (see Corollary 4.3 and Proposition 4.7). Concretely, we show that, given a compact Riemannian surface \mathbb{M} and a subset $\{p_1, \dots, p_m\} \subset \mathbb{M}$, $m \geq 2$, there exist an open Riemann surface \mathcal{R} and a harmonic diffeomorphism $\phi: \mathcal{R} \rightarrow \mathbb{M} - \{p_1, \dots, p_m\}$ such that every end of \mathcal{R} is of hyperbolic conformal type (see Theorem 1.2 of [Ma] for a good setting).

Our strategy to show the harmonic diffeomorphism of Item (i) in Theorem I consists of constructing a maximal graph Σ over $\mathbb{S}^2 - \{p_1, \dots, p_m\}$ in the Lorentzian manifold $\mathbb{S}^2 \times \mathbb{R}_1$, with the conformal type of a circular domain. Then, the projection $\Sigma \rightarrow \mathbb{S}^2 - \{p_1, \dots, p_m\}$ is a surjective harmonic diffeomorphism.

In this direction, we prove the following general existence result:

Theorem II. Let $\mathbb{M} = (\mathbb{M}, \langle \cdot, \cdot \rangle_{\mathbb{M}})$ be a compact Riemannian manifold without boundary of dimension $n \in \mathbb{N}$, $n \geq 2$, and denote by $\mathbb{M} \times \mathbb{R}_1$ the product manifold $\mathbb{M} \times \mathbb{R}$ endowed with the Lorentzian metric $\langle \cdot, \cdot \rangle_{\mathbb{M}} - dt^2$. Let $m \in \mathbb{N}$, $m \geq 2$, and let $\mathfrak{A} = \{(p_i, t_i)\}_{i=1, \dots, m}$ be a subset of $\mathbb{M} \times \mathbb{R}_1$ such that

- $p_i \neq p_j$, and
- $|t_i - t_j| < \text{dist}_{\mathbb{M}}(p_i, p_j)$, for all $\{i, j\} \subset \{1, \dots, m\}$ with $i \neq j$.

Then there exists exactly one entire graph $\Sigma(\mathfrak{A})$ over \mathbb{M} in $\mathbb{M} \times \mathbb{R}_1$ such that

- $\mathfrak{A} \subset \Sigma(\mathfrak{A})$, and
- $\Sigma(\mathfrak{A}) - \mathfrak{A}$ is a spacelike maximal graph over $\mathbb{M} - \{p_i\}_{i=1, \dots, m}$.

Moreover the space \mathcal{G}_m of entire maximal graphs over \mathbb{M} in $\mathbb{M} \times \mathbb{R}_1$ with precisely m singularities, endowed with the topology of uniform convergence, is non-empty, and there exists an $m!$ -sheeted covering, $\mathcal{G}_m \rightarrow \mathcal{G}_m$, where \mathcal{G}_m is an open subset of $(\mathbb{M} \times \mathbb{R})^m$.

Let us point out that our method is different from the one of Collin and Rosenberg [CR] and strongly relies on the theory of maximal hypersurfaces in Lorentzian manifolds. More precisely, it is based on the construction of maximal hypersurfaces with isolated singularities in Lorentzian products $\mathbb{M} \times \mathbb{R}_1$. The study of complete maximal surfaces, with a finite number of singularities and their moduli spaces in the 3-dimensional Minkowski space \mathbb{L}^3 , was developed by Fernández, López and Souam [FLS]. Their study strongly relies on the Weierstrass representation for maximal surfaces in \mathbb{L}^3 . Our approach here relies on a different idea which consists of dealing with the existence problem in Theorem II as a generalized Dirichlet problem. Let us also point out that Klyachin and Miklyukov [KM] have obtained results on the existence of solutions, with a finite number of singularities, to the maximal hypersurface equation in the n -dimensional Minkowski space \mathbb{L}^n with prescribed boundary conditions.

Harmonic maps from Riemann surfaces into \mathbb{S}^2 are related to other natural geometric theories. For instance, the Gauss map of constant mean curvature surfaces in \mathbb{R}^3 is harmonic for the conformal structure induced by the immersion [Ru] (see also [Ke]), whereas the Gauss map of positive constant Gaussian curvature is harmonic for the conformal structure of the second fundamental form [GM]. The latter statement is the key in the proof of Theorem I(ii). More precisely, we show that if the Gauss map of a surface of positive constant curvature in \mathbb{R}^3 is a diffeomorphism onto $\mathbb{S}^2 - \{p\}$, then the conformal structure induced by the second fundamental form of the surface is that of \mathbb{C} .

On the other hand, using the harmonicity of the Gauss map of surfaces of positive constant curvature in \mathbb{R}^3 , harmonic diffeomorphisms from circular domains into domains in \mathbb{S}^2 bounded by a finite family of convex Jordan curves and satisfying a Neumann boundary condition have been recently shown by Gálvez, Hauswirth and Mira [GHM]. It is an open question whether a harmonic diffeomorphism as those in Theorem I(i) can be realized as the Gauss map of either a constant mean curvature or a constant Gaussian curvature surface in \mathbb{R}^3 .

The paper is laid out as follows. In Section 2 we state the necessary notations and preliminaries on harmonic maps between Riemannian manifolds and maximal graphs in Lorentzian product spaces. In Sections 3 and 4 we prove Theorems II and I, respectively. Also in Section 4 we introduce some background on both Riemann surfaces and surfaces of positive constant curvature, for a well understanding of the proofs of items (i) and (ii) in Theorem I. Finally, in Section 5 we discuss the relation between harmonic diffeomorphisms $U \rightarrow \mathbb{S}^2 - \{p_1, \dots, p_m\}$ as those of Theorem I(i) and conformal maximal immersions $U \rightarrow \mathbb{S}^2 \times \mathbb{R}_1$.

2. Preliminaries

Let $M = (M, g)$ and $N = (N, h)$ be smooth Riemannian manifolds. Given a smooth map $f: M \rightarrow N$ and a domain $\Omega \subset M$ with piecewise C^1 boundary $\partial\Omega$, the quantity

$$E_\Omega(f) = \frac{1}{2} \int_\Omega |df|^2 dV_g \quad (2.1)$$

is called the *energy* of f over Ω . Here dV_g denotes the volume element of M , and $|\cdot|$ the norm on (N, h) .

A smooth map $f: M \rightarrow N$ is said to be *harmonic* if it is a critical point of the energy functional, that is, if for any relatively compact domain $\Omega \subset M$ and any smooth variation $F: M \times (-\epsilon, \epsilon) \rightarrow N$ of f supported in Ω (i.e., F is a smooth map, $f_0 = f$ and $f_t|_{M-\Omega} = f|_{M-\Omega}$ for all $t \in (-\epsilon, \epsilon)$, where $f_t := F(\cdot, t): M \rightarrow N$ and $\epsilon > 0$), the first variation $\frac{d}{dt} E_\Omega(f_t)|_{t=0}$ is zero.

If M is 2-dimensional, that is to say, a Riemannian surface, then the energy integral (2.1) is invariant under conformal changes of the metric g , hence so is the harmonicity of f . Therefore, the harmonicity of a map from a Riemann surface to a Riemannian manifold is a well defined notion. On the other hand, the harmonicity of a map is not preserved under conformal changes in the metric of the target manifold.

See the surveys [EL1], [EL2], [HW] for appropriate references.

Remark 2.1. Throughout the paper we denote by $\mathbb{M} = (\mathbb{M}, \langle \cdot, \cdot \rangle_{\mathbb{M}})$ a compact n -dimensional Riemannian manifold without boundary, $n \in \mathbb{N}$, $n \geq 2$.

We denote by $\mathbb{M} \times \mathbb{R}_1$ the Lorentzian product space $\mathbb{M} \times \mathbb{R}$ endowed with the Lorentzian metric

$$\langle \cdot, \cdot \rangle = \pi_{\mathbb{M}}^*(\langle \cdot, \cdot \rangle_{\mathbb{M}}) - \pi_{\mathbb{R}}^*(dt^2),$$

where $\pi_{\mathbb{M}}$ and $\pi_{\mathbb{R}}$ denote the projections from $\mathbb{M} \times \mathbb{R}$ onto each factor. For simplicity, we write

$$\langle \cdot, \cdot \rangle = \langle \cdot, \cdot \rangle_{\mathbb{M}} - dt^2.$$

A smooth immersion $X: \Sigma \rightarrow \mathbb{M} \times \mathbb{R}_1$ of a connected n -dimensional manifold Σ is said to be *spacelike* if X induces a Riemannian metric $X^*(\langle \cdot, \cdot \rangle)$ on Σ .

Let $\Omega \subset \mathbb{M}$ be a connected domain and let $u: \Omega \rightarrow \mathbb{R}$ be a smooth function. Then the map

$$X^u: \Omega \rightarrow \mathbb{M} \times \mathbb{R}_1, \quad X^u(p) = (p, u(p)) \text{ for all } p \in \Omega,$$

determines a smooth graph over Ω in $\mathbb{M} \times \mathbb{R}_1$. The metric induced on Ω by $\langle \cdot, \cdot \rangle$ via X^u is given by

$$\langle \cdot, \cdot \rangle_u := (X^u)^*(\langle \cdot, \cdot \rangle) = \langle \cdot, \cdot \rangle_{\mathbb{M}} - du^2,$$

hence X^u is spacelike if and only if $|\nabla u| < 1$ on Ω , where ∇u denotes the gradient of u in Ω and $|\nabla u|$ denotes its norm, both with respect to the metric $\langle \cdot, \cdot \rangle_{\mathbb{M}}$ in Ω .

In this case the function u is said to be spacelike as well. If u is spacelike, then the mean curvature $H : \Omega \rightarrow \mathbb{R}$ of X^u is given by the equation

$$H = \frac{1}{n} \operatorname{Div} \left(\frac{\nabla u}{\sqrt{1 - |\nabla u|^2}} \right),$$

where Div denotes the divergence operator on Ω with respect to $\langle \cdot, \cdot \rangle_{\mathbb{M}}$.

A smooth function $u : \Omega \rightarrow \mathbb{R}$ and its graph $X^u : \Omega \rightarrow \mathbb{M} \times \mathbb{R}_1$ are said to be *maximal* if u is spacelike and H vanishes identically on Ω .

If $K \subset \mathbb{M}$ is compact then a function $u : K \rightarrow \mathbb{R}$ is said to be smooth (resp., spacelike, maximal) if and only if u extends to an open domain containing K as a smooth (resp., spacelike, maximal) function.

A locally Lipschitz function $u : \Omega \rightarrow \mathbb{R}$ is said to be *weakly spacelike* if and only if $|\nabla u| \leq 1$ a.e. in Ω . In this case the graph X^u is said to be weakly spacelike as well. If u is weakly spacelike then the area of $X^u(\Omega)$ is given by

$$\mathcal{A}(u) := \int_{\Omega} \sqrt{1 - |\nabla u|^2} dV_{\mathbb{M}}, \tag{2.2}$$

where $dV_{\mathbb{M}}$ denotes the volume element of $\langle \cdot, \cdot \rangle_{\mathbb{M}}$. A smooth $u : \bar{\Omega} \rightarrow \mathbb{R}$ is a critical point of (2.2) if and only if u is maximal.

If $u : \Omega \rightarrow \mathbb{R}$ is maximal then $X^u : (\Omega, \langle \cdot, \cdot \rangle_u) \rightarrow (\mathbb{M} \times \mathbb{R}_1, \langle \cdot, \cdot \rangle)$ is a harmonic map. In particular

$$\operatorname{Id} : (\Omega, \langle \cdot, \cdot \rangle_u) \rightarrow (\Omega, \langle \cdot, \cdot \rangle_{\mathbb{M}})$$

is a harmonic diffeomorphism, and

$$u : (\Omega, \langle \cdot, \cdot \rangle_u) \rightarrow \mathbb{R}$$

is a harmonic function.

3. Moduli space of maximal graphs with isolated singularities.

Proof of Theorem II

Throughout this section let $m \in \mathbb{N}$ and let $\mathfrak{X} = \{(p_i, t_i)\}_{i=1}^m$ denote a set of points in $\mathbb{M} \times \mathbb{R}$ such that $p_i \neq p_j$ if $i \neq j$ for all $i, j \in \{1, \dots, m\}$.

We denote by $\mathfrak{M}_{\mathfrak{X}}$ the space of continuous functions $u : \mathbb{M} \rightarrow \mathbb{R}$ such that

- $u(p_i) = t_i$ for all $i = 1, \dots, m$, and
- $\hat{u} := u|_{\mathbb{M} - \{p_i\}_{i=1}^m}$ is maximal.

The following claim trivially follows from the maximum principle for maximal surfaces.

Claim 3.1. *If $m = 1$ then $\mathfrak{M}_{\mathfrak{X}}$ consists of the constant function $u \equiv t_1$.*

From now on in this section assume that $m \geq 2$. The set \mathfrak{X} is said to satisfy the spacelike condition if and only if

$$|t_i - t_j| < \text{dist}_{\mathbb{M}}(p_i, p_j) \quad \text{for all } i, j \in \{1, \dots, m\}, i \neq j,$$

where $\text{dist}_{\mathbb{M}}(\cdot, \cdot)$ means distance in $(\mathbb{M}, \langle \cdot, \cdot \rangle_{\mathbb{M}})$.

From now on assume that \mathfrak{X} satisfies the spacelike condition.

For each $(i, n) \in \{1, \dots, m\} \times \mathbb{N}$ consider an open disk B_i^n in \mathbb{M} satisfying that ∂B_i^n is a smooth Jordan curve, $\overline{B_i^n} \cap \overline{B_j^n} = \emptyset$ if $i \neq j$, $\overline{B_i^{n+1}} \subset B_i^n$, and $\{p_i\} = \bigcap_{n \in \mathbb{N}} B_i^n$. Define $\Delta_n = \mathbb{M} - \bigcup_{i=1}^m B_i^n$, $n \in \mathbb{N}$. Let $\{t_i^n\}_{n \in \mathbb{N}}$ be a sequence of real numbers converging to t_i , $i = 1, \dots, m$.

Consider the function $\varphi_n: \partial \Delta_n \rightarrow \mathbb{R}$ such that

$$\varphi_n|_{\partial B_i^n} = t_i^n, \quad i = 1, \dots, m. \quad (3.1)$$

Since \mathfrak{X} is finite then \mathfrak{X} satisfies the spacelike condition if and only if there exists $\epsilon_{\mathfrak{X}} \in (0, 1)$ such that $|t_i - t_j| < \epsilon_{\mathfrak{X}} \cdot \text{dist}_{\mathbb{M}}(p_i, p_j)$ for all $i \neq j$. It follows that there exists $n_0 \in \mathbb{N}$ such that for each $n \geq n_0$, the function φ_n is $\epsilon_{\mathfrak{X},n}$ -Lipschitz for some $\epsilon_{\mathfrak{X},n} \in (0, 1)$.

It is proved in [Fe], p. 202, that there exists an $\epsilon_{\mathfrak{X},n}$ -Lipschitz extension $\tilde{\varphi}_n$ of φ_n to Δ_n . More precisely, such an extension is given by the formula:

$$\tilde{\varphi}_n(p) = \inf\{\varphi_n(x) + \epsilon_{\mathfrak{X},n} \text{dist}_{\mathbb{M}}(p, x), \quad x \in \partial \Delta_n\}, \quad \text{for } p \in \Delta_n. \quad (3.2)$$

Notice that $\tilde{\varphi}_n$ is smooth near $\partial \Delta_n$. A simple approximation argument then shows that

Claim 3.2. *For all $n \geq n_0$, there exists a smooth spacelike function $\bar{\varphi}_n: \Delta_n \rightarrow \mathbb{R}$ such that $\bar{\varphi}_n|_{\partial B_i^n} = t_i^n$, $i = 1, \dots, m$.*

Then by Theorem 5.1 of [Ge], there exists a maximal function $u_n: \Delta_n \rightarrow \mathbb{R}$ such that

$$u_n|_{\partial B_i^n} = \bar{\varphi}_n|_{\partial B_i^n} = t_i^n, \quad \text{for all } n \geq n_0. \quad (3.3)$$

Notice that the sequence $\{u_n\}_{n \in \mathbb{N}}$ is uniformly bounded (the u_n 's are maximal and there are uniform bounds on the boundary of the Δ_n 's). Moreover $|\nabla u_n| < 1$ on Δ_n , hence Ascoli–Arzela's Theorem and a diagonal argument give that, up to passing to a subsequence,

Claim 3.3. *$\{u_n\}_{n \in \mathbb{N}}$ uniformly converges on compact sets of*

$$\mathbb{M} - \{p_i\}_{i=1}^m = \bigcup_{n \in \mathbb{N}} \Delta_n$$

to a weakly spacelike function $\hat{u}: \mathbb{M} - \{p_i\}_{i=1}^m \rightarrow \mathbb{R}$.

Moreover, from (3.3), the convergence of $\{t_i^n\}_{n \in \mathbb{N}}$ to t_i and the Lipschitz continuity of \hat{u} one has that

Claim 3.4. \hat{u} extends to a weakly spacelike function $u : \mathbb{M} \rightarrow \mathbb{R}$ with $u(p_i) = t_i$ for all $i = 1, \dots, m$.

By the results in [Ba1], §6, \hat{u} is a maximal function except for a set of points $\Lambda \subset \mathbb{M} - \{p_i\}_{i=1}^m$ given by $\Lambda := \{p \in \mathbb{M} - \{p_i\}_{i=1}^m \mid (p, \hat{u}(p)) = \gamma(s_0)\}$ for some $0 < s_0 < 1$, where $\gamma : [0, 1] \rightarrow \mathbb{M} \times \mathbb{R}_1$ is a null geodesic such that $\gamma((0, 1)) \subset X^{\hat{u}}(\mathbb{M} - \{p_i\}_{i=1}^m)$ and $\pi_{\mathbb{M}}(\{\gamma(0), \gamma(1)\}) \subset \{p_i\}_{i=1}^m$.

Since \mathfrak{A} satisfies the spacelike condition then $\Lambda = \emptyset$ and

Claim 3.5. $u \in \mathfrak{M}_{\mathfrak{A}}$.

Now let us show the following

Claim 3.6. $\mathfrak{M}_{\mathfrak{A}}$ consists of exactly one element.

Proof. Consider $u_1, u_2 \in \mathfrak{M}_{\mathfrak{A}}$. By compactness of \mathbb{M} there exists $r_1 \in \mathbb{R}$, $r_1 \geq 0$, such that $r_1 + u_1 \geq u_2$ on \mathbb{M} and the equality holds at a non-empty subset $W \subset \mathbb{M}$. If $r_1 > 0$ then, since $u_1, u_2 \in \mathfrak{M}_{\mathfrak{A}}$, we must have $W \cap (\mathbb{M} - \{p_i\}_{i=1}^m) \neq \emptyset$. By the maximum principle for maximal surfaces $r_1 + u_1 = u_2$ on \mathbb{M} . This contradicts the fact that $u_1, u_2 \in \mathfrak{M}_{\mathfrak{A}}$. Therefore $r_1 = 0$ and so $u_1 \geq u_2$. In a symmetric way we also have $u_2 \geq u_1$. Therefore $u_1 = u_2$ and we are done. \square

At this point notice that the first part of Theorem II in the introduction follows from Claims 3.5 and 3.6. Even more,

Remark 3.7. The first part of Theorem II holds with the same proof if in the sentence

- $\Sigma(\mathfrak{A}) - \mathfrak{A}$ is a spacelike maximal graph over $\mathbb{M} - \{p_i\}_{i=1, \dots, m}$

one changes “maximal” by “constant mean curvature”.

Write $\mathfrak{M}_{\mathfrak{A}} = \{u_{\mathfrak{A}}\}$.

Denote by \mathfrak{G}_m the set of functions $u_{\mathfrak{A}}$ such that \mathfrak{A} consists of m points, \mathfrak{A} satisfies the spacelike condition and $|\nabla u_{\mathfrak{A}}| = 1$ at any point in $\pi_{\mathbb{M}}(\mathfrak{A})$.

Consider a sequence $\{u_{\mathfrak{A}_n}\}_{n \in \mathbb{N} \cup \{0\}} \subset \mathfrak{G}_m$. We say that $\{\mathfrak{A}_n\}_{n \in \mathbb{N}} \rightarrow \mathfrak{A}_0$ if and only if, up to a relabeling, $\{\text{dist}_{\mathbb{M} \times \mathbb{R}}((p_i^n, t_i^n), (p_i^0, t_i^0))\}_{n \in \mathbb{N}} \rightarrow 0$ for all $i = 1, \dots, m$, where $\mathfrak{A}_k = \{(p_i^k, t_i^k)\}_{i=1, \dots, m}$ for all $k \in \mathbb{N} \cup \{0\}$, and $\text{dist}_{\mathbb{M} \times \mathbb{R}}(\cdot, \cdot)$ means distance in $\mathbb{M} \times \mathbb{R}$ with respect to the metric $\langle \cdot, \cdot \rangle_{\mathbb{M}} + dt^2$. Likewise we define the convergence of a sequence of families of m points in \mathbb{M} .

Claim 3.8. $\{u_{\mathfrak{A}_n}\}_{n \in \mathbb{N}}$ uniformly converges to $u_{\mathfrak{A}_0}$ in the C^0 topology in \mathbb{M} if and only if $\{\mathfrak{A}_n\}_{n \in \mathbb{N}} \rightarrow \mathfrak{A}_0$.

Proof. Assume first that $\{u_{\mathfrak{X}_n}\}_{n \in \mathbb{N}}$ uniformly converges to $u_{\mathfrak{X}_0}$ in the C^0 topology in \mathbb{M} . Up to passing to a subsequence, assume that $\{\pi_{\mathbb{M}}(\mathfrak{X}_n)\}_{n \in \mathbb{N}}$ is convergent and let us show that $\{\pi_{\mathbb{M}}(\mathfrak{X}_n)\}_{n \in \mathbb{N}} \rightarrow \pi_{\mathbb{M}}(\mathfrak{X}_0)$. Indeed, suppose for a moment that there exist $p \in \pi_{\mathbb{M}}(\mathfrak{X}_0)$ and an open geodesic disc B centered at p such that, up to passing to a subsequence, $\pi_{\mathbb{M}}(\mathfrak{X}_n) \cap B = \emptyset$ for all $n \in \mathbb{N}$. Reasoning as in the paragraph preceding Claim 3.5, Bartnik’s results [Ba1] give that $u_{\mathfrak{X}_0}$ is smooth and spacelike around p , a contradiction. Then $\pi_{\mathbb{M}}(\mathfrak{X}_0) \subset \lim_{n \rightarrow \infty} \pi_{\mathbb{M}}(\mathfrak{X}_n)$. Since both sets consists of exactly m points then they agree. Since $\{u_{\mathfrak{X}_n}\}_{n \in \mathbb{N}} \rightarrow u_{\mathfrak{X}_0}$ and $\{\pi_{\mathbb{M}}(\mathfrak{X}_n)\}_{n \in \mathbb{N}} \rightarrow \pi_{\mathbb{M}}(\mathfrak{X}_0)$ then $\{\mathfrak{X}_n\}_{n \in \mathbb{N}} \rightarrow \mathfrak{X}_0$ as well.

For the converse assume that $\{\mathfrak{X}_n\}_{n \in \mathbb{N}} \rightarrow \mathfrak{X}_0$. For each $(i, n, k) \in \{1, \dots, m\} \times \mathbb{N} \times \mathbb{N}$ consider an open disc $B_{i,n}^k$ in \mathbb{M} such that $\partial B_{i,n}^k$ is a smooth Jordan curve, $\overline{B_{i,n}^k} \cap \overline{B_{j,n}^k} = \emptyset$ if $i \neq j$, $\overline{B_{i,n}^{k+1}} \subset B_{i,n}^k$, $p_i^n \in B_{i,n}^k$ and

$$\begin{aligned} &\text{for any compact } K \subset \mathbb{M} - \{p_1^0, \dots, p_m^0\} \text{ there} \\ &\text{exists } n_0 \in \mathbb{N} \text{ such that } K \subset \Delta_n^k \text{ for all } n \geq n_0, \end{aligned} \tag{3.4}$$

where $\mathfrak{X}_n = \{(p_1^n, t_i^n), \dots, (p_m^n, t_m^n)\}$ and $\Delta_n^k := \mathbb{M} - \bigcup_{i=1}^m B_{i,n}^k$. Let $u_n^k: \Delta_n^k \rightarrow \mathbb{R}$ be a maximal function satisfying $u_n^k|_{\partial B_{i,n}^k} = t_i^n$. (See the discussion preceding Claim 3.3.) By Claims 3.3, 3.4 and 3.5, the sequence

$$\{u_n^k\}_{k \in \mathbb{N}} \text{ uniformly converges in the } C^0 \text{ topology on } \mathbb{M} \text{ to } u_{\mathfrak{X}_n}. \tag{3.5}$$

Taking into account (3.4), a similar argument gives that the sequence

$$\{u_n^{f(n)}\}_{n \in \mathbb{N}} \text{ uniformly converges in the } C^0 \text{ topology on } \mathbb{M} \text{ to } u_{\mathfrak{X}_0} \tag{3.6}$$

as well, where $f: \mathbb{N} \rightarrow \mathbb{N}$ is any map with $f(n) \geq n$ for all $n \in \mathbb{N}$. Fix $p \in \mathbb{M} - \{p_1^0, \dots, p_m^0\}$ and $\epsilon > 0$. From (3.5), for any $n \in \mathbb{N}$ there exists $k_n \in \mathbb{N}$ such that

$$|u_n^{k_n} - u_{\mathfrak{X}_n}|(p) < \epsilon/2 \quad \text{for all } k \geq k_n, \tag{3.7}$$

where we are assuming that n and k are large enough so that $p \in \Delta_n^k$. Set $v_n := u_n^h$ for $h := \max\{k_n, n\}$. Then (3.6) gives $n_0 \in \mathbb{N}$ such that

$$|v_n - u_{\mathfrak{X}_0}|(p) < \epsilon/2 \quad \text{for all } n \geq n_0. \tag{3.8}$$

Combining (3.7) and (3.8) one has that $|u_{\mathfrak{X}_n} - u_{\mathfrak{X}_0}|(p) < \epsilon$ for all $n \geq n_0$. Since also $\{\mathfrak{X}_n\}_{n \in \mathbb{N}} \rightarrow \mathfrak{X}_0$, we conclude that $\{u_{\mathfrak{X}_n}\}_{n \in \mathbb{N}}$ simply converges to $u_{\mathfrak{X}_0}$. As \mathbb{M} is compact and the $u_{\mathfrak{X}_n}$ are weakly spacelike, this convergence is uniform on \mathbb{M} and we are done. \square

Consider m different points $\{p_1, \dots, p_m\} \subset \mathbb{M}$ and take $t_1 = \dots = t_{m-1} \neq t_m \in \mathbb{R}$ such that $\mathfrak{X} := \{(p_i, t_i)\}_{i=1, \dots, m}$ satisfies the spacelike condition. This is nothing but choosing t_1 and t_m close enough. By Claim 3.5, $u_{\mathfrak{X}}$ is well defined and by the maximum principle for maximal surfaces, $u_{\mathfrak{X}} \in \mathcal{G}_m$. Hence,

Claim 3.9. $\mathcal{G}_m \neq \emptyset$ for any $m \geq 2$.

Let $u \in \mathcal{G}_m$. By definition, a mark in u is an ordering $\mathcal{D} = ((q_1, r_1), \dots, (q_m, r_m))$ of the points in \mathfrak{X} , where $u = u_{\mathfrak{X}}$. Then we say that (u, \mathcal{D}) is a marked function. We denote by \mathcal{G}_m^* the space of marked functions in \mathcal{G}_m . We define the maps

$$\begin{aligned} \mathfrak{s}_1: \mathcal{G}_m^* &\rightarrow \mathcal{G}_m, & \mathfrak{s}_1(u, \mathcal{D}) &= u, \\ \mathfrak{s}_2: \mathcal{G}_m^* &\rightarrow (\mathbb{M} \times \mathbb{R})^m, & \mathfrak{s}_2(u, \mathcal{D}) &= \mathcal{D}. \end{aligned}$$

By Claim 3.6, the map \mathfrak{s}_2 is injective. Moreover,

Claim 3.10. $\mathfrak{s}_2(\mathcal{G}_m^*)$ is an open subset of $(\mathbb{M} \times \mathbb{R})^m$.

Proof. Let $(u, \mathcal{D}) \in \mathcal{G}_m^*$. Write

$$\mathcal{D} = ((q_1, r_1), \dots, (q_m, r_m)) \quad \text{and} \quad \mathfrak{X} = \{(q_1, r_1), \dots, (q_m, r_m)\}.$$

Since $u = u_{\mathfrak{X}}$ then \mathfrak{X} satisfies the spacelike condition. Reason by contradiction and assume that there exists a sequence $\{\mathcal{D}_n = ((q_1^n, r_1^n), \dots, (q_m^n, r_m^n))\}_{n \in \mathbb{N}}$ converging to \mathcal{D} in the metric topology of $(\mathbb{M} \times \mathbb{R})^m$ and $\mathcal{D}_n \notin \mathfrak{s}_2(\mathcal{G}_m^*)$ for all $n \in \mathbb{N}$. Write $\mathfrak{X}_n = \{(q_1^n, r_1^n), \dots, (q_m^n, r_m^n)\}$ and, up to passing to a subsequence, assume that \mathfrak{X}_n satisfies the spacelike condition for all $n \in \mathbb{N}$ (recall that \mathfrak{X} does so). Write $u_n = u_{\mathfrak{X}_n}$, $n \in \mathbb{N}$. By Claim 3.8, $\{u_n\}_{n \in \mathbb{N}}$ uniformly converges to u in the C^0 topology on \mathbb{M} . If, up to passing to a subsequence, $u_n: \mathbb{M} \rightarrow \mathbb{R}$ extends as a spacelike function to a point in $\pi_{\mathbb{M}}(\mathfrak{X}_n)$, that can be assumed to be q_1^n without loss of generality, $n \in \mathbb{N}$, then again Bartnik’s results [Ba1] give that u extends as a spacelike function to q_1 as well, a contradiction. Then $u_n \in \mathcal{G}_m$ (recall that \mathfrak{X}_n consists of m points), hence $(u_n, \mathcal{D}_n) \in \mathcal{G}_m^*$ and $\mathcal{D}_n \in \mathfrak{s}_2(\mathcal{G}_m^*)$, a contradiction. This proves the claim. \square

We set $\overline{\mathcal{G}_m} = \mathfrak{s}_2(\mathcal{G}_m^*)$. We can identify \mathcal{G}_m^* , endowed with the topology induced by the injection \mathfrak{s}_2 into $(\mathbb{M} \times \mathbb{R})^m$, with $\overline{\mathcal{G}_m}$. The permutation group σ_m of order m , acts naturally on \mathcal{G}_m^* as follows: for $\tau \in \sigma_m$ and $(u, \mathcal{D}) \in \mathcal{G}_m^*$ with $\mathcal{D} = ((q_1, r_1), \dots, (q_m, r_m))$, we set $\tau.(u, \mathcal{D}) = (u, \tau(\mathcal{D}))$ where $\tau(\mathcal{D}) = ((q_{\tau(1)}, r_{\tau(1)}), \dots, (q_{\tau(m)}, r_{\tau(m)}))$. This action is clearly free and properly discontinuous and the orbit space is naturally identified to \mathcal{G}_m . By Claim 3.8 the topology induced by the covering map coincides with the topology of C^0 -uniform convergence of graphs on \mathbb{M} .

This completes the proof of Theorem II.

4. Existence or non-existence of harmonic diffeomorphisms. Proof of Theorem I

Throughout this section we assume that $\mathbb{M} = (\mathbb{M}, \langle \cdot, \cdot \rangle_{\mathbb{M}})$ is of dimension $n = 2$, hence, a compact Riemannian surface without boundary.

Let us recall the following classification of Riemann surfaces. A compact Riemann surface (without boundary) is said to be elliptic. An open Riemann surface is said to be *hyperbolic* if it carries non-constant negative subharmonic functions, and it is said to be *parabolic* otherwise. A Riemann surface \mathcal{R} with non-empty boundary is said to be *parabolic* if bounded harmonic functions on \mathcal{R} are determined by their boundary values. Otherwise, \mathcal{R} is said to be *hyperbolic*. (See [AS], [Pe] for a good setting.) For instance, $\mathcal{R}_1 = \{z \in \mathbb{C} \mid 0 < |z| \leq 1\}$ is parabolic whereas $\mathcal{R}_2 = \{z \in \mathbb{C} \mid \alpha < |z| \leq 1\}$, $\alpha \in (0, 1)$, is hyperbolic.

Remark 4.1. An open Riemann surface \mathcal{R} is parabolic if and only if $\mathcal{R} - D$ is parabolic for any open relatively compact disc $D \subset \mathcal{R}$ with smooth boundary.

Indeed, if \mathcal{R} is parabolic then, by [FK], §IV.3.3, the Dirichlet problem has at most one bounded solution on $\mathcal{R} - D$, hence $\mathcal{R} - D$ is parabolic as well. For the converse assume that \mathcal{R} is hyperbolic. Then, by [FK], §IV.3.4, there exists a harmonic function w on $\mathcal{R} - D$ such that $0 < w < 1$ on $\mathcal{R} - \bar{D}$ and $w = 1$ on ∂D , hence $\mathcal{R} - D$ is hyperbolic and we are done.

Let $m \in \mathbb{N}$, $m \geq 2$, $u = u_{\mathfrak{M}} \in \mathcal{G}_m$ and set $\Omega = \mathbb{M} - \pi_{\mathbb{M}}(\mathfrak{A})$.

Recall that $X^u: (\Omega, \langle \cdot, \cdot \rangle_u) \rightarrow \mathbb{M} \times \mathbb{R}_1$ is a conformal harmonic map. Let $p \in \pi_{\mathbb{M}}(\mathfrak{A})$ and let A be an annular end of $(\Omega, \langle \cdot, \cdot \rangle_u)$ corresponding to p . Then A is conformally equivalent to an annulus $A(r, 1) := \{z \in \mathbb{C} \mid r < |z| \leq 1\}$ for some $0 \leq r < 1$. Identify $A \equiv A(r, 1)$ and notice that u extends continuously to $S(r) = \{z \in \mathbb{C} \mid |z| = r\}$ with $u|_{S(r)} = u(p)$. By [Ba2], $X^u(A)$ is tangent to either the upper or the lower light cone at $X^u(p)$ in $\mathbb{M} \times \mathbb{R}_1$. In particular p is either a strict local minimum or a strict local maximum of u . Then, up to a shrinking of A , we can assume that $u|_{S(1)}$ is constant, where $S(1) = \{z \in \mathbb{C} \mid |z| = 1\}$. Since $u|_A$ is harmonic, bounded and non-constant then $r > 0$ and A has hyperbolic conformal type. This proves that

Claim 4.2. $(\Omega, \langle \cdot, \cdot \rangle_u)$ is conformally an open Riemann surface with the same genus as \mathbb{M} and m hyperbolic ends.

In particular, one has the following

Corollary 4.3. Assume \mathbb{M} is a compact Riemannian surface. Let $m \geq 2$ and let $\{p_1, \dots, p_m\} \subset \mathbb{M}$. Then there exist an open Riemann surface \mathcal{R} and a harmonic diffeomorphism $\phi: \mathcal{R} \rightarrow \mathbb{M} - \{p_1, \dots, p_m\}$ such that every end of \mathcal{R} is of hyperbolic type.

By Koebe’s uniformization theorem, any finitely connected planar domain is conformally equivalent to a domain in $\bar{\mathbb{C}}$ whose frontier consists of points and circles. In this setting the corollary above gives Item (i) in Theorem I, that is, one obtains

the following existence result for harmonic diffeomorphism between hyperbolic and parabolic domains in \mathbb{S}^2 .

Corollary 4.4. *Let $m \in \mathbb{N}$, $m \geq 2$, and let $\{p_1, \dots, p_m\} \subset \mathbb{S}^2$.*

Then there exist a circular domain U in $\overline{\mathbb{C}}$ and a harmonic diffeomorphism $\phi: U \rightarrow \mathbb{S}^2 - \{p_1, \dots, p_m\}$.

Let us now show Theorem I (ii).

The proof of Theorem I(ii) fundamentally relies on the theory of surfaces of constant Gaussian curvature in Euclidean space. Before going into the details of the proof, let us state the necessary background on this theory.

Let S be a smooth surface and let $X: S \rightarrow \mathbb{R}^3$ be an immersion with constant Gauss curvature K equal to 1. For convenience we assume that S is simply connected.

Up to changing orientation if necessary, the second fundamental form II_X of X is a positive definite metric. Therefore, II_X induces on S a conformal structure. Denote by \mathcal{S} the Riemann surface with underlying differentiable structure S and conformal structure induced by II_X , and let $z = u + iv$ be a conformal parameter on \mathcal{S} . Then X may be understood as an immersion $X: \mathcal{S} \rightarrow \mathbb{R}^3$ and, following the results by Gálvez and Martínez [GM], §2.1, the equation $K = 1$ implies that

$$X_u = N \times N_v \quad \text{and} \quad X_v = -N \times N_u, \tag{4.1}$$

where $N: \mathcal{S} \rightarrow \mathbb{S}^2$ denotes the unit normal vector field of X . It follows that $N: \mathcal{S} \rightarrow \mathbb{S}^2$ is a harmonic local diffeomorphism.

Conversely, let $N: \mathcal{S} \rightarrow \mathbb{S}^2$ be a harmonic local diffeomorphism. Then the map $X: \mathcal{S} \rightarrow \mathbb{R}^3$ given by (4.1) is an immersion with constant Gauss curvature $K = 1$ (see [GM] again and recall that S is assumed to be simply connected).

On the other hand, in terms of the conformal parameter $z = u + iv$, the first, second and third fundamental forms of $X: \mathcal{S} \rightarrow \mathbb{R}^3$ are given by

$$\begin{aligned} \text{I}_X &= \langle dX, dX \rangle_{\mathbb{R}^3} = Qdz^2 + 2\mu|dz|^2 + \bar{Q}d\bar{z}^2, \\ \text{II}_X &= \langle dX, dN \rangle_{\mathbb{R}^3} = 2\rho|dz|^2, \\ \text{III}_X &= \langle dN, dN \rangle_{\mathbb{R}^3} = -Qdz^2 + 2\mu|dz|^2 - \bar{Q}d\bar{z}^2, \end{aligned} \tag{4.2}$$

where $\langle \cdot, \cdot \rangle_{\mathbb{R}^3}$ denotes the Euclidean metric in \mathbb{R}^3 , Qdz^2 is a holomorphic quadratic differential on \mathcal{S} , and μ and ρ are smooth positive real functions on \mathcal{S} , see [GHM]. Then, as Klotz pointed out in Remark 1 of [KI], there exists an immersion $Y: \mathcal{S} \rightarrow \mathbb{R}^3$ achieving III_X as its first fundamental form, II_X as its second and I_X as its third ones (recall that S is simply connected and observe that III_X is a positive definite metric). Since $X: \mathcal{S} \rightarrow \mathbb{R}^3$ is of constant Gauss curvature $K = 1$, it trivially follows from (4.2) that so is $Y: \mathcal{S} \rightarrow \mathbb{R}^3$.

Now we can prove Theorem I (ii).

Theorem 4.5. *There exists no harmonic diffeomorphism $\phi: \mathbb{D} \rightarrow \mathbb{S}^2 - \{p\}$, $p \in \mathbb{S}^2$.*

Proof. Let \mathcal{S} be a simply connected Riemann surface and let $\varphi: \mathcal{S} \rightarrow \mathbb{S}^2 - \{p\}$ be a harmonic diffeomorphism. To finish it suffices to check that \mathcal{S} is conformally equivalent to the complex plane \mathbb{C} .

By [GM], since $\varphi: \mathcal{S} \rightarrow \mathbb{S}^2 - \{p\}$ is a harmonic (local) diffeomorphism, then, up to replacing φ by $-\varphi$ if necessary, there exists an immersion $X: \mathcal{S} \rightarrow \mathbb{R}^3$ with Gauss map φ , constant curvature $K_X = 1$ and such that the conformal structure of \mathcal{S} is the one induced by the second fundamental form of X , II_X .

Denote by I_X and III_X the first and third fundamental forms of X , respectively. By [KI] there exists another immersion $Y: \mathcal{S} \rightarrow \mathbb{R}^3$ with constant curvature $K_Y = 1$, and such that the first, second and third fundamental forms of Y are given by $\text{I}_Y = \text{III}_X$, $\text{II}_Y = \text{II}_X$ and $\text{III}_Y = \text{I}_X$, respectively, and

$$\text{the conformal structure of } \mathcal{S} \text{ is the one induced by } \text{II}_Y = \text{II}_X. \tag{4.3}$$

Since $\varphi: \mathcal{S} \rightarrow \mathbb{S}^2 - \{p\}$ is a diffeomorphism and $\text{I}_Y = \text{III}_X = \langle d\varphi, d\varphi \rangle_{\mathbb{R}^3} = \varphi^* (\langle \cdot, \cdot \rangle_{\mathbb{S}^2})$ (here $\langle \cdot, \cdot \rangle_{\mathbb{S}^2}$ denotes the canonical metric in \mathbb{S}^2), then $\varphi^{-1}: \mathbb{S}^2 - \{p\} \rightarrow (\mathcal{S}, \text{I}_Y)$ is an isometry. Since obviously $Y: (\mathcal{S}, \text{I}_Y) \rightarrow \mathbb{R}^3$ is an isometric immersion, then

$$Y \circ \varphi^{-1}: \mathbb{S}^2 - \{p\} \rightarrow \mathbb{R}^3 \text{ is an isometric immersion}$$

as well. Following [Po], p. 419, $Y \circ \varphi^{-1}$ is the restriction to $\mathbb{S}^2 - \{p\} \subset \mathbb{R}^3$ of a rigid motion of \mathbb{R}^3 . (An alternate proof of the rigidity of $\mathbb{S}^2 - \{p\}$ in \mathbb{R}^3 can be given using the local structure around the singularities, of immersed surfaces with constant Gaussian curvature in \mathbb{R}^3 having isolated singularities and finite area, see Theorems 5 and 7 of [GHM].) In particular, $Y(\mathcal{S}) \subset \mathbb{R}^3$ is a once-punctured round sphere. Therefore, the conformal structure induced on \mathcal{S} by $\text{II}_Y = \text{II}_X$ is that of \mathbb{C} . This and (4.3) conclude the proof. \square

Remark 4.6. Lemaire [Le] showed that if a harmonic map $\varphi: \mathbb{D} \rightarrow N$ with finite energy satisfies that $\varphi|_{\mathbb{S}^1}$ is constant then φ is constant as well, where N is an arbitrary Riemannian manifold. The above theorem particularly shows that the condition on the energy of φ can be removed if φ is a diffeomorphism and $N = \mathbb{S}^2$.

Finally Theorem I (iii) is a very special instance of the following

Proposition 4.7. *Let \mathcal{R} be a parabolic open Riemann surface, let N be an oriented Riemannian surface and let $\phi: \mathcal{R} \rightarrow N$ be a harmonic local diffeomorphism. Suppose either that N has Gaussian curvature $K_N > 0$ or that $K_N \geq 0$ and N has no flat open subset.*

Then ϕ is either holomorphic or antiholomorphic.

Proof. Assume for instance that ϕ preserves orientation and let us check that ϕ is holomorphic. Let z (resp. ϕ) be a local conformal parameter in \mathcal{R} (resp. in N). The metric on N writes $\rho(\phi)|d\phi|^2$. A conformal metric on \mathcal{R} writes $\lambda(z)|dz|^2$. Following [SY] we consider the following partial energy densities on \mathcal{R} :

$$|\partial\phi|^2 = \frac{\rho(\phi(z))}{\lambda(z)} \left| \frac{\partial\phi}{\partial z} \right|^2 \quad \text{and} \quad |\bar{\partial}\phi|^2 = \frac{\rho(\phi(z))}{\lambda(z)} \left| \frac{\partial\phi}{\partial \bar{z}} \right|^2. \tag{4.4}$$

Denote by $J(\phi)$ the Jacobian of ϕ . We have $J(\phi) = |\partial\phi|^2 - |\bar{\partial}\phi|^2$. By our hypothesis $J(\phi) > 0$, that is, $|\partial\phi| > |\bar{\partial}\phi|$.

Reason by contradiction and assume that ϕ is not holomorphic, that is to say, $|\bar{\partial}\phi|$ is not identically zero on \mathcal{R} . In this case, its zeroes are isolated [SY]. Set $\mathcal{R}^* := \mathcal{R} - \{|\bar{\partial}\phi| = 0\}$. We have

$$\log \frac{|\bar{\partial}\phi|}{|\partial\phi|} < 0 \quad \text{on } \mathcal{R}^*. \tag{4.5}$$

By the Bochner formula (see again Chapter 1, §7, in [SY]):

$$\Delta_{\mathcal{R}} \log \frac{|\bar{\partial}\phi|}{|\partial\phi|} = 2K_N J(\phi). \tag{4.6}$$

Now note that the parabolicity of \mathcal{R} implies that of \mathcal{R}^* (see Claim 4.8 below). Suppose $K_N > 0$. By equations (4.5) and (4.6), $\log \frac{|\bar{\partial}\phi|}{|\partial\phi|}$ is a non-constant negative subharmonic function on the parabolic surface \mathcal{R}^* , which is a contradiction. Suppose now that $K_N \geq 0$. Again, by the equations (4.5) and (4.6), the function $\log \frac{|\bar{\partial}\phi|}{|\partial\phi|}$ is subharmonic and hence constant. From (4.6) we get $K_N J(\phi) \equiv 0$. Since $J(\phi) > 0$, we conclude that $K_N \equiv 0$ on the open set $\phi(\mathcal{R}^*)$, which contradicts our hypothesis.

In the case when ϕ reverses orientation then a parallel argument gives that ϕ is antiholomorphic. This concludes the proof. \square

Since in the setting of Theorem I (iii) the domains $\bar{\mathbb{C}} - \{z_1, \dots, z_m\}$ and $\mathbb{S}^2 - \bigcup_{j=1}^m D_j$ are not conformally equivalent, then the result holds.

For the lack of a reference, we now prove the following well known fact needed in the proof of Proposition 4.7.

Claim 4.8. *Let \mathcal{R} be an open parabolic Riemann surface and let $E \subset \mathcal{R}$ be a closed subset consisting of isolated points.*

Then $\mathcal{R}^ := \mathcal{R} - E$ is an open parabolic Riemann surface.*

Proof. The fact that \mathcal{R}^* is an open Riemann surface is evident. Let us show that it is parabolic. Indeed, consider B an open relatively compact disc in \mathcal{R}^* with smooth boundary and denote by \mathcal{N} the Riemann surface with boundary $\mathcal{N} := \mathcal{R}^* - B$. To

finish it suffices to prove that \mathcal{N} is parabolic (see Remark 4.1). Let $u: \mathcal{N} \rightarrow \mathbb{R}$ be a non-constant bounded harmonic function with $u|_{\partial\mathcal{N}} = 0$. Since E consists of isolated points then u extends harmonically to $\mathcal{N} \cup E = \mathcal{R} - B$. Since $\mathcal{R} - B$ is parabolic (see Remark 4.1 again), $\partial(\mathcal{R} - B) = \partial\mathcal{N}$ and $u|_{\partial\mathcal{N}} = 0$, then u is identically zero on $\mathcal{N} \subset \mathcal{R} - B$. This proves that \mathcal{N} is parabolic and we are done. \square

This completes the proof of Theorem I.

5. Maximal graphs and harmonic diffeomorphisms between surfaces

Let \mathcal{R} be a Riemann surface and let N be a Riemannian surface. A map $X = (f, h): \mathcal{R} \rightarrow N \times \mathbb{R}_1$ is conformal if and only if

$$\left| \frac{\partial f}{\partial x} \right|^2 - \left(\frac{\partial h}{\partial x} \right)^2 = \left| \frac{\partial f}{\partial y} \right|^2 - \left(\frac{\partial h}{\partial y} \right)^2 \quad \text{and} \quad \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle = \frac{\partial h}{\partial x} \frac{\partial h}{\partial y}, \tag{5.1}$$

where $z = x + iy$ is a local conformal parameter on \mathcal{R} and $|\cdot|$ and $\langle \cdot, \cdot \rangle$ denote the norm and metric on N , respectively. If in addition X is harmonic then the above equalities hold if and only if the Hopf differential of $f: \mathcal{R} \rightarrow N$,

$$\Phi_f = \left\langle \frac{\partial f}{\partial z}, \frac{\partial f}{\partial z} \right\rangle dz^2 = \frac{1}{4} \left(\left| \frac{\partial f}{\partial x} \right|^2 - \left| \frac{\partial f}{\partial y} \right|^2 - 2i \left\langle \frac{\partial f}{\partial x}, \frac{\partial f}{\partial y} \right\rangle \right) dz^2,$$

and the one of $h: \mathcal{R} \rightarrow \mathbb{R}$,

$$\Phi_h = \left(\frac{\partial h}{\partial z} \right)^2 dz^2 = \frac{1}{4} \left(\left(\frac{\partial h}{\partial x} \right)^2 - \left(\frac{\partial h}{\partial y} \right)^2 - 2i \frac{\partial h}{\partial x} \frac{\partial h}{\partial y} \right) dz^2,$$

agree.

Furthermore, a conformal harmonic immersion X is spacelike (hence, a conformal maximal immersion) if and only if

$$\left| \frac{\partial f}{\partial x} \right|^2 > \left(\frac{\partial h}{\partial x} \right)^2. \tag{5.2}$$

On the other hand, let $\phi: \mathcal{R} \rightarrow N$ be a harmonic map and denote by Φ_ϕ its Hopf differential. Consider $(\tilde{\mathcal{R}}, \Pi)$ a 2-sheeted covering of \mathcal{R} such that $\tilde{\Phi}_\phi := \Phi_\phi \circ \Pi$ has a well defined square root, and write $\tilde{\Phi}_\phi = (\tilde{\varphi}(z)dz)^2$ on a local conformal parameter $z = x + iy$ on $\tilde{\mathcal{R}}$. Observe that $(\tilde{\mathcal{R}}, \Pi)$ is possibly branched at the zeros of Φ_ϕ .

Consider now $(\hat{\mathcal{R}}, \hat{\Pi})$ a covering of $\tilde{\mathcal{R}}$ such that $\hat{\varphi} := \tilde{\varphi} \circ \hat{\Pi}$ has no real periods, and define

$$X_\phi: \hat{\mathcal{R}} \rightarrow N \times \mathbb{R}_1, \quad X_\phi = (f_\phi, h_\phi),$$

where

$$f_\phi := \phi \circ \tilde{\Pi} \circ \hat{\Pi} \quad \text{and} \quad h_\phi := \Re \int \hat{\varphi} dz.$$

Notice that $(\hat{\mathcal{R}}, \hat{\Pi})$ is infinitely sheeted unless $\tilde{\varphi}$ has no real periods (recall that the periods are additive).

Clearly, the Hopf differentials of f_ϕ and h_ϕ agree, so the above discussion gives that X_ϕ is a conformal harmonic map. Assume in addition that ϕ , and so f_ϕ , is a local harmonic diffeomorphism. From (5.1) and Cauchy–Schwarz inequality one has

$$\left| \frac{\partial f_\phi}{\partial x} \right| \geq \left| \frac{\partial h_\phi}{\partial x} \right|.$$

Assume the equality holds at a point $p \in \hat{\mathcal{R}}$. Then (5.1) gives that, at the point p , $|\partial f_\phi / \partial y| = |\partial h_\phi / \partial y|$ as well and $|\langle \partial f_\phi / \partial x, \partial f_\phi / \partial y \rangle| = |\partial f_\phi / \partial x| \cdot |\partial f_\phi / \partial y|$. This contradicts that f_ϕ is a local diffeomorphism. Therefore $|\partial f_\phi / \partial x| > |\partial h_\phi / \partial x|$ on $\hat{\mathcal{R}}$ and $X_\phi: \hat{\mathcal{R}} \rightarrow \mathcal{N} \times \mathbb{R}_1$ is a possibly branched conformal maximal immersion (see (5.2)).

In this way we have showed the following

Proposition 5.1. *Let \mathcal{R} be a Riemann surface, let N be Riemannian surface and let $\phi: \mathcal{R} \rightarrow N$ be a local harmonic diffeomorphism.*

Then there exist a covering $(\hat{\mathcal{R}}, \Pi)$ of \mathcal{R} and a possibly branched conformal maximal immersion $X_\phi = (f_\phi, h_\phi): \hat{\mathcal{R}} \rightarrow N \times \mathbb{R}_1$ such that $f_\phi = \phi \circ \Pi: \hat{\mathcal{R}} \rightarrow N$.

Let us now focus on the particular case when $U := \mathcal{R} \subset \mathbb{C}$ is a finitely connected circular domain, N is the sphere S^2 with a finite number of points removed and ϕ extends C^1 to the closure \bar{U} of U . Denote by \mathcal{N} the double of \bar{U} (see [St] for details on this construction). Recall that \mathcal{N} is a compact Riemann surface carrying an antiholomorphic involution $\mathcal{J}: \mathcal{N} \rightarrow \mathcal{N}$ having the boundary of \bar{U} as set of fixed points. Let Φ be a smooth quadratic differential on \bar{U} and holomorphic on U . Assume that $\Phi = \varphi(z)dz^2$ with $\varphi(x) \in \mathbb{R}$ for all x for any local conformal parameter $z = x + iy$ on \bar{U} applying a piece of the boundary ∂U of \bar{U} into $\mathbb{R} \subset \mathbb{C}$, then Φ extends holomorphically to \mathcal{N} in the form $\mathcal{J}^* \Phi = \bar{\Phi}$.

Let $\phi: U \rightarrow N$ be a harmonic diffeomorphism extending C^1 to \bar{U} . Obviously ϕ is constant over any connected component of ∂U . Let $z = x + iy$ be a conformal parameter on \bar{U} with $y|_{\partial U} = 0$. Then $\partial \phi / \partial x = 0$ on ∂U , hence the Hopf differential of ϕ can be written on ∂U as

$$(\Phi_\phi)|_{\partial U} = -\frac{1}{4} \left| \frac{\partial \phi}{\partial y} \right|^2 dz^2. \tag{5.3}$$

In particular, Φ_ϕ extends holomorphically to \mathcal{N} with $\mathcal{J}^* \Phi_\phi = \bar{\Phi}_\phi$. This particularly gives that

$$\Phi_\phi \text{ has finitely many zeros on } U. \tag{5.4}$$

Now, as above, we can take a 2-sheeted covering (\tilde{U}, Π) of U such that $\tilde{\Phi}_\phi := \Phi_\phi \circ \Pi$ has a well defined square root. Write $\tilde{\Phi} = (\varphi(z)dz)^2$ in a local conformal parameter z on \tilde{U} . From (5.3) one obtains that $\varphi(z)dz$ has no real periods. Then taking into account (5.4) and following the discussion preceding Proposition 5.1 one has the following

Theorem 5.2. *Let $\phi: U \rightarrow \mathbb{S}^2 - \{p_1, \dots, p_m\}$ be a harmonic diffeomorphism extending C^1 to \bar{U} , where U is a finitely connected circular domain and $\{p_1, \dots, p_m\}$ is a finite subset in \mathbb{S}^2 .*

Then there exist a 2-sheeted covering (\hat{U}, Π) of U and a possibly finitely branched conformal maximal immersion $X_\phi = (f_\phi, h_\phi): \hat{U} \rightarrow \mathbb{S}^2 \times \mathbb{R}_1$ such that $f_\phi = \phi \circ \Pi$.

In the proof of the above theorem, we have used that ϕ extends C^1 to \bar{U} in order to obtain that the Hopf differential Φ_ϕ of ϕ extends holomorphically to the double of U . The authors do not know whether this hypothesis can be removed from the statement of the theorem.

References

- [AS] L. V. Ahlfors and L. Sario, *Riemann surfaces*. Princeton University Press, Princeton, NJ, 1960. [Zbl 0196.33801](#) [MR 0114911](#)
- [Ba1] R. Bartnik, Regularity of variational maximal surfaces. *Acta Math.* **161** (1988), no. 3–4, 145–181. [Zbl 0667.53049](#) [MR 0971795](#)
- [Ba2] R. Bartnik, Isolated singular points of Lorentzian mean curvature hypersurfaces. *Indiana Univ. Math. J.* **38** (1989), no. 4, 811–827. [Zbl 0695.53043](#) [MR 1029678](#)
- [CR] P. Collin and H. Rosenberg, Construction of harmonic diffeomorphisms and minimal graphs. *Ann. of Math. (2)* **172** (2010), 1879–1906. [Zbl 1209.53010](#) [MR 2726102](#)
- [EL1] J. Eells and L. Lemaire, A report on harmonic maps. *Bull. London Math. Soc.* **10** (1978), no. 1, 1–68. [Zbl 0401.58003](#) [MR 1363514](#)
- [EL2] J. Eells and L. Lemaire, Another report on harmonic maps. *Bull. London Math. Soc.* **20** (1988), no. 5, 385–524. [Zbl 0669.58009](#) [MR 1363515](#)
- [FK] H. M. Farkas and I. Kra, *Riemann surfaces*. Grad. Texts in Math. 71, Springer, New York 1980. [Zbl 0475.30001](#) [MR 0583745](#)
- [Fe] H. Federer, *Geometric measure theory*. Grundlehren Math. Wiss. 153, Springer, Berlin 1969. [Zbl 0176.00801](#) [MR 0257325](#)
- [FLS] I. Fernández, F.J. López and R. Souam, The space of complete embedded maximal surfaces with isolated singularities in the 3-dimensional Lorentz–Minkowski space \mathbb{L}^3 . *Math. Ann.* **332** (2005), no. 3, 605–643. [Zbl 1081.53052](#) [MR 2181764](#)
- [GM] J. A. Gálvez and A. Martínez, The Gauss map and second fundamental form of surfaces in \mathbb{R}^3 . *Geom. Dedicata* **81** (2000), no. 1–3, 181–192. [Zbl 0965.53007](#) [MR 1772201](#)

- [GHM] J. A. Gálvez, L. Hauswirth and P. Mira, Surfaces of constant curvature in \mathbb{R}^3 with isolated singularities. *Adv. Math.* **241** (2013), 103–126. [Zbl 06243957](#) [MR 3053706](#)
- [GR] J. A. Gálvez and H. Rosenberg, Minimal surfaces and harmonic diffeomorphisms from the complex plane onto certain Hadamard surface. *Amer. J. Math.* **132** (2010), 1249–1273. [Zbl 1229.53064](#) [MR 2732346](#)
- [Ge] C. Gerhardt, H -surfaces in Lorentzian manifolds. *Comm. Math. Phys.* **89** (1983), no. 4, 523–553. [Zbl 0519.53056](#) [MR 0713684](#)
- [He] E. Heinz, Über die Lösungen der Minimalflächengleichung. *Nachr. Akad. Wiss. Göttingen. Math.-Phys. Kl.* **1952** (1952), 51–56. [Zbl 0048.15401](#) [MR 0054182](#)
- [HW] F. Hélein and J. C. Wood, Harmonic maps. In *Handbook of global analysis*, Elsevier Sci. B. V., Amsterdam 2008, 417–491, 1213. [Zbl 1236.58002](#) [MR 2389639](#)
- [Ke] K. Kenmotsu, Weierstrass formula for surfaces of prescribed mean curvature. *Math. Ann.* **245** (1979), no. 2, 89–99. [Zbl 0402.53002](#) [MR 0552581](#)
- [KI] T. Klotz Milnor, Abstract Weingarten surfaces. *J. Differential Geom.* **15** (1980), no. 3, 365–380 (1981). [Zbl 0464.53002](#) [MR 0620893](#)
- [KM] A. A. Klyachin and V. M. Miklyukov, Existence of solutions with singularities for the maximal surface equation in Minkowski space. *Russian Acad. Sci. Sb. Math.* **80** (1995), 87–104. [Zbl 0620893](#) [MR 1257338](#)
- [Le] L. Lemaire, Applications harmoniques de surfaces riemanniennes. *J. Differential Geom.* **13** (1978), no. 1, 51–78. [Zbl 0388.58003](#) [MR 0520601](#)
- [Ma] V. Markovic, Harmonic diffeomorphisms and conformal distortion of Riemann surfaces. *Comm. Anal. Geom.* **10** (2002), no. 4, 847–876. [Zbl 1024.53044](#) [MR 1925504](#)
- [Pe] J. Pérez, Parabolicity and minimal surfaces. In *Global theory of minimal surfaces*, Clay Math. Proc. 2, Amer. Math. Soc., Providence, RI, 2005, 163–174. [Zbl 1100.53013](#) [MR 2167259](#)
- [Po] A. V. Pogorelov, *Extrinsic geometry of convex surfaces*. Translated from the Russian by Israel Program for Scientific Translations, Transl. Math. Monogr. 35, Amer. Math. Soc., Providence, RI, 1973. [Zbl 0311.53067](#) [MR 0346714](#)
- [Ru] E. A. Ruh, Asymptotic behaviour of non-parametric minimal hypersurfaces. *J. Differential Geom.* **4** (1970), 509–513. [Zbl 0206.50203](#) [MR 0276877](#)
- [SY] R. Schoen and S. T. Yau, Lectures on harmonic maps. Conf. Proc. Lecture Notes Geom. Topology, II, International Press, Cambridge, MA, 1997. [Zbl 0886.53004](#) [MR 1474501](#)
- [St] K. Strebel, *Quadratic differentials*. *Ergeb. Math. Grenzgeb.* (3) 5, Springer, Berlin 1984. [Zbl 0547.30001](#) [MR 0743423](#)

Received October 4, 2011

Antonio Alarcón, Departamento de Geometría y Topología, Universidad de Granada,
18071 Granada, Spain

E-mail: alarcon@ugr.es

Rabah Souam, Institut de Mathématiques de Jussieu, CNRS UMR 7586, Université Paris
Diderot – Paris 7, “Géométrie et Dynamique”, Site Chevaleret, Case 7012, 75205 Paris
Cedex 13, France

E-mail: souam@math.jussieu.fr