

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 89 (2014)

Artikel: Riemann surfaces and totally real tori
Autor: Duval, Julien / Gayet, Damien
DOI: <https://doi.org/10.5169/seals-515673>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Riemann surfaces and totally real tori

Julien Duval and Damien Gayet

Abstract. Given a totally real torus unknotted in the unit sphere S^3 of \mathbb{C}^2 , we prove the following alternative: either the torus is rationally convex and there exists a filling of the torus by holomorphic discs, or its rational hull contains a holomorphic annulus or a pair of holomorphic discs.

Mathematics Subject Classification (2010). 32E20, 58J32.

Keywords. Totally real torus, filling by holomorphic discs, rational convexity.

Introduction

In this paper we address the following question: given a totally real torus in \mathbb{C}^2 , does there always exist a compact Riemann surface in \mathbb{C}^2 with boundary in (or simply *attached to*) the torus?

Recall that (closed connected) surfaces in \mathbb{C}^2 are *totally real* if they are never tangent to a complex line. The only orientable ones are tori. Special cases are *Lagrangian* tori, those on which the standard Kähler form of \mathbb{C}^2 vanishes.

Our question is motivated by geometric function theory (see [15] for background). Given a compact set K in \mathbb{C}^2 , its *polynomial hull* \hat{K} is defined as

$$\hat{K} = \{z \text{ in } \mathbb{C}^2 / |P(z)| \leq \|P\|_K \text{ for every polynomial } P\}.$$

The set K is *polynomially convex* if $\hat{K} = K$. In this case K satisfies Runge theorem. Note that any compact Riemann surface attached to K is contained in \hat{K} . It is therefore tempting to explain the presence of a non trivial hull by Riemann surfaces, at least for nice sets like orientable surfaces (they are not polynomially convex for homological reasons). But quite often a complex tangency of a surface locally gives birth to small holomorphic discs attached to it. Thus the very first global problem arises with totally real orientable surfaces, namely tori.

Note that, in the definitions above, instead of polynomials we could as well work with rational functions without poles on K . This gives rise to the notions of *rational hull* and *rational convexity*. Again an obstruction to rational convexity is the presence

of a compact Riemann surface C attached to K with the additional restriction that ∂C bounds in K .

Here is a bit of history around our question. In 1985 Gromov [10] gave a positive answer for Lagrangian tori, constructing holomorphic discs attached to them. In 1996 by the same method Alexander [1] exhibited for every totally real torus a proper holomorphic disc with all its boundary except one point in the torus. Later on [2] he gave examples of totally real tori without holomorphic discs with full boundary in them, but still admitting holomorphic annuli attached to them.

In the present work we focus¹ on tori in the unit sphere S^3 of \mathbb{C}^2 . They are *unknotted* if they are isotopic to the standard torus in S^3 . We prove the following

Theorem. *Let T be a totally real torus unknotted in S^3 . Then either T is rationally convex and bounds a solid torus foliated by holomorphic discs in the unit ball B , or its rational hull contains a holomorphic annulus or a pair of holomorphic discs attached to T .*

The solid torus is called a *filling* of T which is said in this case *fillable*. The standard torus is an example of the first situation, while the second is illustrated by the following

Example (compare with [2]). Consider the conjugate Hopf fibration

$$\pi: S^3 \subset \mathbb{C}^2 \rightarrow S^2 \subset \mathbb{C} \times \mathbb{R}, \quad (z, w) \mapsto (2zw, |z|^2 - |w|^2).$$

Remark that the fibers of π are circles. Denote by T_γ the preimage by π of an embedded closed curve γ in S^2 . Then T_γ is an unknotted torus in S^3 , totally real if the projection of γ on \mathbb{C} is immersed. Choose this projection as a figure eight which avoids the origin. It follows (see [2]) that every compact Riemann surface with boundary in T_γ is in a fiber of the polynomial $p(z, w) = 2zw$. But T_γ does not separate $p^{-1}(a)$ except if a is the double point of the figure eight. We then get only one holomorphic annulus attached to T . If on the other hand the figure eight intersects itself at the origin we get instead a pair of holomorphic discs attached to T .

The proof of the theorem relies on the technique of filling spheres by holomorphic discs due to Bedford and Klingenberg [5] and Kruzhilin [12] (see also Eliashberg [8]). This is where the restriction to S^3 enters. The spheres come into the picture as approximations of a lift of the torus in a suitable covering. More precisely take a totally real unknotted torus T in S^3 . It divides S^3 in two solid tori. In the same manner its hull \hat{T} separates the unit ball B in two pseudoconvex components. At least one of them has a universal covering which unwinds the corresponding solid torus. Push T slightly in this good component, building a sequence of tori T_n converging

¹following [13] which by the way seems incorrect (see our example)

toward T . We therefore get as a lift of T_n a periodic cylinder sitting in a pseudoconvex boundary. Approximate it by a sphere S_n containing say $2n$ periods of the cylinder.

We are now in position to apply the technique of filling. It provides a sequence of balls bounded by S_n and foliated by holomorphic discs. Single out one of these discs passing through the equator of S_n and call Δ_n its projection downstairs. The alternative reads as follows: either the area of Δ_n remains bounded, or not.

In the former case (the *rationally convex case*) we check that the tori T_n are fillable for large n , and that their fillings converge in some sense to a filling of T . This relies on Gromov compactness theorem.

In the latter (the *non rationally convex case*) we rather look at the limit of Δ_n in terms of currents. Define U as the limit of the normalized currents of integration on Δ_n . Then U is a positive current such that dU bounds in T . Therefore the support of U is contained in the rational hull of T . Moreover a dividing process of U shows that it can be written as an integral of currents of integration over Riemann surfaces. Finally we apply Ahlfors theory of covering surfaces to prove that these Riemann surfaces are holomorphic discs or annuli.

Before entering the details of the proof, we collect some background. In the sequel a limit of a sequence often occurs up to extracting a subsequence, even if not explicitly mentioned. Pseudoconvex domains are also sometimes confused with their closure.

1. Background

a) Filling spheres. Recall the central result of [12] (see also [5]).

Theorem. *Let Ω be a bounded strictly pseudoconvex domain in \mathbb{C}^2 and S a sphere in $\partial\Omega$. Suppose that the complex tangencies of S are elliptic or hyperbolic points. Then S bounds a unique ball Σ in Ω foliated by holomorphic discs.*

This ball Σ is called the *filling* of S . The *complex tangencies* of S are the points where S is tangent to a complex line. Being of elliptic or hyperbolic type (see [5], [12] for the definition) is a generic condition. It can be achieved by a small perturbation localized near the complex points.

The picture looks as follows. Take a sphere in \mathbb{R}^3 endowed with its height function, which is Morse if the sphere is generic. Elliptic points correspond to local maxima and minima of the height, while hyperbolic points translate in saddle points. By Morse theory we have $e - h = 2$ where e and h are respectively the number of elliptic and hyperbolic points. The filling corresponds to the ball bounded by the sphere foliated by the level sets of the height. Therefore all the holomorphic discs of the filling are smooth up to the boundary except those touching a hyperbolic point which have corners.

Another way to describe the complex points of S is via its *characteristic foliation*. This is the foliation generated by the characteristic line field $T_{\mathbb{C}}\partial\Omega \cap TS$ where $T_{\mathbb{C}}\partial\Omega$ is the complex part of $T\partial\Omega$. It is singular precisely at the complex points of S , elliptic points corresponding to foci and hyperbolic to saddle points. The characteristic foliation gives a control on the discs of the filling, in the sense that their boundaries are always transversal to it. This comes from Hopf lemma which asserts that a holomorphic disc contained in Ω is transversal to $\partial\Omega$.

Here are further properties of the filling. First every compact Riemann surface in Ω attached to S is contained in Σ . Next Σ is the envelope of holomorphy of S . Hence Σ is contained in any pseudoconvex domain containing S . Finally if we divide out the sphere S into two half spheres by an equator, at least one of them can be partially filled in the sense of [8]: the surface swept by the boundaries of the discs in Σ contained in the half sphere reaches the equator.

In the sequel we will apply this technique of filling to a sphere in $\partial\tilde{\Omega}$ where $\tilde{\Omega}$ is the universal covering of a pseudoconvex domain Ω which is strictly pseudoconvex where the sphere projects down. The reader can check that all the arguments of [5], [12] apply mutatis mutandis.

b) Geometric function theory. We will use the following facts concerning polynomial convexity (see [15] for this paragraph). Let K be a compact set in S^3 separating the sphere in finitely many components, then its polynomial hull \hat{K} divides B in the same number of components. Moreover by Rossi local maximum principle these components are pseudoconvex domains. We will also rely on the theorem by Alexander describing the polynomial hull of a curve of finite length (with finitely many components): it is a Riemann surface attached to the curve.

We move on to rational convexity. The rational hull $r(K)$ of a compact set K in \mathbb{C}^2 is geometrically defined as the set of points z such that any algebraic curve passing through z meets K . If $K \subset P$ where P is a rational polyhedron, then the algebraic curves can be replaced by analytic curves in P . The usual obstruction to rational convexity is the presence of a compact Riemann surface with boundary in K with the additional restriction that this boundary bounds in K . In our theorem (second situation) the holomorphic annulus or the pair of holomorphic discs will satisfy this condition and therefore be part of $r(T)$. As for the first situation we have the following

Lemma. *A fillable totally real torus in S^3 is rationally convex.*

Proof. Call T the torus and Θ its filling. We first prove that Θ is rationally convex. By Rossi local maximum principle and the Runge property of B it is enough to construct through any point near Θ in the ball B an analytic curve in B (smooth up to S^3) avoiding Θ . We produce them by stability of the filling of T (see [4] for a similar situation). Foliate a neighborhood of T in S^3 by tori, then the fillings of these tori

foliate a neighborhood of Θ in B . Therefore the corresponding holomorphic discs fill out this neighborhood and avoid T if they are not in Θ . At this stage $r(T) \subset \Theta$.

We now prove that $r(T) = T$. According to the first step Θ is a decreasing limit of rational polyhedrons. It is then enough to construct through any point z of $\Theta \setminus T$ an analytic curve in a neighborhood of Θ avoiding T . Take through z a real closed curve in $\Theta \setminus T$ transversal to the holomorphic discs, parametrized by the unit circle. Extend this parametrization as a smooth map f from a thin round annulus in such a way that $\bar{\partial}f$ vanishes to infinite order along the unit circle. By solving an adequate $\bar{\partial}$ -equation perturb now f into a holomorphic map. This map parametrizes a thin holomorphic annulus still passing through z and intersecting Θ near the initial curve, hence avoiding T .

Finally let us recall the analogue in terms of currents of the usual obstructions to polynomial or rational convexity [7]. Let K a compact set in \mathbb{C}^2 and U a positive 1,1-current with compact support. If $\text{supp}(dU) \subset K$ then $\text{supp}(U) \subset \hat{K}$. If moreover $dU = dV$ where V is a current supported by K , then $\text{supp}(U) \subset r(K)$.

c) Ahlfors currents. They are the local version of the currents built from an entire curve in complex hyperbolicity. In our context a current U is an *Ahlfors current* if $U = \lim_{a_n} \frac{[\Delta_n]}{a_n}$, where $[\Delta_n]$ are currents of integration over holomorphic discs Δ_n of area a_n contained in B whose boundary sits mainly in S^3 . Precisely, one has $\text{length}(\partial\Delta_n \setminus S^3) = o(a_n)$. Hence U is a positive 1,1-current with compact support such that $\text{supp}(dU) \subset S^3$. The following lemma (compare with [6]) will be important for the non rationally convex case.

Lemma. *Let U be an Ahlfors current whose support is an analytic curve in B . Then each irreducible component of this curve is a holomorphic disc or annulus.*

Proof. It relies on Ahlfors theory of covering surfaces [14] under the form of the following

Isoperimetric inequality. *Let E be a compact connected Riemann surface with boundary, of negative Euler characteristic. Then there is a constant c such that for any holomorphic disc $f: D \rightarrow E$ we have $\text{area}(f(D)) \leq c \text{ length}(f(\partial D) \setminus \partial E)$.*

Here area and length are computed by means of a given metric on E , taking into account multiplicities.

As in [6] we proceed by contradiction. Let C be a component of the analytic curve which is neither a disc nor an annulus. Then there exists a figure eight e in C such that any component of $C \setminus e$ meets ∂C . In particular e is polynomially convex [15]. We may suppose moreover that e avoids the singularities of C . Thickening e slightly in C we get a disc with two holes E . Identify now a polynomially convex neighborhood V of e to $E \times d$ where d is a small disc. Call π the projection of V

on E . Recall that the current U comes from a sequence of discs (Δ_n) . Shrinking d a bit we may suppose that $\text{length}(\Delta_n \cap (E \times \partial d)) = o(a_n)$. This uses the fact that $U|_V$ does not charge $V \setminus E$ and the coarea formula. Now, as V is polynomially convex, $\Delta_n \cap V$ consists in a union of discs δ_n by the maximum principle. By construction the boundaries of δ_n sit mainly in $\partial E \times d$. We infer that $\text{area}(\pi(\Delta_n \cap V)) = o(a_n)$ by applying the isoperimetric inequality to the maps $\pi: \delta_n \rightarrow E$ and summing up. This contradicts the fact that U charges E .

Remark. Suppose we have an annulus A among the components of the analytic curve. Then the discs Δ_n approximating U satisfy the following additional property: they cannot avoid (for large n) a fixed analytic curve C in B meeting A . Indeed if not we could work out the previous argument in the complement of C , replacing everywhere the polynomial convexity by the convexity with respect to the algebra \mathcal{M} of meromorphic functions in B with poles on C . We would find a \mathcal{M} -convex figure eight in the punctured annulus $A \setminus C$ and proceed as above to reach a contradiction.

We enter now the proof of the theorem.

2. The set up

Let T be an unknotted totally real torus T in S^3 . It divides S^3 into two solid tori ω_i (diffeomorphic to $S^1 \times D^2$) and its polynomial hull \hat{T} separates B into two pseudoconvex domains Ω_i containing ω_i in their closure (§1 b)).

Lemma. *For one of these domains the map $H_1(\omega_i, \mathbb{Z}) \rightarrow H_1(\Omega_i, \mathbb{Z})$ is injective.*

Proof. If not, let γ_i be a generator of $H_1(\omega_i, \mathbb{Z})$. Note first that γ_1 and γ_2 are linked in S^3 , and next that the linking number of two disjoint cycles in S^3 can be computed as the intersection number of the chains they bound in B . Now by assumption $n_i \gamma_i$ bounds a chain in Ω_i for some integer n_i . But Ω_1 and Ω_2 being disjoint this shows that $n_1 \gamma_1$ and $n_2 \gamma_2$ (hence γ_1 and γ_2) are not linked in S^3 . Contradiction.

Let us call simply Ω this good side and ω the corresponding solid torus. We push slightly T inside ω , creating a sequence of tori T_n approximating T .

Consider the universal covering $p: \tilde{\Omega} \rightarrow \Omega$. Because $\pi_1(\omega) \rightarrow \pi_1(\Omega)$ is injective, all the components of $p^{-1}(\omega)$ are diffeomorphic to $\mathbb{R} \times D^2$. Fix one of them and call it $\tilde{\omega}$. Then T_n lifts to a cylinder \tilde{T}_n (diffeomorphic to $\mathbb{R} \times S^1$) inside $\tilde{\omega}$. Let τ be the automorphism of $\tilde{\Omega}$ induced by the action of a generator of $\pi_1(\omega)$. It acts on $\tilde{\omega}$ as a translation on the factor \mathbb{R} and \tilde{T}_n is invariant under this action.

Construct the sphere S_n approximating the cylinder \tilde{T}_n as follows. Pick a disc D in ω (diffeomorphic to $* \times D^2$). Its boundary is a *meridian* of T . Deform D slightly in D_n with boundary in T_n . Choose a lift \tilde{D}_n of D_n in $\tilde{\omega}$. The curves $\tau^{\pm n}(\partial \tilde{D}_n)$

bound an annulus \tilde{A}_n in \tilde{T}_n . The sphere S_n is obtained by smoothing the sphere with corners $\tau^{-n}(\tilde{D}_n) \cup \tilde{A}_n \cup \tau^n(\tilde{D}_n)$. Note that the complex points of S_n can be made generic after a perturbation localized near the caps $\tau^{\pm n}(\tilde{D}_n)$. By construction S_n projects down to the interior of ω where Ω is strictly pseudoconvex. Hence the technics of §1 a) apply. Denote by Σ_n the filling of S_n in $\tilde{\Omega}$. Now the equator $\partial\tilde{D}_n$ divides S_n into two half spheres S_n^{\pm} . At least one of them, say S_n^- , has a partial filling. This means that we may single out a disc $\tilde{\Delta}_n$ of Σ_n touching the equator and whose boundary is entirely contained in S_n^- . Put $\Delta_n = p(\tilde{\Delta}_n)$.

The alternative reads as follows: either the area a_n of Δ_n remains bounded or not. In the former case we will verify by Gromov compactness theorem that T is fillable. This is the *rationally convex case*. In the latter we will consider the Ahlfors current $U = \lim \frac{[\Delta_n]}{a_n}$. By construction its support will be in the rational hull of T and, after a detailed analysis, we will detect holomorphic annuli or discs in it. This is the *non rationally convex case*.

In any case we need to control the boundary of Δ_n . We know by Hopf lemma that $T\partial\Delta_n$ is transversal on T_n to the characteristic line field. Actually we have more. Denote by ω_n the solid torus bounded by T_n in ω . Perturb the ball B in a new strictly convex domain B_n by bumping slightly ω_n out, keeping T_n still in ∂B_n . Note that by construction $\partial\Delta_n \subset B_n$, so $\Delta_n \subset B_n$ by the maximum principle. But tilting the boundary of the domain along T_n translates in rotating the characteristic line field on T_n . We infer that $T\partial\Delta_n$ avoids a full cone field on T_n bounded on one side by the original characteristic line field. As this can be done uniformly in n , we end up with $T\phi_n(\partial\Delta_n \cap T_n)$ avoiding a cone field on T . Here ϕ_n is a diffeomorphism between T_n and T close to identity. We may perturb slightly the characteristic line field of T to push it *inside* this cone field, still keeping its name. We summarize this discussion by saying that $\gamma_n = \phi_n(\partial\Delta_n \cap T_n)$ is *uniformly transversal* to the characteristic foliation \mathcal{C} of T . This actually holds for any disc of Σ_n .

It follows that the length l_n of γ_n is controlled by a_n . For this construct a 1-form β on T whose kernel is the characteristic line field and extend it to \mathbb{C}^2 . Then $l_n \lesssim |\int_{\gamma_n} \beta| \lesssim |\int_{\Delta_n} d\beta| + |\int_{\partial\Delta_n \cap D_n} \beta|$ by the uniform transversality and Stokes theorem. Here \lesssim stands for an estimate up to a multiplicative constant. The first integral on the right is controlled by a_n . The second one is bounded. Indeed note that $\partial\tilde{\Delta}_n$ bounds a disc \tilde{V}_n in S_n^- . Call V_n its projection downstairs. Then $|\int_{\partial\Delta_n \cap D_n} \beta| \leq \int_{\partial D_n \cap V_n} |\beta| + \int_{D_n \cap V_n} |d\beta| \lesssim \text{length}(\partial D) + \text{area}(D)$ by Stokes theorem and the closeness of D_n and \tilde{D} . We end up with an estimate of the form $l_n \leq C(1 + a_n)$.

Conversely a_n is controlled by l_n in the same way. Indeed recall that $a_n = \int_{\Delta_n} \omega$ where ω is the standard Kähler form of \mathbb{C}^2 . Write λ for a primitive of ω . Then $a_n = \int_{\partial\Delta_n} \lambda \lesssim l_n + \int_{\partial\Delta_n \cap D_n} \lambda$ by Stokes theorem and, as before, the last integral is bounded.

The rationally convex case

In this case a_n remains bounded, and so is l_n .

We first check that T_n is fillable. By assumption $\partial\tilde{\Delta}_n$ remains at bounded distance of the equator of S_n . This means that $\tilde{\Delta}_n$ is attached to both S_n and $\tau^{-1}(S_n)$, hence belongs to their fillings (§1 a)). In other words both $\tilde{\Delta}_n$ and $\tau(\tilde{\Delta}_n)$ are part of Σ_n . The discs of Σ_n interpolating between them project down to the desired filling Θ_n of T_n . Note that all the discs Δ'_n of Θ_n have bounded area. Indeed we have $\int_{\Delta'_n} \omega \leq \int_{\Delta_n} \omega + \int_{T_n} |\omega| \lesssim a_n + \text{area}(T)$ by Stokes theorem.

We want now to prove that T is fillable as well. We rely on Gromov compactness theorem [10] (see also [11]). In our context it reads as follows: given a disc Δ'_n in Θ_n , then the sequence (Δ'_n) converges (after extracting a subsequence) toward a finite bunch (with multiplicities) of holomorphic discs Δ' attached to T . These discs do not present self-intersections or mutual intersections in the interior of B . This relies on two facts: intersections of distinct holomorphic curves persist under local deformation, and the convergence does not show accidents inside the ball. Actually an accident means an annulus component of Δ'_n in a fixed small ball converging toward a pair of two discs (its modulus blows up). But all such local components are discs by the maximum principle. Moreover the discs Δ' , if *simple*, are embedded inside B by a knot-theoretic argument [5]. We want to build the filling of T out of these limit discs. The problem is to exhibit sufficiently many such discs, embedded and disjoint in the closed ball. The difficulty takes place at their boundaries. We focus on them.

For a sequence (Δ'_n) as above call $\Gamma' = \cup \partial\Delta'$ the boundary of its limit. By Hopf lemma it is a finite union of immersed curves (with multiplicities). Denote by $\text{Sing}(\Gamma')$ the set of multiple points of Γ' , i.e. its geometric singularities and its multiple components. Similarly put Γ for the boundary of the limit of the original sequence (Δ_n) (after the same extraction). Our first observation is that $\text{Sing}(\Gamma') \subset \Gamma$. Indeed locally at least two strands of $\partial\Delta'_n$ converge at a given point of $\text{Sing}(\Gamma')$: if α is a short piece of the characteristic leaf through this point, it meets $\phi_n(\partial\Delta'_n)$ at least twice. Here again ϕ_n is a diffeomorphism between T_n and T close to identity. In other words α runs from one boundary to the other in the cylinder obtained from T by cutting out $\phi_n(\partial\Delta'_n)$. As $\phi_n(\partial\Delta_n)$ is parallel to these boundaries it always intersects α , and so does Γ . Shrinking α to the initial point concludes.

In particular at each point $q \in \Gamma' \setminus \Gamma$ the convergence of (Δ'_n) is *good*: there exists a unique simple disc Δ' through q in the limit such that Δ'_n converges toward Δ' near q . Our second observation is that this disc does not really depend on (Δ'_n) . If we consider another similar sequence (Δ''_n) converging after the same extraction, such that $q \in (\Gamma' \cap \Gamma'') \setminus \Gamma$, then $\Delta' = \Delta''$. Indeed if not, Δ' and Δ'' would be distinct. But intersections of distinct holomorphic discs attached to a totally real surface and on the same side of a strictly pseudoconvex boundary persist under local deformation. This can be seen by reflecting the discs through the surface to get (pseudo)holomorphic curves in a neighborhood of q and using the positivity of their intersections [16].

Therefore Δ'_n and Δ''_n would still intersect, contradicting their being part of the same filling Θ_n .

According to our previous discussion we focus on $T^* = T \setminus \Gamma$ where all the convergences are good. Pick a countable set Q dense in T^* . Denote by $\Delta_{n,q}$ the disc of Θ_n passing through $\phi_n^{-1}(q)$ for $q \in Q$. By extracting once more we may suppose that all sequences $(\Delta_{n,q})$ converge in Gromov sense. Hence there exists a unique simple disc Δ_q through q in $\lim_{n \rightarrow \infty} \Delta_{n,q}$. We want to extend this construction to T^* .

Pick a point p in T^* . Then the component through p of $\lim_{q \rightarrow p} \Delta_q$ (in Gromov sense) is well defined. Indeed any component through p in $\lim_{q \rightarrow p} \Delta_q$ appears also as a limit of discs in Θ_n : consider discs of the form Δ_{n_k, q_k} for some sequence q_k going to p and n_k rapidly growing. Therefore by the observations above this component is unique and does not depend on any choice. We get a distribution of holomorphic discs Δ_p ($p \in T^*$) whose boundaries are embedded and disjoint (if distinct) in T^* . It turns out that the same holds in the whole T .

Lemma. *The curves $\partial\Delta_p$ are embedded and disjoint (if distinct).*

Proof. We proceed by contradiction. Pick an intersection point s (necessarily in Γ) of two different local branches γ', γ'' of such curves. Note that $\gamma' \cup \gamma''$ cuts out four components in T near s , two of which avoiding the characteristic leaf through s . Call C the union of these two components and put $C^* = C \setminus \Gamma$. Now for all p in C^* the curve $\partial\Delta_p$ is canalized by γ' and γ'' through s . Thus we get a whole family of holomorphic discs Δ_p attached to T with a common point. On the other hand by the maximum principle these discs sit in \hat{T} and even in $\partial\hat{T}$ as limits of discs in $\Theta_n \subset \Omega$. This will be the contradiction.

Let us make this precise. Recall first that we may associate to an immersed holomorphic disc Δ attached to T an even integer, its *Maslov index* $\mu(\Delta)$ (see [3] for background). This index is related to the dimension of the manifold of the holomorphic discs close to Δ and attached to T . If $\mu(\Delta) \leq 0$ this manifold is of dimension 0: Δ does not have any deformation attached to T . If $\mu(\Delta) > 0$ it is of positive dimension $\mu(\Delta) - 1$. Moreover if $\mu(\Delta) = 2$ we get a small 1-parameter family of nearby *locally disjoint* discs attached to T . In particular they cannot pass through a common point. On the other hand if $\mu(\Delta) > 2$ the (at least) 3-parameter family of nearby discs attached to T fills out a whole neighborhood of Δ in B . This forbids Δ to be in $\partial\hat{T}$. To conclude it remains to exhibit a genuine deformation among the family Δ_p passing through s .

What we know already is that Δ_p is the unique component through p in $\lim_{q \rightarrow p} \Delta_q$ for p in C^* . We would like to really have $\lim_{q \rightarrow p} \Delta_q = \Delta_p$. This will be at least the case for Δ_p big enough. For this recall that any holomorphic disc attached to T cannot be too small. This relies for instance on the existence of a basis of strictly pseudoconvex neighborhoods of T . Then according to Lelong theorem the area of

such a disc is bounded from below by some positive constant, say 2ϵ . Now pick p in C^* such that $\text{area}(\Delta_p) \geq \sup_{C^*} \text{area}(\Delta_q) - \epsilon$. As the area is preserved under the convergence in Gromov sense, we infer that there is no other component but Δ_p in $\lim_{q \rightarrow p} \Delta_q$. This concludes.

At this stage we do have a whole smooth family of disjoint embedded holomorphic discs attached to T whose boundaries sweep out at least T^* . To achieve the filling it remains to close this family up on Γ . This goes along the same lines as before. The main point is that if p is in Γ then $\lim_{q \rightarrow p} \Delta_q$ does not present singularities. If it did, as in the first observation above, all the discs Δ_q would pass through this singular point, contradicting the lemma. We leave the details to the reader.

The non rationally convex case

In this case a_n blows up. We want to prove that there exists a Riemann surface (holomorphic annulus or pair of holomorphic discs) attached to T and part of its rational hull. We look at the limit of Δ_n in terms of currents. Consider $\frac{[\Delta_n]}{a_n}$ the normalized current of integration on Δ_n . We get a sequence of positive currents of mass 1 supported in the unit ball. Up to extracting it converges toward an Ahlfors current U . Recall that $\partial\Delta_n = \partial V_n$ where V_n is the projection of the disc \tilde{V}_n bounded by $\partial\tilde{\Delta}_n$ in S_n . Note that a_n is comparable to l_n (§2) and so to the maximal number of sheets of \tilde{V}_n over T_n . Hence $\frac{[V_n]}{a_n}$ converges toward a current V supported on T such that $dV = dU$. Therefore $\text{supp}(U) \subset r(T)$ (§1 b)). We have $dU = \lim \frac{[\gamma_n]}{a_n}$ where $\gamma_n = \phi_n(\partial\Delta_n \cap T_n)$ (§2). As a_n blows up we may even neglect parts of γ_n of bounded length in this limit. To exhibit Riemann surfaces in $r(T)$ we further investigate the current U . We focus first on its boundary.

a) Describing dU . We will prove an integral formula of the form $dU = \int_{\mathcal{G}} [\gamma] d\mu(\gamma)$. Here \mathcal{G} is a compact space of Lipschitz curves in T and μ a positive measure on it, supported on closed curves.

This requires an extra discussion of the characteristic foliation \mathcal{C} . By Denjoy theorem [9] any smooth foliation on T can be perturbed in order to get only a finite number of attracting or repulsive cycles (closed leaves). We may suppose that this holds true for \mathcal{C} as we already perturbed it (§2).

Call c such a characteristic cycle. Observe that the lifts of $\phi_n^{-1}(c)$ cannot be closed in \tilde{T}_n . If it were the case such a lift would divide S_n out into two half spheres, one of which partially fillable. We would thus get a contact between this lift and the boundary of a disc of the filling Σ_n , contradicting the uniform transversality. Hence $p^{-1}(\phi_n^{-1}(c))$ consists in finitely many periodic curves invariant by a power τ^q of τ .

It follows that the number of intersection points between γ_n and c is bounded. Indeed each lift of $\phi_n^{-1}(c)$ cuts at most once $\partial\tilde{\Delta}_n$ by transversality and because $\partial\tilde{\Delta}_n$

separates S_n . Consider now thin tubes along the cycles in T . They divide T in a finite number of annuli. By uniform transversality γ_n cuts the tubes in a bounded number of short arcs. We may neglect them for the computation of dU . Hence the relevant part of γ_n consists in a bounded number of long arcs contained in the annuli.

The crucial observation is that these arcs are *embedded* (up to splitting them into two pieces). To prove this we further analyse the situation upstairs. Call B_n the ball bounded by the sphere S_n in $\tilde{\omega}$ and Ω_n the pseudoconvex domain bounded by $B_n \cup \Sigma_n$ in $\tilde{\Omega}$. Then $\tau^q(S_n^-) \subset \Omega_n$ for large n . This is where the choice of a half sphere enters. Hence its partial filling, as part of its envelope of holomorphy (§1 a)), must also be contained in Ω_n . To be fully correct this argument requires to push slightly S_n off B_n in a new sphere $S'_n \subset \tilde{\omega}$, verify that the partial filling of $\tau^q(S'_n^-)$ is contained in the corresponding pseudoconvex domain Ω'_n and deform back S'_n to S_n . In particular we get that $\tau^q(\tilde{\Delta}_n) \subset \Omega_n$. Hence $\tau^q(\tilde{\Delta}_n)$ remains always on the same side of Σ_n , meaning that $\tau^q(\partial\tilde{\Delta}_n)$ crosses $\partial\tilde{\Delta}_n$ always in the same direction (say entering \tilde{V}_n). Look now at a given lift of $\phi_n^{-1}(A)$ in \tilde{T}_n where A is one of the aforementioned annuli. This is a strip invariant by τ^q . It can be parametrized by $\mathbb{R} \times [0, 1]$ via a diffeomorphism sending the vertical foliation to the characteristic one, τ^q corresponding to the translation by 1. By transversality any component of $\partial\tilde{\Delta}_n$ in the strip is a graph (via the diffeomorphism) with, say, \tilde{V}_n above it. Thus the component and its image by τ^q intersect at most once as the latter crosses the former always bottom up. This allows us to cut the component into two pieces, each of them disjoint from its image by τ^q . Therefore these pieces project down to embedded arcs.

According to this discussion dU is a finite sum of currents of the form $\lim \frac{[\alpha_n]}{a_n}$, where α_n is an embedded arc sitting in an annulus A . We are now in position to prove the integral formula for each such limit. Via the parametrization of the corresponding strip and thanks to the uniform transversality, α_n splits up into a union of graphs of functions from $[0, 1]$ to $[0, 1]$ which are uniformly Lipschitz. Denote by \mathcal{G} the compact space of graphs γ of functions $g: [0, 1] \rightarrow [0, 1]$ such that $\text{Lip}(g) \leq C$ (for some large C). We have $\frac{[\alpha_n]}{a_n} = \int_{\mathcal{G}} [\gamma] d\mu_n(\gamma)$ where μ_n is a positive measure with finite support and bounded mass on \mathcal{G} . Up to extracting μ_n converges toward a positive measure μ on \mathcal{G} . We infer that $\lim \frac{[\alpha_n]}{a_n} = \int_{\mathcal{G}} [\gamma] d\mu(\gamma)$. Moreover the support of μ consists in closed curves (graphs of functions g such that $g(0) = g(1)$). Indeed if the graph of g is in $\text{supp}(\mu)$ then it is certainly the limit of at least two successive graphs (of say g_n and h_n) of α_n (a_n blows up). As $g_n(1) = h_n(0)$ we get $g(0) = g(1)$ in the limit.

b) Describing U . We will prove now an integral formula of the form $U = \int_{\mathcal{P}} W dv(W)$. Here \mathcal{P} is the compact space of positive currents of mass 1 supported in the unit ball and v is a probability measure on it. The point is that $\text{supp}(v)$ consists only in normalized currents of integration on holomorphic discs or

annuli attached to T (or finite sums of them). This formula comes from a division process.

We show first that U can be split up into a sum of four positive currents W of mass at most $\frac{1}{2}$. These currents will be proportional to Ahlfors currents limit of pieces of Δ_n . Precisely $W = \lim \frac{[\delta_n]}{a_n}$ where $\delta_n \subset \Delta_n$, $\text{area}(\delta_n) \leq \frac{a_n}{2}$ and $\text{length}(\partial\delta_n \setminus \partial\Delta_n) = o(a_n)$. In addition we want $\partial\delta_n \cap \partial\Delta_n$ connected.

For this, parametrize Δ_n by the unit disc via a holomorphic map $f_n: D \rightarrow B$ such that the images by f_n of the four half discs cut out in D by \mathbb{R} or $i\mathbb{R}$ have the same area $\frac{a_n}{2}$. Denote by X the cross $(\mathbb{R} \cap D) \cup (i\mathbb{R} \cap D)$. According to the next lemma, we may pick a generic angle θ close to $\frac{\pi}{4}$ such that $\text{length}(f_n(e^{i\theta}X)) = o(a_n)$. The rotated cross $e^{i\theta}X$ divides D out in four quarter discs d . Put $\delta_n = f_n(d)$ and $W = \lim \frac{[\delta_n]}{a_n}$. The currents W have all the desired properties. Here is the precise statement we used.

Lemma. *Let $f_n: D \rightarrow B$ be a sequence of holomorphic discs (piecewise) smooth up to ∂D . Put $a_n = \text{area}(f_n(D))$, $l_n(\theta) = \text{length}(f_n([0, e^{i\theta}]))$ and suppose that a_n blows up. Then $l_n(\theta) = o(a_n)$ for almost all θ (up to extracting a subsequence).*

Proof. We have $l_n(\theta) = \int_0^1 \|f'_n(re^{i\theta})\| dr \leq l + \int_{1/2}^1 \|f'_n(re^{i\theta})\| dr$ for some constant l as $\|f'_n\|$ is uniformly bounded in the disc of radius $\frac{1}{2}$. On the other hand, let $a_n(\theta)$ be the area of the image by f_n of the sector between $[0, 1]$ and $[0, e^{i\theta}]$. Then $\frac{da_n}{d\theta}(\theta) = \int_0^1 \|f'_n(re^{i\theta})\|^2 r dr$. By Cauchy–Schwarz inequality $(l_n(\theta))^2 \leq 2l^2 + 2\ln(2)\frac{da_n}{d\theta}(\theta)$. Integrating, we get $\int_0^{2\pi} (l_n(\theta))^2 d\theta \leq 4\pi l^2 + 2\ln(2)a_n$, so $\lim \int_0^{2\pi} \left(\frac{l_n(\theta)}{a_n}\right)^2 d\theta = 0$. By Fatou's lemma $\int_0^{2\pi} \liminf \left(\frac{l_n(\theta)}{a_n}\right)^2 d\theta = 0$, which concludes.

Iterating this process we may write U as a sum of 4^k positive currents of mass at most 2^{-k} proportional to Ahlfors currents coming from (Δ_n) . Hence $U = \int_{\mathcal{P}} W dv_k(W)$ where v_k is a probability measure supported on these Ahlfors currents. By compactness of \mathcal{P} we may suppose that (v_k) converges toward a probability measure ν on \mathcal{P} and we get our integral formula $U = \int_{\mathcal{P}} W d\nu(W)$. Take now a current W in the support of ν . By construction W is an Ahlfors current as a limit of Ahlfors currents. We will see below that dW is supported on a curve $\gamma \subset T$ of finite length (with finitely many components). Hence $\text{supp}(W) \subset \hat{\gamma}$ which by Alexander theorem (§1 b)) is a Riemann surface. By §1 c) we conclude that W is actually supported in a finite union of holomorphic discs or annuli attached to T .

Let us describe dW . By construction $W = \lim \frac{W_k}{\text{mass}(W_k)}$ where $W_k = \lim \frac{[\delta_{n,k}]}{a_n}$ with $\delta_{n,k} \subset \Delta_n$, $\text{area}(\delta_{n,k}) \leq 2^{-k}a_n$, $\text{length}(\partial\delta_{n,k} \setminus \partial\Delta_n) = o(a_n)$ and $\partial\delta_{n,k} \cap \partial\Delta_n$ connected. We use the notations of the previous paragraph. Recall that we had singled out an annulus A outside thin tubes of the characteristic cycles in T and an arc α_n of γ_n embedded in A . So $\partial\delta_{n,k} \cap \partial\Delta_n$ gives rise to a subarc $\alpha_{n,k}$ of α_n . We check

now that $\lim \frac{[\alpha_{n,k}]}{a_n}$ is supported in a set converging to a curve of finite length (with at most two components). This will conclude as dW is a finite sum of such limits.

Indeed $\alpha_{n,k}$ is built out of graphs in \mathcal{G} and we have $\lim \frac{[\alpha_{n,k}]}{a_n} = \int_{\mathcal{G}} [\gamma] d\mu_k(\gamma)$ for a positive measure $\mu_k \leq \mu$ on \mathcal{G} . Note that we have a partial order on \mathcal{G} given by $\alpha \leq \beta$ if the corresponding functions satisfy $a \leq b$. We may speak of intervals $[\alpha, \beta]$ or $] \alpha, \beta[= [\alpha, \beta] \setminus \{\alpha, \beta\}$. This order is total on the graphs appearing in $\alpha_{n,k}$ (α_n is embedded). Denote by $\eta_{n,k}$ and $\lambda_{n,k}$ the lowest and the highest of these graphs. By compactness of \mathcal{G} we may suppose that $\eta_{n,k}, \lambda_{n,k}$ converge to η_k, λ_k , and that η_k, λ_k converge to η, λ . By construction $\text{supp}(\mu_k) \subset [\eta_k, \lambda_k]$ and, moreover, $\mu_k = \mu$ on $] \eta_k, \lambda_k[$. As the mass of μ_k goes to 0, it follows that μ does not charge $] \eta, \lambda[$. Hence $\text{supp}(\mu_k) \subset [\eta_k, \lambda_k] \setminus] \eta, \lambda[$ which goes to $\{\eta, \lambda\}$. This concludes.

c) End of the argument. At this stage we do have compact Riemann surfaces (holomorphic discs or annuli) attached to T and contained in $r(T)$. We want more. We are looking for a compact Riemann surface C (holomorphic annulus or a pair of holomorphic discs) such that ∂C bounds in T . Here is how we proceed.

Choose a common orientation of the characteristic cycles. Note that the boundaries of our Riemann surfaces are parallel to these cycles. They also inherit a natural orientation from the Riemann surface. We speak of a *positive* boundary if the two orientations agree, or *negative* if not. Call positive (negative) an annulus or a disc with only positive (negative) boundaries, and *opposite* an annulus or a pair of discs with opposite boundaries. We are looking for an opposite annulus or a pair of opposite discs among our Riemann surfaces. Suppose we do not have any.

Recall that dU bounds in T . This implies that our Riemann surfaces cannot be all positive, or all negative. We have three possibilities left: either the presence among them of a positive annulus and a negative annulus, or of a positive annulus and a negative disc, or the converse. By symmetry we may suppose that we have a positive annulus A^+ and a negative Riemann surface (annulus or disc) C^- . Observe now that two disjoint closed curves in T parallel to the characteristic cycles are necessarily linked in S^3 . This can be checked for any pair of disjoint curves in the standard torus, as soon as they are not meridians (i.e. do not bound a disc in the complement of the standard torus).

Hence the boundaries of A^+ and C^- are linked. This implies that A^+ and C^- intersect inside the unit ball. But by construction A^+ is contained in the support of an Ahlfors current coming from (Δ_n) . As A^+ intersects C^- , before the limit Δ_n would have to intersect C^- (§1 c)). This is impossible as $C^- \subset \hat{T}$ and $\Delta_n \subset \Omega$.

References

- [1] H. Alexander, Gromov's method and Bennequin's problem. *Invent. Math.* **125** (1996), 135–148. [Zbl 0853.32003](#) [MR 1389963](#)

- [2] H. Alexander, Disks with boundaries in totally real and Lagrangian manifolds. *Duke Math. J.* **100** (1999), 131–138. [Zbl 0953.32026](#) [MR 1714757](#)
- [3] M. Audin and J. Lafontaine (ed.), *Holomorphic curves in symplectic geometry* Prog. Math. 117, Birkhäuser, Basel 1994. [Zbl 0802.53001](#) [MR 1274924](#)
- [4] E. Bedford, Stability of the polynomial hull of T^2 . *Ann. Scuola Norm. Sup. Pisa* **8** (1981), 311–315 [Zbl 0472.32012](#) [MR 0623939](#)
- [5] E. Bedford and W. Klingenberg, On the envelope of holomorphy of a 2-sphere in \mathbb{C}^2 . *J. Amer. Math. Soc.* **4** (1991), 623–646. [Zbl 0736.32009](#) [MR 1094437](#)
- [6] J. Duval, Singularités des courants d’Ahlfors. *Ann. Sci. Ecole Norm. Sup.* **39** (2006), 527–533. [Zbl 1243.32012](#) [MR 2265678](#)
- [7] J. Duval and N. Sibony, Polynomial convexity, rational convexity, and currents. *Duke Math. J.* **79** (1995), 487–513. [Zbl 0838.32006](#) [MR 1344768](#)
- [8] Y. Eliashberg, Filling by holomorphic discs and its applications. In *Geometry of low dimensional manifolds*, London Math. Soc. Lecture Note Ser. 151, Cambridge University Press, Cambridge 1990, 45–67. [Zbl 0731.53036](#) [MR 1171908](#)
- [9] C. Godbillon, *Dynamical systems on surfaces*. Universitext, Springer, Berlin 1983. [Zbl 0502.58002](#) [MR 0681119](#)
- [10] M. Gromov, Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.* **82** (1985), 307–347. [Zbl 0592.53025](#) [MR 0809718](#)
- [11] S. Ivashkovich and V. Shevchishin, Reflection principle and J -complex curves with boundary on totally real immersions. *Commun. Contemp. Math.* **14** (2002), 65–106. [Zbl 1025.32024](#) [MR 1890078](#)
- [12] N. G. Kruzhilin, Two-dimensional spheres on the boundaries of pseudoconvex domains in \mathbb{C}^2 . *Math. USSR-Izv.* **39** (1992), 1151–1187. [Zbl 0778.32003](#) [MR 1152210](#)
- [13] N. G. Kruzhilin, Holomorphic disks with boundaries in totally real tori in \mathbb{C}^2 . *Math. Notes* **56** (1994), 1244–1248. [Zbl 0842.32011](#) [MR 1330599](#)
- [14] R. Nevanlinna, *Analytic functions*. Grundlehren Math. Wiss. 162, Springer, Berlin 1970. [Zbl 0199.12501](#) [MR 0279280](#)
- [15] E. L. Stout, *Polynomial convexity*. Prog. Math. 261, Birkhäuser, Boston 2007. [Zbl 1124.32007](#) [MR 2305474](#)
- [16] R. Ye, Filling by holomorphic curves in symplectic 4-manifolds. *Trans. Amer. Math. Soc.* **350** (1998), 213–250. [Zbl 0936.53047](#) [MR 1422913](#)

Received November 18, 2011

Julien Duval, Laboratoire de Mathématiques, Université Paris-Sud, 91405 Orsay cedex, France

E-mail: julien.duval@math.u-psud.fr

Damien Gayet, Institut Camille Jordan, Université Claude Bernard, 69622 Villeurbanne cedex, France

E-mail: gayet@math.univ-lyon1.fr

On the values of G -functions

Stéphane Fischler and Tanguy Rivoal

À la mémoire de Philippe Flajolet

Abstract. In this paper we study the set \mathbf{G} of values at algebraic points of analytic continuations of G -functions (in the sense of Siegel). This subring of \mathbb{C} contains values of elliptic integrals, multiple zeta values, and values at algebraic points of generalized hypergeometric functions ${}_pF_q$ with rational coefficients. Its group of units contains non-zero algebraic numbers, π , $\Gamma(a/b)^b$ and $B(x, y)$ (with $a, b \in \mathbb{Z}$ such that $a/b \notin \mathbb{Z}$, and $x, y \in \mathbb{Q}$ such that $B(x, y)$ exists and is non-zero). We prove that for any $\xi \in \mathbf{G}$, both $\operatorname{Re} \xi$ and $\operatorname{Im} \xi$ can be written as $f(1)$, where f is a G -function with rational coefficients of which the radius of convergence can be made arbitrarily large. As an application, we prove that quotients of elements of $\mathbf{G} \cap \mathbb{R}$ are exactly the numbers which can be written as limits of sequences a_n/b_n , where $\sum_{n=0}^{\infty} a_n z^n$ and $\sum_{n=0}^{\infty} b_n z^n$ are G -functions with rational coefficients. This result provides a general setting for irrationality proofs in the style of Apéry for $\zeta(3)$, and gives answers to questions asked by T. Rivoal in “Approximations rationnelles des valeurs de la fonction Gamma aux rationnels : le cas des puissances”, Acta Arith. 142 (2010), no. 4, 347–365.

Mathematics Subject Classification (2010). 11J91, 33E30, 34M35, 34M40.

Keywords. G -function, period, rational approximation, linear differential equation, singularity analysis.

1. Introduction

The purpose of this text is to study the set of values of G -functions at algebraic numbers. Let us recall the following definition, which essentially goes back to Siegel [30].

Definition 1. A G -function f is a formal power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ such that the coefficients a_n are algebraic numbers and there exists $C > 0$ such that:

- (i) the maximum of the moduli of the conjugates of a_n is $\leq C^{n+1}$.
- (ii) there exists a sequence of integers d_n , with $|d_n| \leq C^{n+1}$, such that $d_n a_m$ is an algebraic integer for all $m \leq n$.

- (iii) $f(z)$ satisfies a homogeneous linear differential equation with coefficients in $\overline{\mathbb{Q}}(z)$.¹

Throughout this paper we fix an embedding of $\overline{\mathbb{Q}}$ into \mathbb{C} ; all algebraic numbers and all convergent series are considered in \mathbb{C} .

G -functions occur frequently in analysis, number theory, geometry and physics: for example, algebraic functions over $\overline{\mathbb{Q}}(z)$ which are holomorphic at 0, polylogarithms, Gauss' hypergeometric function with rational parameters, are G -functions. The exponential function is not a G -function but an E -function (that is, it satisfies the requirements of Definition 1 if a_n is replaced with $a_n/n!$ in the expansion of $f(z)$).

In Definition 1, condition (i) ensures that any non-polynomial G -function has finite non-zero radius of convergence at $z = 0$. Condition (iii) implies that in fact the coefficients a_n , $n \geq 0$, all belong to a same number field. Classical references on G -functions are the books [1] and [17].

Siegel's goal was to find conditions ensuring that E and G -functions take irrational or transcendental values at algebraic points: the picture is very well understood for E -functions but largely unknown for G -functions. The main tool to study the nature of values of G -functions is inexplicit Padé-type approximation (see [3], [12], [14], [22]). In an explicit form, Padé approximation is also behind Apéry's celebrated proof [7] of the irrationality of $\zeta(3)$, and similar results in specific cases (see for instance [9], [19]).

In this paper, we study the following set.

Definition 2. Let \mathbf{G} denote the set of all values $f(\alpha)$, where f is a G -function and $\alpha \in \overline{\mathbb{Q}}$. More precisely, all values at α of analytic continuations of f are considered, as soon as they are finite.

This subset of \mathbb{C} is a subring (this can be seen as a consequence of Theorem 1 below). It contains $\overline{\mathbb{Q}}$, and also (see §2.2 for proofs) multiple zeta values, elliptic integrals, and values at algebraic points of generalized hypergeometric functions ${}_pF_q$ with rational coefficients. André proved in [1], p. 123, that the units of the ring of G -functions are exactly the algebraic functions which are holomorphic and don't vanish at the origin. The description of the units of \mathbf{G} is an interesting open problem whose solution is not as simple as for functions, for we show in §2.2 that the group of units of \mathbf{G} contains not only the non-zero algebraic numbers but also π , the values of the Gamma function $\Gamma(a/b)^b$ and that of Euler's Beta function $B(x, y)$ (with $a, b \in \mathbb{Z}$ such that $a/b \notin \mathbb{Z}$, and $x, y \in \mathbb{Q}$ such that $B(x, y)$ exists and is non-zero). On the other hand, there is no explicit interesting number for which we

¹All differential equations considered in this text are homogeneous and consequently we will no longer mention the term "homogeneous".

are able to prove that it is not in \mathbf{G} ;² it is likely that e , Euler's constant γ , $\Gamma(a/b)$ (with a, b integers such that $a/b \notin \mathbb{Z}$) or Liouville numbers do not belong to \mathbf{G} .

A conjecture of Bombieri and Dwork predicts a strong relationship between differential equations satisfied by G -functions and Picard–Fuchs equations satisfied by periods of families of algebraic varieties defined over $\overline{\mathbb{Q}}$. See the precise formulation given by André in [1], p. 7, who proved half of the conjecture in [1], pp. 110–111. Christol [13] also conjectured that globally bounded G -functions are diagonals of rational functions, which are known to satisfy Picard–Fuchs equations. This raises the question of a connection between the set \mathbf{G} and the set \mathcal{P} of periods considered by Kontsevich and Zagier [26]; all elements of \mathcal{P} we have thought of belong also to \mathbf{G} . However $1/\pi$ is conjectured not to belong to \mathcal{P} , so that \mathbf{G} is presumably distinct from \mathcal{P} . However, a natural problem is the determination of the link between \mathbf{G} and $\mathcal{P}[1/\pi]$ (see the discussion at the end of § 2.2).

Our main result is the following.

Theorem 1. *A complex number ξ belongs to \mathbf{G} if, and only if, its real and imaginary parts can be written as $f(1)$, where f is a G -function with rational coefficients of which the radius of convergence can be made arbitrarily large.*

One of the consequences of this theorem is that the set of values of G -functions $\sum_{n=0}^{\infty} a_n z^n$ with $a_n \in \mathbb{Q}$ at points $z \in \mathbb{Q}$ inside the disk of convergence (respectively at points where this series is absolutely convergent, respectively convergent) is equal to $\mathbf{G} \cap \mathbb{R}$.

The main tool in the proof of Theorem 1 is André–Chudnovski–Katz's theorem (stated as Theorem 6 in §4.1 below), which provides for any G -function f and any $\zeta \in \overline{\mathbb{Q}}$ a local basis (g_1, \dots, g_μ) of solutions around ζ of a minimal differential equation satisfied by f . Expanding an analytic continuation of f in this basis yields connection constants $\varpi_1, \dots, \varpi_\mu \in \mathbb{C}$ such that $f(z) = \sum_{j=1}^{\mu} \varpi_j g_j(z)$. As a step towards Theorem 1, we prove the following result which is of independent interest:

Theorem 2. *The connection constants $\varpi_1, \dots, \varpi_\mu$ belong to \mathbf{G} .*

We would like to emphasize that analytic continuation (and its properties encompassed in André–Chudnovski–Katz's theorem) is the main tool in our approach. As the referee pointed out to us, it would be interesting to find a connection with other methods used in similar contexts, including Dèbes–Zannier's [15] or Euler's for accelerating convergent series; however we did not find any. For instance, Euler's binomial transform $\sum_{n \geq 0} (-1)^n a_n = \sum_{n \geq 0} \left(\sum_{k=0}^n (-1)^k \binom{n}{k} a_k \right) 2^{-n-1}$ is involutive and therefore it cannot be used to obtain series with arbitrarily large radius of convergence.

²Since the set \mathbf{G} is countable, there are complex numbers outside \mathbf{G} but the real difficulty is to exhibit such a number by an effective process leading to an analytic expression like a series or an integral for example.

As an application of Theorem 1, we answer questions asked in [28], p. 351, where the second author introduced the notion of rational G -approximations to a real number. This corresponds to assertion (ii) in the next result, which provides a characterization of numbers admitting rational G -approximations.

Given a subring \mathbb{A} of \mathbb{C} , we denote by $\text{Frac}(\mathbb{A})$ the field of fractions of \mathbb{A} , namely the subfield of \mathbb{C} consisting in all elements ξ/ξ' with $\xi, \xi' \in \mathbb{A}$, $\xi' \neq 0$.

Theorem 3. *Let $\xi \in \mathbb{R}^*$. The following statements are equivalent:*

- (i) *We have $\xi \in \text{Frac}(\mathbf{G}) \cap \mathbb{R} = \text{Frac}(\mathbf{G} \cap \mathbb{R})$.*
- (ii) *There exist two sequences $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ of rational numbers such that the series $\sum_{n=0}^{\infty} a_n z^n$ and $\sum_{n=0}^{\infty} b_n z^n$ are G -functions, $b_n \neq 0$ for any n large enough and $\lim_{n \rightarrow +\infty} a_n/b_n = \xi$.*
- (iii) *For any $R \geq 1$ there exist two G -functions $A(z) = \sum_{n=0}^{\infty} a_n z^n$ and $B(z) = \sum_{n=0}^{\infty} b_n z^n$, with rational coefficients and radius of convergence = 1, such that $A(z) - \xi B(z)$ has radius of convergence $> R$.*

Remark. When $\xi \in \mathbf{G}$, we can take $b_n = 1$ in (ii). However, it is not clear to us if this is also the case for other elements $\xi \in \text{Frac}(\mathbf{G})$, in particular because it is doubtful that \mathbf{G} itself is a field.

Apéry has proved [7] that $\zeta(3) \notin \mathbb{Q}$ by constructing sequences $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ essentially as in (iii), such that $b_n \in \mathbb{Z}$ and $\text{lcm}(1, 2, \dots, n)^3 a_n \in \mathbb{Z}$. Since $\zeta(3) = \text{Li}_3(1)$ (where the polylogarithms defined by $\text{Li}_s(z) = \sum_{n=1}^{\infty} \frac{1}{n^s} z^n$, $s \geq 1$, are G -functions), we have $\zeta(3) \in \mathbf{G}$. Theorem 3 provides a general setting for such irrationality proofs and one may wonder if, given a real irrational number $\xi \in \text{Frac}(\mathbf{G})$, there exists a proof *à la Apéry* that ξ is irrational. In particular, this would be a strategy to prove the following conjecture (see §7.2 below):

Conjecture 1. *No $\xi \in \text{Frac}(\mathbf{G})$ can be a Liouville number.*

Our approach does not yield (at least for now) any actual result towards this conjecture, because the denominators of the coefficients of the G -functions we construct grow too fast. It would be interesting to control them in some way.

The paper is organized as follows. We introduce some notation in §2.1, and state slight generalizations of Theorems 1 and 3, namely Theorems 4 and 5. We prove in §2.2 that the numbers mentioned above actually belong to \mathbf{G} . Then we start proving Theorems 4 and 5 by gathering some lemmas in §§2.3 and 2.4. In §3, we prove that the conclusion of Theorem 1 holds for algebraic numbers and their logarithms. In §4, we review some classical results concerning the properties of differential equations satisfied by G -functions (namely Theorem 6, due to André, Chudnovski and Katz). We also prove in this section that connection constants belong to \mathbf{G} , and the conclusion

of Theorem 1 holds for them (see Theorem 7). This result, along with the analytic continuation properties of G -functions deduced from Theorem 6, is used to prove Theorem 4 in §5. In §6, we present the proof of Theorem 5: the main tool is the results of *Singularity Analysis* due to Flajolet and Odlyzko [21], described in details in the book [20]. Finally, we mention in §7 a few problems suggested by our results: what can be said about the case of E -functions and about Diophantine perspectives.

Acknowledgements. We warmly thank Yves André, Daniel Bertrand, Frits Beukers, Gilles Christol, Julien Roques and Michel Waldschmidt for their constructive remarks. We are also indebted to the referee for his pertinent comments that helped us to improve this work, in particular those we present in §7.1. Both authors have been supported by the project HAMOT (ref. ANR 2010-BLAN-0115), and the second author partially by the project Q-DIFF (ref. ANR 2010-JCJC-010501), of the Agence Nationale de la Recherche.

2. Background of the proofs

2.1. Notation and results. In this section we introduce some notation that will be used throughout this text. We also state Theorems 4 and 5, which are slight generalizations of Theorems 1 and 3 respectively.

The letter \mathbb{K} will always stand for a (finite or infinite) algebraic extension of \mathbb{Q} , embedded into $\bar{\mathbb{Q}} \subset \mathbb{C}$.

Definition 3. Given an algebraic extension \mathbb{K} of \mathbb{Q} , we denote by $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$ the set of all values, at points in \mathbb{K} , of multivalued analytic continuations of G -functions with Taylor coefficients at 0 in \mathbb{K} .

For any G -function f with coefficients in \mathbb{K} and any $\alpha \in \mathbb{K}$, we consider all values of $f(\alpha)$ obtained by analytic continuation, as in the definition of \mathbf{G} in the introduction; obviously $\mathbf{G} = \mathbf{G}_{\bar{\mathbb{Q}}}^{\text{a.c.}}$. If α is a singularity of f , then we consider also these values if they are finite. Of course $f(\alpha z)$ is also a G -function with coefficients in \mathbb{K} so that we may restrict ourselves to values at the point 1. By Abel's theorem, $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$ contains all convergent series $\sum_{n=0}^{\infty} a_n \alpha^n$ where $f(z) = \sum_{n=0}^{\infty} a_n z^n$ is a G -function with coefficients in \mathbb{K} and $\alpha \in \mathbb{K}$.

Definition 4. Given an algebraic extension \mathbb{K} of \mathbb{Q} , we denote by $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ the set of all $\xi \in \mathbb{C}$ such that, for any $R \geq 1$, there exists a G -function f with Taylor coefficients at 0 in \mathbb{K} and radius of convergence $> R$ such that $\xi = f(1)$.

For any $R \geq 1$, we denote by $\mathbf{G}_{R,\mathbb{K}}^{\text{cv}}$ the set of all $\xi = f(1)$ where f is a G -function with Taylor coefficients at 0 in \mathbb{K} and radius of convergence $> R$. In this way we have $\mathbf{G}_{\mathbb{K}}^{\text{cv}} = \bigcap_{R \geq 1} \mathbf{G}_{R,\mathbb{K}}^{\text{cv}}$, and also $\mathbf{G}_{R,\mathbb{K}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$ for any $R \geq 1$.

With this notation, Theorem 1 reads $\mathbf{G}_{\mathbb{Q}}^{\text{a.c.}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}} = \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$. Actually we prove that $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$ is independent from \mathbb{K} , so that it is always equal to $\mathbf{G} = \mathbf{G}_{\mathbb{Q}}^{\text{a.c.}}$. Concerning $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$, there is an obvious remark: if $\mathbb{K} \subset \mathbb{R}$ then $\mathbf{G}_{\mathbb{K}}^{\text{cv}} \subset \mathbb{R}$. Apart from this, $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is independent from \mathbb{K} , and equal (up to taking real parts) to \mathbf{G} . Our result reads as follows.

Theorem 4. *Let \mathbb{K} be an algebraic extension of \mathbb{Q} . Then:*

- We have $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}} = \mathbf{G} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$.
- If $\mathbb{K} \not\subset \mathbb{R}$ then $\mathbf{G}_{\mathbb{K}}^{\text{cv}} = \mathbf{G} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$; if $\mathbb{K} \subset \mathbb{R}$ then $\mathbf{G}_{\mathbb{K}}^{\text{cv}} = \mathbf{G} \cap \mathbb{R} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}}$.

In particular this result contains the fact that $\bar{\mathbb{Q}} \cap \mathbb{R} \subset \mathbf{G}_{\mathbb{Q}}^{\text{cv}}$ and $\bar{\mathbb{Q}} \subset \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$; this will be proved in §3.1. Another consequence of this theorem is that the set of values of G -functions $\sum_{n=0}^{\infty} a_n z^n$ with $a_n \in \mathbb{K}$ at points $z \in \mathbb{K}$ inside the disk of convergence (respectively at points where this series is absolutely convergent, respectively convergent) is equal to $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ (so that it is equal to either \mathbf{G} or $\mathbf{G} \cap \mathbb{R}$).

We also generalize Theorem 3 as follows.

Theorem 5. *Let \mathbb{K} be an algebraic extension of \mathbb{Q} , and $\xi \in \mathbb{C}^*$. Then the following statements are equivalent:*

- (i) We have $\xi \in \text{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}})$.
- (ii) There exist two sequences $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ of elements of \mathbb{K} such that $\sum_{n=0}^{\infty} a_n z^n$ and $\sum_{n=0}^{\infty} b_n z^n$ are G -functions, $b_n \neq 0$ for infinitely many n and $a_n - \xi b_n = o(b_n)$.
- (iii) For any $R \geq 1$ there exist two G -functions $A(z) = \sum_{n=0}^{\infty} a_n z^n$ and $B(z) = \sum_{n=0}^{\infty} b_n z^n$, with coefficients $a_n, b_n \in \mathbb{K}$ and radius of convergence = 1, such that $A(z) - \xi B(z)$ has radius of convergence $> R$ and $a_n, b_n \neq 0$ for any n sufficiently large.

When $\mathbb{K} = \mathbb{Q}$, this is a refinement of Theorem 3 because assumption (ii) of Theorem 3 implies assumption (ii) of Theorem 5, and (iii) of Theorem 5 implies (iii) of Theorem 3 (see also Lemma 2 below). The point in assertion (ii) of Theorem 5 is that b_n may vanish for infinitely many n ; by asking $a_n - \xi b_n = o(b_n)$ we require that $a_n = 0$ as soon as $b_n = 0$ and n is sufficiently large.

2.2. Examples and connection to periods. In this section, we prove that the numbers mentioned in the introduction belong to \mathbf{G} , and give some hints on the connection with periods. This section is independent from the rest of the paper, except that we assume here that \mathbf{G} is a ring.

Many examples of G -functions are provided by the generalized hypergeometric series

$$\sum_{n=0}^{\infty} \frac{(\alpha_1)_n (\alpha_2)_n \dots (\alpha_k)_n}{(1)_n (\beta_1)_n \dots (\beta_{k-1})_n} z^n$$

with rational coefficients α 's and β 's, and $(x)_n = x(x+1)\dots(x+n-1)$. Special cases are the polylogarithmic functions $\text{Li}_k(z) = \sum_{n \geq 1} \frac{z^n}{n^k}$ ($k \geq 1$) and $\arctan(z) = \sum_{n \geq 0} (-1)^n \frac{z^{2n+1}}{2n+1}$. We deduce in particular that $\pi = 4 \arctan(1)$ and the values of the Riemann zeta function $\zeta(k) = \text{Li}_k(1)$ are in \mathbf{G} for any integer $k \geq 2$. Catalan's constant $\sum_{n \geq 0} \frac{(-1)^n}{(2n+1)^2}$ is also in \mathbf{G} .

Other examples of G -functions are the multiple polylogarithms

$$\sum_{n_1 > \dots > n_s \geq 1} \frac{z^{n_1}}{n_1^{k_1} \dots n_s^{k_s}}$$

where the k 's are positive integers. This is a consequence of the fact that for $s = 1$, we have a polylogarithm from which we obtain the multiple series by a succession of integrations and multiplications by $1/z$ or $1/(1-z)$; this process does not leave the set of G -functions. As a consequence, multiple zeta values $\zeta(k_1, \dots, k_s) = \sum_{n_1 > \dots > n_s \geq 1} \frac{1}{n_1^{k_1} \dots n_s^{k_s}}$ (with $k_1 \geq 2$) are in \mathbf{G} .

It could seem more surprising that $1/\pi$ is also in \mathbf{G} , a fact proved by each one of the following identities:

$$\frac{1}{\pi} = \sum_{n=0}^{\infty} \frac{\binom{2n}{n}^2}{(1-2n)2^{4n+1}}, \quad \frac{1}{\pi} = \sum_{n=0}^{\infty} \frac{\binom{2n}{n}^3 (42n+5)}{2^{12n+4}}.$$

The first identity is a direct translation of the identity $E(1) = 1$ where $E(k) = \int_0^1 \sqrt{\frac{1-k^2 t^2}{1-t^2}} dt$ is Legendre's complete elliptic function of the second kind. The second identity is due to Ramanujan and it also has an elliptic interpretation. Both series are in fact values of generalized hypergeometric series, hence $1/\pi \in \mathbf{G}$.

In particular, π and the non-zero algebraic numbers are units of \mathbf{G} . These numbers do not span the whole group of units, as we now proceed to prove. Euler's Beta function is defined by

$$B(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} dt$$

for $\text{Re}(x), \text{Re}(y) > 0$. It is well-known that $B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$, which provides the meromorphic continuation of B to \mathbb{C}^2 ; we recall that $\pi = B(\frac{1}{2}, \frac{1}{2})$.

Proposition 1. (i) For all rational numbers x, y such that $B(x, y)$ is defined and non-zero, the number $B(x, y)$ is a unit of \mathbf{G} .

(ii) For any integers $a, b \geq 1$, we have

$$\Gamma\left(\frac{a}{b}\right)^b = (a-1)! \prod_{j=1}^{b-1} B\left(\frac{a}{b}, \frac{ja}{b}\right)$$

and $\Gamma\left(\frac{a}{b}\right)^b$ is a unit of \mathbf{G} .

Remark. a) To sum up, the group of units of \mathbf{G} contains the algebraic numbers and the numbers $B(x, y)$ where $x, y \in \mathbb{Q}$ (as soon as they are defined and non-zero). We don't know if this provides a complete list of generators of this group.

b) Chudnovski proved in 1974 that $\Gamma(1/3)$, respectively $\Gamma(1/4)$, and π are algebraically independent over $\overline{\mathbb{Q}}$. Hence one needs other transcendental generators than π in the group of units of \mathbf{G} .

c) This proposition is a transposition in our context of a discussion in André's book [6], pp. 211–212, where he shows that the numbers $\Gamma(a/b)^b$ are periods (in the geometric sense).

Proof. (i) We first show that $B(x, y) \in \mathbf{G}$ for all rational numbers $0 < x, y \leq 1$. Clearly, $B(x, y)$ is well defined in this case and

$$\begin{aligned} B(x, y) &= \int_0^1 t^{x-1} (1-t)^{y-1} dt = \int_0^1 \sum_{n=0}^{\infty} (-1)^n \binom{y-1}{n} t^{n+x-1} dt \\ &= \sum_{n=0}^{\infty} (-1)^n \binom{y-1}{n} \int_0^1 t^{n+x-1} dt = \sum_{n=0}^{\infty} (-1)^n \frac{\binom{y-1}{n}}{n+x}. \end{aligned}$$

Since $(-1)^n \binom{y-1}{n}$ is positive, permuting the series and integral is licit. Moreover, $\frac{\binom{y-1}{n}}{n+x} = \mathcal{O}(1/n^{y+1})$ so that the final series converges absolutely and is the value at $z = 1$ of a G -function. This proves that $B(x, y) \in \mathbf{G}$ in this case.

From now on, we let $x, y \in \mathbb{Q}$ and we assume that $x, y, x+y \notin \mathbb{Z}$ (otherwise the conclusion is easier to prove). Then $B(x, y)$ is defined and non-zero. There exist two integers M, N such that $0 < x+M, y+N \leq 1$, and the functional equations

$$B(x, y) = \frac{x+y}{x} B(x+1, y), \quad B(x, y) = \frac{x+y}{y} B(x, y+1)$$

yield $B(x, y) = R_{M,N}(x, y) B(x+M, y+N)$ with $R_{M,N}(x, y) \in \mathbb{Q}(x, y)$. Since $B(x+M, y+N)$ is in \mathbf{G} by the previous case, it follows that $B(x, y) \in \mathbf{G}$.

To prove that $1/B(x, y)$ is also in \mathbf{G} , we use the reflection formula $\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin(\pi x)}$ to get

$$\frac{1}{B(x, y)} = \frac{\sin(\pi x) \sin(\pi y)}{\sin \pi(x+y)} \cdot \frac{1-x-y}{\pi} \cdot B(1-x, 1-y).$$

Now $B(1-x, 1-y) \in \mathbf{G}$ by the case above, $\frac{(1-x-y)\sin(\pi x)\sin(\pi y)}{\sin \pi(x+y)}$ is an algebraic number (hence in \mathbf{G}) and $1/\pi \in \mathbf{G}$, so that $\frac{1}{B(x,y)} \in \mathbf{G}$.

(ii) We have

$$\prod_{j=1}^{b-1} B\left(\frac{a}{b}, \frac{ja}{b}\right) = \prod_{j=1}^{b-1} \frac{\Gamma(\frac{a}{b})\Gamma(\frac{aj}{b})}{\Gamma(\frac{a(j+1)}{b})} = \Gamma\left(\frac{a}{b}\right)^{b-1} \frac{\Gamma(\frac{a}{b})}{\Gamma(a)},$$

from which we obtain the claimed identity. Moreover, for any integer $j \geq 1$, $B(\frac{a}{b}, \frac{ja}{b})$ is obviously defined and non-zero, hence is a unit of \mathbf{G} by (i). Thus, this is also the case of $\Gamma(\frac{a}{b})^b$. \square

To conclude this section, we mention some remarks (due to the referee) towards the determination of the link between \mathbf{G} and $\mathcal{P}[1/\pi]$, where \mathcal{P} is the ring of periods (in Kontsevich and Zagier's sense [26]); in particular a natural question is whether $\mathbf{G} = \mathcal{P}[1/\pi]$ or not.

Bombieri–Dwork's conjecture suggests that \mathbf{G} might be contained in $\mathcal{P}[1/\pi]$. Indeed, this conjecture predicts that any G -function is solution of an extension of sub-quotients of Picard–Fuchs equations. It is not clear that such an extension is motivic, but for a Picard–Fuchs equation the G -matrix solution $Y(z)$ is the quotient $P(z)P(0)^{-1}$ of two period matrices. Since the determinant of $P(0)$ is an algebraic number times a power of π (see [2]), the inclusion $\mathbf{G} \subset \mathcal{P}[1/\pi]$ would follow.

Towards the converse inclusion, it is possible to prove that if a one-parameter Picard–Fuchs equation doesn't have 0 as a singularity then the special values of its solutions can be expressed in terms of G -functions which are solutions of the same equation.

In view of this discussion, it would be very interesting to refine Theorem 1 by ensuring that 0 isn't a singularity of the minimal differential equation of the G -function f we construct (such that $f(1)$ is a given $\xi \in \mathbf{G}$). However our proof does not provide this refinement directly and new ideas are necessary to do that.

2.3. General properties of the ring $\mathbf{G}_{\mathbb{K}}^{\text{sv}}$. The set of G -functions satisfies a number of structural properties. It is a ring and even a $\bar{\mathbb{Q}}[z]$ -algebra; it is stable by differentiation and the Hadamard product of two G -functions (obtained by pointwise multiplication of the coefficients) is again a G -function. These properties will be used throughout the text, as well as the fact that algebraic functions over $\bar{\mathbb{Q}}(z)$ which are holomorphic at $z = 0$ are G -functions: this is a consequence of Eisenstein's theorem³ and the fact that an algebraic function over $\bar{\mathbb{Q}}(z)$ satisfies a linear differential equation with coefficients in $\bar{\mathbb{Q}}[z]$.

The following property is useful too:

³which states that for any power series $\sum_{n=0}^{\infty} a_n z^n$ algebraic over $\bar{\mathbb{Q}}(z)$, there exists a positive integer D such that $D^n a_n$ is an algebraic integer for any n .

Lemma 1. Consider a G -function $\sum_{n=0}^{\infty} a_n z^n$. Then the series

$$\sum_{n=0}^{\infty} \bar{a}_n z^n, \quad \sum_{n=0}^{\infty} \operatorname{Re}(a_n) z^n \quad \text{and} \quad \sum_{n=0}^{\infty} \operatorname{Im}(a_n) z^n$$

are also G -functions.

Proof. The series $\sum_{n=0}^{\infty} a_n z^n$ satisfies a linear differential equation $Ly = 0$ with coefficients in $\bar{\mathbb{Q}}[z]$, hence $\sum_{n=0}^{\infty} \bar{a}_n z^n$ satisfies the linear differential equation $\bar{L}y = 0$ where \bar{L} is obtained from L by replacing each coefficient $\sum_{k=0}^d p_k z^k$ with $\sum_{k=0}^d \bar{p}_k z^k$. Furthermore, the moduli of the conjugates of \bar{a}_n and their common denominators obviously grow at most geometrically. Hence, $\sum_{n=0}^{\infty} \bar{a}_n z^n$ is a G -function.

For $\sum_{n=0}^{\infty} \operatorname{Re}(a_n) z^n$ and $\sum_{n=0}^{\infty} \operatorname{Im}(a_n) z^n$, we write $2\operatorname{Re}(a_n) = a_n + \bar{a}_n$, $2i\operatorname{Im}(a_n) = a_n - \bar{a}_n$ and use the fact that the sum of two G -functions is also a G -function. \square

The following lemma includes the easiest properties of $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$; especially (i) will be used very often without explicit reference.

Lemma 2. Let \mathbb{K} be an algebraic extension of \mathbb{Q} .

- (i) $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is a ring and it contains \mathbb{K} .
- (ii) If \mathbb{K} is invariant under complex conjugation then:
 - $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is invariant under complex conjugation.
 - $\mathbf{G}_{\mathbb{K} \cap \mathbb{R}}^{\text{cv}} = \mathbf{G}_{\mathbb{K}}^{\text{cv}} \cap \mathbb{R}$.
 - $\mathbb{R} \cap \operatorname{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}}) = \operatorname{Frac}(\mathbf{G}_{\mathbb{K} \cap \mathbb{R}}^{\text{cv}}) = \operatorname{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}} \cap \mathbb{R})$.
- (iii) $\mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}}[i] = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$, and more generally if $\mathbb{K} \subset \mathbb{R}$ then $\mathbf{G}_{\mathbb{K}(i)}^{\text{cv}} = \mathbf{G}_{\mathbb{K}}^{\text{cv}}[i] = \mathbf{G}_{\mathbb{K}}^{\text{cv}} + i\mathbf{G}_{\mathbb{K}}^{\text{cv}}$.

Proof. (i) The properties of G -functions ensure that the sum and product of two G -functions with coefficients in \mathbb{K} and radii of convergence $> R \geq 1$ are G -functions with coefficients in \mathbb{K} and radii of convergence $> R$. Moreover algebraic constants are G -functions with infinite radius of convergence.

(ii) Using Lemma 1 and the fact that \mathbb{K} is invariant under complex conjugation, if $\sum_{n=0}^{\infty} a_n z^n$ is a G -function with coefficients in \mathbb{K} and radii of convergence $> R \geq 1$ then so is $\sum_{n=0}^{\infty} \bar{a}_n z^n$: this proves that $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is invariant under complex conjugation.

The inclusion $\mathbf{G}_{\mathbb{K} \cap \mathbb{R}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}} \cap \mathbb{R}$ is obvious. Conversely, if $\xi \in \mathbb{R} \cap \mathbf{G}_{\mathbb{K}}^{\text{cv}}$ then for any $R \geq 1$ we have $\xi = \sum_{n=0}^{\infty} a_n$ where $\sum_{n=0}^{\infty} a_n z^n$ is a G -function with coefficients in \mathbb{K} and radius of convergence $> R$. Then $\sum_{n=0}^{\infty} \operatorname{Re}(a_n) z^n$ is also a G -function (by

Lemma 1); it has coefficients in $\mathbb{K} \cap \mathbb{R}$ (because $\operatorname{Re}(a_n) = \frac{1}{2}(a_n + \bar{a}_n)$) and radius of convergence $> R$. Therefore $\xi = \sum_{n=0}^{\infty} \operatorname{Re}(a_n) \in \mathbf{G}_{\mathbb{K} \cap \mathbb{R}}^{\text{cv}}$.

Finally, the inclusion $\operatorname{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}} \cap \mathbb{R}) \subset \mathbb{R} \cap \operatorname{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}})$ is trivial. The converse is trivial too if $\mathbb{K} \subset \mathbb{R}$; otherwise let $\xi, \xi' \in \mathbf{G}_{\mathbb{K}}^{\text{cv}}$ be such that $\xi' \neq 0$ and $\xi/\xi' \in \mathbb{R}$. Multiplying if necessary by a non-real element of \mathbb{K} , we may assume $\xi, \xi' \notin i\mathbb{R}$. Then we have $\xi/\xi' = (\xi + \bar{\xi})/(\xi' + \bar{\xi}') \in \operatorname{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}} \cap \mathbb{R})$.

(iii) Assume $\mathbb{K} \subset \mathbb{R}$. Since $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is a ring and $i^2 = -1 \in \mathbf{G}_{\mathbb{K}}^{\text{cv}}$, we have $\mathbf{G}_{\mathbb{K}}^{\text{cv}}[i] = \mathbf{G}_{\mathbb{K}}^{\text{cv}} + i\mathbf{G}_{\mathbb{K}}^{\text{cv}}$. This is obviously a subset of $\mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$. Conversely, $\mathbb{K}(i)$ is invariant under complex conjugation (because $\mathbb{K} \subset \mathbb{R}$) so that for any $\xi \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$ we have $\operatorname{Re}(\xi) = \frac{1}{2}(\xi + \bar{\xi}) \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}} \cap \mathbb{R} = \mathbf{G}_{\mathbb{K}}^{\text{cv}}$ by (ii). Since $i \in \mathbb{K}(i) \subset \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$ we have $\operatorname{Im}(\xi) = -i(\xi - \operatorname{Re}(\xi)) \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}} \cap \mathbb{R} = \mathbf{G}_{\mathbb{K}}^{\text{cv}}$, using (ii) again. Finally $\xi = \operatorname{Re}(\xi) + i\operatorname{Im}(\xi) \in \mathbf{G}_{\mathbb{K}}^{\text{cv}} + i\mathbf{G}_{\mathbb{K}}^{\text{cv}}$. \square

The following lemma is a consequence of Lemma 7 proved in §3 below; of course the proof of Lemma 7 does not use Lemma 3, hence there is no circularity.

Lemma 3. *Let \mathbb{K} be an algebraic extension of \mathbb{Q} .*

(i) *We have $\bar{\mathbb{Q}} \cap \mathbb{R} \subset \mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$, and $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is a $(\bar{\mathbb{Q}} \cap \mathbb{R})$ -algebra.*

(ii) *If $\mathbb{K} \not\subset \mathbb{R}$ then $\bar{\mathbb{Q}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$, and $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is a $\bar{\mathbb{Q}}$ -algebra.*

Proof. (i) By Lemma 7, we have $\bar{\mathbb{Q}} \cap \mathbb{R} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} \cap \mathbb{R}$; this is equal to $\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$ by Lemma 2. The inclusion $\mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is trivial since $\mathbb{Q} \subset \mathbb{K}$.

(ii) Since $\mathbb{K} \not\subset \mathbb{R}$, there exist $\alpha, \beta \in \mathbb{R}$ such that $\alpha + i\beta \in \mathbb{K}$ and $\beta \neq 0$; since $\alpha - i\beta$ is also algebraic, we have $\alpha, \beta \in \bar{\mathbb{Q}}$. Therefore we can write $i = \frac{1}{\beta}((\alpha + i\beta) - \alpha)$ with $\frac{1}{\beta}, \alpha \in \bar{\mathbb{Q}} \cap \mathbb{R} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$ (by (i)). Since $\mathbf{G}_{\mathbb{K}}^{\text{cv}}$ is a ring which contains $\alpha + i\beta$, this yields $i \in \mathbf{G}_{\mathbb{K}}^{\text{cv}}$, so that (using Lemma 2 and the trivial inclusion $\mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$) $\mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$. Using the inclusion $\bar{\mathbb{Q}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$ proved in Lemma 7, this concludes the proof of (ii). \square

To conclude this section, we state and prove the following lemma, which is very useful for constructing elements of $\mathbf{G}_{R, \mathbb{K}}^{\text{cv}}$. Recall that $\mathbf{G}_{R, \mathbb{K}}^{\text{cv}}$ is the set of all $\xi = f(1)$ where f is a G -function with coefficients in \mathbb{K} and radius of convergence $> R$.

Lemma 4. *Let \mathbb{K} be an algebraic extension of \mathbb{Q} . Let $\zeta \in \mathbb{K}$, and $g(z)$ be a G -function in the variable $\zeta - z$, with coefficients in \mathbb{K} and radius of convergence $\geq r > 0$. Then $g(z_0) \in \mathbf{G}_{R, \mathbb{K}}^{\text{cv}}$ for any $R \geq 1$ and any $z_0 \in \mathbb{K}$ such that $|z_0 - \zeta| < r/R$.*

Proof. Letting $f(z) = g(\zeta + z(z_0 - \zeta))$, we have $f(1) = g(z_0)$ and f is a G -function with coefficients in \mathbb{K} and radius of convergence $> R$. \square

2.4. Miscellaneous lemmas. We gather in this section two lemmas which are neither difficult nor specific to G -functions, but very useful.

Lemma 5. *Let \mathbb{A} be a subring of \mathbb{C} . Let $S \subset \mathbb{N}$ and $T \subset \mathbb{Q}$ be finite subsets. For any $(s, t) \in S \times T$, let $f_{s,t}(z) = \sum_{n=0}^{\infty} a_{s,t,n} z^n \in \mathbb{A}[[z]]$ be a function holomorphic at 0, with Taylor coefficients in \mathbb{A} . Let Ω denote an open subset of \mathbb{C} , with 0 in its boundary, on which a continuous determination of the logarithm is chosen. Then there exist $c \in \mathbb{A}$, $\sigma \in \mathbb{N}$ and $\tau \in \mathbb{Q}$ such that, as $z \rightarrow 0$ with $z \in \Omega$,*

$$\sum_{s \in S} \sum_{t \in T} (\log z)^s z^t f_{s,t}(z) = c (\log z)^\sigma z^\tau (1 + o(1)). \quad (2.1)$$

Proof. Let $T + \mathbb{N} = \{t + n, t \in T, n \in \mathbb{N}\}$. For any $s \in S$ and any $\theta \in T + \mathbb{N}$, let $c_{s,\theta} = \sum_{t \in T} a_{s,t,\theta-t}$ where we let $a_{s,t,\theta-t} = 0$ if $\theta - t \notin \mathbb{N}$. Then the left-hand side of (2.1) can be written, for $z \in \Omega$ sufficiently close to 0, as an absolutely converging series $\sum_{\theta \in T + \mathbb{N}} \sum_{s \in S} c_{s,\theta} (\log z)^s z^\theta$. If $c_{s,\theta} = 0$ for any (s, θ) then (2.1) holds with $c = 0$. Otherwise we denote by τ the minimal value of θ for which there exists $s \in S$ with $c_{s,\theta} \neq 0$, and by σ the largest $s \in S$ such that $c_{s,\tau} \neq 0$. Then (2.1) holds with $c = c_{\sigma,\tau} \in \mathbb{A}$. \square

The following result will be used in the proof of Theorem 5.

Lemma 6. *Let $\omega_1, \dots, \omega_t$ be pairwise distinct complex numbers, with $|\omega_1| = \dots = |\omega_t| = 1$. Let $\kappa_1, \dots, \kappa_t \in \mathbb{C}$ be such that $\lim_{n \rightarrow +\infty} \kappa_1 \omega_1^n + \dots + \kappa_t \omega_t^n = 0$. Then $\kappa_1 = \dots = \kappa_t = 0$.*

Proof. For any $n \geq 0$, let $\delta_n = \det M_n$ where

$$M_n = \begin{pmatrix} \omega_1^n & \omega_2^n & \dots & \omega_t^n \\ \omega_1^{n+1} & \omega_2^{n+1} & \dots & \omega_t^{n+1} \\ \vdots & \vdots & \ddots & \vdots \\ \omega_1^{n+t-1} & \omega_2^{n+t-1} & \dots & \omega_t^{n+t-1} \end{pmatrix}.$$

Let $C_{i,n}$ denote the i -th column of M_n . Since $C_{i,n} = \omega_i^n C_{i,0}$ we have $|\delta_n| = |\omega_1^n \dots \omega_t^n \delta_0| = |\delta_0| \neq 0$ because δ_0 is the Vandermonde determinant built on the pairwise distinct numbers $\omega_1, \dots, \omega_t$. Now assume that $\kappa_j \neq 0$ for some j . Then for computing δ_n we can replace $C_{j,n}$ with $\frac{1}{\kappa_j} \sum_{i=1}^t \kappa_i C_{i,n}$; this implies $\lim_{n \rightarrow +\infty} \delta_n = 0$, in contradiction with the fact that $|\delta_n| = |\delta_0| \neq 0$. \square

3. Algebraic numbers and logarithms as values of G -functions

An important step for us is to show that algebraic numbers are values of G -functions with coefficients in $\mathbb{Q}(i)$ (and, more precisely, that they satisfy the conclusion of

Theorem 1). Despite quite general results in related directions, this fact does not seem to have been proved in the literature in the full form we need. Eisenstein [31] showed that the G -function (of hypergeometric type)

$$\sum_{n=0}^{\infty} (-1)^n \frac{\binom{5n}{n}}{4n+1} a^{4n+1}$$

is a solution of the quintic equation $x^5 + x = a$, provided that $|a| \leq 5^{-5/4}$ (to ensure the convergence of the series). Eisenstein's formula can be proved using Lagrange's inversion formula. More generally, given a polynomial $P(x) \in \mathbb{C}[x]$, it is known that multivariate series can be used to find expressions of the roots of P in terms of its coefficients p_j . For example in [32], it is shown that these roots can be formally expressed as A -hypergeometric series evaluated at rational powers of the p_j 's. (A -hypergeometric series are an example of multivariate G -functions.) It is not clear how such a representation could be used to prove Lemma 7 below: beside the multivariate aspect, the convergence of the series imposes some conditions on the p_j 's and their exponents are not integers in general. Our proof is more in Eisenstein's spirit.

Lemma 7. *Let $\alpha \in \overline{\mathbb{Q}}$, and $Q(X) \in \mathbb{Q}[X]$ be a non-zero polynomial of which α is a simple root. For any $u \in \mathbb{Q}(i)$ such that $Q'(u) \neq 0$, the series*

$$\Phi_u(z) = u + \sum_{n=1}^{\infty} (-1)^n \frac{Q(u)^n}{n!} \frac{\partial^{n-1}}{\partial x^{n-1}} \left(\left(\frac{x-u}{Q(x)-Q(u)} \right)^n \right) \Big|_{x=u} z^n$$

is a G -function with coefficients in $\mathbb{Q}(i)$; it satisfies the equation

$$Q(\Phi_u(z)) = (1-z)Q(u).$$

For any $R \geq 1$, if u is close enough to α then the radius of convergence of Φ_u is $> R$ and $\alpha = \Phi_u(1) \in \mathbf{G}_{R, \mathbb{Q}(i)}^{\text{cv}}$.

Accordingly we have $\overline{\mathbb{Q}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$.

Remarks. a) The proof can be made effective, i.e., given α , Q and R , we can compute $\varepsilon(\alpha, Q, R)$ such that for any $u \in \mathbb{Q}(i)$ with $|\alpha - u| < \varepsilon(\alpha, Q, R)$, we have $\Phi_u(1) = \alpha$ and the radius of convergence of Φ_u is $> R$.

b) Using Lemma 2(ii), we deduce that any real algebraic number is in $\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$.

We also need a similar property for values of the logarithm.

Lemma 8. *Let $\alpha \in \overline{\mathbb{Q}}^*$. For any determination of the logarithm, the number $\log(\alpha)$ belongs to $\mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$.*

3.1. Algebraic numbers

Proof of Lemma 7. If $\deg Q = 1$ then $\Phi_u(z) = u + (\alpha - u)z$ so that Lemma 7 holds trivially. From now on we assume $\deg Q \geq 2$. Then $\frac{Q(X) - Q(u)}{X - u}$ is a non-constant polynomial with coefficients in $\mathbb{Q}(i)$; its value at $X = u$ is $Q'(u) \neq 0$ so that the coefficients of $\Phi_u(z)$ are well-defined and belong to $\mathbb{Q}(i)$. If $Q(u) = 0$ then $\Phi_u(z) = u$ and the result is trivial, so that we may assume $Q(u) \neq 0$ and define the polynomial function

$$z_u(t) = 1 - \frac{Q(t + u)}{Q(u)} \in \mathbb{Q}(i)[t]$$

so that $z_u(0) = 0$ and $z'_u(0) = -\frac{Q'(u)}{Q(u)} \neq 0$. Hence $z_u(t)$ can be locally inverted around $t = 0$ and its inverse $t_u(z) = \sum_{n \geq 1} \phi_n(u) z^n$ is holomorphic at $z = 0$.

The Taylor coefficients of t_u can be computed by means of the Lagrange inversion formula [20], p. 732, which in this case gives $\Phi_u(z) = u + t_u(z)$. By definition of $t_u(z)$, this implies $Q(\Phi_u(z)) = (1 - z)Q(u)$. Therefore Φ_u is an algebraic function hence it is a G -function.

Now let

$$\phi_n(u) = \frac{(-Q(u))^n}{n!} \frac{\partial^{n-1}}{\partial x^{n-1}} \left(\left(\frac{x - u}{Q(x) - Q(u)} \right)^n \right) \Big|_{x=u}$$

denote, for $n \geq 1$, the coefficient of z^n in $\Phi_u(z)$. Then for any $n \geq 1$ we have

$$\phi_n(u) = \frac{Q(u)^n}{2i\pi} \int_{\mathcal{C}} \frac{dz}{(Q(u) - Q(z))^n} \quad (3.1)$$

where \mathcal{C} is a closed path surrounding u but no other roots of the polynomial $Q(X) - Q(u)$. This enables us to get an upper bound on the growth of the coefficients $\phi_n(u)$. Let us denote by $\beta_1(u) = u, \beta_2(u), \dots, \beta_d(u)$ the roots (repeated according to their multiplicities) of the polynomial $Q(X) - Q(u)$, with $d = \deg Q \geq 2$. We take u close enough to α so that $\beta_2(u), \dots, \beta_d(u)$ are also close to the other roots $\alpha_2, \dots, \alpha_d$ of the polynomial $Q(X)$. Since α is a simple root of $Q(X)$, we have $\alpha \notin \{\alpha_2, \dots, \alpha_d\}$. We can then choose the smooth curve \mathcal{C} in (3.1) independent from u such that the distance from \mathcal{C} to any one of $u, \beta_2(u), \dots, \beta_d(u)$ is $\geq \varepsilon > 0$ with ε also independent from u , in such a way that u lies inside \mathcal{C} and $\beta_2(u), \dots, \beta_d(u)$ outside \mathcal{C} .⁴ It follows in particular that, for any $z \in \mathcal{C}$, $|Q(u) - Q(z)| \geq \rho$ for some $\rho > 0$ independent from u . Hence $\max_{z \in \mathcal{C}} \left| \frac{1}{Q(u) - Q(z)} \right| \leq \frac{1}{\rho}$. From the Cauchy integral in (3.1), we deduce that

$$|\phi_n(u)| \leq \frac{|\mathcal{C}|}{2\pi} \cdot \frac{|Q(u)|^n}{\rho^n}, \quad (3.2)$$

⁴We do so because we want to use a curve \mathcal{C} that does not depend of u , whereas the poles of the integrand move with u .

where $|\mathcal{C}|$ is the length of \mathcal{C} . Let $R \geq 1$. Since $Q(u) \rightarrow Q(\alpha) = 0$ as $u \rightarrow \alpha$, we deduce that the radius of convergence of $\Phi_u(z)$ is $> R$ provided that u is sufficiently close to α (namely as soon as $R|Q(u)| < \rho$). Then the series $\Phi_u(1)$ is absolutely convergent and we have

$$|\Phi_u(1) - u| = \left| \sum_{n=1}^{\infty} \phi_n(u) \right| \leq \frac{|\mathcal{C}|}{2\pi} \sum_{n=1}^{\infty} \frac{|Q(u)|^n}{\rho^n} = \mathcal{O}(|Q(u)|). \quad (3.3)$$

Therefore $\Phi_u(1)$ can be made arbitrarily close to u , and accordingly arbitrarily close to α . Now for any z inside the disk of convergence of Φ_u we have $Q(\Phi_u(z)) = (1-z)Q(u)$, so that $\Phi_u(1)$ is a root of $Q(X)$. If it is sufficiently close to α , it has to be α . This completes the proof of Lemma 7. \square

3.2. Logarithms of algebraic numbers

Proof of Lemma 8. Throughout this proof, we will always consider the determination of $\log z$ of which the imaginary part belongs to $(-\pi, \pi]$ (but the result holds for any determination because $i\pi = \log(-1) \in \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$).

Using the formula $\log(\alpha) = n \log(\alpha^{1/n})$ with n sufficiently large, we may assume that α is arbitrarily close to 1; in particular the imaginary part of $\log \alpha$ gets arbitrarily close to 0.

Letting $Q(X)$ denote the minimal polynomial of α , we keep the notation in the proof of Lemma 7, and write $\alpha = \Phi_u(1) = u + u\Psi_u(1)$ where $u \in \mathbb{Q}(i)$ is close enough to α , $\Psi_u(1)$ is in $\mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$ and $\Psi_u(0) = 0$. By Equation (3.2), the radius of convergence at $z = 0$ of the G -function $\Psi_u(z)$ can be taken arbitrarily large provided that $u \in \mathbb{Q}(i)$ is close enough to α . We have

$$\log(\alpha) = \log(\alpha/u) + \log(u) = \log(1 + \Psi_u(1)) + \log(u),$$

because all logarithms in this equality have imaginary parts arbitrarily close to 0. Let $R \geq 1$; we shall prove, if u is close enough to 1, that both $\log(1 + \Psi_u(1))$ and $\log(u)$ belong to $\mathbf{G}_{R, \mathbb{Q}(i)}^{\text{cv}}$.

a) Provided that u is close enough to α , reasoning as in Equation (3.3) we get $|\Psi_u(z)| < 1$ for all z in a disk of center 0 and radius $> R$. Hence for such a u , the radius of convergence of the Taylor series of $\log(1 + \Psi_u(z))$ at $z = 0$ is $> R \geq 1$. To see that it is a G -function with coefficients in $\mathbb{Q}(i)$, we observe that $\frac{d}{dz} \log(1 + \Psi_u(z)) = \frac{\Psi'_u(z)}{1 + \Psi_u(z)}$ is an algebraic function holomorphic at the origin: its Taylor series is a G -function $\sum_{n=0}^{\infty} a_n z^n \in \mathbb{Q}(i)[[z]]$. Therefore $\log(1 + \Psi_u(z)) = \sum_{n=0}^{\infty} \frac{a_n}{n+1} z^{n+1} \in \mathbb{Q}(i)[[z]]$; this is a G -function because the set of G -functions is stable under Hadamard product and both $\sum_{n=0}^{\infty} a_n z^{n+1}$ and $\sum_{n=0}^{\infty} \frac{1}{n+1} z^{n+1}$ are G -functions. Whence, $\log(1 + \Psi_u(1)) \in \mathbf{G}_{R, \mathbb{Q}(i)}^{\text{cv}}$.

b) It remains to prove that $\log(u) \in \mathbf{G}_{R, \mathbb{Q}(i)}^{\text{cv}}$ for any $u \in \mathbb{Q}(i)$ sufficiently close to 1. Let $a, b \in \mathbb{Q}$ be such that $u = a + ib$. Then we have

$$\log(u) = \frac{1}{2} \log(a^2 + b^2) + i \arctan\left(\frac{b}{a}\right).$$

Now $\log(1 + z) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} z^n$ and $\arctan(z) = \sum_{n=0}^{\infty} \frac{(-1)^n}{2n+1} z^{2n+1}$ are G -functions with rational coefficients and radius of convergence = 1, and we may assume that $|a^2 + b^2 - 1| < 1/R$ and $|b/a| < 1/R$. Then $\log(u) \in \mathbf{G}_{R, \mathbb{Q}(i)}^{\text{cv}}$ (see Lemma 4). \square

4. Analytic continuation and connection constants

4.1. Properties of differential equations of G -functions. Let \mathbb{K} be an algebraic extension of \mathbb{Q} , and $f(z) = \sum_{n=0}^{\infty} a_n z^n \in \mathbb{K}[[z]]$ be a G -function with coefficients $a_n \in \mathbb{K}$. Let L be a minimal differential equation with coefficients in $\mathbb{K}[z]$ of which $f(z)$ is a solution. We denote by $\xi_1, \dots, \xi_p \in \mathbb{C}$ the singularities of L (throughout this paper, we will consider only points at finite distance). For any $i \in \{1, \dots, p\}$, let Δ_i be a closed broken line from ξ_i to the point at infinity; we assume $\Delta_i \cap \Delta_j = \emptyset$ for any $i \neq j$, and let $\mathcal{D} = \mathbb{C} \setminus (\Delta_1 \cup \dots \cup \Delta_p)$: this is a simply connected open subset of \mathbb{C} . In most cases we shall take for Δ_i a closed half-line starting at ξ_i .

The differential equation $Ly = 0$ has holomorphic solutions on \mathcal{D} , and these solutions make up a \mathbb{C} -vector space of dimension equal to the order of L ; a basis of this vector space will be referred to as a *basis of solutions of L* .

Let ζ be a singularity of L . Then for any sufficiently small open disk D centered at ζ , the intersection $D \cap \mathcal{D}$ is equal to D with a ray removed; let us choose a determination of the logarithm of $\zeta - z$, denoted by $\log(\zeta - z)$, for $z \in D \cap \mathcal{D}$ (in such a way that it is holomorphic in z). If $\zeta \in \mathcal{D}$ is not a singularity of L , the function $\log(\zeta - z)$ will cancel out in what follows.

We shall use the following theorem (see [4], p. 719, for a discussion).

Theorem 6 (André, Chudnovski, Katz). *Let \mathbb{K} denote an algebraic extension of \mathbb{Q} . Consider a minimal differential equation L of order μ , with coefficients in $\mathbb{K}[z]$ and admitting a solution at $z = 0$ which is a G -function in $\mathbb{K}[[z]]$. Let \mathcal{D} , ξ_1, \dots, ξ_p be as above. Then L is fuchsian with rational exponents at each of its singularities, and for each point $\zeta \in \mathcal{D} \cup \{\xi_1, \dots, \xi_p\}$ there is a basis of solutions $(g_1(z), \dots, g_\mu(z))$ of L , holomorphic on \mathcal{D} , with the following properties:*

- *There exists an open disk D centered at ζ and functions $F_{s,t,j}(z)$, holomorphic at 0, such that for any $j \in \{1, \dots, \mu\}$ and any $z \in D \cap \mathcal{D}$:*

$$g_j(z) = \sum_{s \in S_j} \sum_{t \in T_j} (\log(\zeta - z))^s (\zeta - z)^t F_{s,t,j}(\zeta - z)$$

where $S_j \subset \mathbb{N}$ and $T_j \subset \mathbb{Q}$ are finite subsets.

- If $\zeta \in \mathbb{K}$ then the functions $F_{s,t,j}(z)$ are G -functions with coefficients in \mathbb{K} .
- If ζ is not a singularity of L then $S_j = T_j = \{0\}$ for any j , so that $g_1(z), \dots, g_\mu(z)$ are holomorphic at $z = \zeta$.

This theorem is usually stated in a more precise form, namely

$$(g_1(z), \dots, g_\mu(z)) = (f_1(\zeta - z), f_2(\zeta - z), \dots, f_\mu(\zeta - z)) \cdot (\zeta - z)^{C_\zeta}$$

where the functions $f_j(z)$ are holomorphic at 0 and C_ζ is an upper triangular matrix, and a similar formulation holds for the singularity at infinity, where one replaces $\zeta - z$ by $1/z$. However this precise version won't be used in this paper.

4.2. Statement of the theorem on connection constants. Let \mathbb{K} , f , L and \mathcal{D} be as in §4.1. Let (g_1, \dots, g_μ) denote a basis of the \mathbb{C} -vector space of holomorphic solutions on \mathcal{D} of the differential equation $Ly = 0$; here μ is the order of L . Since $f \in \mathbb{K}[[z]]$ satisfies $Lf = 0$ and is holomorphic on a small open disk centered at 0, it can be analytically continued to \mathcal{D} and expanded in the basis (g_1, \dots, g_μ) :

$$f(z) = \sum_{j=1}^{\mu} \varpi_j g_j(z) \quad (4.1)$$

for any $z \in \mathcal{D}$, where $\varpi_1, \dots, \varpi_\mu \in \mathbb{C}$ are called *connection constants*.

The following theorem⁵ is an important ingredient in the proof of Theorems 4 and 5.

Theorem 7. Let \mathbb{K} denote an algebraic extension of \mathbb{Q} . Consider a minimal differential equation L of order μ , with coefficients in $\mathbb{K}[z]$ and admitting a solution at $z = 0$ which is a G -function $f \in \mathbb{K}[[z]]$. Let \mathcal{D} , ξ_1, \dots, ξ_p be as above, $\zeta \in \mathbb{K} \cap (\mathcal{D} \cup \{\xi_1, \dots, \xi_p\})$ and (g_1, \dots, g_μ) be a basis of solutions given by Theorem 6. Then the connection constants $\varpi_1, \dots, \varpi_\mu$ defined by Equation (4.1) belong to $\mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$.

The following corollary is a consequence of Theorem 7 and Lemma 5 (applied with $\mathbb{A} = \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$). It is used in the proof of Theorem 5.

Corollary 1. Let \mathbb{K} , f , \mathcal{D} , ζ be as in Theorem 7. Then there exist $c \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$, $\sigma \in \mathbb{N}$ and $\tau \in \mathbb{Q}$ such that, as $z \rightarrow \zeta$ with $z \in \mathcal{D}$,

$$f(z) = c (\log(\zeta - z))^\sigma (\zeta - z)^\tau (1 + o(1)).$$

⁵As the proof shows, Theorem 7 holds under slightly weaker assumptions: it applies to any G -operator L such that $Lf = 0$, and also to $\zeta = \infty$.

4.3. Wronskian of fuchsian equations. Given a linear differential equation L with coefficients in $\bar{\mathbb{Q}}(z)$, of order μ and with a basis of solutions f_1, f_2, \dots, f_μ , the wronskian $W = W(f_1, \dots, f_\mu)$ is the determinant

$$W(z) = \begin{vmatrix} f_1(z) & f_2(z) & \cdots & f_\mu(z) \\ f_1^{(1)}(z) & f_2^{(1)}(z) & \cdots & f_\mu^{(1)}(z) \\ \vdots & \vdots & \cdots & \vdots \\ f_1^{(\mu-1)}(z) & f_2^{(\mu-1)}(z) & \cdots & f_\mu^{(\mu-1)}(z) \end{vmatrix}.$$

The wronskian can be defined in a more intrinsic way as follows. We write L as

$$y^{(\mu)}(z) + a_{\mu-1}(z)y^{(\mu-1)}(z) + \cdots + a_1(z)y'(z) + a_0(z)y(z) = 0$$

where $a_j(z) \in \bar{\mathbb{Q}}(z)$, $j = 1, \dots, \mu - 1$. Then $W(z)$ is a solution of the linear equation

$$y'(z) = -a_{\mu-1}(z)y(z), \quad (4.2)$$

hence $W(z) = v_0 \exp\left(-\int a_{\mu-1}(z)dz\right)$. The value of the constant v_0 is determined by the solutions f_1, f_2, \dots, f_μ .

Lemma 9. *Let $\mathbb{K}, f, L, \mathcal{D}, \zeta, g_1, \dots, g_\mu$ be as in Theorem 7. Then the wronskian $W(z) = W(g_1, \dots, g_\mu)(z)$ is an algebraic function over $\bar{\mathbb{Q}}(z)$, and its zeros and singularities lie among the poles of $a_{\mu-1}(z)$.*

Proof. Since the differential equation (4.2) is fuchsian, Equation (5.1.16) in [24], p. 148, yields $W(z) = v \prod_{j=1}^J (z - p_j)^{-r_j}$ where $p_1, \dots, p_J \in \bar{\mathbb{Q}}$ are the poles of $a_{\mu-1}(z)$ (which are simple because L is fuchsian), $r_1, \dots, r_J \in \mathbb{Q}$ (because L has rational exponents at its singularities), and $v \in \mathbb{C}^*$. It remains to prove that v is algebraic.

With this aim in view, we compute the determinant $W(z)$ for $z \in \mathcal{D}$ sufficiently close to ζ by means of the expansions of g_1, \dots, g_μ and their derivatives. This yields

$$W(z) = \sum_{s \in S} \sum_{t \in T} (\log(\zeta - z))^s (\zeta - z)^t F_{s,t}(\zeta - z)$$

where $S \subset \mathbb{N}$ and $T \subset \mathbb{Q}$ are finite subsets, and the $F_{s,t}(z)$ are G -functions with coefficients in \mathbb{K} . Now Lemma 5 provides $c \in \mathbb{K}$, $\sigma \in \mathbb{N}$ and $\tau \in \mathbb{Q}$ such that, as $z \rightarrow \zeta$ with $z \in \mathcal{D}$:

$$W(z) = c (\log(\zeta - z))^\sigma (\zeta - z)^\tau (1 + o(1)).$$

On the other hand we also have $\prod_{j=1}^J (z - p_j)^{-r_j} = \tilde{c}(\zeta - z)^{\tilde{\tau}}(1 + o(1))$ for some $\tilde{c} \in \bar{\mathbb{Q}}^*$ and $\tilde{\tau} \in \mathbb{Q}$. Since the quotient is a constant, namely v , taking limits as $z \rightarrow \zeta$ yields $\sigma = 0$, $\tau = \tilde{\tau}$ and $v = c/\tilde{c} \in \bar{\mathbb{Q}}$. This concludes the proof of Lemma 9. \square

4.4. Proof of Theorem 7. Let $R \geq 1$. For any $\xi \in (\mathcal{D} \setminus \{0, \zeta\}) \cap \mathbb{K}(i)$, let $r_\xi > 0$ be the distance of ξ to the border $\Delta_1 \cup \dots \cup \Delta_p$ of \mathcal{D} (with the notation of §4.1), and D_ξ be the open disk centered at ξ of radius r_ξ/R . Since ξ is not a singularity of L , there is a basis $g_{1,\xi}(z), \dots, g_{\mu,\xi}(z)$ of solutions of $Ly = 0$ consisting in G -functions in the variable $\xi - z$ with coefficients in $\mathbb{K}(i)$ (by Theorem 6); these G -functions have radii of convergence $\geq r_\xi$, so that $g_{j,\xi}(z) \in \mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$ for any $z \in D_\xi \cap \mathbb{K}(i)$ and any j (see Lemma 4).

Let $r_0 > 0$ be the radius of convergence of the G -function $f(z)$, and D_0 denote the open disk centered at 0 with radius r_0/R . Finally, for any $j \in \{1, \dots, \mu\}$ we let $g_{j,\zeta}(z) = g_j(z)$; by assumption there exists $r_\zeta > 0$ such that

$$g_{j,\zeta}(z) = \sum_{s \in S_j} \sum_{t \in T_j} (\log(\zeta - z))^s (\zeta - z)^t F_{s,t,j}(\zeta - z)$$

for any $z \in \mathcal{D}$ such that $|z - \zeta| < r_\zeta$, where $S_j \subset \mathbb{N}$ and $T_j \subset \mathbb{Q}$ are finite subsets and the $F_{s,t,j}$ are G -functions with coefficients in \mathbb{K} and radii of convergence $\geq r_\zeta$. Then we let D_ζ be the open disk centered at ζ with radius r_ζ/R , so that for any $z \in D_\zeta \cap \mathbb{K}(i)$ and any j we have $g_{j,\zeta}(z) \in \mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$ by Lemmas 4, 7 and 8.

Following a smooth injective compact path from 0 to ζ inside $\mathcal{D} \cup \{0, \zeta\}$, we can find $s - 2$ points $\xi_2, \dots, \xi_{s-1} \in (\mathcal{D} \setminus \{0, \zeta\}) \cap \mathbb{K}(i)$ (with $s \geq 3$) such that $D_{k-1} \cap D_k \neq \emptyset$ for any $k \in \{2, \dots, s\}$, where we let $D_k = D_{\xi_k}$ and $\xi_1 = 0$, $\xi_s = \zeta$.

As in the beginning of §4.2, we have connection constants $\varpi_{j,2} \in \mathbb{C}$ such that

$$f(z) = \sum_{j=1}^{\mu} \varpi_{j,2} g_{j,\xi_2}(z) \quad (4.3)$$

for any $z \in \mathcal{D}$. In the same way, for any $z \in \mathcal{D}$, any $k \in \{3, \dots, s\}$ and any $j \in \{1, \dots, \mu\}$ we have

$$g_{j,\xi_{k-1}}(z) = \sum_{\ell=1}^{\mu} \varpi_{j,k,\ell} g_{\ell,\xi_k}(z). \quad (4.4)$$

Obviously the connection constants $\varpi_j \in \mathbb{C}$ in Theorem 7 are obtained by making products of the vector $(\varpi_{j,2})_{1 \leq j \leq \mu}$ and the matrices $(\varpi_{j,k,\ell})_{1 \leq j, \ell \leq \mu}$ (for $k \in \{3, \dots, s\}$), because $g_{j,\xi_s}(z) = g_j(z)$. Since $\mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$ is a ring and $R \geq 1$ can be any real number, Theorem 7 follows from the fact that all constants $\varpi_{j,2}$ and $\varpi_{j,k,\ell}$ in (4.3) and (4.4) belong to $\mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$. We will prove it now for (4.4); the proof is similar for (4.3).

Let $k \in \{3, \dots, s\}$ and $j \in \{1, \dots, \mu\}$. We differentiate $\mu - 1$ times Equation (4.4), so that we get the μ equations

$$g_{j,\xi_{k-1}}^{(s)}(z) = \sum_{\ell=1}^{\mu} \varpi_{j,k,\ell} g_{\ell,\xi_k}^{(s)}(z), \quad s = 0, \dots, \mu - 1.$$

We choose $z = \rho_k \in D_{k-1} \cap D_k \cap \mathbb{K}(i)$ outside the poles of $a_{\mu-1}(z)$ (with the notation of §4.3). Doing so yields a system of μ linear equations in the μ unknowns $\varpi_{j,k,\ell}$, $\ell = 1, \dots, \mu$, which can be solved using Cramer's rule because the determinant of the system (namely $W(\rho_k)$, where $W(z)$ is the wronskian of L built on the basis of solutions $g_{1,\xi_k}(z), \dots, g_{\mu,\xi_k}(z)$) does not vanish, by Lemma 9. Using again Lemma 9, we have $W(\rho_k) \in \overline{\mathbb{Q}}^\star$ and therefore $\frac{1}{W(\rho_k)} \in \overline{\mathbb{Q}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$ by Lemma 7. Now Cramer's rule yields the following expression for $\varpi_{j,k,\ell}$:

$$\frac{1}{W(\rho_k)} \begin{vmatrix} g_{1,\xi_k}(\rho_k) & \cdots & g_{\ell-1,\xi_k}(\rho_k) & g_{j,\xi_{k-1}}(\rho_k) & g_{\ell+1,\xi_k}(\rho_k) & \cdots & g_{\mu,\xi_k}(\rho_k) \\ g_{1,\xi_k}^{(1)}(\rho_k) & \cdots & g_{\ell-1,\xi_k}^{(1)}(\rho_k) & g_{j,\xi_{k-1}}^{(1)}(\rho_k) & g_{\ell+1,\xi_k}^{(1)}(\rho_k) & \cdots & g_{\mu,\xi_k}^{(1)}(\rho_k) \\ \vdots & \cdots & \vdots & \vdots & \vdots & \cdots & \vdots \\ g_{1,\xi_k}^{(\mu-1)}(\rho_k) & \cdots & g_{\ell-1,\xi_k}^{(\mu-1)}(\rho_k) & g_{j,\xi_{k-1}}^{(\mu-1)}(\rho_k) & g_{\ell+1,\xi_k}^{(\mu-1)}(\rho_k) & \cdots & g_{\mu,\xi_k}^{(\mu-1)}(\rho_k) \end{vmatrix}.$$

Since $\rho_k \in D_{k-1} \cap D_k$, the entries in this determinant belong to the ring $\mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$ (as noticed above), so that $\varpi_{j,k,\ell} \in \mathbf{G}_{R,\mathbb{K}(i)}^{\text{cv}}$. This concludes the proof of Theorem 7.

5. Proof of Theorem 4

The main part in the proof of Theorem 4 is to prove that $\mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$; this will be done below. We deduce Theorem 4 from this inclusion as follows, by Lemmas 2 and 3. If $\mathbb{K} \not\subset \mathbb{R}$, we have

$$\mathbf{G}_{\mathbb{K}}^{\text{a.c.}} \subset \mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$$

and Theorem 4 follows. If $\mathbb{K} \subset \mathbb{R}$, we have:

$$\mathbf{G}_{\mathbb{K}}^{\text{cv}} \subset \mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}} \cap \mathbb{R} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}} \cap \mathbb{R} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{cv}}$$

so that $\mathbf{G}_{\mathbb{K}}^{\text{cv}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}}$. The inclusion $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}} \subset \mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$ is trivial; let us prove that $\mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}} \subset \mathbf{G}_{\mathbb{K}}^{\text{a.c.}}$. Let $\xi_1, \xi_2 \in \mathbf{G}_{\mathbb{Q}}^{\text{cv}}$, and f, g, h be G -functions with rational coefficients and radii of convergence > 2 such that $f(1) = \xi_1$, $g(1) = \xi_2$, and $h(1) = \sqrt[4]{2}$. Then $k(z) = f(z) + g(z)h(z)\sqrt[4]{1 - \frac{z}{2}}$ is a G -function with coefficients in $\mathbb{Q} \subset \mathbb{K}$, and $\xi_1 + i\xi_2$ is the value at 1 of an analytic continuation of k (obtained after a small loop around $z = 2$). This concludes the proof that $\mathbf{G}_{\mathbb{K}}^{\text{a.c.}} = \mathbf{G}_{\mathbb{Q}}^{\text{cv}} + i\mathbf{G}_{\mathbb{Q}}^{\text{cv}}$ if $\mathbb{K} \subset \mathbb{R}$.

The rest of the section is devoted to the proof that $\mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}} \subset \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$. Let $\xi \in \mathbf{G}_{\overline{\mathbb{Q}}}^{\text{a.c.}}$; we may assume $\xi \neq 0$. There exists a G -function $f(z) = \sum_{n=0}^{\infty} a_n z^n$ with coefficients $a_n \in \overline{\mathbb{Q}}$, and $z_0 \in \overline{\mathbb{Q}}$, such that ξ is one of the values at z_0 of the multivalued analytic continuation of f . Replacing $f(z)$ with $f(z_0 z)$, we may assume $z_0 = 1$.

Let L denote the minimal differential equation satisfied by f , and ξ_1, \dots, ξ_p be the singularities of L . To keep the notation simple (and because the general case can be proved along the same lines), we shall assume that there is an open subset $\mathcal{D} \subset \mathbb{C}$ (as in §4.1) such that $1 \in \mathcal{D} \cup \{\xi_1, \dots, \xi_p\}$ and $\xi = f(1)$, where f denotes the analytic continuation of the G -function $\sum a_n z^n$ to \mathcal{D} . If 1 is a singularity of L then $f(1)$ is the (necessarily finite) limit of $f(z)$ as $z \rightarrow 1$, $z \in \mathcal{D}$.

The coefficients a_n ($n \geq 0$) belong to a number field $\mathbb{K} = \mathbb{Q}(\beta)$ for some primitive element β of degree d say. We can assume without loss of generality that \mathbb{K} is a Galois extension of \mathbb{Q} , i.e., that all Galois conjugates of β are in \mathbb{K} . There exist d sequences of rational numbers $(u_{j,n})_{n \geq 0}$, $j = 0, \dots, d-1$, such that, for all $n \geq 0$, $a_n = \sum_{j=0}^{d-1} u_{j,n} \beta^j$ and thus (at least formally)

$$f(z) = \sum_{n=0}^{\infty} a_n z^n = \sum_{j=0}^{d-1} \beta^j \sum_{n=0}^{\infty} u_{j,n} z^n. \quad (5.1)$$

The power series $U_j(z) = \sum_{n=0}^{\infty} u_{j,n} z^n$ are G -functions (see [17], Proposition VIII.1.4, p. 266), so that Equation (5.1) holds as soon as $|z|$ is sufficiently small. Moreover U_j has rational coefficients, so that it satisfies a differential equation with coefficients in $\mathbb{Q}[z]$ (see for instance [17], Proposition VIII.2.1 (iv), p. 268). We let L_j denote a minimal one, of order μ_j . Let \mathcal{S}_j denote the set of singularities of L_j , and $\mathcal{S} = \mathcal{S}_0 \cup \dots \cup \mathcal{S}_{d-1}$. Let Γ denote a compact broken line without multiple points from 0 to 1 inside $\mathcal{D} \cup \{0, 1\}$. Since \mathcal{S} is a finite set, we may assume that $\Gamma \cap \mathcal{S} \subset \{0, 1\}$ and find a (small) simply connected open subset $\Omega \subset \mathbb{C}$ such that $\Gamma \setminus \{0, 1\} \subset \Omega \subset \mathcal{D} \setminus \{1\}$ and $\Omega \cap \mathcal{S} = \emptyset$. If Γ and Ω are chosen appropriately, it is possible to construct $\mathcal{D}_0, \dots, \mathcal{D}_{d-1}$ as in §4.1 (with respect to L_0, \dots, L_{d-1}) such that $\Omega \subset \mathcal{D}_0 \cap \dots \cap \mathcal{D}_{d-1}$. Since Ω is simply connected and $1 \notin \Omega$, we choose a continuous determination of $\log(1-z)$ for $z \in \Omega$. Now Equation (5.1) holds in a neighborhood of 0, and 0 lies in the closure of Ω so that, by analytic continuation,

$$f(z) = \sum_{j=0}^{d-1} \beta^j U_j(z) \text{ for any } z \in \Omega. \quad (5.2)$$

We shall now expand this equality around the point 1, which lies also in the closure of Ω . For any $j \in \{0, \dots, d-1\}$, let $(g_{j,1}, \dots, g_{j,\mu_j})$ denote a basis of solutions of the differential equation $L_j y = 0$ provided by Theorem 6 with $\zeta = 1$. Then Theorem 7 gives $\varpi_{j,1}, \dots, \varpi_{j,\mu_j} \in \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$ such that $U_j(z) = \varpi_{j,1} g_{j,1}(z) + \dots + \varpi_{j,\mu_j} g_{j,\mu_j}(z)$ for any $z \in \Omega$. Since $\beta^j \in \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$ by Lemma 7, Equation (5.2) yields finite subsets $S \subset \mathbb{N}$ and $T \subset \mathbb{Q}$ such that, for $z \in \Omega$ sufficiently close to 1,

$$f(z) = \sum_{s \in S} \sum_{t \in T} (\log(1-z))^s (1-z)^t F_{s,t}(1-z)$$

where the functions $F_{s,t}(z)$ are holomorphic at 0 and have Taylor coefficients at 0 in $\mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$. Then Lemma 5 gives $c \in \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$, $\sigma \in \mathbb{N}$ and $\tau \in \mathbb{Q}$ such that $f(z) = c (\log(1-z))^\sigma (1-z)^\tau (1+o(1))$ as $z \rightarrow 1$ with $z \in \Omega$. Since $\lim_{z \rightarrow 1} f(z) = \xi \neq 0$, we have $\sigma = \tau = 0$ and $\xi = c \in \mathbf{G}_{\mathbb{Q}(i)}^{\text{cv}}$. This concludes the proof of Theorem 4.

6. Rational approximations to quotients of values of G -functions

This section is devoted to the proof of Theorem 5: in §6.1 we prove that (i) \Rightarrow (iii), and in §6.2 that (ii) \Rightarrow (i). Since (iii) obviously implies (ii), this will conclude the proof.

6.1. Construction of rational approximants. Assume that assertion (i) holds. Let $\xi_1, \xi_2 \in \mathbf{G}_{\mathbb{K}}^{\text{cv}} \setminus \{0\}$ be such that $\xi = \xi_1/\xi_2$. Let $R \geq 1$, and $U(z) = \sum_{n=0}^{\infty} u_n z^n$, $V(z) = \sum_{n=0}^{\infty} v_n z^n$ be G -functions with coefficients in \mathbb{K} and radii of convergence $> R$, such that $U(1) = \sum_{n=0}^{\infty} u_n = \xi_1$ and $V(1) = \sum_{n=0}^{\infty} v_n = \xi_2$.

For any $n \geq 0$, let $a_n = \sum_{k=0}^n u_k$ and $b_n = \sum_{k=0}^n v_k$, $A(z) = \sum_{n=0}^{\infty} a_n z^n$ and $B(z) = \sum_{n=0}^{\infty} b_n z^n$. Then $A(z) = U(z) \sum_{n=0}^{\infty} z^n = \frac{U(z)}{1-z}$ and $B(z) = \frac{V(z)}{1-z}$ are G -functions with coefficients in \mathbb{K} and radii of convergence $= 1$. Moreover $\lim_{n \rightarrow +\infty} a_n = \xi_1$ and $\lim_{n \rightarrow +\infty} b_n = \xi_2$ so that $a_n, b_n \neq 0$ for any n sufficiently large, and

$$|a_n - \xi b_n| = |(a_n - \xi_1) - \xi(b_n - \xi_2)| \leq \sum_{k=n+1}^{\infty} |u_k| + |\xi| \sum_{k=n+1}^{\infty} |v_k| = \mathcal{O}(R^{-n})$$

because $u_n, v_n = \mathcal{O}(R^{-n})$ as $n \rightarrow +\infty$ and we may assume $R \geq 2$. Therefore $A(z) - \xi B(z)$ has radius of convergence $\geq R$, thereby concluding the proof that (i) \Rightarrow (iii).

6.2. Application of Singularity Analysis. Let us prove that (ii) \Rightarrow (i) in Theorem 5.

Let $A(z) = \sum_{n=0}^{\infty} a_n z^n$ and $B(z) = \sum_{n=0}^{\infty} b_n z^n$ be G -functions with coefficients in \mathbb{K} , such that $b_n \neq 0$ for infinitely many n and $a_n - \xi b_n = o(b_n)$. Since $\xi \neq 0$, we have $a_n \neq 0$ for infinitely many n : none of $A(z)$ and $B(z)$ is a polynomial. Therefore these G -functions have finite positive radii of convergence, say ρ and $\tilde{\rho}$ respectively.

Let us denote by L the minimal differential equation over $\mathbb{K}[z]$ satisfied by $A(z)$, and by $\rho\zeta_1, \dots, \rho\zeta_q$ the pairwise distinct singularities of $A(z)$ of modulus ρ (so that $|\zeta_1| = \dots = |\zeta_q| = 1$). Then we have $q \geq 1$, and all $\rho\zeta_i$ are singularities of L and are algebraic numbers.

Let $\theta_0 \in (-\pi/2, \pi/2)$ and $\Delta_0 = \{z \in \mathbb{C}, z = 1 \text{ or } \arg(z-1) \equiv \theta_0 \pmod{2\pi}\}$. For any $i \in \{1, \dots, q\}$, let $\Delta_i = \rho\zeta_i \Delta_0 = \{\rho\zeta_i z, z \in \Delta_0\}$. Denoting by $\xi_1 = \rho\zeta_1$,

..., $\xi_q = \rho\zeta_q$, ξ_{q+1} , ..., ξ_p the singularities of L , we may assume (by choosing θ_0 properly) that Δ_1 , ..., Δ_q and some appropriate half-lines Δ_{q+1} , ..., Δ_p satisfy the assumptions made at the beginning of §4.1, so that we can take $\mathcal{D} = \mathbb{C} \setminus (\Delta_1 \cup \dots \cup \Delta_p)$. Choosing arbitrary determinations for $\log(\rho\zeta_i)$ ($i = 1, \dots, q$), and also a continuous one for $\log z$ when $z \in \mathbb{C} \setminus \Delta_0$, we may define $\log(\rho\zeta_i - z)$ to be $\log(\rho\zeta_i) + \log\left(1 - \frac{z}{\rho\zeta_i}\right)$ for $z \in \mathcal{D}$ sufficiently close to $\rho\zeta_i$ (because $\frac{1}{\rho\zeta_i}\Delta_i = \Delta_0$). For any $i \in \{1, \dots, q\}$, Corollary 1 yields $c_i \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}} \setminus \{0\}$, $\sigma_i \in \mathbb{N}$ and $\tau_i \in \mathbb{Q}$ such that

$$\begin{aligned} A(z) &= c_i (\log(\rho\zeta_i - z))^{\sigma_i} (\rho\zeta_i - z)^{\tau_i} (1 + o(1)) \\ &= c_i (\rho\zeta_i)^{\tau_i} \left(\log\left(1 - \frac{z}{\rho\zeta_i}\right) \right)^{\sigma_i} \left(1 - \frac{z}{\rho\zeta_i}\right)^{\tau_i} (1 + o(1)) \end{aligned}$$

as $z \rightarrow \rho\zeta_i$ with $z \in \mathcal{D}$. Replacing $A(z)$ and $B(z)$ with their ℓ -th derivatives from the beginning, where ℓ is a sufficiently large integer, we may assume $\tau_1 < 0$ (because $\rho\zeta_1$ is a singularity of $A(z)$). Let $\tau = \min(\tau_1, \dots, \tau_q) < 0$, and σ denote the maximal value of σ_i among those indices i such that $\tau_i = \tau$. Let $g(z) = (\log(1 - z))^{\sigma} (1 - z)^{\tau}$ for $z \in \mathbb{C} \setminus \Delta_0$, and $d_i = c_i (\rho\zeta_i)^{\tau_i}$ if $(\sigma_i, \tau_i) = (\sigma, \tau)$, $d_i = 0$ otherwise. Then $(d_1, \dots, d_q) \neq (0, \dots, 0)$ and, for any $i \in \{1, \dots, q\}$, we have $d_i \in \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$ (by Lemma 7, because $\rho\zeta_i \in \bar{\mathbb{Q}}$). Finally,

$$A(z) = d_i g\left(\frac{z}{\rho\zeta_i}\right) + o\left(g\left(\frac{z}{\rho\zeta_i}\right)\right) \quad (6.1)$$

as $z \rightarrow \rho\zeta_i$ with $z \in \mathcal{D}$. We have checked all assumptions of Theorem VI.5 (§VI.5, p. 398) of [20] (see also [21]). This result enables one to transfer this estimate (6.1) around the singularities on the circle of convergence into an asymptotic estimate for the coefficients of $A(z)$, namely

$$a_n = \frac{(-1)^{\sigma}}{\Gamma(-\tau)} \cdot \frac{(\log n)^{\sigma}}{\rho^n n^{\tau+1}} \cdot (\chi_n + o(1)), \quad \text{with } \chi_n = \sum_{i=1}^q d_i \zeta_i^{-n}. \quad (6.2)$$

Remark. Equation (6.2), the proof of which is based on Singularity Analysis, seems to be interesting for itself (and not only as a step in the proof of Theorem 5).

The same arguments with $B(z)$ provide $\tilde{\rho}, \tilde{\sigma}, \tilde{\tau}, \tilde{\zeta}_1, \dots, \tilde{\zeta}_{\tilde{q}}, \tilde{d}_1, \dots, \tilde{d}_{\tilde{q}}$ such that

$$b_n = \frac{(-1)^{\tilde{\sigma}}}{\Gamma(-\tilde{\tau})} \cdot \frac{(\log n)^{\tilde{\sigma}}}{\tilde{\rho}^n n^{\tilde{\tau}+1}} \cdot (\tilde{\chi}_n + o(1)), \quad \text{with } \tilde{\chi}_n = \sum_{i=1}^{\tilde{q}} \tilde{d}_i \tilde{\zeta}_i^{-n}. \quad (6.3)$$

Let $\mathcal{N}_0 = \{n \in \mathbb{N}, b_n = 0\}$ and $\mathcal{N} = \mathbb{N} \setminus \mathcal{N}_0$. By assumption \mathcal{N} is infinite, and $a_n = 0$ for any $n \in \mathcal{N}_0$ sufficiently large. In what follows, we assume implicitly

that \mathcal{N}_0 is infinite (otherwise the proof is the same, and even easier since everything works as if $\mathcal{N}_0 = \emptyset$ and $\mathcal{N} = \mathbb{N}$).

By Equations (6.2) and (6.3), we have as $n \rightarrow +\infty$ with $n \in \mathcal{N}$,

$$\frac{a_n}{b_n} = (-1)^{\sigma-\tilde{\sigma}} \frac{\Gamma(-\tilde{\tau})}{\Gamma(-\tau)} \cdot \frac{\chi_n + o(1)}{\tilde{\chi}_n + o(1)} \cdot \left(\frac{\tilde{\rho}}{\rho}\right)^n n^{\tilde{\tau}-\tau} (\log n)^{\sigma-\tilde{\sigma}}. \quad (6.4)$$

Now the left-hand side tends to $\xi \neq 0$ as $n \rightarrow +\infty$ with $n \in \mathcal{N}$. If $(\rho, \sigma, \tau) \neq (\tilde{\rho}, \tilde{\sigma}, \tilde{\tau})$ then $\left|\frac{\chi_n + o(1)}{\tilde{\chi}_n + o(1)}\right|$ tends to 0 or $+\infty$ as $n \rightarrow +\infty$ with $n \in \mathcal{N}$. Since both χ_n and $\tilde{\chi}_n$ are bounded, this implies that χ_n or $\tilde{\chi}_n$ tends to 0 as $n \rightarrow +\infty$ with $n \in \mathcal{N}$. Since $\chi_n = o(1)$ and $\tilde{\chi}_n = o(1)$ as $n \rightarrow \infty$ with $n \in \mathcal{N}_0$ (using (6.2) and (6.3), because $a_n = b_n = 0$ for $n \in \mathcal{N}_0$ sufficiently large), we have $\lim_{n \rightarrow +\infty} \chi_n = 0$ or $\lim_{n \rightarrow +\infty} \tilde{\chi}_n = 0$. By Lemma 6 this implies $d_1 = \dots = d_q = 0$ or $\tilde{d}_1 = \dots = \tilde{d}_{\tilde{q}} = 0$, which is a contradiction.

Therefore we have $(\rho, \sigma, \tau) = (\tilde{\rho}, \tilde{\sigma}, \tilde{\tau})$ in Equation (6.4), so that $\frac{a_n}{b_n} = \frac{\chi_n + o(1)}{\tilde{\chi}_n + o(1)}$ as $n \rightarrow +\infty$ with $n \in \mathcal{N}$. Therefore $\frac{\chi_n - \xi \tilde{\chi}_n + o(1)}{\tilde{\chi}_n + o(1)} = \frac{a_n}{b_n} - \xi$ tends to 0 as $n \rightarrow +\infty$ with $n \in \mathcal{N}$. Since $\tilde{\chi}_n$ is bounded, we deduce $\lim_{n \rightarrow +\infty} \chi_n - \xi \tilde{\chi}_n = 0$ (using the fact that $\chi_n = o(1)$ and $\tilde{\chi}_n = o(1)$ as $n \rightarrow \infty$ with $n \in \mathcal{N}_0$). Writing $\chi_n - \xi \tilde{\chi}_n = \sum_{j=1}^t \kappa_j \omega_j^n$ where $\{\omega_1, \dots, \omega_t\} = \{\zeta_1^{-1}, \dots, \zeta_q^{-1}, \tilde{\zeta}_1^{-1}, \dots, \tilde{\zeta}_{\tilde{q}}^{-1}\}$ with $\omega_1, \dots, \omega_t$ pairwise distinct, Lemma 6 yields $\kappa_1 = \dots = \kappa_t = 0$. Reordering the ζ_j 's and the ω_k 's if necessary, we may assume that $d_1 \neq 0$ and $\omega_1 = \zeta_1^{-1}$. Then $\kappa_1 = d_1 - \xi \tilde{d}_i$ if there is a (necessarily unique) i such that $\omega_1 = \tilde{\zeta}_i^{-1}$, and $\kappa_1 = d_1$ otherwise. Since $\kappa_1 = 0 \neq d_1$, there is such an i and it satisfies $\tilde{d}_i \neq 0$ and $\xi = d_1 / \tilde{d}_i \in \text{Frac}(\mathbf{G}_{\mathbb{K}(i)}^{\text{cv}})$. If $\mathbb{K} \not\subset \mathbb{R}$ then $\mathbf{G}_{\mathbb{K}}^{\text{cv}} = \mathbf{G}_{\mathbb{K}(i)}^{\text{cv}}$ by Theorem 4; otherwise we have $\xi \in \mathbb{R} \cap \text{Frac}(\mathbf{G}_{\mathbb{Q}}^{\text{cv}}) = \text{Frac}(\mathbf{G}_{\mathbb{Q} \cap \mathbb{R}}^{\text{cv}}) = \text{Frac}(\mathbf{G}_{\mathbb{K}}^{\text{cv}})$ by Theorem 4 and Lemma 2. In both cases, this concludes the proof of Theorem 5.

7. Perspectives

7.1. Other classes of arithmetic power series. It is natural to wonder if the results presented in this paper can be adapted to other classes of arithmetic power series. The most natural class is that of E -functions, also introduced by Siegel in [30]. The definition of these functions (see the Introduction) is formally similar to that of G -functions, but of course the presence of $n!$ at the denominator of the Taylor coefficients changes drastically the properties of E -functions. An E -function is entire and André proved in Theorem 4.3 of [4] that any E -function is solution of a linear differential equation with polynomial coefficients (not necessarily minimal) whose singularities are 0 (a regular singularity with rational exponents) and infinity (an irregular singularity in general). Like the set of G -functions, the set of E -functions enjoys certain stability properties; for instance, it is a ring.

Let us denote by \mathbf{E} as the set of all values of E -functions at algebraic points. This is the analogue of \mathbf{G} and it is a ring; it would be interesting to prove a result on \mathbf{E} analogous to Theorem 1. However we do not even know what a reasonable conjecture would be in this respect; what is clear is that the situation is really different, as the following result shows (we are indebted to the referee for suggesting its proof to us).

Proposition 2. *Let f be an E -function with coefficients in $\mathbb{Q}(i)$, and $\alpha \in \bar{\mathbb{Q}}$ be such that $f(1) = \alpha$ or $f(1) = e^\alpha$. Then $\alpha \in \mathbb{Q}(i)$.*

Proof. Let $\phi(z)$ denote either α or $e^{\alpha z}$, with $\alpha \in \bar{\mathbb{Q}}$; assume there exists an E -function f with coefficients in $\mathbb{Q}(i)$ such that $f(1) = \phi(1)$. Replacing $f(z)$ with $f(z) - \beta$ or $f(z)e^{-\beta z}$ for a suitable $\beta \in \mathbb{Q}(i)$, we may assume that α has zero trace over $\mathbb{Q}(i)$. Now there exist $\bar{\mathbb{Q}}(z)$ -linearly independent E -functions f_1, \dots, f_n with coefficients in $\mathbb{Q}(i)$ such that $f_1(1) = \phi(1)$ and the vector $\underline{f} = {}^t(f_1, \dots, f_n)$ is a solution of the differential system $\underline{y}' = A\underline{y}$ where A is an $n \times n$ matrix with entries in $\mathbb{Q}(i)(z)$. Modifying f_1, \dots, f_n if necessary as in the proof of Theorem 1.5 of [11], we may assume that 1 is not a pole of an entry of A . Using Beukers' version of Siegel–Shidlovskii's theorem (namely Theorem 1.3 of [11]), the relation $f_1(1) = \phi(1)$ can be lifted to $P_1(z)f_1(z) + \dots + P_n(z)f_n(z) = P_0(z)\phi(z)$ with $P_0, \dots, P_n \in \bar{\mathbb{Q}}[z]$ such that $P_0(1) = P_1(1) = 1$ and $P_2(1) = \dots = P_n(1) = 0$.

If $\phi(z) = \alpha$, taking the trace over $\mathbb{Q}(i)$ yields $Q_0, \dots, Q_n \in \mathbb{Q}(i)[z]$ such that $Q_1(z)f_1(z) + \dots + Q_n(z)f_n(z) = Q_0(z)$ with $Q_1(1) = 1$, $Q_2(1) = \dots = Q_n(1) = 0$, and $Q_0(1) = 0$ since α has zero trace. Therefore $f_1(1) = 0$, and $\alpha = 0$.

If $\phi(z) = e^{\alpha z}$, we take the norm over $\mathbb{Q}(i)$ of the relation $P_1(z)f_1(z) + \dots + P_n(z)f_n(z) = P_0(z)e^{\alpha z}$. Letting d denote the degree of a finite Galois extension of $\mathbb{Q}(i)$ which contains α and all coefficients of P_0, \dots, P_n , this provides (since α has zero trace) a relation $\sum_{\kappa} Q_{\kappa}(z)f_{\kappa}(z) = Q_0(z)$ where $Q_0 \in \mathbb{Q}(i)[z]$, $\kappa = (\kappa_1, \dots, \kappa_n) \in \mathbb{N}^n$ is such that $\kappa_1 + \dots + \kappa_n = d$, $f_{\kappa}(z) = f_1(z)^{\kappa_1} \dots f_n(z)^{\kappa_n}$, and $Q_{\kappa}(z) \in \bar{\mathbb{Q}}[z]$ is such that $Q_{\kappa}(1) = 0$ for $\kappa \neq (d, 0, \dots, 0)$ and $Q_{(d, 0, \dots, 0)}(1) = 1$. Taking $z = 1$ yields $f_1(1)^d = Q_0(1) \in \mathbb{Q}(i)$ hence $e^\alpha \in \bar{\mathbb{Q}}$, so that $\alpha = 0$.

This concludes the proof of Proposition 2. \square

The possibility of a result analogous to Theorem 3 is also uncertain. It is easy to describe the limits of sequences A_n/B_n where $A_n, B_n \in \bar{\mathbb{Q}}$, $B_n \neq 0$ for all large enough n and $\sum_{n=0}^{\infty} A_n z^n$ and $\sum_{n=0}^{\infty} B_n z^n$ are E -functions. This is simply $\text{Frac}(\mathbf{G})$, because the series $\sum_{n=0}^{\infty} n! A_n z^n$ and $\sum_{n=0}^{\infty} n! B_n z^n$ are G -functions, and conversely if $\sum_{n=0}^{\infty} a_n z^n$ is a G -function, then $\sum_{n=0}^{\infty} \frac{a_n}{n!} z^n$ is an E -function. This can hardly be the analogue we seek. We now observe that given an E -function $f(z) = \sum_{n=0}^{\infty} A_n z^n$, the sequence p_n/q_n , with $p_n = \sum_{k=0}^n A_k$ and $q_n = 1$, tends to $f(1)$, but $\sum_{n=0}^{\infty} p_n z^n = \frac{f(z)}{1-z}$ is not an E -function and $\sum_{n=0}^{\infty} z^n = \frac{1}{1-z}$ is a G -function. Hence a result analogous to Theorem 3 and involving \mathbf{E} might be achieved by considering simultaneously E and G -functions. It is also possible that similar

questions might be easier to answer in the larger class of *arithmetic Gevrey series* introduced by André in [4], [5].

7.2. Possible applications to irrationality questions. The Diophantine theory of E -functions is well understood after the works of many authors, among which we may cite Siegel [30] and Shidlovskii [29], and more recently André [5] and Beukers [11]. An E -function essentially takes transcendental values at all non-zero algebraic points, and the algebraic points where it may take an algebraic value are fully controlled *a priori*.

This is far from being true for a non-algebraic G -function. There are many examples in the literature of G -functions taking algebraic values at some algebraic points without an obvious reason, see for example [10]. After the pioneering works of Galochkin [22] and Bombieri [12], it is known that, given a transcendental G -function f , if α is a non-zero algebraic number of modulus $\leq c$, then $f(\alpha)$ cannot be an algebraic number of degree $\leq d$. Here, $c > 0$ and $d \geq 1$ are explicit quantities that depend on f and on the degree and height of α . A typical example is that if $\alpha = 1/q$ is the inverse of an integer, then $f(\alpha)$ is an irrational number provided that $|q| \geq Q$ is sufficiently large in terms of f . An important issue is that the constant c is usually much smaller than the radius of convergence of f : the point where the value is taken has to be very close to 0.

On the contrary, a few results are known in which such a restriction is not necessary. One of them is Wolfart's theorem [33] on transcendence of values of Gauss' hypergeometric function at algebraic points. Another, more related to the present paper, is Apéry's proof of the irrationality of $\zeta(3)$; it involves evaluating a G -function on the border of its disk of convergence. The starting point of his method is given by Theorem 5: he constructs two sequences $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$ of rational numbers, whose generating functions are G -functions⁶, such that a_n/b_n tends to $\zeta(3)$. To prove irrationality, more is needed, i.e., one also has to find a suitable common denominator D_n of a_n and b_n , and then prove that the linear form $D_n a_n + D_n b_n \zeta(3) \in \mathbb{Z} + \mathbb{Z} \zeta(3)$ tends to 0 without being equal to 0. (In this case, $D_n = \text{lcm}(1, 2, \dots, n)^3$.) The growth of D_n is usually the main problem in attempts at proving irrationality in Apéry's style. Indeed, there exist many examples of values $f(\alpha)$ of a G -function f at an algebraic point α having approximations in the sense of Theorem 3 (iii) (see [28] for references), but the growth of the relevant denominators D_n prevents one to prove irrationality when the modulus of α is too close to the radius of convergence of f . For instance, this approach has failed so far to establish the irrationality of $\zeta(5)$ or of Catalan's constant $G = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2}$.

In the following proposition, we explain in details how the growth of D_n , the radii of convergence and the irrationality exponent $\mu(\xi)$ of ξ are connected. Recall that $\mu(\xi)$ is the supremum of the set of real numbers μ such that, for infinitely many

⁶This was apparently first observed by Dwork in [16]; see also [18], §1.10, for references.

fractions p/q , $|\xi - p/q| < q^{-\mu}$. In particular ξ is said to be a Liouville number if $\mu(\xi) = +\infty$.

Proposition 3. *Let $\xi \in \mathbf{G} \cap \mathbb{R}$. Let $A(z) = \sum_{n=0}^{\infty} a_n z^n$ and $B(z) = \sum_{n=0}^{\infty} b_n z^n$ be G -functions, with rational coefficients and radii of convergence $= r > 0$, such that $A(z) - \xi B(z)$ has a finite radius of convergence, which is $\geq R > r$. Let $C \geq 1$ be such that a_n and b_n have a common denominator $\leq C^{n(1+o(1))}$ (as $n \rightarrow +\infty$). Then:*

- If $C < R$ then $\xi \notin \mathbb{Q}$ and $\mu(\xi) \leq 1 - \frac{\log(C/r)}{\log(C/R)}$.
- Necessarily $C \geq \sqrt{Rr}$.

This proposition is analogous to the other ones used to bound $\mu(\xi)$ from above when small linear forms $a_n \xi - b_n$ are available; the main difference here is that we do not assume $\lim_{n \rightarrow \infty} |a_n \xi - b_n|^{1/n}$ to exist. We hope this proposition can be used to make some progress towards Conjecture 1 stated in the introduction; of course the difficult point is to construct the G -functions with a control upon the denominators of a_n and b_n (so that C is not too large).

We have considered here only the case of one number ξ , but G -functions also arise in proofs of linear independence, in the same way as in Apéry's, for instance concerning the irrationality [8], [27] of $\zeta(s)$ for infinitely many odd $s \geq 3$.

Proof of Proposition 3. The second assertion follows from the first one because $\mu(\xi) \geq 2$ for any $\xi \in \mathbb{R} \setminus \mathbb{Q}$. Let us prove the first one.

Let $p_n = D_n a_n \in \mathbb{Z}$ and $q_n = D_n b_n \in \mathbb{Z}$, where n is sufficiently large and $D_n \in \mathbb{Z}$ is such that $1 \leq D_n \leq C^n$ (increasing C slightly if necessary). Decreasing R slightly if necessary, we may assume that the radius of convergence of $A(z) - \xi B(z)$ is $> R$, so that $|q_n \xi - p_n| \leq (C/R)^n$ for any n sufficiently large. Since $C < R$ and $q_n \xi - p_n \neq 0$ for infinitely many n (because $A(z) - \xi B(z)$ has a finite radius of convergence), this implies $\xi \notin \mathbb{Q}$. Moreover there exists a non-trivial linear recurrence relation $P_0(n)u_n + P_1(n)u_{n+1} + \dots + P_r(n)u_{n+r} = 0$, with coefficients $P_j(n) \in \mathbb{Z}[n]$, satisfied by both sequences $(a_n)_{n \geq 0}$ and $(b_n)_{n \geq 0}$. We claim that for any n sufficiently large, the vectors $(p_n, q_n), (p_{n+1}, q_{n+1}), \dots, (p_{n+r}, q_{n+r})$ span the \mathbb{Q} -vector space \mathbb{Q}^2 . Using Lemma 3.2 in [23], this implies $\mu(\xi) \leq 1 - \frac{\log(C/r')}{\log(C/R)}$ for any $r' < r$, because $|p_n|, |q_n| \leq (C/r')^n$ for any n sufficiently large. To prove the claim we argue by contradiction, and assume (permuting $(p_n)_{n \geq 0}$ and $(q_n)_{n \geq 0}$ if necessary) that for some $\lambda \in \mathbb{Q}$ we have $q_k = \lambda p_k$ for any $k \in \{n, n+1, \dots, n+r\}$. Then the sequence $(b_i - \lambda a_i)_{i \geq n}$ satisfies the above-mentioned recurrence relation, and its first $r+1$ terms vanish. If n is sufficiently large then $P_r(i) \neq 0$ for any $i \geq n+r+1$ (because we may assume P_r to be non-zero), so that $q_i - \lambda p_i = b_i - \lambda a_i = 0$ for any $i \geq n$. Since $\lim_{i \rightarrow +\infty} q_i \xi - p_i = 0$ and $p_i \neq 0$ for infinitely many n , we deduce $\lambda \xi = 1$, in contradiction with the fact that $\xi \notin \mathbb{Q}$. \square

Bibliography

- [1] Y. André, *G-functions and geometry*. Aspects Math. E13. Friedr. Vieweg & Sohn, Braunschweig 1989. [Zbl 0688.10032](#) [MR 0990016](#)
- [2] Y. André, Théorie des motifs et interprétation géométrique des valeurs p -adiques de G -fonctions. In *Number theory*, Papers from the Séminaire de théorie des nombres de Paris 1992/1993, London Math. Soc. Lecture Notes Ser. 215, Cambridge University Press, Cambridge 1995, 37–60. [Zbl 0848.11034](#) [MR 1345171](#)
- [3] Y. André, G -fonctions et transcendance. *J. Reine Angew. Math.* **476** (1996), 95–125. [Zbl 0848.11033](#) [MR 1401697](#)
- [4] Y. André, Séries Gevrey de type arithmétique, I. Théorèmes de pureté et de dualité. *Ann. Math.* **151** (2000), 705–740. [Zbl 1037.11049](#) [MR 1765707](#)
- [5] Y. André, Séries Gevrey de type arithmétique, II. Transcendance sans transcendance. *Ann. Math.* **151** (2000), 741–756. [Zbl 1037.11050](#) [MR 1765708](#)
- [6] Y. André, *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*. Panor. Synthèses 17, Soc. Math. de France, Paris 2004. [Zbl 1060.14001](#) [MR 2115000](#)
- [7] R. Apéry, Irrationalité de $\zeta(2)$ et $\zeta(3)$. *Astérisque* **61** (1979), 11–13. [Zbl 0401.10049](#)
- [8] K. Ball and T. Rivoal, Irrationalité d’une infinité de valeurs de la fonction zêta aux entiers impairs. *Invent. Math.* **146** (2001), no. 1, 193–207. [Zbl 1058.11051](#) [MR 1859021](#)
- [9] F. Beukers, Padé-approximations in number theory. In *Padé approximation and its applications, Amsterdam 1980*, Lecture Notes in Math. 888, Springer, Berlin 1981, 90–99. [Zbl 0478.10016](#) [MR 0649087](#)
- [10] F. Beukers, Algebraic values of G -functions. *J. Reine Angew. Math.* **434** (1993), 45–65. [Zbl 0753.11024](#) [MR 1195690](#)
- [11] F. Beukers, A refined version of the Siegel-Shidlovskii theorem. *Ann. of Math.* **163** (2006), no. 1, 369–379. [Zbl 1133.11044](#) [MR 2195138](#)
- [12] E. Bombieri, On G -functions. In *Recent progress in analytic number theory*, Vol. 2, Academic Press Inc., London 1981, 1–67. [Zbl 0461.10031](#) [MR 0637359](#)
- [13] G. Christol, Fonctions hypergéométriques bornées. In *Groupe de travail d’analyse ultramétrique* 14 (1986–1987), Exposé no. 8, 16 p.
- [14] D. V. Chudnovsky and G. V. Chudnovsky, Applications of Padé approximations to Diophantine inequalities in values of G -functions. In *Number theory* (New York, 1983/84), Lecture Notes in Math. 1135, Springer, Berlin 1985, 9–51. [Zbl 0561.10016](#) [MR 0803349](#)
- [15] P. Dèbes and U. Zannier, Hilbert’s irreducibility theorem and G -functions. *Math. Ann.* **309** (1997), no. 3, 491–503. [Zbl 0892.11024](#) [MR 1474204](#)
- [16] B. Dwork, On Apéry’s differential operator. In *Groupe de travail d’analyse ultramétrique* 7-8 (1979–1981), Exposé no. 25, 6 pages. [Zbl 0493.12031](#) [MR 0628166](#)
- [17] B. Dwork, G. Gerotto and F. J. Sullivan, *An introduction to G -functions*. Ann. Math. Studies 133, Princeton University Press, Princeton, NJ, 1994. [Zbl 0830.12004](#) [MR 1274045](#)
- [18] S. Fischler, Irrationalité de valeurs de zêta (d’après Apéry, Rivoal, ...). Séminaire Bourbaki 2002–2003, exposé no. 910 (nov. 2002), *Astérisque* **294** (2004), 27–62. [Zbl 1101.11024](#) [MR 2111638](#)

- [19] S. Fischler and T. Rivoal, Approximants de Padé et séries hypergéométriques équilibrées. *J. Math. Pures Appl.* **82** (2003), no. 10, 1369–1394. [Zbl 1064.11053](#) [MR 2020926](#)
- [20] P. Flajolet and R. Sedgewick, *Analytic combinatorics*. Cambridge University Press, Cambridge 2009. [Zbl 1165.05001](#) [MR 2483235](#)
- [21] P. Flajolet and A. Odlyzko, Singularity analysis of generating functions. *SIAM J. Discrete Math.* **3** (1990), 216–240. [Zbl 0712.05004](#) [MR 1039294](#)
- [22] A. I. Galochkin, Lower bounds of polynomials in the values of a certain class of analytic functions. *Mat. Sb.* **95** (137) (1974), 396–417, 471 (in Russian). [Zbl 0311.10035](#) [MR 0357338](#)
- [23] M. Hata, Legendre type polynomials and irrationality measures. *J. Reine Angew. Math.* **407** (1990), 99–125. [Zbl 0692.10034](#) [MR 1048530](#)
- [24] E. Hille, *Ordinary differential equations in the complex domain*. Pure and Applied Mathematics, Monographs and Texts, Wiley, 1976. [Zbl 0343.34007](#) [MR 0499382](#)
- [25] N. M. Katz, Nilpotent connections and the monodromy theorem: Applications of a result of Turrittin. *Inst. Hautes Études Sci. Publ. Math.* **39** (1970), 175–232. [Zbl 0221.14007](#) [MR 0291177](#)
- [26] M. Kontsevich and D. Zagier, Periods. In *Mathematics Unlimited – 2001 and beyond*, Springer, Berlin 2001, 771–808. [Zbl 1039.11002](#) [MR 1852188](#)
- [27] T. Rivoal, La fonction zêta de Riemann prend une infinité de valeurs irrationnelles aux entiers impairs. *C. R. Acad. Sci. Paris Sér. I Math.* **331** (2000), no. 4, 267–270. [Zbl 0973.11072](#) [MR 1787183](#)
- [28] T. Rivoal, Approximations rationnelles des valeurs de la fonction Gamma aux rationnels : le cas des puissances. *Acta Arith.* **142** (2010), no. 4, 347–365. [Zbl 1264.11057](#) [MR 2640064](#)
- [29] A. B. Shidlovskii, Transcendental numbers. *De Gruyter Stud. Math.* **12**, Berlin 1989. [Zbl 0689.10043](#) [MR 1033015](#)
- [30] C. Siegel, Über einige Anwendungen diophantischer Approximationen. *Abhandlungen Akad. Berlin* **1** (1929), 70p. [Zbl 56.0180.05](#)
- [31] J. Stillwell, Eisenstein’s footnote. *Math. Intelligencer* **17** (1995), no. 2, 58–62. [Zbl 0870.12001](#) [MR 1336074](#)
- [32] B. Sturmfels, Solving algebraic equations in terms of \mathcal{A} -hypergeometric series. *Discrete Math.* **210** (2000), 171–181. [Zbl 0963.33010](#) [MR 1731613](#)
- [33] J. Wolfart, Werte hypergeometrischer Funktionen. *Invent. Math.* **92** (1988), 187–216. [Zbl 0649.10022](#) [MR 0931211](#)

Received January 9, 2012

Stéphane Fischler, Equipe d’Arithmétique et de Géométrie Algébrique, Université Paris-Sud, Campus d’Orsay, Bâtiment 425, 91405 Orsay, France
E-mail: stephane.fischler@math.u-psud.fr

Tanguy Rivoal, CNRS et Université Grenoble 1, 100 rue des maths, BP 74, 38402 St Martin d’Hères cedex, France
E-mail: tanguy.rivoal@ujf-grenoble.fr