

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 89 (2014)

Artikel: A -structure on Lagrangian Grassmannians
Autor: Albers, Peter / Frauenfelder, Urs / Solomon, Jake P.
DOI: <https://doi.org/10.5169/seals-515691>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 17.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A Γ -structure on Lagrangian Grassmannians

Peter Albers, Urs Frauenfelder and Jake P. Solomon

Abstract. For n odd the Lagrangian Grassmannian of \mathbb{R}^{2n} is a Γ -manifold.

Mathematics Subject Classification (2010). 53Dxx, 55Mxx.

Keywords. Lagrangian Grassmannian, Gamma manifolds.

1. Introduction and statement of the result

We denote by $(\mathbb{R}^{2n}, \omega)$ the standard symplectic vector space. The (unoriented) Lagrangian Grassmannian \mathcal{L} is the space of all Lagrangian subspaces of \mathbb{R}^{2n} . It is a homogeneous space

$$\mathcal{L} \cong \mathrm{U}(n)/\mathrm{O}(n),$$

see [AG01], [MS98]. Every Lagrangian subspace can be identified with the fixed point set of a linear orthogonal anti-symplectic involution. Using this identification, we define a smooth map

$$\Theta: \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$$

by

$$(R, S) \mapsto RSR,$$

which we think of as a product. On every space there are products such as constant maps and projections to one factor. In [Hop41] Hopf introduced the notion of Γ -manifolds which rules these trivial products out. The purpose of this paper is to prove that the above product gives the Lagrangian Grassmannian \mathcal{L} the structure of a Γ -manifold for n odd.

Definition 1.1. A closed, connected, orientable manifold M carries the structure of a Γ -manifold if there exists a map

$$\Theta: M \times M \rightarrow M$$

such that the maps

$$x \mapsto \Theta(x, y_0) \quad \text{and} \quad y \mapsto \Theta(x_0, y)$$

have non-zero mapping degree for one and thus all pairs $(x_0, y_0) \in M \times M$.

It is well known that \mathcal{L} is orientable if and only if n odd, see [Fuk68]. The main result of this article is the following theorem.

Theorem 1.2. *If n is odd, then (\mathcal{L}, Θ) is a Γ -manifold.*

Using Hopf's theorem ([Hop41], Satz 1), we get a new proof of the following corollary due to Fuks [Fuk68].

Corollary 1.3 ([Fuk68]). *For n odd, the rational cohomology ring of \mathcal{L} is an exterior algebra on generators of odd degree.*

Remark 1.4. The cohomology ring of the oriented and unoriented Lagrangian Grassmannian was computed by Borel and Fuks for all n , see [Bor53a], [Bor53b], [Fuk68]. A nice summary of these results can be found in Chapter 22 of the book by Vassilyev [Vas88].

The above situation fits into the following general framework. It is well known that \mathcal{L} embeds into $U(n)$ as the set $U(n) \cap \text{Sym}(n)$, i.e. the symmetric unitary matrices. Indeed the image of a Lagrangian subspace $\Lambda \subset \mathbb{C}^n$ is the symmetric unitary matrix $A_\Lambda := uu^t \in U(n) \cap \text{Sym}(n)$ where $a \in U(n)$ maps \mathbb{R}^n onto Λ . The unique orthogonal anti-symplectic involution with fixed point set Λ is then the map $A_\Lambda \circ \tau$ where $\tau: \mathbb{C}^n \rightarrow \mathbb{C}^n$ is complex conjugation. Thus, the Lagrangian Grassmannian \mathcal{L} can be interpreted as the fixed point set of the involutive anti-isomorphism $A \mapsto A^T$ of $U(n)$. On any Lie group G we can define a new product: $(g, h) \mapsto gh^{-1}g$. If $I: G \rightarrow G$ is an involutive anti-isomorphism then this new product restricts to a product on the fixed point set $\text{Fix}(I)$. This is precisely the situation for the Lagrangian Grassmannian, namely the map Θ corresponds under the embedding of \mathcal{L} into $U(n)$ to $(g, h) \mapsto gh^{-1}g$.

For general Lie groups this new product does not always give rise to a Γ -structure for various reasons. For example, if we take $G = O(n)$ resp. $G = U(n)$ and $I(A) := A^{-1}$, then $\text{Fix}(I)$ can be identified with $\bigcup_k G(k, n)$, the union of all real resp. complex Grassmannians, which is not connected. Another example is $G = SU(n)$ with $I =$ transposition. Then for $n = 2$ we can identify $\text{Fix}(I) \cong S^2$. But by Hopf's theorem ([Hop41], Satz 1) S^2 is not a Γ -manifold.

Acknowledgements. Parts of this article were written during a visit of the first two authors at the Forschungsinstitut für Mathematik (FIM), ETH Zürich. The authors thank the FIM for its stimulating working atmosphere.

This work is supported by the SFB 878 – Groups, Geometry and Actions (PA), by the Alexander von Humboldt Foundation (UF), by Israel Science Foundation grant 1321/2009 and by Marie Curie grant No. 239381 (JPS).

2. Proof of Theorem 1.2

We recall that the (unoriented) Lagrangian Grassmannian \mathcal{L} is the homogeneous space

$$\mathcal{L} \cong \mathrm{U}(n)/\mathrm{O}(n),$$

see [AG01], [MS98]. Since n is odd, \mathcal{L} is a closed connected orientable manifold [Fuk68]. The space \mathcal{L} is naturally identified with the space of linear orthogonal anti-symplectic involutions of \mathbb{R}^{2n} with the standard symplectic structure. Using this identification, we define the map

$$\Theta: \mathcal{L} \times \mathcal{L} \rightarrow \mathcal{L}$$

by $(R, S) \mapsto RSR$. In order to prove Theorem 1.2, it suffices to show for one choice of basepoint R_0 that the mapping degrees of

$$S \mapsto \Theta(R_0, S) \quad \text{and} \quad S \mapsto \Theta(S, R_0)$$

are non-zero. Since

$$S \mapsto \Theta(R_0, S) = R_0SR_0 \mapsto \Theta(R_0, R_0SR_0) = R_0R_0SR_0R_0 = S,$$

the first map is an involution and therefore has mapping degree ± 1 . The non-trivial case is to compute the mapping degree of

$$\Theta_0(S) := \Theta(S, R_0) = SR_0S.$$

Theorem 1.2 follows immediately from the following proposition.

Proposition 2.1. *The mapping degree of Θ_0 equals*

$$\deg \Theta_0 = 2^{m+1}$$

where $n = 2m + 1$.

Proof. Identify $\mathbb{R}^{2n} = \mathbb{C}^n$ in the standard way. Denote by $\tau: \mathbb{C}^n \rightarrow \mathbb{C}^n$ the map given by complex conjugation of all coordinates simultaneously. It is a standard fact, see for instance [MS98], that an orthogonal symplectic map $\mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ corresponds to a unitary map $\mathbb{C}^n \rightarrow \mathbb{C}^n$. It follows that an orthogonal anti-symplectic map

$R: \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ can be written as the composition $A \circ \tau: \mathbb{C}^n \rightarrow \mathbb{C}^n$ for A a unitary linear map. The condition $R^2 = \text{Id}$ then translates to $A\bar{A} = \text{Id}$. So, we identify

$$\mathcal{L} = \{A \in \text{U}(n) \mid A\bar{A} = \text{Id}\}.$$

Under this identification, the map Θ is given by

$$\Theta(A, B) = A\bar{B}A.$$

Let B_0 be the unitary matrix corresponding to R_0 . Then the map Θ_0 is given by

$$\Theta_0(A) = \Theta(A, B_0) = A\bar{B}_0A.$$

In the following, we take $B_0 = B$, the diagonal matrix with entries $b_{jk} = e^{i\theta_j} \delta_{jk}$ where

$$0 < \theta_j < 2\pi, \quad \theta_1 < \theta_2 < \dots < \theta_n.$$

For this choice of B_0 , we show that Id is a regular value of Θ_0 and compute the signed cardinality of $\Theta_0^{-1}(\text{Id})$.

Indeed, if $\Theta_0(A) = \text{Id}$, then $A\bar{B}A = \text{Id}$, and therefore $\bar{A}B = A$. Throughout this paper, we do *not* use the Einstein summation convention. Letting a_{jk} denote the matrix entries of A , we have

$$\bar{a}_{jk}e^{i\theta_k} = a_{jk}.$$

Write $a_{jk} = r_{jk}e^{i\psi_{jk}}$, where $r_{jk} \in \mathbb{R}$ and $0 \leq \psi_{jk} < \pi$. So,

$$e^{i2\psi_{jk}} = a_{jk}/\bar{a}_{jk} = e^{i\theta_k},$$

and therefore $\psi_{jk} = \theta_k/2$. Writing the unitary condition for A in terms of r_{jk} and ψ_{jk} , we have

$$\delta_{jl} = \sum_k a_{jk}\bar{a}_{lk} = \sum_k r_{jk}r_{lk}e^{i(\psi_{jk}-\psi_{lk})} = \sum_k r_{jk}r_{lk}.$$

Thus r_{jk} is an orthogonal matrix. Furthermore, the condition $A\bar{A} = \text{Id}$ translates to

$$\delta_{jl} = \sum_k a_{jk}\bar{a}_{kl} = \sum_k r_{jk}r_{kl}e^{i(\theta_k-\theta_l)/2}.$$

In particular, taking $j = l$, we obtain

$$1 = \sum_k r_{jk}r_{kj} \cos((\theta_k - \theta_j)/2).$$

Writing $r'_{jk} = \cos((\theta_k - \theta_j)/2)r_{jk}$, we can reformulate the preceding equation in terms of the inner product of the row and column vectors $r'_{j\cdot}$ and $r_{\cdot j}$. Namely,

$$r'_{j\cdot} \cdot r_{\cdot j} = 1. \tag{2.1}$$

On the other hand, since r_{jk} is unitary, $|r_{.j}| = 1$ and

$$|r'_{.j}|^2 = \sum_k r_{jk}^2 \cos^2((\theta_k - \theta_j)/2) \leq \sum_k r_{jk}^2 = |r_{.j}|^2 = 1,$$

with equality only if $r_{jk} = 0$ when $k \neq j$. Applying Cauchy–Schwartz to equation (2.1), we have

$$1 \leq |r'_{.j}| |r_{.j}| = |r'_{.j}| \leq 1.$$

Thus equality must hold, and the matrix r_{jk} is diagonal. Moreover, orthogonality implies that $r_{jk} = \pm \delta_{jk}$. Summing up, $A \in \Theta_0^{-1}(\text{Id})$ if and only if we have $A = A^\epsilon$, where

$$\epsilon = (\epsilon_1, \dots, \epsilon_n), \quad \epsilon_k \in \{0, 1\},$$

and A^ϵ is the matrix with elements

$$a_{jk}^\epsilon = e^{i(\theta_k/2 + \epsilon_k \pi)}.$$

In particular, $\Theta_0^{-1}(\text{Id})$ has unsigned cardinality 2^n .

It remains to show that Id is a regular value and compute the signs. Let $\text{Sym}(n)$ denote the space of real $n \times n$ symmetric matrices. It is easy to see that the tangent space to \mathcal{L} at $A = \text{Id}$ is given by

$$T_{\text{Id}} \mathcal{L} = \{T \in \mathfrak{u}(n) \mid T + \bar{T} = 0\} = \{iQ \mid Q \in \text{Sym}(n)\}.$$

Recall that $U(n)$ acts on \mathcal{L} by $A \mapsto UA\bar{U}^{-1}$. Thus, if $A = U\bar{U}^{-1}$, we have an isomorphism

$$\kappa_U : T_{\text{Id}} \mathcal{L} \xrightarrow{\sim} T_A \mathcal{L}$$

given by $T \mapsto UT\bar{U}^{-1}$. Since \mathcal{L} is a $U(n)$ homogeneous space, the isomorphism κ_U preserves orientation. Moreover, for $T \in T_{A^\epsilon} \mathcal{L}$ we have

$$d\Theta_0|_{A^\epsilon}(T) = T\bar{B}A^\epsilon + A^\epsilon\bar{B}T = T\bar{A}^\epsilon + \bar{A}^\epsilon T.$$

If $U^\epsilon \in U(n)$ satisfies

$$A^\epsilon = U^\epsilon(\bar{U}^\epsilon)^{-1},$$

then A^ϵ is a regular point of Θ_0 if the linear map

$$\alpha^\epsilon = d\Theta_0|_{A^\epsilon} \circ \kappa_U : T_{\text{Id}} \mathcal{L} \rightarrow T_{\text{Id}} \mathcal{L}$$

is invertible, and in that case the sign of A^ϵ is $\text{sign det}(\alpha^\epsilon)$. Explicitly,

$$\begin{aligned} \alpha^\epsilon(T) &= U^\epsilon T(\bar{U}^\epsilon)^{-1} \bar{A}^\epsilon + \bar{A}^\epsilon U^\epsilon T(\bar{U}^\epsilon)^{-1} \\ &= U^\epsilon T(U^\epsilon)^{-1} + \bar{U}^\epsilon T(\bar{U}^\epsilon)^{-1} \\ &= U^\epsilon T(U^\epsilon)^{-1} - \bar{U}^\epsilon \bar{T}(\bar{U}^\epsilon)^{-1} \\ &= 2i \text{Im}(U^\epsilon T(U^\epsilon)^{-1}). \end{aligned}$$

Writing $T = iQ$, we can think of α^ϵ as the map $\text{Sym}(n) \rightarrow \text{Sym}(n)$ given by

$$\alpha^\epsilon(Q) = 2 \operatorname{Re}(U^\epsilon Q (U^\epsilon)^{-1}).$$

For convenience, we take U^ϵ to be the unitary linear map given by

$$u_{jk}^\epsilon = e^{i(\theta_k/4 + \epsilon_k \pi/2)} \delta_{jk}.$$

Then, denoting by q_{jk} the matrix elements of Q , we have

$$\begin{aligned} \alpha^\epsilon(Q)_{jk} &= 2 \operatorname{Re} \left(e^{i((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2)} \right) q_{jk} \\ &= 2 \cos \left((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2 \right) q_{jk}. \end{aligned}$$

Since Q is a symmetric matrix, it is determined by q_{jk} for $j \leq k$. Thus

$$\det(\alpha^\epsilon) = \prod_{j \leq k} 2 \cos \left((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2 \right) q_{jk}.$$

We need to show that this determinant does not vanish and compute its sign. For $j = k$, clearly $\cos \left((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2 \right) = 1$. For $j < k$, by assumption, $0 < \theta_j < \theta_k < 2\pi$, so

$$-\frac{\pi}{2} < \frac{\theta_j - \theta_k}{4} < 0.$$

It follows that for all $j \leq k$, we have $\cos \left((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2 \right) \neq 0$. Therefore $\det(\alpha^\epsilon) \neq 0$ for all ϵ and Id is a regular value. Moreover,

$$\cos \left((\theta_j - \theta_k)/4 + (\epsilon_j - \epsilon_k)\pi/2 \right) < 0 \quad \text{if and only if} \quad \epsilon_j = 0, \epsilon_k = 1.$$

Let Υ_n be the set of all binary sequences $\epsilon = (\epsilon_1, \dots, \epsilon_n)$. For $\epsilon \in \Upsilon_n$ define $\text{sign}(\epsilon)$ to be the number modulo 2 of pairs $j < k$ such that $\epsilon_j = 0$ and $\epsilon_k = 1$. The upshot of the preceding calculations is that

$$\text{sign} \det(\alpha^\epsilon) = \text{sign}(\epsilon),$$

therefore

$$\deg \Theta_0 = \sum_{\epsilon \in \Upsilon_n} (-1)^{\text{sign}(\epsilon)}.$$

A combinatorial argument given below in Lemma 2.2 then implies the theorem. \square

Lemma 2.2. For $n = 2m + 1$, we have

$$d_n := \sum_{\epsilon \in \Upsilon_n} (-1)^{\text{sign}(\epsilon)} = 2^{m+1}.$$

Proof. Let M_n denote the number of $\epsilon \in \Upsilon_n$ such that $\text{sign}(\epsilon) = 0$. Then

$$d_n = M_n - (2^n - M_n) = 2M_n - 2^n.$$

For $\epsilon \in \Upsilon_n$ denote by $\text{par}(\epsilon)$ the parity of ϵ , or in other words the number modulo 2 of j such that $\epsilon_j = 1$. Let P_n denote the number of $\epsilon \in \Upsilon_n$ such that $\text{sign}(\epsilon) + \text{par}(\epsilon) = 0$. By analyzing what happens when we adjoin either 1 or 0 to the beginning of a sequence $\epsilon \in \Upsilon_{n-1}$, we find that

$$M_n = M_{n-1} + P_{n-1}, \quad P_n = (2^{n-1} - P_{n-1}) + M_{n-1}.$$

Iterating these recursions twice, we obtain

$$M_n = M_{n-2} + P_{n-2} + 2^{n-2} - P_{n-2} + M_{n-2} = 2M_{n-2} + 2^{n-2}.$$

Clearly $M_1 = 2$, so $d_1 = 2$. Using the preceding recursion for M_n , we obtain

$$d_n = 2(2M_{n-2} + 2^{n-2}) - 2^n = 2(2M_{n-2} - 2^{n-2}) = 2d_{n-2}.$$

The lemma follows by induction. □

References

- [AG01] V. I. Arnol'd and A. B. Givental, Symplectic geometry. In *Dynamical systems, IV*, Encyclopaedia Math. Sci. 4, second ed., Springer, Berlin 2001, 1–138. [Zbl 0780.58016](#) [MR 1866631](#)
- [Bor53a] A. Borel, La cohomologie mod 2 de certains espaces homogènes. *Comment. Math. Helv.* **27** (1953), 165–197. [Zbl 0052.40301](#) [MR 0057541](#)
- [Bor53b] A. Borel, Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts. *Ann. of Math. (2)* **57** (1953), 115–207. [Zbl 0052.40001](#) [MR 0051508](#)
- [Fuk68] D. B. Fuks, The Maslov–Arnol'd characteristic classes. *Dokl. Akad. Nauk SSSR* **178** (1968), 303–306; English transl. *Soviet Math. Dokl.* **9** (1968), 96–99. [Zbl 0175.20304](#) [MR 0225340](#)
- [Hop41] H. Hopf, Über die Topologie der Gruppen-Mannigfaltigkeiten und ihre Verallgemeinerungen. *Ann. of Math. (2)* **42** (1941), 22–52. [Zbl 67.0747.01](#) [MR 0004784](#)
- [MS98] D. McDuff and D. A. Salamon, *Introduction to symplectic topology*. Second ed., Oxford Math. Monogr., The Clarendon Press, Oxford University Press, New York 1998. [Zbl 0844.58029](#) [MR 1698616](#)
- [Vas88] V. A. Vassilyev, *Lagrange and Legendre characteristic classes*. Adv. Stud. Contemp. Math. 3, Gordon and Breach Science Publishers, New York 1988. [Zbl 0715.53001](#) [MR 1065996](#)

Received December 13, 2012

Peter Albers, Mathematisches Institut, Westfälische Wilhelms-Universität Münster,
Germany

E-mail: peter.albers@wwu.de

Urs Frauenfelder, Department of Mathematics and Research Institute of Mathematics, Seoul
National University, Korea

E-mail: frauenf@snu.ac.kr

Jake P. Solomon, Institute of Mathematics, Hebrew University of Jerusalem, Israel

E-mail: jake@math.huji.ac.il