Zeitschrift: Commentarii Mathematici Helvetici

Herausgeber: Schweizerische Mathematische Gesellschaft

Band: 90 (2015)

Heft: 2

Artikel: Finite-dimensionality and cycles on powers of K3 surfaces

Autor: Yin, Qizheng

DOI: https://doi.org/10.5169/seals-658060

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Finite-dimensionality and cycles on powers of K3 surfaces

Qizheng Yin

Abstract. For a K3 surface S, consider the subring of $CH(S^n)$ generated by divisor and diagonal classes (with \mathbb{Q} -coefficients). Voisin conjectures that the restriction of the cycle class map to this ring is injective. We prove that Voisin's conjecture is equivalent to the finite-dimensionality of S in the sense of Kimura–O'Sullivan. As a consequence, we obtain examples of S whose Hilbert schemes satisfy the Beauville–Voisin conjecture.

Mathematics Subject Classification (2010). 14C15, 14C25, 14J28.

Keywords. Chow groups, *K*3 surfaces, finite-dimensional motives, Beauville–Voisin conjecture.

1. Introduction

Let S be a smooth projective K3 surface over a field k. For $n \ge 1$, consider the \mathbb{Q} -subalgebra of the Chow ring $CH(S^n)$ with \mathbb{Q} -coefficients generated by (pull-backs of) divisor classes on S and the diagonal class on $S \times S$. We denote it by $R(S^n)$. Regarding its structure, Voisin made the following conjecture ([15], Conjecture 1.6).

Conjecture 1. For $n \ge 1$, the restriction of the cycle class map $cl: CH(S^n) \to H(S^n)$ to $R(S^n)$ is injective.

The case n=1 is the well-known result of Beauville and Voisin ([2], Theorem 1). Voisin also proved Conjecture 1 for $n \le 2b_{\rm tr} + 1$, where $b_{\rm tr} = 22 - \rho$ is the rank of the transcendental part of $H^2(S)$ ([15], Proposition 2.2). As is remarked in [16], Section 5.1, Conjecture 1 turns out to be rather strong. Notably it implies the finite-dimensionality of S in the sense of Kimura–O'Sullivan ([8]; see also Section 2.4), a conjecture that is widely open even for K3 surfaces.

The aim of this short note is to prove the converse: that finite-dimensionality suffices to deduce Conjecture 1.

Theorem. Conjecture 1 holds for S if and only if S is finite-dimensional.

More precisely, we prove that the relations found in [2] (see also Section 2.2) plus the one predicted by the finite-dimensionality of S generate all relations in $\operatorname{cl}(R(S^n))$. To achieve this, we reduce the problem to a manageable algebraic form, whose solution has long been known to algebraists (Hanlon and Wales [6]).

Further, let k be algebraically closed. As is shown by Voisin ([15], Proposition 2.5), Conjecture 1 implies the following conjecture for the Hilbert schemes $S^{[n]}$ of S (Conjecture 1.3 in *loc. cit.*; stated for $k = \mathbb{C}$ and often referred to as the Beauville–Voisin conjecture).

Conjecture 2. Let X be an irreducible holomorphic symplectic variety (hyper-Kähler manifold). Then the restriction of the cycle class map $cl: CH(X) \to H(X)$ to the \mathbb{Q} -subalgebra generated by divisor classes and Chern classes of the tangent bundle is injective.

We thus obtain an immediate consequence.

Corollary. If S is finite-dimensional, then Conjecture 2 holds for $S^{[n]}$ and for all $n \ge 1$.

Finally, we refer to [11] in characteristic 0 and [9] in positive characteristic for *K*3 surfaces known to be finite-dimensional. Among them are Kummer surfaces¹, surfaces of Picard rank 19, 20 and 22 (supersingular), and some sporadic cases of even Picard rank.

Notation and conventions. Throughout, Chow groups CH and Picard groups Pic are taken with \mathbb{Q} -coefficients. We fix a Weil cohomology theory H (e.g. singular cohomology when $k = \mathbb{C}$, or ℓ -adic cohomology in general). For a set A, we denote by $\mathbb{Q}A$ the free \mathbb{Q} -linear span of A.

Acknowledgements. Thanks to Mehdi Tavakol for explaining the result of Hanlon and Wales, to Claire Voisin and Rahul Pandharipande for useful discussions, and to the referee for comments and suggestions. This work was carried out in the group of Pandharipande at ETH Zürich, supported by grant ERC-2012-AdG-320368-MCSK.

2. Preliminaries

2.1. Let S be a smooth projective K3 surface over k. Denote by $o \in \mathrm{CH}^2(S)$ the distinguished class on S as in [2], Theorem 1 (e.g. the class of any point on a rational curve in S). Take a basis $\{L^s\}_{1 \leq s \leq \rho}$ of $\mathrm{Pic}(S)$, and write $l^s = c_1(L^s) \in \mathrm{CH}^1(S)$. For convenience we assume $\{l^s\}$ to be orthogonal. Further, denote by $\delta = [\Delta] \in \mathrm{CH}^2(S \times S)$ the class of the diagonal.

¹For Kummer surfaces, one can also prove Conjectures 1 and 2 by applying results on abelian varieties (*e.g.* [10], Corollary 9.4). We refer to [4] for a similar argument proving Conjecture 2 for generalized Kummer varieties.

Consider the projections $\operatorname{pr}_i : S^n \to S$ for $1 \le i \le n$, and $(\operatorname{pr}_i, \operatorname{pr}_j) : S^n \to S \times S$ for $1 \le i, j \le n$ and $i \ne j$. We write

$$o_i = \operatorname{pr}_i^*(o) \in \operatorname{CH}^2(S^n), \quad l_i^s = \operatorname{pr}_i^*(l^s) \in \operatorname{CH}^1(S^n),$$

and $\delta_{i,j} = (\operatorname{pr}_i, \operatorname{pr}_j)^*(\delta) \in \operatorname{CH}^2(S^n).$

The ring $R(S^n)$ is defined to be the \mathbb{Q} -subalgebra of $CH(S^n)$ generated by $\{o_i\}, \{l_i^s\}$ and $\{\delta_{i,j}\}.$

2.2. Relations in $R(S^n)$ **.** The following set of relations summarizes the main results of $[2]^2$ (namely Theorem 1 and Proposition 3.2 in *loc. cit.*)

$$o_i \cdot o_i = 0$$
, $l_i^s \cdot o_i = 0$, and $l_i^s \cdot l_i^s = \deg(l^s \cdot l^s)o_i$; (1)

$$\delta_{i,j} \cdot o_i = o_i \cdot o_j$$
, $\delta_{i,j} \cdot l_i^s = l_i^s \cdot o_j + o_i \cdot l_j^s$, and $\delta_{i,j} \cdot \delta_{i,j} = 24o_i \cdot o_j$; (2)

$$\delta_{i,j} \cdot \delta_{i,k} = \delta_{i,j} \cdot o_k + \delta_{i,k} \cdot o_j + \delta_{j,k} \cdot o_i - o_i \cdot o_j - o_i \cdot o_k - o_j \cdot o_k. \tag{3}$$

Note that (1), (2) and (3) involve 1, 2 and 3 factors of S^n respectively.

As we will see, it is both meaningful and convenient to replace $\delta_{i,j}$ (as generator of $R(S^n)$) by

$$\tau_{i,j} = \delta_{i,j} - o_i - o_j - \sum_{s=1}^{\rho} \frac{l_i^s \cdot l_j^s}{\deg(l^s \cdot l^s)} \in \mathrm{CH}^2(S^n).$$

Here τ stands for "transcendental". The relations above now appear in an even simpler form.

2.3. Lemma. In $R(S^n)$ we have relations

$$o_i \cdot o_i = 0$$
, $l_i^s \cdot o_i = 0$, and $l_i^s \cdot l_i^s = \deg(l^s \cdot l^s)o_i$; (4)

$$\tau_{i,j} \cdot o_i = 0, \quad \tau_{i,j} \cdot l_i^s = 0, \quad \text{and} \quad \tau_{i,j} \cdot \tau_{i,j} = b_{\text{tr}} o_i \cdot o_j;$$
 (5)

$$\tau_{i,j} \cdot \tau_{i,k} = \tau_{j,k} \cdot o_i, \tag{6}$$

where $b_{tr} = 22 - \rho$ is the rank of the transcendental part of $H^2(S)$.

²By lifting to characteristic 0, the results of [2] remain valid in positive characteristic.

506 Q. Yin CMH

Proof. The calculation is straightforward and we only do (6). By (2), (3) and (5), we get

$$\begin{aligned} \tau_{i,j} \cdot \tau_{i,k} &= \tau_{i,j} \cdot \left(\delta_{i,k} - o_i - o_k - \sum_{s=1}^{\rho} \frac{l_i^s \cdot l_k^s}{\deg(l^s \cdot l^s)} \right) \\ &= \tau_{i,j} \cdot \delta_{i,k} - \tau_{i,j} \cdot o_k \\ &= \left(\delta_{i,j} - o_i - o_j - \sum_{s=1}^{\rho} \frac{l_i^s \cdot l_j^s}{\deg(l^s \cdot l^s)} \right) \cdot \delta_{i,k} \\ &- \left(\delta_{i,j} - o_i - o_j - \sum_{s=1}^{\rho} \frac{l_i^s \cdot l_j^s}{\deg(l^s \cdot l^s)} \right) \cdot o_k \\ &= \delta_{i,j} \cdot \delta_{i,k} - o_i \cdot o_k - \delta_{i,k} \cdot o_j - \delta_{i,j} \cdot o_k \\ &+ o_i \cdot o_k + o_j \cdot o_k - \left(\sum_{s=1}^{\rho} \frac{l_j^s \cdot l_k^s}{\deg(l^s \cdot l^s)} \right) \cdot o_i \\ &= \delta_{j,k} \cdot o_i - o_i \cdot o_j - o_i \cdot o_k - \left(\sum_{s=1}^{\rho} \frac{l_j^s \cdot l_k^s}{\deg(l^s \cdot l^s)} \right) \cdot o_i = \tau_{j,k} \cdot o_i. \quad \Box \end{aligned}$$

2.4. Finite-dimensionality. We refer to [1], Chapitre 4 for the definition of Chow motives over k. A motive M is said to be finite-dimensional if M can be decomposed into $M^{\text{odd}} \oplus M^{\text{even}}$ satisfying $S^{N_1}(M^{\text{odd}}) = 0$ and $\wedge^{N_2}(M^{\text{even}}) = 0$ for some $N_1, N_2 > 0$. Here S and \wedge are the symmetric and exterior powers respectively. More precisely, if M is finite-dimensional, one can take $N_1 = \dim H^{\text{odd}}(M) + 1$ and $N_2 = \dim H^{\text{even}}(M) + 1$.

It is conjectured that all Chow motives are finite-dimensional ([8], Conjecture 7.1), although this is proven only for the subcategory generated by the motives of curves (Theorem 4.2 in *loc. cit.*). The motive of a K3 surface is believed to be in this category (e.g. over $\mathbb C$ by applying the Kuga–Satake construction and the Lefschetz standard conjecture in addition). However, as we discussed, even its finite-dimensionality remains unknown in general.

2.5. Back to the K3 surface S. We now interpret what it means for S to be finite-dimensional. By [7], Section 7.2, the motive of S (denoted by h(S)) admits a decomposition

$$h(S) = h^0(S) \oplus h^2_{\mathrm{alg}}(S) \oplus h^2_{\mathrm{tr}}(S) \oplus h^4(S) = \mathbb{1} \oplus \mathbb{L}^{\oplus \rho} \oplus h^2_{\mathrm{tr}}(S) \oplus \mathbb{L}^{\otimes 2}.$$

Here \mathbb{I} is the unit motive and \mathbb{L} is the Lefschetz motive, both of which are (evenly) finite-dimensional. The only part that remains unclear is the motive $h_{tr}^2(S)$, which is defined exactly by the projector $\tau = \tau_{1,2} \in CH^2(S \times S)$. We have

dim $H(h_{tr}^2(S)) = \dim H^2(h_{tr}^2(S)) = b_{tr}$. It follows that S is finite-dimensional if and only if $\wedge^{b_{tr}+1}h_{tr}^2(S) = 0$. In down-to-earth terms, this means

$$\sum_{g \in \mathfrak{S}_{b_{\text{tr}}+1}} \operatorname{sgn}(g) \prod_{i=1}^{b_{\text{tr}}+1} \tau_{i,b_{\text{tr}}+1+g(i)} = 0 \text{ in } R^{b_{\text{tr}}+1} \left(S^{2(b_{\text{tr}}+1)} \right), \tag{7}$$

where \mathfrak{S} stands for the symmetric group and sgn the signature. Note that (7) holds in $H(S^{2(b_{tr}+1)})$, so Conjecture 1 implies the finite-dimensionality of S (first observed in [16], Section 5.1).

The group $\mathfrak{S}_{2(b_{tr}+1)}$ acts on $S^{2(b_{tr}+1)}$ by permutations. It then acts on (7) and produces more relations in $R(S^{2(b_{tr}+1)})$.

3. Proof of the theorem

- **3.1.** Using the relations (4), (5) and (6), it is not difficult to see that $R^{2n}(S^n) = \mathbb{Q}\{\prod_{i=1}^n o_i\}$. Then for $0 \le m \le 2n$, consider the pairing between $R^m(S^n)$ and $R^{2n-m}(S^n)$. We will show that by assuming (7) and its permutations, the pairing is already perfect. This means there cannot be more relations in cl $(R(S^n))$ than in $R(S^n)$, which proves the theorem.
- **3.2.** The first step is essentially the same as in [15], proof of Lemma 2.3. By applying (4), (5) and (6), one observes that $R(S^n)$ is linearly spanned by monomials in $\{o_i\}, \{l_i^s\}$ and $\{\tau_{i,j}\}$ with no repeated index, *i.e.* each index $i \in \{1, ..., n\}$ appears at most once. From now on, we view $\{o_i\}, \{l_i^s\}$ and $\{\tau_{i,j}\}$ as abstract variables. Denote by $\operatorname{Mon}^m(n)$ the set of all formal monomials with no repeated index and with image in $R^m(S^n)$ (when m = 0 we set $\operatorname{Mon}^0(n) = \{1\}$).

We introduce the following symbol for an element in $Mon^{m}(n)$

$$\tau_{I,\alpha} \cdot l_{J,\beta} \cdot o_K$$
.

Here I, J and K are pairwise disjoint subsets of $\{1, \ldots, n\}$ satisfying |I| + |J| + 2|K| = m and |I| even, α is a partition of I into pairs and $\beta \in \{1, \ldots, \rho\}^J$. We set $\tau_{I,\alpha}$ to be the product of $\tau_{i,j}$'s corresponding to pairs in α , $l_{J,\beta} = \prod_{j \in J} l_j^{\beta(j)}$ and $o_K = \prod_{k \in K} o_k$ (again, we set $\tau_{\emptyset,\alpha} = l_{\emptyset,\beta} = o_{\emptyset} = 1$). There is a bijection between $\operatorname{Mon}^m(n)$ and $\operatorname{Mon}^{2n-m}(n)$ given by

$$\tau_{I,\alpha} \cdot l_{J,\beta} \cdot o_K \longleftrightarrow \tau_{I,\alpha} \cdot l_{J,\beta} \cdot o_{(I \cup J \cup K)} c.$$

For $0 \le m \le 2n$, consider the pairing

$$\mathbb{Q}\operatorname{Mon}^{m}(n) \times \mathbb{Q}\operatorname{Mon}^{2n-m}(n) \to \mathbb{Q}\operatorname{Mon}^{2n}(n) = \mathbb{Q}\left\{\prod_{i=1}^{n} o_{i}\right\}$$
(8)

formally defined by the same recipes (4), (5) and (6).

We have the following observation.

3.3. Lemma. Let $\tau_{I,\alpha} \cdot l_{J,\beta} \cdot o_K$ and $\tau_{I',\alpha'} \cdot l_{J',\beta'} \cdot o_{K'}$ be elements in $\mathsf{Mon}^m(n)$ and $\mathsf{Mon}^{2n-m}(n)$ respectively. Then the pairing of the two can be non-zero only if $I' = I, J' = J, K' = (I \cup J \cup K)^{\complement}$ and $\beta' = \beta$.

Proof. Suppose the pairing is non-zero. We have

$$2n = |I| + |J| + 2|K| + |I'| + |J'| + 2|K'| \le 2|I \cup I'| + 2|J \cup J'| + 2|K| + 2|K'| \le 2n.$$

Here the first inequality is obvious and the second follows from the fact that $I \cup I'$, $J \cup J'$, K and K' are pairwise disjoint, which in turn follows from (4), (5) and (6). Therefore the two inequalities are both equalities, which implies I' = I, J' = J and $K' = (I \cup J \cup K)^{\complement}$. Further, the assumption that $\{l^s\}$ is an orthogonal basis implies $\beta' = \beta$.

3.4. Lemma 3.3 shows that after a suitable ordering of the bases, the pairing matrix of (8) is block diagonal. Moreover, the diagonal blocks correspond to the pairing of elements in $\mathrm{Mon}^d(d)$ for some $d \leq \min\{m, 2n - m\}$ that consist solely of $\tau_{i,j}$'s. We denote by $\mathrm{Mon}^d_{\tau}(d) \subset \mathrm{Mon}^d(d)$ the subset of all such elements, *i.e.* monomials of the form $\tau_{\alpha} = \tau_{\{1,\ldots,d\},\alpha}$ where α is a partition of $\{1,\ldots,d\}$ into pairs (d even).

We are left to consider the pairing

$$\mathbb{Q}\operatorname{Mon}_{\tau}^{d}(d) \times \mathbb{Q}\operatorname{Mon}_{\tau}^{d}(d) \to \mathbb{Q}\operatorname{Mon}^{2d}(d) = \mathbb{Q}\left\{\prod_{i=1}^{d} o_{i}\right\}. \tag{9}$$

It turns out that the matrix of (9) has been studied in detail by Hanlon and Wales [6]. It is denoted by $T_r(x)$ with r = d/2 and $x = b_{tr}$. Here we only cite (and translate) what is relevant to our problem, namely Theorem 3.1 in *loc. cit*..

3.5. Proposition. The symmetric group \mathfrak{S}_d acts on $\mathbb{Q} \operatorname{Mon}_{\tau}^d(d)$ by permutations. We have

$$\mathbb{Q}\operatorname{Mon}_{\tau}^d(d) \simeq \bigoplus_{\lambda \in \Lambda_d} V_{\lambda},$$

where Λ_d is the set of partitions of $\{1, \ldots, d\}$ whose parts are all even, and V_{λ} is the irreducible representation of \mathfrak{S}_d associated to λ . All V_{λ} 's are eigenspaces of the matrix $T_{d/2}(b_{tr})$. The eigenvalue is 0 if and only if λ contains at least $b_{tr} + 1$ parts.

3.6. For λ a partition of $\{1, \ldots, d\}$, we recall the definition of V_{λ} via Specht modules (see [5], Problem 4.47). Define a tabloid $\{T\}$ to be an equivalence class of Young tableaux T associated to λ , two being equivalent if the rows are the same

up to order. The group \mathfrak{S}_d acts by permutations on the set of such tabloids, denoted by Tab(λ). For each tableau T, define

$$E_T = \sum_{g \in Q_T} \operatorname{sgn}(g) \{g(T)\} \in \mathbb{Q} \operatorname{Tab}(\lambda),$$

where $Q_T \subset \mathfrak{S}_d$ is the column stabilizer of T. Then V_{λ} is the \mathbb{Q} -linear span of all E_T 's in \mathbb{Q} Tab(λ). A basis of V_{λ} is given by the E_T 's with standard tableaux T.

When $\lambda \in \Lambda_d$, we locate V_λ inside $\mathbb{Q}\operatorname{Mon}_{\tau}^d(d)$ as follows. Take a tableau T associated to λ and denote by $T_i \subset \{1, \ldots, d\}$ the i-th row of T, with $|T_i|$ even. For each T_i consider the sum $\sum_{\alpha_i} \tau_{T_i,\alpha_i}$, where α_i runs through all partitions of T_i into pairs. Altogether we assign to T the product

$$\phi(T) = \prod_{i} \left(\sum_{\alpha_i} \tau_{T_i, \alpha_i} \right) \in \mathbb{Q} \operatorname{Mon}_{\tau}^d(d).$$

By construction ϕ descends to the tabloids and the resulting map $\phi: \mathbb{Q} \operatorname{Tab}(\lambda) \to \mathbb{Q} \operatorname{Mon}_{\tau}^d(d)$ is \mathfrak{S}_d -equivariant. Restricting to V_{λ} we get a morphism of \mathfrak{S}_d -modules $\phi|_{V_{\lambda}}: V_{\lambda} \to \mathbb{Q} \operatorname{Mon}_{\tau}^d(d)$, which is injective since it is non-zero and V_{λ} is irreducible. We identify V_{λ} with its image.

3.7. End of proof. By Proposition 3.5, we know that V_{λ} lies in the kernel (*i.e.* radical) of (9) if and only if λ contains at least $b_{tr} + 1$ parts. The first occurrence is when $d = 2(b_{tr} + 1)$ and $\lambda = (2, ..., 2)$. Take for example the following standard tableau T, with $Q_T \simeq \mathfrak{S}_{b_{tr}+1} \times \mathfrak{S}_{b_{tr}+1}$.


$$\begin{array}{c|cccc}
1 & b_{tr} + 2 \\
2 & b_{tr} + 3 \\
\vdots & \vdots \\
b_{tr} + 1 & 2b_{tr} + 2
\end{array}$$

A direct calculation gives

$$\phi(E_T) = (b_{tr} + 1)! \sum_{g \in \mathfrak{S}_{b_{tr} + 1}} \operatorname{sgn}(g) \prod_{i=1}^{b_{tr} + 1} \tau_{i, b_{tr} + 1 + g(i)}, \tag{10}$$

which is exactly $(b_{tr} + 1)!$ times the left-hand side of (7). The other $\phi(E_T)$'s are given by permuting the indices on the right-hand side of (10).

The situation is similar as long as λ contains at least $b_{tr}+1$ parts. We draw a standard tableau T of such a λ , with the length of the first and second columns $e \geq b_{tr}+1$ (lengths of other columns do not matter). Here $Q_{\lambda} \simeq \mathfrak{S}_e \times \mathfrak{S}_e \times \cdots$.



By writing \mathfrak{S}_e as the union of cosets $\{g\mathfrak{S}_{b_{tr}+1}\}$, it is not difficult to see that $\phi(E_T)$ is generated by various pull-backs and permutations of the right-hand side of (10). The same holds for the other $\phi(E_T)$'s again by permutations.

We conclude that the kernel of (9) is entirely generated by the right-hand side of (10) and its permutations. Then by the assumption (7) those classes vanish in $R(S^{2(b_{tr}+1)})$. It follows that the kernel of (9) vanishes in $R(S^d)$, and that the pairing between $R^m(S^n)$ and $R^{2n-m}(S^n)$ is perfect.

- **3.8. Final remarks.** (i) A 1-dimensional analogue of our result concerns a hyperelliptic curve C. There the ring $R(C^n)$ contains (pull-backs of) the canonical class of C and the diagonal class on $C \times C$. It is proven that all relations in $R(C^n)$ are generated by the vanishing of the Faber–Pandharipande cycle in $R^2(C^2)$, the vanishing of the Gross–Schoen cycle in $R^2(C^3)$, and one relation corresponding to the vanishing of the motive $S^{2g+2}h^1(C)$ (unconditional). This is the work of Tavakol ([12]; see also [13]), which inspired the present note. It would also be interesting to see if there are higher-dimensional analogues.
- (ii) As is remarked in [16], Section 5.1, it might be the case that Conjecture 2 for $S^{[n]}$ also implies Conjecture 1, *i.e.* the two are equivalent. Our result shows that it suffices to deduce the relation (7) from Conjecture 2. By the work of de Cataldo and Migliorini on the decomposition of $CH(S^{[n]})$ ([3], Theorem 5.4.1), one can express the left-hand side of (7) as a homologically trivial class in $CH(S^{[n]})$ for some very large n ($n \ge 1 + 2 + \cdots + (b_{tr} + 1)$). It remains to see if this class is generated by divisor classes and Chern classes of the tangent bundle. The computation feels like a "reverse engineering" of [15], proof of Proposition 2.6.
- (iii) Further, observe that the relations (4), (5), (6) and (7) hold in cohomology for any smooth projective surface of Albanese dimension 0. The same argument then shows that for such a surface S, all relations in cl $(R(S^n))$ are generated by (4), (5), (6) and (7). We refer to [17] and [14] for recent applications of this result.

References

[1] André, Y. *Une introduction aux motifs (motifs purs, motifs mixtes, périodes)*. Panoramas et Synthèses, 17. Société Mathématique de France, Paris, 2004. xii+261 pp. Zbl 1060.14001 MR 2115000

- [2] Beauville, A. and C. Voisin. On the Chow ring of a *K*3 surface. *Journal of Algebraic Geometry* **13** (2004), no. 3, 417–426. Zbl 1069.14006 MR 2047674
- [3] de Cataldo, M. and L. Migliorini. The Chow groups and the motive of the Hilbert scheme of points on a surface. *Journal of Algebra* **251** (2002), no. 2, 824–848. Zbl 1033.14004 MR 1919155
- [4] Fu, L. Beauville–Voisin conjecture for generalized Kummer varieties. Preprint, arXiv:1309.4977, 2013. To appear in International Mathematics Research Notices.
- [5] Fulton, W. and J. Harris. *Representation theory. A first course*. Graduate Texts in Mathematics, 129. Readings in Mathematics. Springer-Verlag, New York, 1991. xvi+551 pp. Zbl 0744.22001 MR 1153249
- [6] Hanlon, P. and D. Wales. On the decomposition of Brauer's centralizer algebras. *Journal of Algebra* 121 (1989), no. 2, 409–445. Zbl 0695.20026 MR 992775
- [7] Kahn, B. J. Murre, and C. Pedrini. On the transcendental part of the motive of a surface. Algebraic cycles and motives. Vol. 2, 143–202, London Mathematical Society Lecture Note Series, 344, Cambridge Univ. Press, Cambridge, 2007. Zbl 1130.14008 MR 2187153
- [8] Kimura, S.-I. Chow groups are finite dimensional, in some sense. *Mathematische Annalen* **331** (2005), no. 1, 173–201. Zbl 1067.14006 MR 2107443
- [9] Liedtke, C. Supersingular K3 Surfaces are Unirational. Preprint, arXiv:1304.5623, 2013. To appear in Inventiones Mathematicae.
- [10] Moonen, B. On the Chow motive of an abelian scheme with non-trivial endomorphisms. Preprint, arXiv:1110.4264, 2011. To appear in Journal für die Reine und Angewandte Mathematik.
- [11] Pedrini, C. On the finite dimensionality of a K3 surface. *Manuscripta Mathematica* **138** (2012), no. 1-2, 59–72. Zbl 1278.14012 MR 2898747
- [12] Tavakol, M. The tautological ring of the moduli space $\mathcal{M}_{2,n}^{rt}$. Preprint, arXiv:1101.5242, 2011. To appear in *International Mathematics Research Notices*.
- [13] Tavakol, M. *Tautological classes on the moduli space of hyperelliptic curves with rational tails.* Preprint, arXiv:1406.7403, 2014.
- [14] Vial, C. On the motive of hyperkaehler varieties. Preprint, arXiv:1406.1073, 2014. To appear in Journal für die Reine und Angewandte Mathematik.
- [15] Voisin, C. On the Chow ring of certain algebraic hyper-Kähler manifolds. *Pure and Applied Mathematics Quarterly* **4** (2008), no. 3, part 2, 613–649. Zbl 1165.14012 MR 2435839
- [16] Voisin, C. Chow rings, decomposition of the diagonal and the topology of families. Annals of Mathematics Studies, 187. Princeton University Press, Princeton, NJ, 2014. viii+163 pp. Zbl 1288.14001 MR 3186044
- [17] Voisin, C. Some new results on modified diagonals. Preprint, arXiv:1405.6957, 2014. To appear in Geometry & Topology.

Received May 12, 2014; revised December 14, 2014

Q. Yin, ETH Zürich, Departement Mathematik, Rämistrasse 101, 8092 Zürich, Switzerland E-mail: yin@math.ethz.ch