

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 91 (2016)
Heft: 1

Artikel: Asymptotic equivalence of symplectic capacities
Autor: Gluskin, Efim D. / Ostrover, Yaron
DOI: <https://doi.org/10.5169/seals-630554>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 26.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Asymptotic equivalence of symplectic capacities

Efim D. Gluskin and Yaron Ostrover*

Abstract. A long-standing conjecture states that all normalized symplectic capacities coincide on the class of convex subsets of \mathbb{R}^{2n} . In this note we focus on an asymptotic (in the dimension) version of this conjecture, and show that when restricted to the class of centrally symmetric convex bodies in \mathbb{R}^{2n} , several symplectic capacities, including the Ekeland–Hofer–Zehnder capacity, the displacement energy capacity, and the cylindrical capacity, are all equivalent up to a universal constant.

Mathematics Subject Classification (2010). 53D35, 52A23, 52A20, 46B07, 46B20.

Keywords. Symplectic capacities, convex bodies, asymptotic behaviour.

1. Introduction

Consider the space \mathbb{R}^{2n} equipped both with the standard symplectic form $\omega = dp \wedge dq$, and with the standard inner product $\langle \cdot, \cdot \rangle$. Note that under the usual identification between \mathbb{R}^{2n} and \mathbb{C}^n , these two structures are the real and the imaginary parts, respectively, of the standard Hermitian inner product in \mathbb{C}^n . Moreover, one has that $\omega(v, u) = \langle v, Ju \rangle$, where J is the standard complex structure in $\mathbb{R}^{2n} \simeq \mathbb{C}^n$. Symplectic capacities, whose axiomatic definition below is due to Ekeland and Hofer [4], are numerical invariants which roughly speaking measure the symplectic size of sets. More precisely, let $B^{2n}(r)$ stand for the Euclidean open ball of radius r , and $Z^{2n}(r)$ for the cylinder $B^2(r) \times \mathbb{C}^{n-1}$.

Definition 1.1. A symplectic capacity on $(\mathbb{R}^{2n}, \omega)$ associates to each subset $U \subset \mathbb{R}^{2n}$ a number $c(U) \in [0, \infty]$ such that the following hold:

- (P1) $c(U) \leq c(V)$ whenever $U \subseteq V$ (monotonicity),
- (P2) $c(\psi(U)) = |\alpha|c(U)$ for $\psi \in \text{Diff}(\mathbb{R}^{2n})$ such that $\psi^*\omega = \alpha\omega$ for $0 \neq \alpha \in \mathbb{R}$ (conformality),
- (P3) $0 < c(B^{2n}(r))$, and $c(Z^{2n}(r)) < \infty$ (nontriviality).

*The second-named author was partially supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme, starting grant No. 637386, and by the ISF grant No. 1274/14.

Moreover, a symplectic capacity is said to be normalized if in addition it satisfies

$$(P4) \quad c(B^{2n}(r)) = c(Z^{2n}(r)) = \pi r^2 \text{ (normalization).}$$

Note that property (P2) implies that c is a symplectic invariant which scales like a two-dimensional invariant, and (P3) that symplectic capacities significantly differ from any volume related invariants. The first examples of symplectic capacities were constructed by Gromov in [9], where he developed and used pseudoholomorphic curve techniques to prove a striking symplectic rigidity result, nowadays known as Gromov's "non-squeezing theorem". It states that one cannot map a ball inside a thinner cylinder by a symplectic embedding. More precisely, the theorem asserts that if $r < 1$, there is no symplectic embedding of the unit ball B^{2n} into the cylinder $Z^{2n}(r)$. This naturally leads to the definition of two normalized symplectic capacities: the Gromov width, given by $\underline{c}(U) = \sup\{\pi r^2 \mid B^{2n}(r) \xrightarrow{s} U\}$; and the cylindrical capacity, $\bar{c}(U) = \inf\{\pi r^2 \mid U \xrightarrow{s} Z^{2n}(r)\}$. Here \xrightarrow{s} stands for symplectic embedding. It is not hard to verify that these two capacities are the smallest and largest possible normalized symplectic capacities, respectively.

Shortly after Gromov's work [9] many other symplectic capacities were constructed, reflecting different geometrical and dynamical properties. Among these are the Hofer–Zehnder capacity [13, 14], the Ekeland–Hofer capacities [4, 5], the displacement energy [11], the Floer–Hofer capacity [7, 8], spectral capacities [6, 19, 26], and more recently, Hutchings' embedded contact homology (ECH) capacities [15]. These quantities play an important role in symplectic geometry, and their properties, interrelations, and applications to symplectic topology and Hamiltonian dynamics are intensively studied (see e.g., [3] and [17] for two excellent surveys).

In the two-dimensional case, Siburg [23] showed that any symplectic capacity of a compact connected domain with smooth boundary $\Omega \subset \mathbb{R}^2$ equals its Lebesgue measure. In higher dimensions symplectic capacities do not coincide in general. A theorem by Hermann [10] states that for any $n \geq 2$ there is a bounded star-shaped domain $S \subset \mathbb{R}^{2n}$ with cylindrical capacity $\bar{c}(S) \geq 1$, and arbitrarily small Gromov width $\underline{c}(S)$. Still, for a large class of sets in \mathbb{R}^{2n} , including ellipsoids, polydiscs, and convex Reinhardt domains, all normalized symplectic capacities coincide [10]. In [25] Viterbo showed that for any bounded convex set K of \mathbb{R}^{2n} one has $\bar{c}(K) \leq 4n^2 \underline{c}(K)$. Moreover, it was conjectured [10, 12, 25] that:

Conjecture 1.2. *For any convex body K in \mathbb{R}^{2n} one has $\underline{c}(K) = \bar{c}(K)$.*

Here, by a convex body we mean a compact convex subset of \mathbb{R}^{2n} with non-empty interior. The above conjecture is particularly challenging due to the scarcity of examples of convex domains for which capacities have been computed. Moreover, an affirmative answer to Conjecture 1.2 would in particular implies Viterbo's volume-capacity conjecture [25], and it was recently shown that the latter would in turn settle a 70-years old question in convex geometry known as the Mahler conjecture. For more information regarding these applications of Conjecture 1.2 see [1] and [20].

A somewhat more modest question in the same direction (c.f. Problem 1.4 in [10], Problem 8 in [3], and Section 5 in [20]) is whether Conjecture 1.2 above holds asymptotically in the dimension, i.e.,

Question 1.3. Is there is an absolute constant $A > 0$ such that for every convex body K in \mathbb{R}^{2n} one has

$$\bar{c}(K) \leq A c(K).$$

Here we will give a partial answer to this question. Before we state our main result we wish to recall the definition of the Ekeland–Hofer–Zehnder capacity. The restriction of the symplectic form ω to a smooth closed hypersurface $\mathcal{S} \subset \mathbb{R}^{2n}$ canonically defines a 1-dimensional subbundle, $\ker(\omega|_{\mathcal{S}})$, whose integral curves comprise the characteristic foliation of \mathcal{S} . In other words, a closed characteristic of \mathcal{S} is an embedded circle in \mathcal{S} tangent to the canonical line bundle

$$\mathfrak{C}_{\mathcal{S}} = \{(x, \xi) \in T\mathcal{S} \mid \omega(\xi, \eta) = 0 \text{ for all } \eta \in T_x\mathcal{S}\}.$$

Recall that the symplectic action of a closed curve γ is defined by $A(\gamma) = \int_{\gamma} \lambda$, where $\lambda = pdq$ is the Liouville 1-form. The action spectrum of \mathcal{S} is

$$\mathcal{L}(\mathcal{S}) = \{|A(\gamma)|; \gamma \text{ is a closed characteristic on } \mathcal{S}\}.$$

In [4] and [14] it was proved that for a smooth convex body $K \subset \mathbb{R}^{2n}$, the two aforementioned Hofer–Zehnder and Ekeland–Hofer capacities coincide, and are given by the minimal action over all closed characteristics on the boundary of the body K , i.e.,

$$c_{\text{EH}}(K) = c_{\text{HZ}}(K) = \min \mathcal{L}(\partial K). \quad (1)$$

We remark that although the above definition of closed characteristics, as well as the equalities in (1), were given only for the class of convex bodies with smooth boundary, they can naturally be generalized to the class of convex sets in \mathbb{R}^{2n} with nonempty interior (see e.g., [2]). In what follows, we refer to the coinciding Ekeland–Hofer and Hofer–Zehnder capacities on this class as the Ekeland–Hofer–Zehnder capacity, and denote it by c_{EHZ} .

Our first result in the note is the following. Recall that a convex body $K \subset \mathbb{R}^n$ is said to be centrally symmetric if $K = -K$.

Theorem 1.4. *For every centrally symmetric convex body K in \mathbb{R}^{2n} ,*

$$\bar{c}(K) \leq 4c_{\text{EHZ}}(K).$$

Remark 1.5. Other symplectic capacities, like the spectral capacities c_{σ} , which are based on a choice of an action selector σ , and the displacement energy d , are known to be bigger than or equal to the Hofer–Zehnder capacity (see e.g., Section 2.3.4 in [3]). Thus, it follows from Theorem 1.4 that on the class of symmetric convex sets in \mathbb{R}^{2n} , the normalized symplectic capacities c_{EHZ} , d , c_{σ} and \bar{c} , all coincide up to an absolute constant.

In fact, we prove a slightly stronger result than Theorem 1.4 which shows that for a centrally symmetric convex body $K \subset \mathbb{R}^{2n}$, the aforementioned symplectic capacities are all equivalent to yet another quantity associated with the body K . More precisely, for a convex body $K \subset \mathbb{R}^{2n}$ with $0 \in \text{Int}(K)$, we denote by $K^\circ = \{y \in \mathbb{R}^{2n} \mid \langle x, y \rangle \leq 1, \text{ for every } x \in K\}$ the polar body of K ¹. Moreover, we denote

$$\|J\|_{K^\circ \rightarrow K} := \sup_{v, u \in K^\circ} \langle Jv, u \rangle.$$

To explain the reason for this notation, we remark that when the convex body K is centrally symmetric, $\|J\|_{K^\circ \rightarrow K}$ is the operator norm of the complex structure J , when the latter is considered as a linear map between the normed spaces $J : (\mathbb{R}^{2n}, \|\cdot\|_{K^\circ}) \rightarrow (\mathbb{R}^{2n}, \|\cdot\|_K)$, i.e.,

$$\|J\|_{K^\circ \rightarrow K} = \sup_{v, u \in K^\circ} \langle Jv, u \rangle = \sup_{v: \|v\|_{K^\circ} \leq 1} \|Jv\|_K.$$

Here we use the standard identification between normed spaces and centrally symmetric convex bodies, i.e., for a non-empty centrally symmetric convex body K in \mathbb{R}^{2n} we denote by $\|\cdot\|_K$ the norm on \mathbb{R}^{2n} induced by K , that is,

$$\|\cdot\|_K = \inf\{r : x \in rK\}.$$

Theorem 1.6. *For every centrally symmetric convex body K in \mathbb{R}^{2n} ,*

$$\frac{1}{\|J\|_{K^\circ \rightarrow K}} \leq c_{\text{EHZ}}(K) \leq \bar{c}(K) \leq \frac{4}{\|J\|_{K^\circ \rightarrow K}}. \tag{2}$$

Remark 1.7. In fact, in the proof of Theorem 1.6 we use the centrally symmetric assumption on the body K only for the right-most inequality of (2). The first two inequalities on the left-hand side hold for every convex body K in \mathbb{R}^{2n} .

Note that Theorem 1.4 follows immediately from Theorem 1.6. Moreover, we wish to emphasize that Theorem 1.6 provides in many cases an efficient way to approximate the numerical value of the capacities $c_{\text{EHZ}}(K)$ and $\bar{c}(K)$ (for centrally symmetric convex bodies), as the quantity $\|J\|_{K^\circ \rightarrow K}$ is a-priori much easier to compute than the above mentioned symplectic capacities.

Another by-product of Theorem 1.6, which may be of independent interest, concerns the equivalence of the cylindrical capacity and the Gromov width capacity with their linearized versions \bar{c}_{lin} and $\underline{c}_{\text{lin}}$ respectively. The definitions of these two quantities are given in Definitions 2.4 and 3.1 below. It turns out that for centrally symmetric convex bodies in \mathbb{R}^{2n} , the cylindrical capacity \bar{c} is asymptotically equivalent to its linearized version \bar{c}_{lin} , while surprisingly enough, this is false for the Gromov width capacity.

¹As a matter of fact, the polar body K° should be defined as a subset of the dual space of \mathbb{R}^{2n} . However, since we have fixed a scalar product in our setting, we will identify the latter space with \mathbb{R}^{2n} itself.

More precisely,

Theorem 1.8. *For every centrally symmetric convex body K in \mathbb{R}^{2n} ,*

$$\bar{c}(K) \leq \bar{c}_{\text{lin}}(K) \leq 4\bar{c}(K).$$

On the other hand, there exist a centrally symmetric convex body \tilde{K} in \mathbb{R}^{2n} such that

$$\underline{c}_{\text{lin}}(\tilde{K}) \leq \pi, \text{ while } \underline{c}(\tilde{K}) \geq \sqrt{\frac{n}{2}}.$$

Note that an immediate corollary from Theorem 1.8 is that the linearized versions of the Gromov width and the cylindrical capacity are not asymptotically equivalent.

Notation. We denote by \mathcal{K}^n the class of convex bodies of \mathbb{R}^n , i.e., compact convex sets with non-empty interior. For $K \in \mathcal{K}^n$, we denote by $h_K : \mathbb{R}^n \rightarrow \mathbb{R}$ its support function given by $h_K(u) = \sup\{\langle x, u \rangle : x \in K\}$. Also, we denote by $g_K : \mathbb{R}^n \rightarrow \mathbb{R}$ the gauge function $g_K(x) = \inf\{r \mid x \in rK\}$ associated with K . Note that when K is centrally symmetric, i.e., $K = -K$, the gauge function $g_K(x)$ is a norm, and is denoted by $\|x\|_K$. Furthermore, when $0 \in \text{int}(K)$, one has that $h_K = g_{K^\circ}$, where $K^\circ = \{y \in \mathbb{R}^n \mid \langle x, y \rangle \leq 1, \text{ for every } x \in K\}$ is the polar body of K . The Euclidean norm will be denoted by $|\cdot|$. Finally, we denote by \mathbb{S}^n the unit sphere in \mathbb{R}^{n+1} , i.e., $\mathbb{S}^n = \{x \in \mathbb{R}^{n+1} \mid |x| = 1\}$.

Acknowledgements. The authors are grateful to Shiri Artstein-Avidan and Boaz Klartag for many stimulating discussions on various topics related to symplectic geometry and convexity.

2. Proof of Theorem 1.6

Note first that there is no loss of generality in assuming that in addition to being compact and with non-empty interior, all convex bodies considered also have a smooth boundary, and contain the origin in their interior. Indeed, affine translations in \mathbb{R}^{2n} are symplectomorphisms, which accounts for the assumption that the origin is in the interior. Secondly, once Theorem 1.6 is proved for smooth convex domains, the general case follows by standard approximation arguments, as symplectic capacities are continuous on the class of convex bodies with respect to the Hausdorff distance (see e.g. [18, p. 376]).

Moreover, in what follows we will make repeated use of the following well-known geometric observation from convex geometry.

Lemma 2.1. *Let g_K be the gauge function associated with a smooth convex body K . Then, when restricted to the boundary ∂K , the gradient ∇g_K is a surjective map $\nabla g_K : \partial K \rightarrow \partial K^\circ$.*

A proof of Lemma 2.1 can be found e.g., in Subsection 1.7.1 of [22]. We turn now to the proof of Theorem 1.6, and start with the following proposition.

Proposition 2.2. *For every smooth convex body $K \in \mathcal{K}^{2n}$,*

$$\frac{1}{\|J\|_{K^\circ \rightarrow K}} \leq c_{\text{EHZ}}(K).$$

To prove Proposition 2.2 we first need some preparation. Recall (see e.g., Chapter 1 of [13]) that the classical geometric problem of finding closed characteristics on ∂K has the following dynamical interpretation. If the boundary ∂K is represented as a regular energy surface $\{x \in \mathbb{R}^{2n} \mid H(x) = 1\}$ of a smooth Hamiltonian function $H : \mathbb{R}^{2n} \rightarrow \mathbb{R}$, then the restriction to ∂K of the Hamiltonian vector field X_H , defined by $i_{X_H}\omega = -dH$, is a section of the line bundle $\mathfrak{S}_{\partial K}$. Thus, the images of the periodic solutions of the classical Hamiltonian equation $\dot{x} = X_H(x) = J\nabla H(x)$ on ∂K are precisely the closed characteristics of ∂K . In particular, the closed characteristics do not depend (up to parametrization) on the choice of the Hamiltonian function. Indeed, if the energy surface can be represented as a regular level set of some other function $F : \mathbb{R}^{2n} \rightarrow \mathbb{R}$, then $X_H = \alpha X_F$ on ∂K for some scalar function $\alpha \neq 0$, and the corresponding Hamiltonian equations have the same solutions up to parametrization. Finally, note that for a smooth convex body K the gauge function g_K is a *defining function* for K , i.e., $K = g_K^{-1}([0, 1])$, $\partial K = g_K^{-1}(1)$, and 1 is a regular value of g_K .

Lemma 2.3. *Let $\gamma : [0, T] \rightarrow \partial K$ be a solution of the Hamiltonian equation $\dot{\gamma} = J\nabla g_K(\gamma)$, with $\gamma(0) = \gamma(T)$. Then there exist $t_0 \in [0, T]$ such that*

$$g_K(\gamma(t_0) - \gamma(0)) \geq 1.$$

Proof of Lemma 2.3. It follows immediately from the assumptions that

$$0 = \int_0^T \dot{\gamma}(t) dt = \int_0^T J\nabla g_K(\gamma(t)) dt = J \int_0^T \nabla g_K(\gamma(t)) dt.$$

From this one can conclude that

$$\int_0^T \langle \nabla g_K(\gamma(t)), \gamma(0) \rangle dt = 0.$$

In particular, this implies that there exists $t_0 \in [0, T]$ such that

$$\langle \nabla g_K(\gamma(t_0)), \gamma(0) \rangle \leq 0. \tag{3}$$

Next, from Lemma 2.1 it follows that $\nabla g_K(\gamma(t_0)) \in \partial K^\circ$, and we obtain that

$$\begin{aligned} g_K(\gamma(t_0) - \gamma(0)) &= \sup\{\langle \gamma(t_0) - \gamma(0), u \rangle \mid u \in K^\circ\} \\ &\geq \langle \gamma(t_0) - \gamma(0), \nabla g_K(\gamma(t_0)) \rangle. \end{aligned} \tag{4}$$

Finally, from Euler's homogeneous function theorem it follows that for every $x \in \partial K$, one has $\langle x, \nabla g_K(x) \rangle = g_K(x) = 1$, and hence the combination of this fact together with inequalities (3) and (4) completes the proof of the lemma. \square

Proof of Proposition 2.2. Let $\gamma : [0, T] \rightarrow \partial K$ be a closed characteristic on the boundary ∂K , i.e., a solution of the Hamiltonian equation $\dot{\gamma} = J\nabla g_K(\gamma)$, with $\gamma(0) = \gamma(T)$. Note that

$$A(\gamma) = \frac{1}{2} \int_0^T \langle J\dot{\gamma}(t), \dot{\gamma}(t) \rangle dt = \frac{1}{2} \int_0^T \langle \dot{\gamma}(t), \nabla g_K(\gamma(t)) \rangle dt = \frac{T}{2}. \quad (5)$$

It follows from Lemma 2.3, the subadditivity property of g_K , and the definition of γ , that there exists $t_0 \in [0, T]$ such that

$$1 \leq g_K \left(\int_0^{t_0} \dot{\gamma}(t) dt \right) \leq \int_0^{t_0} g_K(\dot{\gamma}(t)) dt = \int_0^{t_0} g_K(J\nabla g_K(\gamma(t))) dt. \quad (6)$$

On the other hand, it follows from the definition of an operator norm that

$$\int_0^{t_0} g_K(J\nabla g_K(\gamma(t))) dt \leq \int_0^{t_0} \|J\|_{K^\circ \rightarrow K} g_{K^\circ}(\nabla g_K(\gamma(t))) dt. \quad (7)$$

The combination of (6), (7), and Lemma 2.1 gives

$$1 \leq \int_0^{t_0} \|J\|_{K^\circ \rightarrow K} g_{K^\circ}(\nabla g_K(\gamma(t))) dt = \int_0^{t_0} \|J\|_{K^\circ \rightarrow K} dt, \quad (8)$$

and thus we obtain that

$$\frac{1}{\|J\|_{K^\circ \rightarrow K}} \leq t_0. \quad (9)$$

Note that since $\gamma(0) = \gamma(T)$, repeating the same arguments as above (this time, integrating in (6), (7), and (8) between t_0 and T) we obtain also that

$$\frac{1}{\|J\|_{K^\circ \rightarrow K}} \leq T - t_0. \quad (10)$$

From (5) it follows that $\min\{t_0, T - t_0\} \leq T/2 = A(\gamma)$, and since, by definition, the capacity $c_{\text{EHZ}}(K)$ is defined to be the minimal action of closed characteristics on the boundary ∂K , we conclude from (9) and (10) that,

$$\frac{1}{\|J\|_{K^\circ \rightarrow K}} \leq c_{\text{EHZ}}(K).$$

This completes the proof of the proposition. \square

To describe the second ingredient in the proof of Theorem 1.6 we need to introduce one more definition. It is known (see e.g., Appendix C in [21]) that for a Lebesgue measurable set $\mathcal{U} \subset \mathbb{R}^{2n}$,

$$\bar{c}(\mathcal{U}) = \inf_{\varphi} \text{Area}(\pi(\varphi(\mathcal{U}))),$$

where π is the orthogonal projection to the complex line

$$E = \{z \in \mathbb{C}^n \mid z_j = 0 \text{ for } j \neq 1\},$$

and the infimum is taken over all symplectic embeddings φ of \mathcal{U} into \mathbb{R}^{2n} . Recall that with our notations, under the natural identification $\mathbb{R}^{2n} \simeq \mathbb{C}^n$ one has that $z_j = q_j + ip_j$. Thus, a natural way to “linearize” the cylindrical capacity \bar{c} is as follows. Let $\text{ISp}(2n)$ be the affine symplectic group, defined as the semi-direct product $\text{Sp}(2n) \times \text{T}(2n)$ of the linear symplectic group and the group of translations in \mathbb{R}^{2n} .

Definition 2.4. The linearized cylindrical capacity \bar{c}_{lin} of a set $\mathcal{U} \subset \mathbb{R}^{2n}$ is defined as

$$\bar{c}_{\text{lin}}(\mathcal{U}) = \inf_S \text{Area}(\pi(S(\mathcal{U}))),$$

where the infimum is taken over all affine symplectic maps $S \in \text{ISp}(2n)$.

The second main ingredient in the proof of Theorem 1.6 is the following:

Proposition 2.5. *For every centrally symmetric convex body $K \in \mathcal{K}^{2n}$,*

$$\bar{c}(K) \leq \bar{c}_{\text{lin}}(K) \leq \frac{4}{\|J\|_{K^\circ \rightarrow K}}. \quad (11)$$

To establish Proposition 2.5 we shall need the following geometric observation. For $v \in \mathbb{R}^{2n}$, we denote by K_v the section $K \cap \{v\}^\perp$, and by $\|\cdot\|_{K_v^\circ}$ the semi-norm defined by

$$\|w\|_{K_v^\circ} = \sup\{\langle w, y \rangle \mid y \in K_v\}.$$

Lemma 2.6. *For a symmetric convex body $K \in \mathcal{K}^{2n}$, a linear symplectic map $S \in \text{Sp}(2n)$, and the orthogonal projection π to the complex line*

$$E = \{z \in \mathbb{C}^n \mid z_j = 0 \text{ for } j \neq 1\}$$

defined above, one has

$$\text{Area}(\pi(S(K))) \leq 4\|S^T e\|_{K^\circ} \|S^T J e\|_{K_v^\circ}, \quad (12)$$

where S^T stands for the transpose of the matrix S , e is a unit vector parallel to the q_1 -axis, and $v = S^T e$.

Proof of Lemma 2.6. The lemma follows from a much more general result by Rogers and Shephard [24], which states that for every symmetric convex body $K \subset \mathbb{R}^n$ one has

$$\text{Vol}_n(K) \leq \left(\text{Vol}_k(\pi_E(K)) \text{Vol}_{n-k}(K \cap E^\perp) \right) \leq \binom{n}{k} \text{Vol}_n(K), \tag{13}$$

for every k -dimensional subspace E of \mathbb{R}^n , where π_E stands for the orthogonal projection on the subspace E . We remark that we use only the case where $n = 2$ and $k = 1$, in which inequality (13) is an elementary geometric fact. It can be easily checked that the right-hand side of (12) exactly equals the product of the length of the projection of $\pi(SK)$ to the q_1 -axis, and the length of the intersection of $\pi(SK)$ with the p_1 -axis. \square

We are now in a position to prove Proposition 2.5.

Proof of Proposition 2.5. Note that, by definition, for every measurable set $\mathcal{U} \subset \mathbb{R}^{2n}$ one has $\bar{c}(\mathcal{U}) \leq \bar{c}_{\text{lin}}(\mathcal{U})$, and hence the left-hand side inequality in (11) holds. Next, we recall the easily verified fact that for any $v, w \in \mathbb{R}^{2n}$ such that $\omega(v, w) = 1$, there exists a linear symplectic map $S \in \text{Sp}(2n)$ such that $v = S^T e$ and $w = S^T J e$, where as before, e is a unit vector parallel to the q_1 -axis. From this fact, Lemma 2.6, and Definition 2.4 it follows that for a centrally symmetric convex body $K \in \mathcal{K}^{2n}$

$$\begin{aligned} \bar{c}_{\text{lin}}(K) &\leq 4 \inf_{v \in \mathbb{S}^{2n-1}} \inf_{w : \langle Jv, w \rangle = 1} \|v\|_{K^\circ} \|w\|_{K_v^\circ} \\ &= 4 \inf_{v \in \mathbb{S}^{2n-1}} \|v\|_{K^\circ} \inf_{w : \langle Jv, w \rangle = 1} \|w\|_{K_v^\circ}. \end{aligned} \tag{14}$$

We focus now on the second infimum on the right-hand side of (14). Note that for a fixed vector $v \in \mathbb{S}^{2n-1}$, the equality $\langle Jv, w \rangle = 1$ is equivalent to $\langle Jv, w - Jv \rangle = 0$. Denoting $z := w - Jv$, we can write

$$\inf_{w : \langle Jv, w \rangle = 1} \|w\|_{K_v^\circ} = \inf_{z : z \perp Jv} \|Jv + z\|_{K_v^\circ}. \tag{15}$$

This quantity measures the distance, with respect to the semi-metric induced by $\|\cdot\|_{K_v^\circ}$, between the vector Jv and the subspace $\{Jv\}^\perp$ orthogonal to it. By duality,

$$\inf_{z : z \perp Jv} \|Jv + z\|_{K_v^\circ} = \text{dist}_{\|\cdot\|_{K_v^\circ}}(Jv, \{Jv\}^\perp) = \sup_u \langle u, Jv \rangle, \tag{16}$$

where the supremum is taken over all vectors u such that $u \in \text{span}\{Jv\}$ and $\|u\|_{K_v} \leq 1$. Note that we have used the fact that $(K_v^\circ)^\circ = K_v$. Next, we use

the fact that Jv is orthogonal to v (and hence in particular $\|Jv\|_{K_v} < \infty$) to deduce from (15) and (16) that

$$\inf_{w : \langle Jv, w \rangle = 1} \|w\|_{K_v^\circ} = \sup_{\substack{u \in \text{span}\{Jv\}, \\ \|u\|_{K_v} \leq 1}} \langle u, Jv \rangle \leq \frac{\langle Jv, Jv \rangle}{\|Jv\|_{K_v}} = \frac{1}{\|Jv\|_K}. \quad (17)$$

From the combination of (14) and (17) we conclude that

$$\bar{c}_{\text{lin}}(K) \leq 4 \inf_{v \neq 0} \frac{\|v\|_{K^\circ}}{\|Jv\|_K} = \frac{4}{\|J\|_{K^\circ \rightarrow K}}, \quad (18)$$

which completes the proof of the proposition. \square

Remark 2.7. For a general convex body K in \mathbb{R}^{2n} (not necessarily centrally symmetric), the same proof as the one above will give the following bound:

$$\frac{1}{4} \bar{c}_{\text{lin}}(K - K) \leq \bar{c}_{\text{lin}}(K) \leq \frac{1}{\|J\|_{(K-K)^\circ \rightarrow (K-K)}}. \quad (19)$$

On the other hand, from Proposition 2.2 it follows that

$$c_{\text{EHZ}}(K) \geq \sup_v \frac{1}{\|J\|_{(K-v)^\circ \rightarrow (K-v)}}, \quad (20)$$

where the supremum is taken over all $v \in \mathbb{R}^{2n}$ such that $v \in \text{Int}(K)$. We remark that although the upper bound for $\bar{c}_{\text{lin}}(K)$ in (19), and the lower bound for $c_{\text{EHZ}}(K)$ in (20) seem not too far away, we do not expect them to be asymptotically equivalent in general.

Proof of Theorem 1.6. For a smooth symmetric convex body K , the proof follows immediately from Propositions 2.2 and 2.5. The general case (i.e., without the smoothness assumption) follows by a standard approximation argument, as indicated at the beginning of this section. \square

3. Linearized symplectic capacities

In this section we prove Theorem 1.8. We recall first the following definition.

Definition 3.1. The linearized Gromov width $\underline{c}_{\text{lin}}$ of a set $\mathcal{U} \subset \mathbb{R}^{2n}$ is defined as

$$\underline{c}_{\text{lin}}(\mathcal{U}) = \sup_S \{\pi r^2 \mid SB^{2n}(r) \subset \mathcal{U}\},$$

where the supremum is taken over all affine symplectic maps $S \in \text{ISp}(2n)$.

The following is the main ingredient in the proof of Theorem 1.8.

Proposition 3.2. *Let $Q = [-1, 1]^{2n}$ be the standard cube in \mathbb{R}^{2n} . Then, for every orthogonal transformation $O \in O(2n)$ one has $\underline{c}_{\text{lin}}(OQ) \leq \pi$. Moreover, there is a rotation $\tilde{O} \in O(2n)$ for which $\underline{c}(\tilde{O}Q) \geq \sqrt{n/2}$.*

Proof of Proposition 3.2. Note first that for every orthogonal transformation $O \in O(2n)$,

$$\underline{c}_{\text{lin}}(OQ) \leq \sup_L \{ \pi r^2 \mid LB^{2n}(r) \subseteq OQ \}, \quad (21)$$

where the supremum is taken over all affine volume-preserving linear maps L of \mathbb{R}^{2n} . It is straightforward to check that the largest ellipsoid contained in the cube Q is the unit-ball $B^{2n}(1)$, and hence $\underline{c}_{\text{lin}}(OQ) \leq \pi$ for every orthogonal transformation $O \in O(2n)$.

For the second part of the proposition, consider the Lagrangian splitting $\mathbb{R}^n(q) \times \mathbb{R}^n(p)$ of \mathbb{R}^{2n} , and the following configuration: $B_\infty^n(\alpha) \times B_1^n(\beta) \subset \mathbb{R}^{2n}$, where

$$B_\infty^n(\alpha) = \{(q_1, \dots, q_n) \in \mathbb{R}^n(q) \mid \max\{|q_1|, \dots, |q_n|\} < \alpha\},$$

$$B_1^n(\beta) = \{(p_1, \dots, p_n) \in \mathbb{R}^n(p) \mid \sum_{i=1}^n |p_i| < \beta\}.$$

Note that $B_\infty^n(1) \times B_1^n(1)$ is the product of a hypercube and its dual body, the cross-polytope. It is known (see e.g., §4 of [16]) that for every $\varepsilon > 0$, the ball $B^{2n}(r)$ symplectically embeds (via a non-linear symplectomorphism) into the product $B_\infty^n(1) \times B_1^n(\beta(1 + \varepsilon))$, for a parameter β such that $\text{Vol}(B_\infty^n(1) \times B_1^n(\beta)) = \text{Vol}(B^{2n}(r))$. In particular, this implies that for an orthogonal transformation O of \mathbb{R}^{2n} , one has the following lower bound

$$\underline{c}(OQ) \geq \sup\{4r \mid B_\infty^n(1) \times B_1^n(r) \subseteq OQ\}.$$

Thus, to complete the proof of the proposition it is enough to find an orthogonal transformation $O \in O(2n)$ such that

$$O(B_\infty^n(1) \times B_1^n(\tilde{r})) \subseteq Q,$$

where $\tilde{r} \geq \sqrt{n/2}$. In particular, it is enough to find an orthogonal transformation O' of $\mathbb{R}^n(p)$ such that $O'(B_1^n(\tilde{r})) \subseteq [-1, 1]^n \subset \mathbb{R}^n(p)$, with \tilde{r} as above. The fact that such a transformation exists is well known to experts. For completeness we will give

an explicit construction². We define the elements O'_{kj} of the matrix O' by

$$\sqrt{n} O'_{kj} = \begin{cases} \sqrt{2} \sin\left(\frac{kj}{n} 2\pi\right) & \text{for } 1 \leq k < \frac{n}{2} \text{ and } 1 \leq j \leq n, \\ (-1)^j & \text{for } k = \frac{n}{2} \text{ and } 1 \leq j \leq n, \\ \sqrt{2} \cos\left(\frac{kj}{n} 2\pi\right) & \text{for } \frac{n}{2} < k < n \text{ and } 1 \leq j \leq n, \\ 1 & \text{for } k = n \text{ and } 1 \leq j \leq n. \end{cases} \quad (22)$$

It is a straightforward computation (based on the orthonormality of the standard Fourier basis) to check that the matrix O' defined by (22) is indeed an orthogonal transformation. Moreover, denote by $\{e_i\}_{i=1}^n$ the standard basis of $\mathbb{R}^n(p)$. Note that $B_1^n(1) = \text{Conv}\{\pm e_i\}$. It follows immediately from the definition of the matrix O' that

$$\|O'e_i\|_\infty := \max_{1 \leq j \leq n} |(O'e_i)_j| \leq \frac{\sqrt{2}}{\sqrt{n}},$$

where $(O'e_i)_j$ stands for the j -th component of the vector $O'e_i \in \mathbb{R}^n(p)$. This implies in particular that

$$O'(B_1^n(\sqrt{n})) = O'(\text{Conv}\{\pm \sqrt{n}e_i\}) = \text{Conv}\{\pm O' \sqrt{n}e_i\} \subseteq [-\sqrt{2}, \sqrt{2}]^n,$$

which completes the proof of Proposition 3.2. \square

Proof of Theorem 1.8. Note that an immediate corollary from Propositions 2.2 and 2.5 is that the cylindrical capacity \bar{c} is asymptotically equivalent to its linearized version \bar{c}_{lin} for symmetric convex domains in \mathbb{R}^{2n} , i.e., for every symmetric convex body $K \in \mathcal{K}^{2n}$,

$$\bar{c}(K) \leq \bar{c}_{\text{lin}}(K) \leq 4c_{\text{EHZ}}(K) \leq 4\bar{c}(K).$$

This establishes the first part of Theorem 1.8. The second part follows from Proposition 3.2. \square

References

- [1] S. Artstein-Avidan, R. N. Karasev and Y. Ostrover, From symplectic measurements to the Mahler conjecture, *Duke Math. Jour.*, **163** (2014), 2003–2022. Zbl 06345977 MR 3263026
- [2] S. Artstein-Avidan and Y. Ostrover, Bounds for Minkowski billiard trajectories in convex bodies, *Intern. Math. Res. Not. (IMRN)*, (2014), no. 1, 165–193. Zbl 06340328 MR 3158530

²The first named author learned this example from R.S. Ismagilov around 1976.

- [3] T. Cieliebak, H. Hofer, J. Latschev and F. Schlenk, Quantitative symplectic geometry, in *Dynamics, ergodic theory, and geometry*, 1–44, Math. Sci. Res. Inst. Publ., 54, Cambridge Univ. Press, Cambridge, 2007. Zbl 1143.53341 MR 2369441
- [4] I. Ekeland and H. Hofer, Symplectic topology and Hamiltonian dynamics, *Math. Z.*, **200** (1989), 355–378. Zbl 0641.53035 MR 0978597
- [5] I. Ekeland and H. Hofer, Symplectic topology and Hamiltonian dynamics. II, *Math. Z.*, **203** (1990), 553–567. Zbl 0729.53039 MR 1044064
- [6] U. Frauenfelder, V. Ginzburg and F. Schlenk, Energy capacity inequalities via an action selector, in *Geometry, spectral theory, groups, and dynamics*, 129–152, Contemp. Math., 387, Amer. Math. Soc., Providence, RI, 2005. Zbl 1094.53079 MR 2179791
- [7] A. Floer and H. Hofer, Symplectic homology, I. Open sets in \mathbb{C}^n , *Math. Z.*, **215** (1994), no. 1, 37–88. Zbl 0810.58013 MR 1254813
- [8] A. Floer, H. Hofer and K. Wysocki, Applications of symplectic homology. I, *Math. Z.*, **217** (1994), no. 4, 577–606. Zbl 0869.58012 MR 1306027
- [9] M. Gromov, Pseudoholomorphic curves in symplectic manifolds, *Invent. Math.*, **82** (1985), no. 2, 307–347. Zbl 0592.53025 MR 0809718
- [10] D. Hermann, *Non-equivalence of symplectic capacities for open sets with restricted contact type boundary*, Prépublication d’Orsay numéro 32 (29/4/1998).
- [11] H. Hofer, On the topological properties of symplectic maps, *Proc. Roy. Soc. Edinburgh Sect. A*, **115** (1990), 25–38. Zbl 0713.58004 MR 1059642
- [12] H. Hofer, Symplectic capacities, in *Geometry of low-dimensional manifolds, 2 (Durham, 1989)*, 15–34, London Math. Soc. Lecture Note Ser., 151, Cambridge Univ. Press, Cambridge, 1990. Zbl 0729.58026 MR 1171906
- [13] H. Hofer and E. Zehnder, *Symplectic Invariants and Hamiltonian Dynamics*, Birkhauser Advanced Texts, Birkhauser Verlag, 1994. Zbl 0805.58003 MR 1306732
- [14] H. Hofer and E. Zehnder, A new capacity for symplectic manifolds, in *Analysis, et cetera*, 405–427, Academic Press, Boston, MA, 1990. Zbl 0702.58021 MR 1039354
- [15] M. Hutchings, Quantitative embedded contact homology, *J. Diff. Geom.*, **88** (2011), 231–266. Zbl 1238.53061 MR 2838266
- [16] J. Latschev, D. McDuff and F. Schlenk, The Gromov width of 4-dimensional tori, *Geom. Topol.*, **17** (2013), 2813–2853. Zbl 1277.57024 MR 3190299
- [17] D. McDuff, *Symplectic topology today*, 2014 AMS Joint Mathematics Meeting.

- [18] D. McDuff and D. Salamon, *Introduction to Symplectic Topology*, 2nd edition, Oxford University Press, Oxford, England 1998. Zbl 1066.53137 MR 1698616
- [19] Y-G. Oh, Chain level Floer theory and Hofer's geometry of the Hamiltonian diffeomorphism group, *Asian J. Math.*, **6** (2002), no. 4, 579–624. Zbl 1038.53084 MR 1958084
- [20] Y. Ostrover, When symplectic topology meets Banach space geometry, in *Proceedings of the International Congress of Mathematicians (Seoul, 2014)* II, 959–981, Jang, S.Y.; Kim, Y.R., Lee, D.-W.; Yie, I. (Eds.), Kyung Moon SA Co. Ltd, Seoul, Korea 2014.
- [21] F. Schlenk, *Embedding Problems in Symplectic Geometry*, de Gruyter Expositions in Mathematics, 40, Berlin, 2005. Zbl 1073.53117 MR 2147307
- [22] R. Schneider, *Convex Bodies: the Brunn-Minkowski Theory*, Second expanded edition, Encyclopedia of Mathematics and its Applications, 151, Cambridge University Press, Cambridge, 2014. Zbl 1287.52001 MR 3155183
- [23] K. F. Siburg, Symplectic capacities in two dimensions, *Manuscripta Math.*, **78** (1993), no. 2, 149–163. Zbl 0791.58043 MR 1202158
- [24] C. A. Rogers and G. C. Shephard, Convex bodies associated with a given convex body, *J. London Soc.*, **33** (1958), 270–281. Zbl 0083.38402 MR 0101508
- [25] C. Viterbo, Metric and isoperimetric problems in symplectic geometry, *J. Amer. Math. Soc.*, **13** (2000), no. 2, 411–431. Zbl 0964.53050 MR 1750956
- [26] C. Viterbo, Symplectic topology as the geometry of generating functions, *Math. Ann.*, **292** (1992), 685–710. Zbl 0735.58019 MR 1157321

Received September 10, 2015

E. D. Gluskin, School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel
E-mail: gluskin@post.tau.ac.il

Y. Ostrover, School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel
E-mail: ostrover@post.tau.ac.il