Zeitschrift:	Commentarii Mathematici Helvetici
Herausgeber:	Schweizerische Mathematische Gesellschaft
Band:	91 (2016)
Heft:	1
Artikel:	On Serre's injectivity question and norm principle
Autor:	Bhaskhar, Nivedita
DOI:	https://doi.org/10.5169/seals-630555

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 20.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

On Serre's injectivity question and norm principle

Nivedita Bhaskhar*

Abstract. Let k be a field of characteristic not 2. We give a positive answer to Serre's injectivity question for any smooth connected reductive k-group whose Dynkin diagram contains connected components only of type A_n , B_n or C_n . We do this by relating Serre's question to the norm principles proved by Barquero and Merkurjev. We give a scalar obstruction defined up to spinor norms whose vanishing will imply the norm principle for the non-trialitarian D_n case and yield a positive answer to Serre's question for connected reductive k-groups whose Dynkin diagrams contain components of (non-trialitarian) type D_n too. We also investigate Serre's question for quasi-split reductive k-groups.

Mathematics Subject Classification (2010). 14L35, 20G10, 20G15.

Keywords. Principal homogeneous spaces, Serre's question, zero cycles, norm principles, spinor norm, Galois cohomology.

1. Introduction

Let k be a field. Then the following question of Serre, which is open in general, asks

Question 1.1 (Serre, [13, p. 233]). Let G be any connected linear algebraic group over a field k. Let L_1, L_2, \ldots, L_r be finite field extensions of k of degrees d_1, d_2, \ldots, d_r respectively such that $gcd_i(d_i) = 1$. Then is the following sequence exact?

$$1 \to \mathrm{H}^{1}(k, G) \to \prod_{i=1}^{r} \mathrm{H}^{1}(L_{i}, G).$$

The classical result that the index of a central simple algebra divides the degrees of its splitting fields answers Serre's question affirmatively for the group PGL_n. Springer's theorem for quadratic forms answers it affirmatively for the (albeit sometimes disconnected) group O(q) and Bayer–Lenstra's theorem [2], for the groups of isometries of algebras with involutions. Jodi Black [3] answers Serre's question positively for absolutely simple simply connected and adjoint *k*-groups of classical type. In this paper, we use and extend Jodi's result to connected reductive *k*-groups whose Dynkin diagram contains connected components only of type A_n , B_n or C_n .

^{*}The author acknowledges support from the NSF-FRG grant 1463882.

Theorem 1.2. Let k be a field of characteristic not 2. Let G be a connected reductive k-group whose Dynkin diagram contains connected components only of type A_n , B_n or C_n . Then Serre's question has a positive answer for G.

We also investigate Serre's question for reductive k-groups whose derived subgroups admit quasi-split simply connected covers. More precisely, we give a uniform proof for the following :

Theorem 1.3. Let k be a field of characteristic not 2. Let G be a connected quasi-split reductive k-group whose Dynkin diagram does not contain connected components of type E_8 . Then Serre's question has a positive answer for G.

We relate Serre's question for G with the norm principles of other closely related groups following a series of reductions previously used by Barquero and Merkurjev to prove the norm principles for reductive groups whose Dynkin diagrams do not contain connected components of type D_n , E_6 or E_7 [1]. We also give a scalar obstruction defined up to spinor norms whose vanishing will imply the norm principle for the (non-trialitarian) D_n case and yield a positive answer to Serre's question for connected reductive k-groups whose Dynkin diagrams contain components of this type also.

In the next section, we begin with some lemmata and preliminary reductions. In Section 3, we introduce intermediate groups \hat{G} and \tilde{G} and relate Serre's question for G to Serre's question for \hat{G} and \tilde{G} via the norm principle. In Section 4, we investigate the norm principle for (non-trialitarian) type D_n groups and find the scalar obstruction whose vanishing will imply the norm principle for the (non-trialitarian) D_n case. In the final section, we use the reduction techniques used in Sections 2 and 3 to discuss Serre's question for connected reductive k-groups whose derived subgroups admit quasi-split simply connected covers.

2. Preliminaries

We work over the base field k of characteristic not 2. By a k-group, we mean a smooth connected linear algebraic group defined over k. And mostly, we will restrict ourselves to reductive groups. We say that a k-group G satisfies SQ if Serre's question has a positive answer for G.

2.1. Reduction to characteristic 0. Let *G* be a connected reductive *k*-group whose Dynkin diagram contains connected components only of type A_n , B_n , C_n or (non-trialitarian) D_n . Without loss of generality we may assume that *k* is of characteristic 0 [7, p. 47]. We give a sketch of the reduction argument for the sake of completeness.

Suppose that the characteristic of k is p > 0. Let L_1, L_2, \ldots, L_r be finite field extensions of k of degrees d_1, d_2, \ldots, d_r respectively such that $gcd_i(d_i) = 1$ and

let ξ be an element in the kernel of

$$\mathrm{H}^{1}(k,G) \to \prod_{i=1}^{r} \mathrm{H}^{1}(L_{i},G).$$

By a theorem of Gabber, Liu and Lorenzini [5, Thm. 9.2] which was pointed out to us by O. Wittenberg, we note that any torsor under a smooth group scheme G/kwhich admits a zero-cycle of degree 1 also admits a zero-cycle of degree 1 whose support is étale over k. Thus without loss of generality we can assume that the given coprime extensions L_i/k are in fact separable.

By [10, Thms. 1 & 2], there exists a complete discrete valuation ring R with residue field k and fraction field K of characteristic zero. Let S_i denote corresponding étale extensions of R with residue fields L_i and fraction fields K_i .

There exists a smooth *R*-group scheme \tilde{G} with special fiber *G* and connected reductive generic fiber \tilde{G}_K . Now given any torsor $t \in H^1(k, G)$, there exists a torsor $\tilde{t} \in H^1_{\text{ét}}(R, \tilde{G})$ specializing to *t* which is unique upto isomorphism. This in turn gives a torsor \tilde{t}_K in $H^1(K, \tilde{G}_K)$ by base change, thus defining a map $i_k : H^1(k, G) \to$ $H^1(K, \tilde{G}_K)$ [6, p. 29]. It clearly sends the trivial element to the trivial element. The map *i* also behaves well with the natural restriction maps, i.e., it fits into the following commutative diagram :

$$\begin{array}{c} \mathrm{H}^{1}(k,G) \xrightarrow{i_{k}} \mathrm{H}^{1}(K,\tilde{G}_{K}) \\ \downarrow & \downarrow \\ \prod \mathrm{H}^{1}(L_{i},G) \xrightarrow{\prod i_{L_{i}}} \prod \mathrm{H}^{1}(K_{i},\tilde{G}_{K}). \end{array}$$

Let $\tilde{\xi}$ denote the torsor in $\mathrm{H}^{1}_{\mathrm{\acute{e}t}}(R, \tilde{G})$ corresponding to ξ as above. Therefore $\tilde{\xi}_{K} := i_{k}(\xi)$ is in the kernel of

$$\mathrm{H}^{1}(K, \tilde{G}_{K}) \to \prod_{i=1}^{\prime} \mathrm{H}^{1}(K_{i}, \tilde{G}_{K}).$$

Suppose that \tilde{G}_K satisfies SQ. Then $\tilde{\xi}_K$ is trivial. However by [12], the natural map $\mathrm{H}^1_{\mathrm{\acute{e}t}}(R, \tilde{G}) \to \mathrm{H}^1(K, \tilde{G}_K)$ is injective and hence $\tilde{\xi}$ is trivial in $\mathrm{H}^1_{\mathrm{\acute{e}t}}(R, \tilde{G})$. This implies that its specialization, ξ , is trivial in $\mathrm{H}^1(k, G)$.

Thus from here on, we assume that the base field k has characteristic 0.

2.2. Lemmata.

Lemma 2.1. Let k-groups G and H satisfy SQ. Then $G \times_k H$ also satisfies SQ.

Proof. Let L/k be a field extension. Then the map

$$\mathrm{H}^{1}(k, G \times_{k} H) \to \mathrm{H}^{1}(L, G \times_{k} H)$$

is precisely the product of the maps

$$\mathrm{H}^{1}(k,G) \to \mathrm{H}^{1}(L,G)$$
 and $\mathrm{H}^{1}(k,H) \to \mathrm{H}^{1}(L,H)$.

This immediately shows that if G and H satisfy SQ, so does $G \times_k H$.

Lemma 2.2. Let $1 \rightarrow Q \rightarrow H \rightarrow G \rightarrow 1$ be a central extension of a k-group G by a quasi-trivial torus Q. Then H satisfies SQ if and only if G satisfies SQ.

Proof. Let L_i be field extensions of k such that $gcd[L_i : k] = 1$. Since Q is quasitrivial, $H^1(L, Q) = \{1\} \forall L/k$. From the long exact sequence in cohomology, we have the following commutative diagram.

From the above diagram, it is clear that if G satisfies SQ, so does H.

Conversely, assume that H satisfies SQ. Let $a \in H^1(k, G)$ become trivial in $\prod H^1(L_i, G)$. Then $\delta_k(a)$ becomes trivial in each $H^2(L_i, Q)$. Hence the corestriction $\operatorname{Cor}_{L_i/k}(\delta_k(a)) = \delta_k(a)^{d_i}$ becomes trivial in $H^2(k, Q)$ where $d_i = [L_i : k]$. Since $\operatorname{gcd}_i(d_i) = 1$, this implies that $\delta_k(a)$ is itself trivial in $H^2(k, Q)$. Therefore a comes from an element $b \in H^1(k, H)$ which is trivial in $\prod H^1(L_i, H)$. (The fact that $H^1(L_i, Q) = \{1\}$ guarantees that b is trivial in $H^1(L_i, H)$.) Since Hsatisfies SQ by assumption, b is trivial in $H^1(k, H)$ which implies the triviality of ain $H^1(k, G)$.

Lemma 2.3. Let *E* be a finite separable field extension of *k* and let *H* be an *E*-group satisfying SQ. Then the *k*-group $R_{E/k}(H)$ also satisfies SQ.

Proof. Set $G = R_{E/k}(H)$ and let ξ be an element in the kernel of $H^1(k, G) \rightarrow \prod_{i=1}^{r} H^1(L_i, G)$ where $gcd_i[L_i:k] = 1$.

Since char(k) = 0, $L_i \otimes_k E$ is an étale *E*-algebra and hence isomorphic to $E_{1,i} \times E_{2,i} \times \cdots \times E_{n_i,i}$ where each $E_{j,i}$ is a separable field extension of *E*. Thus $\sum_{j=1}^{n_i} [E_{j,i} : E] = [L_i : k]$ and therefore gcd $[E_{j,i} : E] = 1$ where $1 \le i \le r$ and $1 \le j \le n_i$.

By Eckmann–Faddeev–Shapiro, we have a natural bijection of pointed sets

$$H^{1}(k,G) \simeq H^{1}(E,H),$$
$$H^{1}(L_{i},G) \simeq \prod_{j=1}^{n_{i}} H^{1}(E_{j,i},H)$$

Thus we have that ξ is in the kernel of $H^1(E, H) \to \prod_{i \le r, j \le n_i} H^1(E_{j,i}, H)$. Since *H* satisfies *SQ*, we see that ξ is trivial.

Vol. 91 (2016)

3. Serre's question and norm principles

3.1. Intermediate groups \hat{G} and \tilde{G} . Notations are as in Section 5 of [1].

Let G be our given connected reductive k-group whose Dynkin diagram contains connected components only of type A_n , B_n , C_n or (non-trialitarian) D_n and let G' denote its derived subgroup. Let Z(G) = T and $Z(G') = \mu$.

Let $\rho : \mu \hookrightarrow S$ be an embedding of μ into a quasi-trivial torus S. We denote the cofibre product $e(G', \rho) = \frac{G' \times S}{\mu}$ by \hat{G} . This *k*-group is called an *envelope* of G'.

$$\begin{array}{c} \mu \xrightarrow{\delta} G' \\ \downarrow^{\rho} \qquad \downarrow \\ S \xrightarrow{\gamma} \hat{G} \end{array}$$

Now the quasi-trivial torus $S = Z(\hat{G})$ and \hat{G} fit into an exact sequence as follows:

$$1 \to S \to \hat{G} \to G'^{ad} \to 1 \tag{(*)}$$

where G'^{ad} corresponds to the adjoint group of G'. We now recall the following result of Jodi Black which addresses Serre's question for adjoint groups of classical type.

Theorem 3.1 (Jodi Black, [3, Thm. 0.2]). Let k be a field of characteristic different from 2 and let J be an absolutely simple algebraic k-group which is not of type E_8 and which is either a simply connected or adjoint classical group or a quasi-split exceptional group. Then Serre's question has a positive answer for J.

Since every adjoint group of classical type is a product of Weil restrictions of absolutely simple adjoint groups, the above theorem, along with Lemmata 2.1 and 2.3, implies that G'^{ad} satisfies SQ. Applying Lemma 2.2 to the exact sequence (*) above, we see that \hat{G} satisfies SQ. Let us chose such an envelope \hat{G} of G' which satisfies SQ.

Define an intermediate abelian group \tilde{T} to be the cofibre product $\frac{T \times S}{\mu}$.

$$\begin{array}{ccc} \mu & \longrightarrow & T \\ \downarrow^{\rho} & & \downarrow^{\alpha} \\ S & \stackrel{\nu}{\longrightarrow} & \tilde{T} \end{array}$$

Let the algebraic group \tilde{G} be the cofibre product defined by the following diagram:

$$\begin{array}{ccc} G' \times T & \stackrel{m}{\longrightarrow} & G \\ & & \downarrow^{id \times \alpha} & & \downarrow^{\beta} \\ G' \times \tilde{T} & \stackrel{\epsilon}{\longrightarrow} & \tilde{G}. \end{array}$$

Then we have the following commutative diagram with exact rows [1, Prop. 5.1]. Note that each row is a central extension of \tilde{G} .

$$1 \longrightarrow \mu \xrightarrow{\delta, \nu \rho} G' \times \tilde{T} \xrightarrow{\epsilon} \tilde{G} \longrightarrow 1 \qquad (**)$$

 $1 \longrightarrow S \xrightarrow{\gamma, \nu} \hat{G} \times \tilde{T} \longrightarrow \tilde{G} \longrightarrow 1 \qquad (* * *)$

Since \tilde{T} is abelian, the existence of the co-restriction map shows that \tilde{T} satisfies SQ. Since \hat{G} satisfies SQ, we can apply Lemmata 2.1 and 2.2 to (* * *) to see that \tilde{G} satisfies SQ.

3.2. Norm principle and weak norm principle. Let $f : G \to T$ be a map of k-groups where T is an abelian k-group. Then we have norm maps $N_{L/k}: T(L) \to T(k)$ for any separable field extension L/k.

$$\begin{array}{ccc} G(L) & \xrightarrow{f(L)} & T(L) \\ & & & & \downarrow^{N_{L/k}} \\ G(k) & \xrightarrow{f(k)} & T(k) \end{array}$$

We say that the *norm principle* holds for $f : G \to T$ if for all separable field extensions L/k,

$$N_{L/k}$$
(Image $f(L)$) \subseteq Image $f(k)$.

That is, we say that the *norm principle* holds for $f : G \to T$ if given any separable field extension L/k and any $t \in T(L)$ such that

t
$$\in$$
 (Image $f(L) : G(L) \to T(L)$),
then $N_{L/k}(t) \in$ (Image $f(k) : G(k) \to T(k)$).

Note that the norm principle holds for any algebraic group homomorphism between abelian groups.

We say that the *weak norm principle* holds for $f : G \to T$ if given any $t \in T(k)$ such that

$$t \in (\text{Image } f(L) : G(L) \to T(L)),$$

$$t^{[L:k]} = N_{L/k}(t) \in (\text{Image } f(k) : G(k) \to T(k)).$$

It is clear that if the norm principle holds for f, then so does the weak norm principle.

then

150

CMH

3.3. Relating Serre's question and norm principle. The deduction of SQ for *G* from \hat{G} and \tilde{G} follows via the (weak) norm principles.

Let $\beta : G \to \tilde{G}$ be the embedding of k-groups with the cokernel P isomorphic to the torus $\frac{S}{\mu}$ where \tilde{G} and G are as in Section 3.1. Thus we have the following exact sequence:

$$1 \to G \xrightarrow{\beta} \tilde{G} \xrightarrow{\pi} P \to 1.$$

Lemma 3.2. If the weak norm principle holds for $\pi : \tilde{G} \to P$, then G satisfies SQ.

Proof. From the long exact sequence of cohomology, we have the following commutative diagram:

Let $a \in H^1(k, G)$ become trivial in $\prod H^1(L_i, G)$. As \tilde{G} satisfies SQ, $\beta_k(a)$ becomes trivial in $H^1(k, \tilde{G})$. Hence $a = \delta_k(b)$ for some $b \in P(k)$ and $\delta_{L_i}(b)$ is trivial in $H^1(L_i, G)$. Therefore, there exist $c_i \in \tilde{G}(L_i)$ such that $\pi_{L_i}(c_i) = b$.

Showing that G satisfies SQ, i.e. that a is trivial, is equivalent to showing

$$b \in (\operatorname{Image} \pi_k : \tilde{G}(k) \to P(k)).$$

However $b \in (\text{Image } \pi_{L_i} : \tilde{G}(L_i) \to P(L_i))$. Since the weak norm principle holds for $\pi : \tilde{G} \to P, b^{d_i} \in \text{Image } (\pi_k : \tilde{G}(k) \to P(k))$ where $[L_i : k] = d_i$ for each *i*. As $\text{gcd}_i(d_i) = 1$, this means $b \in \text{Image } (\pi_k : \tilde{G}(k) \to P(k))$.

We recall now the norm principle of Merkurjev and Barquero for reductive groups of classical type.

Theorem 3.3 (Barquero–Merkurjev, [1]). Let G be a reductive group over a field k. Assume that the Dynkin diagram of G does not contain connected components D_n , $n \ge 4$, E_6 or E_7 . Let T be any commutative k-group. Then the norm principle holds for any group homomorphism $G \rightarrow T$.

This shows that the norm principle and hence the weak norm principle holds for the map $\pi : \tilde{G} \to P$ for reductive k-groups G as in the main theorem (Theorem 1.2). Thus we have concluded the proof for the following:

Theorem 1.2. Let k be a field of characteristic not 2. Let G be a connected reductive k-group whose Dynkin diagram contains connected components only of type A_n , B_n or C_n . Then Serre's question has a positive answer for G.

4. Obstruction to norm principle for (non-trialitarian) D_n

4.1. Preliminaries. Let (A, σ) be a central simple algebra of degree 2n over k and let σ be an orthogonal involution. Let $C(A, \sigma)$ denote its Clifford algebra which is a central simple algebra over its center, Z/k, the discriminant extension. Let i denote the non-trivial automorphism of Z/k and let σ denote the canonical involution of $C(A, \sigma)$.

Recall that, depending on the parity of $n, \underline{\sigma}$ is either an involution of the second kind (when *n* is odd) or of the first kind (when *n* is even). Let $\underline{\mu}$: Sim (C (A, σ), $\underline{\sigma}$) $\rightarrow R_{Z/k}\mathbb{G}_m$ denote the multiplier map sending similitude *c* to $\underline{\sigma}(c)c$.

Let $\Omega(A, \sigma)$ be the *extended Clifford group*. Note that this has center $R_{Z/k}\mathbb{G}_m$ and is an *envelope* of Spin (A, σ) [1, Ex. 4.4]. We recall below the map $\varkappa : \Omega(A, \sigma)(k) \to Z^*/k^*$ as defined in [9, p. 182].

Given $\omega \in \Omega(A, \sigma)(k)$, let $g \in GO^+(A, \sigma)(k)$ be some similitude such that $\omega \rightsquigarrow gk^*$ under the natural surjection $\Omega(A, \sigma)(k) \rightarrow PGO^+(A, \sigma)(k)$.

Let $h = \mu(g)^{-1}g^2 \in O^+(A, \sigma)(k)$ and let $\gamma \in \Gamma(A, \sigma)(k)$ be some element in the *special Clifford group* which maps to h under the vector representation $\chi' : \Gamma(A, \sigma)(k) \to O^+(A, \sigma)(k)$. Then $\omega^2 = \gamma z$ for some $z \in Z^*$ and $\varkappa(\omega) = zk^*$.

Note that the map \varkappa has Γ (A, σ) (k) as kernel. Also if $z \in Z^*$, then $\varkappa(z) = z^2 k^*$.

By following the reductions in [1], it is easy to see that one needs to investigate whether the norm principle holds for the canonical map

$$\Omega (\mathbf{A}, \sigma) \to \frac{\Omega (\mathbf{A}, \sigma)}{\left[\Omega (\mathbf{A}, \sigma), \Omega (\mathbf{A}, \sigma)\right]}.$$

We will need to investigate the norm principle for two different maps depending on the parity of n.

The map μ_* for *n* odd. Let $U \subset \mathbb{G}_m \times R_{Z/k} \mathbb{G}_m$ be the algebraic subgroup defined by

$$U(k) = \{ (f, z) \in k^* \times Z^* | f^4 = N_{Z/k}(z) \}.$$

Recall the map $\mu_* : \Omega(A, \sigma) \to U$ defined in [9, p. 188] which sends

$$\omega \rightsquigarrow \left(\underline{\mu}(\omega), ai(a)^{-1}\underline{\mu}(\omega)^2\right),$$

where $\omega \in \Omega(A, \sigma)(k)$ and $\varkappa(\omega) = a k^*$. This induces the following exact sequence [9, p. 190]

$$1 \to \operatorname{Spin} (\mathbf{A}, \sigma) \to \Omega (\mathbf{A}, \sigma) \xrightarrow{\mu_*} U \to 1.$$

Since the semisimple part of $\Omega(A, \sigma)$ is Spin (A, σ) , the above exact sequence shows that it suffices to check the norm principle for the map μ_* .

The map μ for *n* even. Recall the following exact sequence induced by restricting μ to Ω (A, σ) [9, p. 187]

$$1 \to \operatorname{Spin}(\mathcal{A}, \sigma) \to \Omega(\mathcal{A}, \sigma) \xrightarrow{\underline{\mu}} R_{Z/k} \mathbb{G}_m \to 1.$$

Since the semisimple part of $\Omega(A, \sigma)$ is Spin (A, σ) , the above exact sequence shows that it suffices to check the norm principle for the map μ .

4.2. An obstruction to being in the image of μ_* for *n* odd. Given $(f, z) \in U(k)$, we would like to formulate an obstruction which prevents (f, z) from being in the image $\mu_* (\Omega (A, \sigma) (k))$. Note that for $z \in Z^*$, $\mu_*(z) = (N_{Z/k}(z), z^4)$ and hence the algebraic subgroup $U_0 \subseteq U$ defined by

$$U_0(k) = \{ (N_{Z/k}(z), z^4) | z \in Z^* \}$$

has its *k*-points in the image $\mu_* (\Omega(A, \sigma)(k))$.

Let $\mu_{n[Z]}$ denote the kernel of the norm map $R_{K/k}\mu_n \xrightarrow{N} \mu_n$ where K/k is a quadratic extension. Note that $\mu_{4[Z]}$ is the center of Spin (A, σ) as *n* is odd. Also recall that [9, Prop. 30.13, p. 418]

$$\mathrm{H}^{1}\left(k,\mu_{4[Z]}\right) \cong \frac{U(k)}{U_{0}(k)}.$$

Thus, we can construct the map $S : PGO^+(A, \sigma)(k) \to H^1(k, \mu_{4[Z]})$ induced by the following commutative diagram with exact rows:

$$1 \longrightarrow Z^{*} \longrightarrow \Omega(A, \sigma)(k) \xrightarrow{\chi'} PGO^{+}(A, \sigma)(k) \longrightarrow 1$$
$$\downarrow^{\mu_{*}} \qquad \qquad \downarrow^{\mu_{*}} \qquad \qquad \downarrow^{S}$$
$$1 \longrightarrow U_{0}(k) \longrightarrow U(k) \longrightarrow H^{1}(k, \mu_{4[Z]}) \longrightarrow 1$$

The map *S* also turns out to be the connecting map from PGO⁺ (A, σ) (*k*) \rightarrow H¹ (*k*, $\mu_{4[Z]}$) [9, Prop. 13.37, p. 190] in the long exact sequence of cohomology corresponding to the exact sequence

$$1 \rightarrow \mu_{4[Z]} \rightarrow \text{Spin}(A, \sigma) \rightarrow \text{PGO}^+(A, \sigma) \rightarrow 1.$$

Since the maps $\mu_* : Z^* \to U_0(k)$ and $\chi' : \Omega(A, \sigma)(k) \to PGO^+(A, \sigma)(k)$ are surjective, an element $(f, z) \in U(k)$ is in the image $\mu_*(\Omega(A, \sigma)(k))$ if and only if its image $[f, z] \in H^1(k, \mu_{4[Z]})$ is in the image $S(PGO^+(A, \sigma)(k))$.

Therefore we look for an obstruction preventing [f, z] from being in the image $S(\text{PGO}^+(A, \sigma)(k))$. Recall the following commutative diagram with exact rows and columns:



The long exact sequence of cohomology induces the following commutative diagram (Figure 1) with exact columns [9, Prop. 13.36, p. 189], where



Figure 1. Spinor norms and S for *n* odd

 $\mu: \operatorname{PGO}^+(A, \sigma)(k) \to \frac{k^*}{k^{*2}} \text{ is induced by the multiplier map } \mu: \operatorname{GO}^+(A, \sigma) \to \mathbb{G}_m$ $i: \frac{k^*}{k^{*2}} \to \operatorname{H}^1\left(k, \mu_{4[Z]}\right) = \frac{U(k)}{U_0(k)} \text{ is the map sending } fk^{*2} \to [f, f^2]$ $j: \frac{U(k)}{U_0(k)} = \operatorname{H}^1\left(k, \mu_{4[Z]}\right) \to \frac{k^*}{k^{*2}} \text{ is the map sending } [f, z] \to \operatorname{N}(z_0)k^{*2},$ where $z_0 \in Z^*$ is such that $z_0i(z_0)^{-1} = f^{-2}z$.

Definition 4.1. We call an element $(f, z) \in U(k)$ to be *special* if there exists a $[g] \in PGO^+(A, \sigma)(k)$ such that $j([f, z]) = \mu([g])$.

Vol. 91 (2016)

Let $(f, z) \in U(k)$ be a special element and let $[g] \in PGO^+(A, \sigma)(k)$ be such that $j([f, z]) = \mu([g])$. From the discussion above, it is clear that (f, z) is in the image $\mu_*(\Omega(A, \sigma)(k))$ if and only if [f, z] is in the image $S(PGO^+(A, \sigma)(k))$.

Thus $S([g])[f, z]^{-1}$ is in kernel j = Image i and hence there exists some $\alpha \in k^*$ such that

$$[f, z] = S([g])[\alpha, \alpha^2] \in \frac{U(k)}{U_0(k)}$$

Note that if g is changed by an element in $O^+(A, \sigma)(k)$, then α changes by a spinor norm by Figure 1 above. Thus given a special element, we have produced a scalar $\alpha \in k^*$ which is well defined upto spinor norms.

$$[f, z] \in S \left(\text{PGO}^+ (A, \sigma) (k) \right) \iff [\alpha, \alpha^2] \in S \left(\text{PGO}^+ (A, \sigma) (k) \right)$$
$$\iff (\alpha, \alpha^2) \in \mu_* \left(\Omega (A, \sigma) (k) \right).$$

This happens if and only if there exists $w \in \Omega(A, \sigma)(k)$ such that

$$\alpha = \underline{\mu}(w)$$

$$\alpha^{2} = \varkappa(w)i(\varkappa(w))^{-1}\underline{\mu}(w)^{2}$$

This implies $\varkappa(w) \in k^*$ and hence $w \in \Gamma(A, \sigma)(k)$. Thus α is a spinor norm, being the similarity of an element in the special Clifford group. Also note if α is a spinor norm, then $\alpha = \mu(\gamma)$ for some $\gamma \in \Gamma(A, \sigma)(k)$ and $\mu_*(\gamma) = (\mu(\gamma), \mu(\gamma)^2)$.

Thus a special element (f, z) is in the image of μ_* if and only if the produced scalar α is a spinor norm. We call the class of α in $\frac{k^*}{\operatorname{Sn}(A,\sigma)}$ to be the scalar obstruction preventing the *special* element $(f, z) \in U(k)$ from being in the image $\mu_* (\Omega(A, \sigma)(k))$.

4.3. An obstruction to being in the image of μ for *n* even. Given $z \in Z^*$, we would like to formulate an obstruction which prevents *z* from being in the image $\mu(\Omega(A, \sigma)(k))$. Note that for $z \in Z^*$, $\mu(z) = z^2$ and hence the subgroup Z^{*2} is in the image $\mu(\Omega(A, \sigma)(k))$.

Like in the case of odd *n*, we can construct the map $S : \text{PGO}^+(A, \sigma)(k) \to \frac{Z^*}{Z^{*2}}$ induced by the following commutative diagram with exact rows [9, Def. 13.32, p. 187]:

Again by the surjectivity of the maps, $\underline{\mu} : Z^* \to Z^{*2}$ and $\chi' : \Omega(A, \sigma)(k) \to PGO^+(A, \sigma)(k)$, an element $z \in Z^*$ is in the image $\underline{\mu}(\Omega(A, \sigma)(k))$ if and only

if its image $[z] \in \frac{Z^*}{Z^{*2}}$ is in the image $S(\text{PGO}^+(A, \sigma)(k))$. Therefore we look for an obstruction preventing [z] from being in the image $S(\text{PGO}^+(A, \sigma)(k))$. And as before, we arrive at the the following commutative diagram (Figure 2) with exact rows and columns [9, Prop. 13.33, p. 188], where



Figure 2. Spinor norms and S for *n* even

 $\mu: \operatorname{PGO}^+(A, \sigma)(k) \to \frac{k^*}{k^{*2}} \text{ is induced by the multiplier map } \mu: \operatorname{GO}^+(A, \sigma) \to \mathbb{G}_m$ $i: \frac{k^*}{k^{*2}} \to \frac{Z^*}{Z^{*2}} \text{ is the inclusion map}$

 $j: \frac{Z^*}{Z^{*2}} \to \frac{k^*}{k^{*2}}$ is induced by the norm map from $Z^* \to k^*$.

Definition 4.2. We call an element $z \in Z^*$ to be *special* if there exists a $[g] \in PGO^+(A, \sigma)(k)$ such that $j([z]) = \mu([g])$.

Let $z \in Z^*$ be a special element and let $[g] \in PGO^+(A, \sigma)(k)$ be such that $j([z]) = \mu([g])$. As before a *special* element $z \in Z^*$ is in the image $\underline{\mu}(\Omega(A, \sigma)(k))$ if and only if [z] is in the image $S(PGO^+(A, \sigma)(k))$.

Thus $S([g])[z]^{-1}$ is in kernel j = Image i and hence there exists some $\alpha \in k^*$ such that

$$[z] = S([g])[\alpha] \in \frac{Z^*}{Z^{*2}}.$$

Note that if g is changed by an element in $O^+(A, \sigma)(k)$, then α changes by a spinor norm by Figure 2 above. Thus given a special element, we have produced a scalar $\alpha \in k^*$ which is well defined up to spinor norms.

$$[z] \in S \left(\mathsf{PGO}^+ (\mathsf{A}, \sigma) (k) \right) \iff [\alpha] \in S \left(\mathsf{PGO}^+ (\mathsf{A}, \sigma) (k) \right)$$
$$\iff (\alpha) \in \mu \left(\Omega \left(\mathsf{A}, \sigma \right) (k) \right).$$

Since $\alpha \in k^*$ also, this is equivalent to α being a spinor norm [9, Prop. 13.25, p. 184].

We call the class of α in $\frac{k^*}{\operatorname{Sn}(A,\sigma)}$ to be the scalar obstruction preventing the *special* element $z \in Z^*$ from being in the image $\underline{\mu} (\Omega (A, \sigma) (k))$.

4.4. Scharlau's norm principle for μ : **GO**⁺(**A**, σ) $\rightarrow \mathbb{G}_m$. Let μ : **GO**⁺(**A**, σ) $\rightarrow \mathbb{G}_m$ denote the multiplier map and let L/k be a separable field extension of finite degree. Let $g_1 \in GO^+(A, \sigma)(L)$ be such that $\mu(g_1) = f_1 \in L^*$. Let f denote $N_{L/k}(f_1)$. We would like to show that f is in the image μ (GO⁺(A, σ)(k)).

Note that by a generalization of Scharlau's norm principle ([9, Prop. 12.21]; [3, Lemma 4.3]) there exists a $\tilde{g} \in \text{GO}(A, \sigma)(k)$ such that $f = \mu(\tilde{g})$. However we would like to find a *proper* similitude $g \in \text{GO}^+(A, \sigma)(k)$ such that $\mu(g) = f$.

We investigate the cases when the algebra A is non-split and split separately.

Case I: *A* is non-split. Note that $g_1 \in \text{GO}^+(A, \sigma)(L)$. If $\tilde{g} \in \text{GO}^+(A, \sigma)(k)$, we are done. Hence assume $\tilde{g} \notin \text{GO}^+(A, \sigma)(k)$. By a generalization of Dieudonné's theorem [9, Thm. 13.38, p. 190], we see that the quaternion algebras

$$B_1 = (Z, f_1) = 0 \in Br(L),$$

 $B_2 = (Z, f) = A \in Br(k).$

Since A is non-split, $B_2 \neq 0 \in Br(k)$. However co-restriction of B_1 from L to k gives a contradiction, because

$$0 = \operatorname{Cor} B_1 = (Z, \operatorname{N}_{L/k}(f_1)) = B_2 \in \operatorname{Br}(k).$$

Hence $\tilde{g} \in \mathrm{GO}^+(\mathrm{A},\sigma)(k)$.

Case II: *A* is split. Since *A* is split, A = End V where (V, q) is a quadratic space and σ is the adjoint involution for the quadratic form *q*. Again, if $\tilde{g} \in \text{GO}^+(A, \sigma)(k)$, we are done. Hence assume $\tilde{g} \notin \text{GO}^+(A, \sigma)(k)$. That is

$$\det(\tilde{g}) = -f^{2n/2} = -(f^n).$$

Since A is of even degree (2n) and split, there exists an isometry¹ h of determinant -1. Set $g = \tilde{g}h$. Then $det(g) = f^n$ where $\mu(g) = f$. Thus we have found a suitable $g \in GO^+(A, \sigma)(k)$ which concludes the proof of the following:

Theorem 4.3. The norm principle holds for the map $\mu : \mathrm{GO}^+(\mathrm{A}, \sigma) \to \mathbb{G}_m$.

4.5. Spinor obstruction to norm principle for non-trialitarian D_n . Let L/k be a separable field extension of finite degree. And let $w_1 \in \Omega(A, \sigma)(L)$ be such that for

n odd :
$$\mu_*(w_1) = \theta$$
 which is equal to $(f_1, z_1) \in U(L)$,
n even : $\mu(w_1) = \theta$ which is equal to $z_1 \in (R_{Z/k}\mathbb{G}_m)(L)$.

¹Since V is of even dimension 2n, h can be chosen to be a hyperplane reflection for instance

We would like to investigate whether $N_{L/k}(\theta)$ is in the image of $\mu_*(\Omega(A, \sigma)(k))$ (resp. $\underline{\mu}(\Omega(A, \sigma)(k))$) when *n* is odd (resp. even) in order to check if the norm principle holds for the map $\mu_*: \Omega(A, \sigma) \to U$ (resp. $\mu: \Omega(A, \sigma) \to R_{Z/k}\mathbb{G}_m$).

Let $[g_1] \in PGO^+(A, \sigma)(L)$ be the image of w_1 under the canonical map $\chi' : \Omega(A, \sigma)(L) \to PGO^+(A, \sigma)(L)$. Clearly θ is *special* and let $g_1 \in GO^+(A, \sigma)(L)$ be such that $\mu([g_1]) = j([\theta])$.

By Theorem 4.3, there exists a $g \in \text{GO}^+(A, \sigma)(k)$ such that²

$$\mu([g]) = \mathcal{N}_{L/k} \left(j\left[\theta\right] \right) = j\left(\left[\mathcal{N}_{L/k} \theta \right] \right).$$

Hence $N_{L/k}(\theta)$ is special.

By Subsection 4.2 (resp. 4.3), $N_{L/k}(\theta)$ is in the image of μ_* (resp $\underline{\mu}$) if and only if the scalar obstruction $\alpha \in \frac{k^*}{\operatorname{Sn}(A,\sigma)}$ defined for $N_{L/k}(\theta)$ vanishes. Thus we have a spinor norm obstruction given below.

Theorem 4.4 (Spinor norm obstruction). Let L/k be a finite separable extension of fields. Let f denote the map μ_* (resp μ) in the case when n is odd (resp. even). Given $\theta \in f(\Omega(A, \sigma)(L))$, there exists scalar obstruction $\alpha \in k^*$ such that

$$N_{L/k}(\theta) \in f(\Omega(\mathbf{A},\sigma)(k)) \iff \alpha = 1 \in \frac{k*}{\mathrm{Sn}(\mathbf{A},\sigma)}.$$

Thus the norm principle for the canonical map

$$\Omega\left(\mathbf{A},\sigma\right) \rightarrow \frac{\Omega\left(\mathbf{A},\sigma\right)}{\left[\Omega\left(\mathbf{A},\sigma\right),\Omega\left(\mathbf{A},\sigma\right)\right]}$$

and hence for non-trialitarian D_n holds if and only if the scalar obstructions are spinor norms.

5. Quasi-split groups

Let G be a connected reductive k-group whose Dynkin diagram does not contain connected components of type E_8 and let G' denote its derived subgroup. Let G^{sc} denote the simply connected cover of G'. Then one has the exact sequence $1 \rightarrow C \rightarrow G^{sc} \rightarrow G' \rightarrow 1$, where C is a finite k-group of multiplicative type, central in G^{sc} . Assuming that G^{sc} is quasi-split, we would like to show that G satisfies SQ by following the reduction techniques used in Sections 2 and 3.

Lemma 5.1. Let G be a connected reductive k-group. If G^{sc} is quasi-split, then there exists an extension $1 \rightarrow Q \rightarrow H \xrightarrow{\psi} G \rightarrow 1$, where Q is a quasi-trivial k-torus, central in reductive k-group H with H' simply connected and quasi-split.

²The map j commutes with $N_{L/k}$ in both cases.

Proof. Recall that there is a central extension (called a *z*-extension) of *G* by a quasitrivial torus Q such that H' is semisimple and simply connected ([11, Prop. 3.1] and [4, Lemma 1.1.4]).

$$1 \to Q \to H \xrightarrow{\psi} G \to 1.$$

The restriction $\psi|_{H'}: H' \to G$ yields the fact that H' is the simply connected cover of G' and hence is quasi-split.

Lemmata 2.2 and 5.1 imply that we can restrict ourselves to connected reductive k-groups G such that G' is simply connected and quasi-split.

Lemma 5.2. Let H be any reductive k-group such that its derived subgroup H' is semisimple simply connected and quasi-split. Let T denote the k-torus H/H'. Then the natural exact sequence $1 \rightarrow H' \rightarrow H \xrightarrow{\phi} T \rightarrow 1$ induces surjective maps $\phi(L) : H(L) \rightarrow T(L)$ for all field extensions L/k. In particular, the norm principle holds for $\phi : H \rightarrow T$.

Proof. There exists a quasi-trivial maximal torus Q_1 of H' defined over k [8, Lem. 6.7]. Let $Q_1 \subset Q_2$, where Q_2 is a maximal torus of H defined over k. The proof of [8, Lem. 6.6] shows that $\phi|_{Q_2} : Q_2 \to T$ is surjective and that $Q_2 \cap H'$ is a maximal torus of H'. Since $Q_2 \cap H' \subseteq Q_1$, we get the following extension of k-tori

$$1 \to Q_1 \to Q_2 \to T \to 1$$

Since Q_1 is quasitrivial, $H^1(L, Q_1) = 0$ for any field extension L/k which gives the surjectivity of $\phi(L) : Q_2(L) \to T(L)$ and hence of $\phi(L) : H(L) \to T(L)$. \Box

Let \hat{G} be an envelope of G' defined using an embedding of $\mu = Z(G')$ into a quasi-trivial torus S. Note that G' is assumed to be simply connected and quasi-split and is also the derived subgroup of \hat{G} by construction.

$$\begin{array}{c} \mu \xrightarrow{\delta} G' \\ \downarrow^{\rho} \qquad \downarrow \\ S \xrightarrow{\gamma} \hat{G} \end{array}$$

Thus, we get an exact sequence $1 \to G' \to \hat{G} \to \hat{G}/G' \to 1$ to which we can apply Lemma 5.2 to conclude that the norm principle holds for the canonical map $\hat{G} \to \frac{\hat{G}}{[\hat{G},\hat{G}]}$.

Constructing the intermediate group \tilde{G} as in Section 3.1, we see that the norm principle also holds for the natural map $\tilde{G} \rightarrow \tilde{G}/G$ [1, Prop. 5.1]. Then using Theorem 3.1 [3], Lemma 3.2, and a remark from Gopal Prasad that G^{sc} is quasi-split if and only if G is quasi-split, we can conclude that Theorem 1.3 (restated below) holds. **Theorem 1.3.** Let k be a field of characteristic not 2. Let G be a connected quasi-split reductive k-group whose Dynkin diagram does not contain connected components of type E_8 . Then Serre's question has a positive answer for G.

Acknowledgements. The author thanks Professors A. S. Merkurjev, R. Parimala, O. Wittenberg and the anonymous referee for their many valuable suggestions and critical comments.

References

- P. Barquero and A. Merkurjev, *Norm Principle for Reductive Algebraic Groups*, J. Proceedings of the International Colloquium on Algebra, Arithmetic and Geometry TIFR, Mumbai, 2000.
- [2] E. Bayer-Fluckiger and H. W. Lenstra Jr., Forms in odd degree extensions and self-dual normal bases, *Amer. J. Math.*, **112** (1990), no. 3, 359–373. Zbl 0729.12006 MR 1055648
- [3] J. Black, Zero cycles of degree one on principal homogeneous spaces, *Journal* of Algebra, **334** (2011), 232–246. Zbl 1271.11046 MR 2787661
- [4] M. Borovoi and B. Kunyavskiĭ, Formulas for the unramified Brauer group of a principal homogeneous space of a linear algebraic group, *Journal of Algebra*, 225 (2000), no. 2, 804–821. Zbl 1054.14510 MR 1741563
- [5] O. Gabber, Q. Liu and D. Lorenzini, The index of an algebraic variety, *Inventiones mathematicae*, **192** (2013), no. 3, 567–626. Zbl 1268.13009 MR 3049930
- [6] S. Garibaldi, A. Merkurjev and J.-P. Serre, *Cohomological invariants in Galois cohomology*, University Lecture Series, 28, American Mathematical Society, Providence, RI, 2003. Zbl 1159.12311 MR 1999383
- [7] P. Gille, Serre's conjecture II: a survey, in *Quadratic forms, linear algebraic groups, and cohomology*, 41–56, Springer, New York, 2010. Zbl 1239.11046 MR 2648719
- [8] D. Harari and T. Szamuely, Local-global questions for tori over *p*-adic function fields, *preprint*, 2013. arXiv:1307.4782
- M.-A. Knus, A. Merkurjev, M. Rost and J.-P. Tignol, *The book of involutions*, American Mathematical Society Colloquium Publications, 44, AMS, 1998. Zbl 0955.16001 MR 1632779
- [10] S. MacLane, Subfields and automorphism groups of *p*-adic fields, Annals of Mathematics, 40 (1939), 423–442. Zbl 0021.00501 MR 1503470

Vol. 91 (2016)

- [11] J. S. Milne and K. Y. Shih, Conjugates of Shimura varieties, in *Hodge cycles*, *motives*, *and Shimura varieties*, 280–356, Springer, Berlin Heidelberg, 1981.
- [12] Y. Nisnevich, Rationally trivial principal homogeneous spaces and arithmetic of reductive group schemes over Dedekind rings, C. R. Acad. Sci. Paris, Série I, 299 (1984), no. 1, 5–8. Zbl 0587.14033 MR 0756297
- [13] J.-P. Serre, Cohomologie galoisienne: progrès et problèmes, Séminaire Bourbaki (1993/94), Astérisque, 227 (1995), Exp. No. 783, 4, 229–257. Zbl 0837.12003 MR 1321649

Received November 1, 2015

N. Bhaskhar, Department of Mathematics and Computer Science, Emory University, Atlanta, GA 30322, USA

E-mail: nbhaskh@emory.edu