

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 94 (2019)
Heft: 1

Artikel: Generalized triangle groups, expanders, and a problem of Agol and Wise
Autor: Lubotzky, Alexander / Fox Manning, Jason / Wilton, Henry
DOI: <https://doi.org/10.5169/seals-823091>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Generalized triangle groups, expanders, and a problem of Agol and Wise

Alexander Lubotzky, Jason Fox Manning and Henry Wilton

Abstract. Answering a question asked by Agol and Wise, we show that a desired stronger form of Wise’s malnormal special quotient theorem does not hold. The counterexamples are generalizations of triangle groups, built using the Ramanujan graphs constructed by Lubotzky–Phillips–Sarnak.

Mathematics Subject Classification (2010). 20F67; 05C75, 20E26, 20F65.

Keywords. Expanders, virtually special groups, combinatorial Dehn filling.

1. Introduction

Consider the following notorious question in geometric group theory (see for example [9, 5.3.B], [5, Question 1.15]).

Question 1.1. *Is every hyperbolic group residually finite?*

“Dehn filling” is a powerful technique for constructing hyperbolic groups. A *group pair* (G, \mathcal{P}) is a group G together with a finite collection \mathcal{P} of subgroups of G . The subgroups \mathcal{P} are referred to as *peripheral groups* of the pair. A *Dehn filling* of a group pair is a quotient

$$G(N_1, \dots, N_n) := G / \langle\langle \cup_i N_i \rangle\rangle$$

where, for each i , the subgroup N_i is normal in P_i . If each N_i is finite index in P_i , the filling is said to be *peripherally finite* or *PF*. We say that a property Π holds for *all sufficiently long* Dehn fillings of (G, \mathcal{P}) if there is a finite subset $B \subseteq G \setminus 1$ so that, whenever $N_i \cap B = \emptyset$ for all i , the corresponding Dehn filling $G(N_1, \dots, N_n)$ has Π . If all sufficiently long Dehn fillings either satisfy Π or are not PF, we say Π holds for *all sufficiently long PF Dehn fillings*. The archetypal Dehn filling theorem is a far-reaching generalization of Thurston’s famous Hyperbolic Dehn Filling theorem to the group-theoretic context. In the context of PF fillings of hyperbolic groups, it has the following consequence (see Definition 2.6 for the definition of almost malnormal).

Theorem 1.2 ([15], cf. [8]). *Let G be hyperbolic, and let $\mathcal{P} = \{P_1, \dots, P_n\}$ be an almost malnormal collection of quasiconvex subgroups. All sufficiently long PF Dehn fillings*

$$G(N_1, \dots, N_n) := G / \langle\langle \cup_i N_i \rangle\rangle$$

are infinite and hyperbolic.

Even if one starts with a residually finite hyperbolic group (even a free group) G , there is no reason to believe that the resulting Dehn fillings $G(N_1, \dots, N_n)$ should be residually finite. (Compare [11, Theorem 8.1], in which it is shown that an infinite hyperbolic proper quotient G of a linear group Γ need not be linear; this applies, for instance, if G is a Dehn filling of Γ .) Theorem 1.2 therefore seems like a promising candidate for constructing non-residually-finite hyperbolic groups.

These considerations made the work of Wise and his coauthors on *virtually special groups* all the more surprising. A nonpositively curved cube complex X is *special* if there is a locally isometric immersion to the Salvetti complex associated to some right-angled Artin group. A group is *special* if it is the fundamental group of a compact special cube complex. A group is *virtually special* if it has a special subgroup of finite index. Virtually special groups have numerous attractive properties. For example, they are virtually subgroups of right-angled Artin groups, which are linear. It follows that virtually special groups are linear, and therefore residually finite. In addition, an infinite virtually special group has a subgroup of finite index with infinite abelianization.

One of the most important theorems about virtually special groups is Wise's Malnormal Special Quotient Theorem, which can be thought of as a Dehn filling result. In order to state it, we need one additional piece of terminology about Dehn fillings of a group pair (G, \mathcal{P}) .

We say that a property Π holds for *a positive fraction of all* Dehn fillings if, for each i , there is a subgroup $\dot{P}_i < P_i$ of finite index so that, whenever $N_i < \dot{P}_i$ for all i , the corresponding Dehn filling $G(N_1, \dots, N_n)$ has Π . The Malnormal Special Quotient theorem can now be stated as follows [18] (cf. [1]).

Theorem 1.3 (Wise's Malnormal Special Quotient Theorem). *Let G be hyperbolic and virtually special, and let $\mathcal{P} = \{P_1, \dots, P_n\}$ be an almost malnormal collection of quasiconvex subgroups. A positive fraction of all PF Dehn fillings*

$$G(N_1, \dots, N_n) := G / \langle\langle \cup_i N_i \rangle\rangle$$

are hyperbolic and virtually special.

Thus, remarkably, in the context of virtually special groups, Dehn fillings can be performed that preserve residual finiteness. This was one of the most important ingredients in Agol's celebrated proof of the Virtual Haken conjecture [2].

Nevertheless, the Malnormal Special Quotient Theorem does not completely rule out the possibility of constructing a non-residually finite hyperbolic group using Dehn

filling, since it only applies to a *positive fraction* of all possible Dehn fillings. As a result of Theorem 1.2, all sufficiently long PF Dehn fillings of a virtually special group are infinite and hyperbolic, but only a positive fraction of them are guaranteed to be virtually special (and hence residually finite). One is therefore led to wonder whether the Malnormal Special Quotient Theorem can be given such a form. This led Ian Agol [3, Problem 14] and Daniel Wise [19, Problem 13.16] to ask the following question in their 2014 ICM talks.

Question 1.4. *Let G be hyperbolic and virtually special, and let $\mathcal{P} = \{P_1, \dots, P_n\}$ be an almost malnormal collection of quasiconvex subgroups. Are all sufficiently long PF Dehn fillings*

$$G(N_1, \dots, N_n) := G / \langle\langle \cup_i N_i \rangle\rangle$$

virtually special?

The purpose of the current note is to show that this question has a *negative* answer in some simple situations, meaning that the Malnormal Special Quotient Theorem is in some sense as strong as it can be.

Our examples will be *k-fold triangle groups* (discussed at length in Sections 2 and 3). We briefly give the definition now. Let $k \geq 2$, and let G be a free product of three copies of \mathbb{Z}/k .¹ The collection $\mathcal{P} = \{P_{12}, P_{13}, P_{23}\}$ consists of three two-fold free products of copies of \mathbb{Z}/k obtained each by omitting one of the copies. Fix a surjection $G \rightarrow \mathbb{Z}/k$ taking each free factor isomorphically to the target, and let G_0 be the kernel. The collection $\mathcal{P}_0 = \{P_{12,0}, P_{13,0}, P_{23,0}\}$ is the collection of intersections of elements of \mathcal{P} with G_0 . Each $P_{ij,0}$ can be identified with a free group F_{k-1} on $k-1$ generators and with a subgroup of index k in $\mathbb{Z}/k * \mathbb{Z}/k$.

In Section 2, we define, for normal subgroups of finite index in $P_{ij,0}$ notions of *rotundness* (large girth for some associated graph), and *expansiveness* (good expansion for the associated graph). See Definition 2.10 for the precise definitions. For the group G_0 just described, we prove:

Theorem 1.5. *If, for each $i \in \{1, 2, 3\}$, the subgroup $N_{ij,0} \triangleleft P_{ij,0} \cong F_{k-1}$ is rotund and expansive, then $G_0(N_{12,0}, N_{13,0}, N_{23,0})$ is hyperbolic and has property (T).*

In fact (see Theorem 2.11) rotundness alone suffices for hyperbolicity; that some lower bound on girth suffices can also be seen from Theorem 1.2.

In Section 4, we use the Ramanujan graphs constructed in [12] to show the following proposition (note that the group $P_{ij,0}$ is free of rank $k-1$).

Proposition 1.6. *There exists $k \geq 18$ and, for each $1 \leq i < j \leq 3$, a sequence $\{K_{ij,n}\}_{n \in \mathbb{N}}$ of normal, rotund, expansive subgroups of $P_{ij,0}$ so that*

$$\bigcap_{n \in \mathbb{N}} K_{ij,n} = \{1\}.$$

¹By \mathbb{Z}/k we mean the cyclic group of order k .

Note that each group $P_{ij,0}$ in the statement of the Proposition is free of rank $k - 1$, and that the resulting graphs are k -valent. The possible k include $p + 1$ for any prime $p \geq 17$ so that $p \equiv 1 \pmod{4}$.

Corollary 1.7. *The answer to Question 1.4 is “no”.*

Proof. Fix a k as in Proposition 1.6. The pair (G_0, \mathcal{P}_0) we have just described satisfies:

- (1) G_0 is free, hence hyperbolic.
- (2) The elements of \mathcal{P}_0 are quasiconvex.
- (3) \mathcal{P}_0 is a malnormal collection.

Suppose the answer were “yes”. Then for some j , the quotient $G_0(K_{12,j}, K_{13,j}, K_{23,j})$ is an infinite virtually special group; in particular it has a finite index subgroup with infinite abelianization [4]. This contradicts property (T). \square

It is interesting to point out that the solution to this group theoretic problem relies essentially on number theory, via the construction in [12].

Question 1.8. *Are the examples from Corollary 1.7 virtually torsion-free? Residually finite? Linear?*

1.1. Conventions. We use the notation $A \triangleleft B$ to indicate that A is finite index in B , and $A \triangleleft\!\!\triangleleft B$ to indicate that A is a finite index normal subgroup of B . If G is a group and $S \subset G$, we use the notation $\langle\langle S \rangle\rangle$ to denote the normal closure of S in G .

Acknowledgements. The first author was supported by ERC, NSF, and BSF grants. The second and third authors are grateful to the Mathematical Sciences Research Institute in Berkeley, California, where this project was started during a special semester on Geometric Group Theory in Fall 2016, funded under NSF grant DMS-1440140. The second author is partially funded by NSF grant DMS-1462263. The third author is partially funded by EPSRC Standard Grant number EP/L026481/1. This paper was completed while the third author was participating in the *Non-positive curvature, group actions and cohomology* programme at the Isaac Newton Institute, funded by EPSRC Grant number EP/K032208/1.

2. k -fold triangle groups

In this section, we describe a generalization of the classical triangle groups which we will use to prove the main result of the paper (Corollary 1.7). To motivate, let us recall first the classical (hyperbolic) triangle groups.

Let $l \geq m \geq r \geq 2$ be integers, so that $\frac{1}{l} + \frac{1}{m} + \frac{1}{r} < 1$. Then there exists an essentially unique hyperbolic triangle with angles $\frac{\pi}{l}$, $\frac{\pi}{m}$, and $\frac{\pi}{r}$. The group generated

by reflections in the sides of this triangle is a cocompact lattice in $SO(2, 1)$, with group presentation given by the Poincaré polyhedron theorem:

$$\Delta(l, m, r) = \langle x_1, x_2, x_3 \mid x_1^2 = x_2^2 = x_3^2 = (x_1x_2)^l = (x_2x_3)^m = (x_1x_3)^r = 1 \rangle \quad (2.1)$$

The group of orientation-preserving elements in $\Delta(l, m, r)$ has index 2, and the following presentation:

$$\Delta_0(l, m, r) = \langle a_1, a_2, a_3 \mid a_1^l = a_2^m = a_3^r = a_1a_2a_3 = 1 \rangle \quad (2.2)$$

Both $\Delta(l, m, r)$ and $\Delta_0(l, m, r)$ are often called triangle groups. Sometimes Δ_0 is called an *ordinary* triangle group or a *von Dyck* group.

Let us propose a generalization of the triangle groups, generated by elements of order k instead of involutions. We first fix parent groups G and G_0 , which generalize the orbifold fundamental groups of a mirrored ideal triangle and a pair of pants, respectively. We also specify some peripheral subgroups.

Definition 2.1 (The parent groups). Fix $k \geq 2$. Let G be the free product of three copies of \mathbb{Z}/k ,

$$G = \langle x_1, x_2, x_3 \mid x_1^k = x_2^k = x_3^k \rangle.$$

For $i < j$, let $P_{ij} = \langle x_i, x_j \rangle < G$. Let G_0 be the kernel of the map $G \rightarrow \mathbb{Z}/k$ taking x_i to $\bar{1}$ for each i . For $i < j$ let $P_{ij,0} = P_{ij} \cap G_0$. Then G_0 is free of rank $2k - 2$, and each $P_{ij,0}$ is a free factor of rank $k - 1$.

Definition 2.2 (Ordinary triangle groups). Let $L \triangleleft P_{12,0}$, $M \triangleleft P_{13,0}$, and $R \triangleleft P_{23,0}$. Let K_0 be the normal closure of $L \cup M \cup R$ in G_0 , and define the (*ordinary*) k -fold triangle group:

$$G_0(L, M, R) = G_0/K_0.$$

We will often omit the word “ordinary”. Notice that the 2-fold triangle groups are the classical triangle groups, where we take $L = \langle (x_1x_2)^l \rangle$, $M = \langle (x_1x_3)^m \rangle$, and $R = \langle (x_2x_3)^r \rangle$.

Likewise, for L, M, R normal subgroups in the $P_{ij} < G$, we can define analogues of the triangle groups $\Delta(l, m, r)$.

Definition 2.3 (Extended triangle groups). Suppose that $L \triangleleft P_{12}$, $M \triangleleft P_{13}$, and $R \triangleleft P_{23}$. Let K be the normal closure of $L \cup M \cup R$ in G . Then we define the *extended* k -fold triangle group:

$$G(L, M, R) = G/K.$$

We next note that, if the subgroups L, M, R are normal subgroups of both the P_{ij} and the $P_{ij,0}$, then the ordinary triangle groups and extended triangle groups are related as one expects.

Lemma 2.4. *Suppose that $L \triangleleft P_{12}$ and $L < P_{12,0}$, that $M \triangleleft P_{13}$ and $M < P_{13,0}$, and that $R \triangleleft P_{23}$ and $R < P_{23,0}$. Then the normal closure of $L \cup M \cup R$ in G_0 is equal to K .*

Proof. Clearly $K_0 = \langle\langle L \cup M \cup R \rangle\rangle_{G_0} < K = \langle\langle L \cup M \cup R \rangle\rangle_G$. But we can write $K_0 = K_{L,0}K_{M,0}K_{R,0}$, and $K = K_LK_MK_R$, where $K_{L,0} = \langle\langle L \rangle\rangle_{G_0}$, $K_L = \langle\langle L \rangle\rangle_G$, and so on. Thus it suffices to show, for example, that $K_{L,0} = K_L$, that the normal closures of L in G and G_0 coincide. Since $\langle x_1 \rangle$ maps onto G/G_0 , it is enough to show that $x_1^{-1}K_{L,0}x_1 = K_{L,0}$. Consider a generator $z = x^{-1}yx$ of $K_{L,0}$, where $y \in L$ and $x \in G_0$. Then $x_1^{-1}zx_1 \in G_0$, since $G_0 \triangleleft G$ and $x_1^{-1}yx_1 \in L$, since $L \triangleleft P_{12} = \langle x_1, x_2 \rangle < G$. Thus $x_1^{-1}zx_1 \in K_{L,0}$. Since z was arbitrary, we have shown that $K_{L,0}$ is normal in G , and so equal to K_L , as desired. \square

Corollary 2.5. *If the ordinary and extended triangle groups are both defined, then $G_0(L, M, R)$ is a subgroup of index k in $G(L, M, R)$.*

2.1. Malnormality.

Definition 2.6. Let H be a group, and \mathcal{Q} a collection of subgroups of H . Then \mathcal{Q} is *malnormal* if whenever $h \in H$, $Q, Q' \in \mathcal{Q}$, and $Q \cap hQ'g^{-1}$ is nontrivial, then $Q = Q'$ and $h \in Q$.

The collection is *almost malnormal* if whenever $h \in H$, $Q, Q' \in \mathcal{Q}$, and $Q \cap hQ'h^{-1}$ is infinite, then $Q = Q'$ and $h \in Q$.

We make the following observation, whose (easy) proof is left to the reader.

Lemma 2.7. *With the notation of Definition 2.1, the collection $\mathcal{P} = \{P_{12}, P_{13}, P_{23}\}$ is almost malnormal in G , and $\mathcal{P}_0 = \{P_{12,0}, P_{13,0}, P_{23,0}\}$ is malnormal in G_0 .*

Both G and G_0 are virtually free, and thus virtually special locally quasiconvex. In particular, the pairs (G, \mathcal{P}) and (G_0, \mathcal{P}_0) both satisfy the hypotheses of Theorem 1.3 and Question 1.4.

2.2. Geometric conditions on graphs and triangle groups.

Definition 2.8. Let Γ be a graph. The *girth* of Γ is the length of the shortest circuit in Γ .

Definition 2.9. If Γ is a connected k -regular graph, we define the *Laplacian* in terms of the adjacency matrix A :

$$\delta = I - \frac{1}{k}A.$$

With this normalization, the spectrum of δ always contains 0 and lies in the interval $[0, 2]$. We define $\lambda_1(\Gamma)$ to be the smallest positive eigenvalue of δ .

For each pair $i < j$, the group P_{ij} acts on the regular k -valent tree T_k with quotient equal to a single edge. For definiteness we fix a planar embedding of this tree, and an oriented edge e_0 . Let x_i act on this tree by rotating around $i(e_0)$ and let x_j act by rotating around $t(e_0)$. In this way we make T_k into a Bass–Serre tree for the P_{ij} , considered as a free product. Any finite index subgroup N of P_{ij} acts on T_k with a finite quotient graph $N \backslash T_k$.

Definition 2.10. Let $N \triangleleft P_{ij,0} < P_{ij} \cong (\mathbb{Z}/k) * (\mathbb{Z}/k)$. We say that N is *rotund* if $\text{girth}(N \backslash T_k) > 6$. We say that N is *expansive* if $\lambda_1(N \backslash T_k) > \frac{1}{2}$.

These characterizations of subgroups as rotund or expansive depend on the particular action of $P_{ij,0}$ on T_k given. Here is a more precise version of Theorem 1.5.

Theorem 2.11. Let $G_0, P_{ij,0}, L, M, R$ be as in Definition 2.2.

- (1) If L, M, R are rotund, then $G_0(L, M, R)$ is hyperbolic.
- (2) If L, M, R are rotund and expansive, then $G_0(L, M, R)$ has property (T).

Theorem 2.11 will be proved in Section 3. In Section 4 we will produce many examples of rotund expansive L, M, R , proving Proposition 1.6.

3. Triangular complexes of groups

In this section, we give the geometric framework necessary to understand why the groups discussed in the last section answer Question 1.4. In particular, we will prove Theorem 2.11.

The virtually free group G can usefully be thought of as a complex of groups in two ways, both shown in Figure 1.




Figure 1. G as a graph of finite cyclic groups, and as a triangle of groups.

On the left, we see G as the fundamental group of a graph (a tripod) of finite groups \mathcal{G} . On the right, we see G as a triangle of groups \mathcal{D} , with cyclic edge groups, and vertex groups equal to the peripheral groups. Both complexes of groups are

developable in the sense of [6, III.ℳ]. This means in the second case that there is an action of G on a simply connected complex (the *development*) with quotient \mathcal{D} , and the complex-of-groups data can be recovered from the action. Likewise \mathcal{G} is the quotient of a Bass–Serre tree T by the natural action of G . Here is a way to recover the development in this case: Each $P_{ij} \in \mathcal{P}$ has a minimal invariant subtree $T_{P_{ij}}$ in $\tilde{\mathcal{D}}$. The development $\tilde{\mathcal{D}}$ of \mathcal{D} , is homeomorphic to a complex which is obtained from T by coning off each translate of any $T_{P_{ij}}$. The link of a vertex in $\tilde{\mathcal{D}}$ can be identified with the Bass–Serre tree of $\mathbb{Z}/k * \mathbb{Z}/k$ (see Lemma 4.2 below).

Likewise, the free group G_0 is the fundamental group of a graph \mathcal{G}_0 and a complex of groups \mathcal{D}_0 , both shown in Figure 2. Here the vertex groups of \mathcal{D}_0 are the elements of \mathcal{P}_0 . The development of the complex of groups is also $\tilde{\mathcal{D}}$. The link of a vertex of \mathcal{D}_0 is a graph with two vertices joined by k edges.




Figure 2. G_0 is the fundamental group of the graph on the left, and also the fundamental group of the complex of groups with underlying complex Y on the right. The case $k = 3$ is shown.

Now fix a k -fold triangle group $G_0(L, M, R)$ as in Definition 2.2. We obtain a complex of groups structure $\mathcal{D}_0(L, M, R)$ for $G_0(L, M, R)$ in terms of the one for G_0 , by replacing the vertex groups (elements of \mathcal{P}_0) with their finite quotients $P_{12,0}/L$, $P_{13,0}/M$, and $P_{23,0}/R$.

Bridson and Haefliger give a criterion which implies that a given complex of groups is developable.

Theorem 3.1 ([6, Theorem III.ℳ.4.17]). *If a complex of groups is non-positively curved it is developable. Moreover, if the local developments are $CAT(-1)$ then the development is $CAT(-1)$.*

To say that a complex of groups is non-positively curved is precisely to say that the *local developments* are non-positively curved. This condition depends on how we metrize the cells of the complex. In our case, we can metrize the triangles as hyperbolic triangles with some constant angle θ . The local development at a vertex marked by a group $P_{ij,0}/N$ where $N \triangleleft P_{ij,0} \triangleleft \mathbb{Z}/k * \mathbb{Z}/k$ is the hyperbolic cone on the graph $N \setminus T_k$, where this graph has been metrized so each edge has length θ .

If $\theta \cdot \text{girth}(N \setminus T_k) \geq 2\pi$, this local development is locally $\text{CAT}(-1)$. In particular it will satisfy the nonpositive curvature hypothesis in Theorem 3.1.

Proposition 3.2. *Suppose that $L, M,$ and R are rotund.*

- (1) $\mathcal{D}_0(L, M, R)$ is developable, and the development is contractible;
- (2) $G_0(L, M, R)$ is hyperbolic; and
- (3) The link of any vertex of the development of $\mathcal{D}_0(L, M, R)$ is isomorphic to $N \setminus T_k$ where $N \in \{L, M, R\}$.

Proof. The local development of $\mathcal{D}_0(L, M, R)$ at a vertex is as described in item (3). Thus if $\mathcal{D}_0(L, M, R)$ is in fact developable, item (3) will follow.

Let $n \geq 7$ be the minimum girth of the graphs $N \setminus T_k$ where $N \in \{L, M, R\}$. Metrizing the triangles of $\mathcal{D}_0(L, M, R)$ by equilateral hyperbolic triangles with angle $2\pi/n$, we can verify the conditions of Theorem 3.1 as discussed above, and see that $\mathcal{D}_0(L, M, R)$ is developable, establishing the first part of item (1). The development X is moreover locally (and hence globally) $\text{CAT}(-1)$ and thus contractible. Moreover the group $G_0(L, M, R)$ acts properly cocompactly on X . Thus $G_0(L, M, R)$ is hyperbolic, establishing item (2). □

To deduce property (T) when the normal subgroups are expansive, we need the following criterion.

Theorem 3.3 ([7, Corollary 1]). *Let $\Gamma \curvearrowright Z$ properly and cocompactly, where Z is a contractible simplicial 2-complex so that for every vertex v of Z , the link Z_v of v is connected and satisfies $\lambda_1(Z_v) > \frac{1}{2}$. Then Γ has Property (T).*

Proposition 3.4. *If $L, M,$ and R are rotund and expansive, then $G_0(L, M, R)$ has property (T).*

Proof. Since $L, M,$ and R are rotund, the group $G_0(L, M, R)$ acts properly and cocompactly on the development $\mathcal{D}_0(L, M, R)$, which is a contractible complex with each link isomorphic to $N \setminus T_k$ for $N \in \{L, M, R\}$. Since L, M, R are expansive, we have $\lambda_1(\text{link}(v)) > \frac{1}{2}$ for each vertex v . We can thus apply Theorem 3.3 to conclude that $G_0(L, M, R)$ has property (T). □

Propositions 3.2 and 3.4 together imply Theorem 2.11.

4. Finding good expanders

In this section we prove Proposition 1.6, about the existence of the expanders we need. The proposition is phrased in terms of a subgroup N , normal and finite index in $P_0 \triangleleft P \cong \mathbb{Z}/k * \mathbb{Z}/k$, where P_0 is defined to be the kernel of the map $\mathbb{Z}/k * \mathbb{Z}/k \rightarrow \mathbb{Z}/k$ taking each generator to $\bar{1} \in \mathbb{Z}/k$. In applying the results of this section we identify P_0 with one of the $P_{ij,0}$ described before. For T_k equal to the

Bass–Serre tree associated to the free splitting of P , we are interested in the girth and first eigenvalue of the graphs $N \setminus T_k$. We proceed by identifying P_0 with a certain arithmetic subgroup of $\mathrm{PGL}_2(\mathbb{Q}_p)$. (We should emphasize that none of the results of the next two subsections are really new, but we want to include enough of the ideas from [12, 13] so that the reader gets the flavor of what is going on.)

4.1. The setup. If T is a tree, we let $\mathrm{Aut}^+(T) < \mathrm{Aut}(T)$ be the subgroup of index at most two consisting of those ϕ which move a point (hence every point) an even distance. Note that $\mathrm{Aut}^+(T)$ acts on T without inversions, and $\Gamma \setminus T$ is bipartite for any $\Gamma < \mathrm{Aut}^+(T)$. The first lemma is an easy corollary of the fact that T is contractible.

Lemma 4.1. *Let T be a locally finite tree and D, Δ two discrete torsion-free subgroups of $\mathrm{Aut}^+(T)$ so that the graphs $D \setminus T$ and $\Delta \setminus T$ are isomorphic. Then D and Δ are conjugate in $\mathrm{Aut}(T)$.*

The second lemma is also standard.

Lemma 4.2. *Let $k \in \mathbb{Z}^+$, and let $P = \mathbb{Z}/k * \mathbb{Z}/k$, and let T be the Bass–Serre tree for the free splitting of P . (In other words vertices are in one-to-one correspondence with left cosets of the free factors, and if A_1 is the first and A_2 the second free factor, each gA_1 is connected to gA_2 by an edge.) Let P_0 be the kernel of the map $P \rightarrow \mathbb{Z}/k$ which is the identity on each free factor.*

(1) T is a k -regular tree.

(2) $P_0 \setminus T$ consists of 2 vertices connected by k edges.

Let p be a prime and T_{p+1} be the regular $(p+1)$ -valent tree, and let \mathbb{Q}_p be the field of p -adic numbers. Then $\mathrm{PGL}_2(\mathbb{Q}_p)$ acts on the Bruhat–Tits tree T_{p+1} , as explained in [16].

Our goal now is to find an *arithmetic* subgroup Δ of

$$\mathrm{PSL}_2(\mathbb{Q}_p) < \mathrm{PGL}_2(\mathbb{Q}_p) < \mathrm{Aut}(T_{p+1}),$$

so that the quotient $\Delta \setminus T_{p+1} \cong P_0 \setminus T$, the graph from Lemma 4.2, with $k = p+1$. Lemma 4.1 implies that Δ is conjugate to P_0 in $\mathrm{Aut}(T_{p+1})$.

We will take suitable *congruence* subgroups $\Delta(i)$ of Δ ; the isomorphism taking Δ to P_0 will take these congruence subgroups to the groups L, M, R specified in Theorem 2.11. The fact that the graphs $\Delta(i) \setminus T$ are Ramanujan will come from [13, Section 7.3, Theorem 7.3.12], using the solution of Deligne to the Ramanujan–Peterson conjecture. We will see that in fact, one can choose the subgroups $\Delta(i)$ to be nested with $\bigcap_i \Delta(i) = \{1\}$, so that the girth of $\Delta(i) \setminus T$ goes to infinity.

4.2. Constructing Δ . Fix p prime, with $p \equiv 1 \pmod{4}$ and $p \geq 17$.

Recall the classical four square theorem of Jacobi [10, Theorem 386]:

Theorem 4.3 (Jacobi). *Let $n \in \mathbb{Z}^+$, and define*

$$r_4(n) = \#\{(x_0, x_1, x_2, x_3) \in \mathbb{Z}^4 \mid \sum x_j^2 = n\}.$$

Then

$$r_4(n) = 8 \sum_{d \mid n, 4 \nmid d} d.$$

In particular since p is prime, $r_4(p) = 8(p + 1)$. We are assuming $p \equiv 1 \pmod{4}$, so for any four integers whose squares sum to p , exactly three are even. Thus if we take

$$S = \{(x_0, x_1, x_2, x_3) \in \mathbb{Z}^4 \mid x_0 > 0 \text{ odd}, \sum x_j^2 = p\}, \tag{4.1}$$

then $\#S = p + 1$.

Next we claim (again using $p \equiv 1 \pmod{4}$) that there exists $\epsilon \in \mathbb{Z}_p \subseteq \mathbb{Q}_p$ so that $\epsilon^2 = -1$. Indeed it is well-known that such an ϵ exists in \mathbb{Z}/p , and by the Hensel Lemma it can be lifted to \mathbb{Z}_p . For every $\alpha \in S$, associate the matrix

$$\tilde{\alpha} = \begin{pmatrix} x_0 + \epsilon x_1 & x_2 + \epsilon x_3 \\ -x_2 + \epsilon x_3 & x_0 - \epsilon x_1 \end{pmatrix} \in M_2(\mathbb{Q}_p). \tag{4.2}$$

Note that $\det(\tilde{\alpha}) = p$; we abuse notation by also thinking of $\tilde{\alpha}$ as an element of $\text{PGL}_2(\mathbb{Q}_p)$. Let Γ be the subgroup generated by $\tilde{S} = \{\tilde{\alpha} \mid \alpha \in S\}$.

Lemma 4.4. Γ is a discrete cocompact subgroup of $\text{PGL}_2(\mathbb{Q}_p)$.

This is actually a special case of Theorem 7.3.12 of [13]. Let us explain this special case in some detail.

Let H be the Hamiltonian quaternion algebra; for a commutative ring R , we have

$$H(R) = \{a_0 + a_1i + a_2j + a_3k \mid a_i \in R\},$$

the associative R -algebra generated by symbols i, j, k , satisfying the relations $i^2 = j^2 = k^2 = -1$ and $ij = k = -ji$. Let $H^*(R)$ be the group of invertible elements in $H(R)$.

Since $\mathbb{Z}[\frac{1}{p}]$ is discrete in $\mathbb{R} \times \mathbb{Q}_p$, there are discrete embeddings

$$H(\mathbb{Z}[\frac{1}{p}]) \hookrightarrow H(\mathbb{R}) \times H(\mathbb{Q}_p)$$

and

$$H^*(\mathbb{Z}[\frac{1}{p}])/\text{center} \hookrightarrow (H^*(\mathbb{R})/\text{center}) \times (H^*(\mathbb{Q}_p)/\text{center}).$$

Now $H^*(\mathbb{R})/\text{center} = \mathbb{RP}^3$ is compact, while $H(\mathbb{Q}_p) \cong M_2(\mathbb{Q}_p)$ (i.e., H splits over \mathbb{Q}_p , hence $H^*(\mathbb{Q}_p)/\text{center} \cong PGL_2(\mathbb{Q}_p)$); the map

$$a_0 + a_1i + a_2j + a_3k \mapsto \begin{pmatrix} a_0 + \epsilon a_1 & a_2 + \epsilon a_3 \\ -a_2 + \epsilon a_3 & a_0 - \epsilon a_1 \end{pmatrix}$$

gives the explicit isomorphism). Since $H^*(\mathbb{R})/\text{center}$ is compact, the projection of $\Gamma_0 = H^*(\mathbb{Z}[\frac{1}{p}])/\text{center}$ to $H^*(\mathbb{Q}_p)/\text{center} = PGL_2(\mathbb{Q}_p)$ gives a discrete subgroup! (This despite the fact that $\mathbb{Z}[\frac{1}{p}]$ projected to \mathbb{Q}_p is dense.)

Now, our Γ is inside the projection, since every $\alpha \in S$ is invertible as an element of $H(\mathbb{Z}[\frac{1}{p}])$; indeed $\|\alpha\| = \alpha \cdot \bar{\alpha} = p$ is invertible in $\mathbb{Z}[\frac{1}{p}]$, and $\alpha^{-1} = \frac{\bar{\alpha}}{\|\alpha\|}$. (In general $\alpha \in H(R)$ is invertible if and only if $\|\alpha\|$ is invertible in the ring R .) One can easily see from (4.1) and (4.2) that $\Gamma \subseteq \Gamma_0(2)$, the mod 2 congruence subgroup of Γ_0 . This all explains why Γ is discrete. But it is also cocompact; in fact $\Gamma = \Gamma_0(2)$, and every $\tilde{\alpha} \in \tilde{S}$ takes the root of the tree (which is the equivalence class of the lattice $\mathbb{Z}_p^2 \subseteq \mathbb{Q}_p^2$; see [16] for this model of the Bruhat–Tits tree) to a sublattice of index p (since $\det(\alpha) = p$) and there are exactly $p + 1$ such sublattices — representing the $p + 1$ neighbors of the root vertex. From this one deduces that Γ acts *transitively* on the vertices of the tree T_{p+1} . In fact Γ acts simply transitively, and is therefore a free group on $\frac{p+1}{2}$ generators. (Note $\tilde{\alpha} = \tilde{\alpha}^{-1}$ where $\bar{\alpha}$ is the quaternionic conjugate, and so the image of S is a symmetric subset of $PGL_2(\mathbb{Q}_p)$.) Thus T_{p+1} can be identified with the Cayley graph $\text{Cay}(\Gamma, \tilde{S})$. In particular $\Gamma \backslash T_{p+1}$ is a bouquet of $k/2$ circles, and hence compact.

Now let $\Delta = \Gamma \cap \text{Aut}^+(T_{p+1})$; this is an index-2 subgroup of Γ which preserves the 2-coloring of the tree. Because $\Gamma = \Gamma_0(2)$ is free of rank $\frac{p+1}{2}$, the rank of Δ is, by the Nielsen–Schreier Theorem,

$$2\left(\frac{p+1}{2} - 1\right) + 1 = p = k - 1,$$

and there are two orbits of vertices. In particular, there is an isomorphism Ψ from Δ to $P_0 = \ker(\mathbb{Z}/k * \mathbb{Z}/k \rightarrow \mathbb{Z}/k)$ and an equivariant isomorphism from the tree T_{p+1} to the Bass–Serre tree of $\mathbb{Z}/k * \mathbb{Z}/k$. In particular, we can find rotund or expansive subgroups of P_0 by specifying them in Δ , which we now do.

Let $q \neq p$ be a prime or prime power, so that $(\frac{p}{q}) = -1$, i.e., p is not a quadratic residue mod q . As explained in [13, 7.3.12], in this case $\Gamma_0(2q)$ (the mod $2q$ congruence subgroup) preserves the coloring of the tree, so it lies inside Δ .

Moreover, by [13, 7.3.12], the quotients $\Gamma_0(2q) \backslash T$ have the following properties:

- (1) They are k -regular *Ramanujan* graphs, i.e., $\lambda_1(\Gamma_0(2q) \backslash T) \geq 1 - \frac{2\sqrt{k-1}}{k}$.
- (2) The girth of $\Gamma_0(2q) \backslash T$ is at least $\frac{4}{3} \log_p(q)$.

So for fixed p and $q \rightarrow \infty$ we are finished. (Take for example $p = 17$, $q = 3^l$ and $l \rightarrow \infty$ for a nested family.)

Remark 4.5. The graphs $\Gamma_0(2q)\backslash T$, for q a prime congruent to 1 mod 4, are really the same as the Ramanujan graphs presented in Lubotzky–Phillips–Sarnak [12]. But one can make use of many other examples, e.g., for $k = p^e + 1$, those constructed by Morgenstern [14].

4.3. Other constructions. Another source of examples is provided by a result communicated to us by Varjù [17].

Theorem 4.6 (Varjù). *There is an absolute constant $c > 0$ such that the following holds. Let $k \geq 3$. Let p be a sufficiently large prime so that $k \geq 3$ is a divisor of $(p + 1)/2$, and let $L = PSL_2(p)$. Then there are subgroups $H_1, H_2 \cong \mathbb{Z}/k$ of L so that the following inequalities hold. Let K be the kernel of a homomorphism from $P_{ij} \cong \mathbb{Z}/k * \mathbb{Z}/k$ to L that takes the first factor to H_1 and the second factor to H_2 , and let $\Gamma = K\backslash T_k$.*

- (1) $\text{girth } \Gamma \geq c \frac{\log |\Gamma|}{\log(k-1)}$, and
- (2) $\lambda_1(\Gamma) > 1 - (k - 1)^{-c}$.

By taking k large enough that $1 - (k - 1)^{-c} > \frac{1}{2}$, and letting p tend to infinity, we obtain examples which are *extended k -fold triangle groups*. Applying Theorem 2.11, these give many more examples to show the answer to Question 1.4 is “no”. In Varjù’s theorem, the subgroups H_1 and H_2 are obtained as random unipotent conjugates of a fixed cyclic subgroup.

Remark 4.7. Another construction of negatively curved triangle complexes with prescribed links is provided by [7, Theorem 2]. It is possible that these can also be thought of as Dehn fillings of virtually free groups, in which case these would provide another route to answering Question 1.4.

References

- [1] I. Agol, D. Groves, and J. F. Manning, An alternate proof of Wise’s malnormal special quotient theorem, *Forum Math. Pi*, **4** (2016), e1, 54pp. Zbl 1380.20047 MR 3456181
- [2] I. Agol, The virtual Haken conjecture. With an appendix by Agol, Daniel Groves, and Jason Manning, *Doc. Math.*, **18** (2013), 1045–1087. Zbl 1286.57019 MR 3104553
- [3] I. Agol, Virtual properties of 3-manifolds, *Proceedings of the International Congress of Mathematicians (Seoul, 2014) Vol. 1*, 141–170, Kyung Moon Sa, Seoul, 2014. Zbl 1379.57001 MR 3728467
- [4] Y. Antolín and A. Minasyan, Tits alternatives for graph products, *J. Reine Angew. Math.*, **704** (2015), 55–83. Zbl 1368.20046 MR 3365774
- [5] M. Bestvina, Questions in geometric group theory. Available at: <http://www.math.utah.edu/~bestvina/eprints/questions-updated.pdf>
- [6] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 319, Springer-Verlag, Berlin, 1999. Zbl 0988.53001 MR 1744486

- [7] W. Ballmann and J. Świątkowski, On L^2 -cohomology and property (T) for automorphism groups of polyhedral cell complexes, *Geom. Funct. Anal.*, **7** (1997), no. 4, 615–645. Zbl 0897.22007 MR 1465598
- [8] D. Groves and J. F. Manning, Dehn filling in relatively hyperbolic groups, *Israel J. Math.*, **168** (2008), 317–429. Zbl 1211.20038 MR 2448064
- [9] M. Gromov, Word hyperbolic groups, in *Essays in Group Theory*, S. M. Gersten (ed.), 75–264, Mathematical Sciences Research Institute Publications, 8, Springer-Verlag, New York, 1987. Zbl 0634.20015 MR 919826
- [10] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*. Fifth edition, The Clarendon Press, Oxford University Press, New York, 1979. Zbl 0423.10001 MR 568909
- [11] M. Kapovich, Representations of polygons of finite groups, *Geom. Topol.*, **9** (2005), 1915–1951 (electronic). Zbl 1163.20029 MR 2175160
- [12] A. Lubotzky, R. Phillips, and P. Sarnak, Ramanujan graphs, *Combinatorica*, **8** (1988), no. 3, 261–277. Zbl 0661.05035 MR 963118
- [13] A. Lubotzky, *Discrete groups, expanding graphs and invariant measures*. With an appendix by Jonathan D. Rogawski, Progress in Mathematics, 125, Birkhäuser Verlag, Basel, 1994. Zbl 0826.22012 MR 1308046
- [14] M. Morgenstern, Existence and explicit constructions of $q + 1$ regular Ramanujan graphs for every prime power q , *J. Combin. Theory Ser. B*, **62** (1994), no. 1, 44–62. Zbl 0814.68098 MR 1290630
- [15] D. V. Osin, Peripheral fillings of relatively hyperbolic groups, *Invent. Math.*, **167** (2007), no. 2, 295–326. Zbl 1116.20031 MR 2270456
- [16] J.-P. Serre, *Trees*. Translated from the French by John Stillwell, Springer-Verlag, Berlin-New York, 1980. Zbl 0548.20018 MR 607504
- [17] P. P. Varjú, Expansion of coset graphs of $\mathrm{PSL}_2(F_p)$, to appear in *Israel J. Math.* arXiv:1810.09925
- [18] D. T. Wise, The structure of groups with a quasiconvex hierarchy, to appear in *Annals of Mathematics Studies*, Princeton University Press, Princeton, NJ.
- [19] D. T. Wise, The cubical route to understanding groups, *Proceedings of the International Congress of Mathematicians (Seoul, 2014) Vol. 2*, 1075–1099, Kyung Moon Sa, Seoul, 2014. Zbl 1373.20053 MR 3728653

Received February 27, 2017

A. Lubotzky, Institute of Mathematics, Hebrew University,
Jerusalem 91904, Israel

E-mail: alex.lubotzky@mail.huji.ac.il

J. F. Manning, Department of Mathematics, Cornell University,
310 Malott Hall, Ithaca, NY 14853, USA

E-mail: jfmanning@math.cornell.edu

H. Wilton, Centre for Mathematical Sciences, University of Cambridge,
Wilberforce Road, Cambridge CB3 0WB, UK

E-mail: h.wilton@maths.cam.ac.uk