

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 94 (2019)
Heft: 2

Artikel: The maximum number of strokes for genus two Riemann surfaces with abelian differentials
Autor: Judge, Chris / Parlier, Hugo
DOI: <https://doi.org/10.5169/seals-846785>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The maximum number of systoles for genus two Riemann surfaces with abelian differentials

Chris Judge* and Hugo Parlier**

Abstract. In this article, we provide bounds on systoles associated to a holomorphic 1-form ω on a Riemann surface X . In particular, we show that if X has genus two, then, up to homotopy, there are at most 10 systolic loops on (X, ω) and, moreover, that this bound is realized by a unique translation surface up to homothety. For general genus g and a holomorphic 1-form ω with one zero, we provide the optimal upper bound, $6g - 3$, on the number of homotopy classes of systoles. If, in addition, X is hyperelliptic, then we prove that the optimal upper bound is $6g - 5$.

Mathematics Subject Classification (2010). 32G15; 30F10, 53C22.

Keywords. Systoles, translation surfaces, abelian differentials.

1. Introduction

The *systolic length* of a length space (X, d) is the infimum of the lengths of non-contractible loops in X . If a non-contractible loop γ achieves this infimum, then we will call γ a *systole*. The systolic length and systoles have received a great deal of attention beginning with work of Loewner who is credited [12] with proving that among unit area Riemannian surfaces of genus one, the unit area hexagonal torus has the largest systolic length, $\sqrt{2/\sqrt{3}}$, and is the unique such surface that achieves this value.

The hexagonal torus has another extremal property: Among all Riemannian surfaces of genus one, it has the maximum number of distinct homotopy classes of systoles, three. With respect to this property, the hexagonal torus is not the unique extremal among all genus one Riemannian surfaces, but it is the unique — up to homothety — extremal among quotients of \mathbb{C} by lattices Λ equipped with the metric $|dz|^2$.

The form dz on \mathbb{C}/Λ is an example of a holomorphic 1-form on a Riemann surface. More generally, given a holomorphic 1-form ω on a Riemann surface X ,

*Research partially supported by a Simons collaboration grant.

**Research partially supported by Swiss National Science Foundation grant number PP00P2_153024.

one integrates $|\omega|$ over arcs to obtain a length metric d_ω on X . On the complement of the zero set of ω the metric is locally Euclidean, and each zero of order n is a conical singularity with angle $2\pi \cdot (n + 1)$.

The length space (X, d_ω) determined by (X, ω) is a basic object of study in the burgeoning field of Teichmüller dynamics. See, for example, the recent surveys of [5] and [16].

In this paper we prove the following.

Theorem 1.1. *Let ω be a holomorphic 1-form on a closed Riemann surface X of genus two. The number of distinct homotopy classes of systolic minimizers on (X, d_ω) is at most 10. Moreover, up to homothety, there is a unique metric space of the form (X, d_ω) for which there exist exactly 10 distinct homotopy classes of systoles.*

In other words, among the unit area surfaces (X, d_ω) of genus two, there exists a unique surface $(X_{10}, d_{\omega_{10}})$ that achieves the maximum number of systolic homotopy classes. The surface obtained by multiplying the unit area metric $d_{\omega_{10}}$ by $\sqrt{4\sqrt{3}}$ is described in Figure 1. The surface $(X_{10}, d_{\omega_{10}})$ has two conical singularities each of angle 4π corresponding to the vertices of the polygon pictured in Figure 1. In other words, the 1-form ω_{10} has simple zeros corresponding to these vertices. Four of the ten systolic homotopy classes consist of geodesics that lie in an embedded Euclidean cylinders. Each of the other six systolic homotopy classes has a unique geodesic representative that necessarily passes through one of the two zeros of ω_{10} . It is interesting to note that some of the latter systoles intersect twice. Both intersections necessarily occur at zeros of ω_{10} . Indeed, if two curves intersect twice and one of the intersection points is a smooth point of the Riemannian metric, then a standard perturbation argument produces a curve of shorter length.




Figure 1. A pair (X, ω) that has ten systoles: By identifying parallel sides with the same letters, we obtain a Riemann surface X . The one form dz in the plane defines a holomorphic 1-form on X .

Perhaps surprisingly, $(X_{10}, d_{\omega_{10}})$ does not maximize the systolic length among all unit area, genus two surfaces of the form (X, d_{ω}) . To discuss this, it will be convenient to introduce the *systolic ratio*: the square of the systolic length divided by the area of the surface. A surface maximizes the systolic length among unit area surfaces if and only if it maximizes systolic ratio among all surfaces.

A genus two surface (X, d_{ω}) that has ten systoles has systolic ratio equal to $1/\sqrt{3} = .57735\dots$. On the other hand, the surface described in Figure 2 has systolic ratio equal to

$$\frac{2 \cdot (\sqrt{13} - 3)^2}{\sqrt{3} \cdot (1 - \frac{3}{4}(\sqrt{13} - 3)^2)} = .58404\dots \tag{1.1}$$

We believe that this surface has maximal systolic ratio.

Conjecture 1.2. *The supremum of the systolic ratio over surfaces (X, d_{ω}) of genus two equals the constant in (1.1). Moreover, up to homothety, the surface described in Figure 2 is the unique surface that achieves this systolic ratio.*




Figure 2. A surface (X, d_{ω}) whose systolic ratio equals the constant in (1.1). The surface is obtained from gluing parallel sides of two isometric cyclic hexagons in \mathbb{C} . Each hexagon has a rotational symmetry of order 3. The 1-form ω corresponds to dz in the plane.

By the Riemann–Roch theorem, the total number of zeros, including multiplicities, of a holomorphic 1-form on a Riemann surface of genus g equals $2g - 2$. In particular, a 1-form ω on a genus two Riemann surface X consists of either two simple zeros or one double zero. Thus, we have a partition of the moduli space

of pairs (X, ω) into the stratum, $\mathcal{H}(1, 1)$, of those for which d_ω has two conical singularities of angle 4π and the complementary stratum, $\mathcal{H}(2)$, those for which d_ω has a single conical singularity of angle 6π .

In order to prove Theorem 1.1, we study each stratum separately. It turns out that the stratum $\mathcal{H}(2)$ is considerably easier to analyse. Indeed, for $\mathcal{H}(2)$ we are able to prove sharp bounds on both the systolic ratio and on the number of systolic homotopy classes. This is due to the fact that if there is only one zero, then each homotopy class of systoles may be represented by a single saddle connection.

Theorem 1.3. *If $(X, \omega) \in \mathcal{H}(2)$, then (X, d_ω) has at most 7 homotopy classes of systoles, and the systolic ratio of (X, d_ω) is at most $2/(3\sqrt{3}) = .3849\dots$ Furthermore, either inequality is an equality if and only if (X, d_ω) is tiled by an equilateral triangle.¹*

The unique surface that attains both optimal bounds is illustrated in Figure 3.




Figure 3. The genus two surface (X, d_ω) that achieves the optimal bounds in $\mathcal{H}(2)$.

To prove the optimal systolic bounds for a holomorphic 1-form with one zero, we adapt the argument that Fejes Tóth used to prove that a hexagonal lattice gives the optimal disc packing of the plane [4]. This method of proof extends to higher genus surfaces equipped with holomorphic one forms that have exactly one zero.

Theorem 1.4. *If $(X, \omega) \in \mathcal{H}(2g - 2)$, then the systolic ratio of (X, d_ω) is at most $\frac{4}{(4g-2)\sqrt{3}}$. Equality is achieved if and only if the surface is tiled by equilateral triangles whose vertices lie at the zero of ω .*

Theorem 1.4 has been independently observed by Boissy and Geninska [1].

As indicated above, when ω has only one zero, each systole is homotopic to a saddle connection of the same length. Smillie and Weiss [14] provided an upper bound on the length ℓ_0 of the shortest saddle connection for surfaces (X, d_ω) of genus g and area 1. In particular, they showed that $\ell_0 \leq \sqrt{1/\pi \cdot (2g - 2 + n)}$ where n is the number of zeros of ω .

We also identify optimal bounds for the number of homotopy classes of systoles of surfaces in $\mathcal{H}(2g - 2)$, and show that the optimal bounds are not attained by hyperelliptic surfaces in these strata. A condensed version of these results is the following (see Proposition 3.1 and Theorem 3.3):

¹A surface (X, d_ω) is tiled by an equilateral triangle T if there exists a triangulation of X such that each triangle is isometric to T and each vertex is a zero of ω .

Theorem 1.5. *If ω is a holomorphic 1-form on X that has exactly one zero, then (X, d_ω) has at most $6g - 3$ homotopy classes of systoles. If in addition ω is hyperelliptic, then (X, d_ω) has at most $6g - 5$ homotopy classes of systoles. Both bounds are realized.*

The bulk of the present paper verifies Theorem 1.1 for the stratum $\mathcal{H}(1, 1)$. The proof begins in §4 where we show that each nonseparating systole is homotopic to a systole that passes through exactly two Weierstrass points. Such a systole is divided by the Weierstrass points into two geodesic arcs that are interchanged by the hyperelliptic involution τ . We regard each such “systolic Weierstrass arc” as an arc on $X/\langle\tau\rangle$ that joins the two corresponding angle π cone points. If a Weierstrass arc misses the angle 4π cone point on $X/\langle\tau\rangle$ that corresponds to the zeros of ω , then we will call it “direct.” Otherwise, the arc will be called “indirect.” In §5 we show that for each angle π cone point c there are at most two direct systolic Weierstrass arcs that have an endpoint at c , and hence there are at most six homotopy classes of systoles that correspond to direct Weierstrass arcs.

The angle 4π cone point c^* divides each indirect systolic Weierstrass arc into two subarcs that we call “prongs.” Observe that if some prong has length $\ell \leq \text{sys}(X)/4$, then each of the other prongs emanating from c^* has length $\text{sys}(X)/2 - \ell$. In §6 we show that if all of the prongs have the same length — necessarily $\text{sys}(X)/4$ — then there are at most four prongs, and if there is a “short” prong of length $\ell < \text{sys}(X)/4$, then there are at most five prongs. In the former case, we obtain at most six indirect systolic Weierstrass arcs and in the latter case, we obtain at most five indirect systolic Weierstrass arcs.²

In §7 we show that there is at most one systole that is a separating curve. Moreover, we show that if the surface has a systole which is a separating curve, then the surface has either no prongs or exactly two prongs of equal length. It follows that a surface with a separating systole has at most eight homotopy classes of separating systoles.

In §8, we show that if there are exactly four prongs of equal length, then the surface has at most ten homotopy classes of systoles, and if there are ten, then the surface is homothetic to the surface described in Figure 1. In §9, we show that if one of the prongs is shorter than the others, then there are at most none homotopy classes of systoles. This finishes the proof of Theorem 1.1 in the case of surfaces from the stratum $\mathcal{H}(1, 1)$.

Although the questions that we address in this paper regarding systoles have not been systematically studied previously in the context of translation surfaces, they have been studied in the context of hyperbolic and general Riemannian surfaces. As hinted at above, smooth surfaces have systoles that intersect at most once, and from this one can deduce that there are at most 12 homotopy classes of systole in genus two (see for instance [9]). This bound is sharp. Indeed, among hyperbolic surfaces of genus two, there is a unique surface, called the Bolza surface, with exactly 12 systoles. It can be

²Note that a particular prong can lie in more than one systolic Weierstrass arc.

obtained by gluing opposite edges of a regular hyperbolic octagon with all angles $\frac{\pi}{4}$. This same surface is also optimal (again among hyperbolic surfaces) for systolic ratio, a result of Jenni [7]. There are bounds on these quantities in higher genus, but these bounds are not optimal. Interestingly, Katz and Sabourau [8] showed that among CAT(0) genus two surfaces, the optimal surface is an explicit flat surface with cone point singularities, conformally equivalent to the Bolza surface. This singular surface can not be optimal among all Riemannian surfaces however, as by a result of Sabourau, the optimal surface in genus two necessarily has a region with positive curvature [13]. The optimal systolic ratio among all Riemannian surfaces is still not known.

Acknowledgements. We are grateful to the referee for a careful reading of the paper and valuable comments. We thank Marston Conder for the examples in Remark 3.2. We thank Carlos Matheus Santos and Gabriela Weitze-Schmithüsen for kindly pointing out some mistakes as well as some missing references in earlier versions. H. P. acknowledges support from U.S. National Science Foundation grants DMS 1107452, 1107263, 1107367 RNMS: Geometric structures And Representation varieties (the GEAR Network). C.J. acknowledges support from the Simons Foundation.

2. Facts concerning the geometry of (X, d_ω)

We collect here some relevant facts about the geometry of the surface (X, d_ω) sometimes called a “translation surface.” Much of this material can be found in, for example, [2, 6, 10].

2.1. Integrating the 1-form. By integrating the holomorphic 1-form ω along a piecewise differentiable path $\alpha: [a, b] \rightarrow X$, we obtain a path in $\bar{\alpha}: [a, b] \rightarrow \mathbb{C}$ defined by

$$\bar{\alpha}(t) = \int_{\alpha|_{[a,t]}} \omega. \quad (2.1)$$

Since ω is closed, if two paths α, β in X are homotopic rel endpoints, then $\bar{\alpha}$ and $\bar{\beta}$ are homotopic rel endpoints. Thus, if $U \subset X$ is simply connected neighborhood of a point x , then

$$\mu_{x,U}(y) := \int_{\alpha_y} \omega \quad (2.2)$$

is independent of the path α_y joining x to y . Note that $\mu_{x,U}$ is a holomorphic map from U into \mathbb{C} . If x is not a zero of ω , then it follows from the inverse function theorem that there exists a neighborhood U so that $\mu_{x,U}$ is a biholomorphism onto its image.

2.2. The metric. The norm, $|\omega|$, of ω defines an arc length element on X . We will let $\ell_\omega(\alpha)$ denote the length of a path on X , and we will let d_ω denote the metric obtained by taking the infimum of lengths of paths joining two points.

If x is not a zero of ω and U is a simply connected neighborhood of x , then $\mu_{x,U}$ is a local isometry from U into \mathbb{C} equipped with its usual Euclidean metric $|dz|^2$. If, in addition, U is star convex at x , then $\mu_{x,U}$ is an isometry onto its image.

If x is a zero of ω of order k , then there exists a neighborhood V of x and a chart $\nu: V \rightarrow \mathbb{C}$ such that $\omega = (k+1) \cdot \nu^*(z^k dz) = \nu^*(d(z^{k+1}))$ and $\nu(x) = 0$. If V is sufficiently small, the map ν is an isometry from (V, d_ω) to $(\nu(V), d_{d(z^{k+1})})$. In turn, the map $z \mapsto z^{k+1}$ is a local isometry from $(\nu(V) - \{0\}, d_{d(z^{k+1})})$ to a neighborhood of the origin with the Euclidean metric $|dz|^2$. Since the branched covering $z \mapsto z^{k+1}$ has degree $k+1$, the arc length of the boundary of an ϵ -neighborhood of x is $2\pi(k+1) \cdot \epsilon$. Therefore, we refer to x as a *cone point* of angle $2\pi(k+1)$, and the set of zeros, denoted Z_ω , will be regarded as the set of cone points of (X, d_ω) .

2.3. Universal cover, developing map and holonomy. Let $p: \tilde{X} \rightarrow X$ be the universal covering map, and let $\tilde{\omega} = p^*(\omega)$. If we let $d_{\tilde{\omega}}$ be the associated metric on \tilde{X} , then p is a local isometry from $(\tilde{X}, d_{\tilde{\omega}})$ onto (X, d_ω) . Since \tilde{X} is simply connected, we may fix $\tilde{x}_0 \in \tilde{X}$ and integrate $\tilde{\omega}$ as in (2.2) to obtain a map $\text{dev}: \tilde{X} \rightarrow \mathbb{C}$ called the *developing map*. The restriction of dev to $\tilde{X} - Z_{\tilde{\omega}}$ is a local biholomorphism and a local isometry. Each zero of $\tilde{\omega}$ is a branch point whose degree equals the order of the zero. If C is the closure of a convex subset of $\tilde{X} - Z_{\tilde{\omega}}$, then the restriction of dev to C is injective.

Let $x_0 = p(\tilde{x}_0)$, and consider loops α in X based at x_0 . The assignment $\alpha \rightarrow \bar{\alpha}$ defines a homomorphism, hol , from $\pi_1(X, x_0)$ to the additive group \mathbb{C} . Moreover, for each $[\alpha] \in \pi_1(X, x_0)$ and $\tilde{x} \in \tilde{X}$ we have

$$\text{dev}([\alpha] \cdot \tilde{x}) = \text{dev}(\tilde{x}) + \text{hol}([\alpha]) \quad (2.3)$$

where $\alpha \cdot \tilde{x}$ denotes action by covering transformations. See, for example, [6].

2.4. Geodesics. If a geodesic γ on (X, d_ω) passes through a zero of ω , then γ will be called *indirect* and otherwise *direct*. If γ is a direct simple geodesic loop, then, since Z_ω is finite, for sufficiently small $\epsilon > 0$, the ϵ -tubular neighborhood, N , of γ is disjoint from Z_ω . Each lift $\tilde{N} \subset \tilde{X}$ of N is convex and hence the restriction of the developing map to \tilde{N} is an isometry onto $\text{dev}(\tilde{N})$. Since \tilde{N} is stabilized by the cyclic subgroup $\langle \gamma \rangle$ of the deck group generated by γ , it follows from (2.3) that $\text{dev}(\tilde{N})$ is the convex hull of two parallel lines, and, moreover, the map dev determines an isometry from N to $\text{dev}(\tilde{N})/\langle \text{hol}(\gamma) \rangle$. In particular, N is isometric to a Euclidean cylinder $[0, w] \times \mathbb{R}/\ell\mathbb{Z}$ where $\ell = |\text{hol}(\gamma)|$ and w is the distance between the parallel lines. If $Z_\omega \neq \emptyset$, then the union of all Euclidean cylinders embedded in $X - Z_\omega$ that

contain γ is a cylinder called the *maximal cylinder* associated to γ . Each component of the frontier of a maximal cylinder consists of finitely many indirect geodesics.

Proposition 2.1. *If ω has at least one zero, then each homotopy class of loops is represented by a geodesic loop that passes through a zero of ω .*

Proof. Since X is compact, a homotopy class of simple loops has a geodesic representative γ . If γ does not pass through a zero, then γ lies in a maximal cylinder. The boundary of the maximal cylinder contains a geodesic representative that passes through a zero. \square

Proposition 2.2. *If two direct simple geodesic loops are homotopic, then they lie in the closure of the same maximal cylinder.*

Proof. Because the angle at each cone point $\tilde{z} \in Z_{\tilde{\omega}}$ is greater than 2π , the length space $(\tilde{X}, d_{\tilde{\omega}})$ is CAT(0). If two geodesic loops γ and γ' are homotopic, then they have lifts that are asymptotic in $(\tilde{X}, d_{\tilde{\omega}})$. By the flat strip theorem [3], the convex hull of the two lifts is isometric to a strip $[0, w] \times \mathbb{R}$. Thus, since each cone point has angle larger than 2π , the interior I of the convex hull contains no cone points. The developing map restricted to I is an isometry onto a strip in \mathbb{C} , and, moreover, it induces an isometry from $I/\langle g \rangle$ to the cylinder $\text{dev}(I)/\langle \text{hol}(g) \rangle$ where g is the deck transformation associated to the common homotopy class of γ and γ' . Since the lifts are boundary components of I , the loops γ and γ' lie in the boundary of the cylinder $\text{dev}(I)/\langle \text{hol}(g) \rangle$. \square

2.5. The Delaunay cell decomposition. The Delaunay decomposition is well-known in the context of complete constant curvature geometries. Thurston observed that the construction also applies to constant curvature metrics with conical singularities [15].

We will first describe the Delaunay decomposition of the universal cover \tilde{X} . Given $\tilde{x} \in \tilde{X} - Z_{\tilde{\omega}}$, let $D_{\tilde{x}}$ be the largest open disk centered at \tilde{x} that does not intersect $Z_{\tilde{\omega}}$. Since $D_{\tilde{x}}$ is convex, the restriction of dev to the closure $\bar{D}_{\tilde{x}}$ is an isometry onto a closed Euclidean disk in \mathbb{C} . Since $Z_{\tilde{\omega}}$ is discrete, the intersection $Z_{\tilde{\omega}} \cap \bar{D}_{\tilde{x}}$ is finite. Let \mathcal{V} be the set of $\tilde{x} \in \tilde{X} - Z_{\tilde{\omega}}$ such that $Z_{\tilde{\omega}} \cap \bar{D}_{\tilde{x}}$ contains at least three points. Because three points determine a circle, the set \mathcal{V} is discrete.

For each $\tilde{x} \in \mathcal{V}$, let $P_{\tilde{x}}$ denote the convex hull of $Z_{\tilde{\omega}} \cap \bar{D}_{\tilde{x}}$. It is isometric to a convex polygon in the plane. Again, because three points determine a circle, if $\tilde{x}, \tilde{y} \in \mathcal{V}$ and $\tilde{x} \neq \tilde{y}$, then the set $Z_{\tilde{\omega}} \cap \bar{D}_{\tilde{x}} \cap \bar{D}_{\tilde{y}}$ consists of at most two points, and hence $P_{\tilde{x}} \cap P_{\tilde{y}}$ is either empty, a point, or a geodesic arc lying in both the boundary of $P_{\tilde{x}}$ and the boundary of $P_{\tilde{y}}$. The interior of $P_{\tilde{x}}$ is called a *Delaunay 2-cell* and the boundary edges are called *Delaunay edges*. The vertex set of this decomposition of \tilde{X} is the set of zeros of $\tilde{\omega}$.

The deck group of the universal covering map p permutes the cells of the Delaunay decomposition, and so we obtain a decomposition of X . Note the restriction of p to each 2-cell P is an isometry onto its image. Indeed, if not then there exists a covering

transformation γ , a lift \tilde{P} of P , and $\tilde{x} \in \tilde{P}$ such that $\gamma \cdot \tilde{x} \in \tilde{P}$. Since \tilde{P} is convex, it follows that for some vertex $\tilde{z} \in \tilde{P}$, we would have $\gamma \cdot \tilde{z} \in \tilde{P}$. But γ maps $Z_{\tilde{\omega}}$ to itself.

Our interest in the Delaunay decomposition stems from the following.

Proposition 2.3. *If α is a shortest non-null homotopic arc with both endpoints in Z_{ω} , then α is a Delaunay edge.*

Proof. Since the universal covering map p preserves the length of arcs, it suffices to prove that the analogous statement holds for the universal cover \tilde{X} . Because α is a shortest arc, if m is the midpoint of α , then the largest disc D centered at m has diameter equal to $\ell(\alpha)$ and $\bar{D} \cap Z_{\tilde{\omega}}$ consists of exactly two points, the endpoints z and z' of α . The circle $\text{dev}(\partial D)$ belongs to the pencil of circles containing $\text{dev}(z)$ and $\text{dev}(z')$. Since X is compact, by varying over this pencil, we find a disk D' so that $\bar{D}' \cap Z_{\tilde{\omega}}$ contains z, z' , and at least one other point. The center c of D' belongs to \mathcal{V} and α is a boundary edge of the polygon P_c . \square

Proposition 2.4. *Let ω be a holomorphic 1-form on a closed surface of genus g . If ω has v zeros, then the Delaunay decomposition of X has at most $6g - 6 + 3 \cdot v$ edges and the number of 2-cells is at most $4g - 4 + 2 \cdot v$. Equality holds if and only if each 2-cell is a triangle.*

Proof. By dividing the Delaunay 2-cells (convex polygons) into triangles, we obtain a triangulation with v vertices. By Euler's formula and the fact that there are 3 oriented edges for each triangle, we find that each triangulation has $6g - 6 + 3v$ edges and $4g - 4 + 2 \cdot v$ triangles. \square

3. Systoles of 1-forms in $\mathcal{H}(2g - 2)$

In this section, we consider holomorphic 1-forms with a single zero. In the first part of the section we give the optimal bound on the number of homotopy classes of systoles of such surfaces as well as the optimal bound for the hyperelliptic surfaces with one zero. In the second part, we provide the optimal estimate on the systolic ratio of such surfaces.

3.1. Bounds on the number of systoles.

Proposition 3.1. *If ω is a holomorphic 1-form on X that has exactly one zero, then (X, d_{ω}) has at most $6g - 3$ homotopy classes of systoles.*

Proof. By Proposition 2.1, each homotopy class of systoles contains a representative that passes through the zero. Proposition 2.3 implies that each such systole is a Delaunay edge. By Proposition 2.4, there are at most $6g - 3$ Delaunay edges and hence at most $6g - 3$ homotopy classes of systoles. \square

The bound in Proposition 3.1 is sharp if the genus g of X is at least 3. For example, if $g = 3, 4, 5$, then consider the surfaces described in Figures 4, 5, and 6.




Figure 4. Genus 3 example that saturates bound in Proposition 3.1. Glue the edges of the polygon according to the labels. Each edge is a systole, the 1-form ω has exactly one zero, and no two Delaunay edges are homotopic.




Figure 5. Genus 4 example that saturates bound in Proposition 3.1.




Figure 6. Genus 5 example that saturates bound in Proposition 3.1.

More generally, given a holomorphic 1-form ω_g on a surface X_g of genus g with one zero that achieves the bound $6g - 3$, one can construct a holomorphic 1-form ω_{g+3} with one zero on a surface X_{g+3} of genus $g + 3$ that achieves the bound $6(g + 3) - 3$. Indeed, remove the interior of a Delaunay edge from $(X_g, d\omega_g)$ to obtain a surface X'_g with “figure eight” boundary consisting of two segments F_-, F_+ each corresponding to the Delaunay edge. Let (Y_2, dz) be the genus two surface with two boundary components G_-, G_+ that is described in Figure 7. By gluing F_\pm to G_\pm , we obtain the desired (X_{g+3}, ω_{g+3}) .




Figure 7. Glue solid edges of the polygon that have the same label to obtain the Delaunay triangulation associated a holomorphic 1-form on a surface of genus two having two boundary components.

Remark 3.2. The problem of constructing surfaces that saturate the bound in Proposition 3.1 is equivalent to the problem of constructing two fixed-point free elements σ, τ in the symmetric group $S_{2g-1} = \text{Sym}(\{1, \dots, 2g-1\})$ such that $\sigma \cdot \tau$ has no fixed points and the commutator $[\sigma, \tau]$ is a $(2g-1)$ -cycle. Indeed, let P_1, \dots, P_g be $2g-1$ disjoint copies of the convex hull of $\{0, 1, e^{\pi i/3}, 1 + e^{\pi i/3}\}$. Given $\sigma, \tau \in S_{2g-1}$, glue the left side of P_i to the right side of $P_{\sigma(i)}$ and the top side of P_i to the bottom side of $P_{\tau(i)}$ to obtain a surface with a holomorphic 1-form ω . If $[\sigma, \tau]$ is an n -cycle, then it follows that ω has one zero, and if σ, τ , and $\sigma \cdot \tau$ have no fixed points, then it follows that (X, d_ω) has no cylinder with girth equal to the systole. Thus, by Proposition 2.2, no two systolic edges are homotopic.

Conversely, suppose that a holomorphic 1-form surface saturates the bound, then the necessarily equilateral Delaunay triangles can be paired to form parallelograms as above that are glued according to permutations σ and τ . One verifies that σ and τ satisfy the desired properties.

The surface constructed in Figure 4 corresponds to the pair

$$\sigma = (12345), \quad \tau = (15243),$$

the surface constructed in Figure 5 corresponds to the pair

$$\sigma = (1234567), \quad \tau = (1364527),$$

and the surface in Figure 6 corresponds to

$$\sigma = (123456789), \quad \tau = (146379285).$$

We thank Marston Condon for finding these examples for us.

If the genus of the surface is two, then one can show that the maximum number of homotopy classes of systoles is $7 = 6g - 5$. More generally, the following is true.

Theorem 3.3. *Let ω be a holomorphic 1-form on a surface with a hyperelliptic involution τ . If ω has exactly one zero, then (X, d_ω) has at most $6g - 5$ homotopy classes of systoles. Moreover, (X, d_ω) has exactly $6g - 5$ homotopy classes of systoles if and only if each Delaunay edge is a systole and there exist exactly four Delaunay 2-cells each of which have two edges that are preserved by the hyperelliptic involution.*

For each $g \geq 2$, the bound given in Theorem 3.3 is achieved by, for example, the surface described in Figure 8.

Proof. Each homotopy class of systole is represented by at least one systolic Delaunay edge. Since ω has exactly one zero, z_0 , the number of Delaunay edges is at most $6g - 3$. Thus, we wish to show that if there are $6g - 3$ or $6g - 4$ systolic Delaunay edges, then there exist at least two homotopic pairs of systolic edges, and that, if there are $6g - 5$ systolic edges, then there is at least one pair of homotopic edges.




Figure 8. Glue edges of the polygon that have the same label to obtain the Delaunay triangulation associated to a holomorphic 1-form on a surface of genus g . The surface is hyperelliptic, the 1-form ω has exactly one zero, and there are exactly $6g - 5$ homotopy classes of systoles.

$6g - 3$ systolic edges: Suppose that there are exactly $6g - 3$ systolic Delaunay edges. Then each Delaunay 2-cell is an equilateral triangle and by Proposition 2.4 there are $4g - 2$ such cells. Since τ is an isometry, it preserves the Delaunay partition. In particular, since z_0 is the unique 0-cell, we have $\tau(z_0) = z_0$, and since an equilateral triangle has no (orientation preserving) involutive isometry, the involution τ has no fixed points on the interior of each 2-cell. Thus, the remaining $2g + 1$ fixed points of τ lie on 1-cells. In particular, τ fixes exactly $2g + 1$ Delaunay edges.

Suppose that T is a 2-cell with two fixed edges. Then $T \cup \tau(T)$ is a cylinder whose boundary components are the “third” edges of T and $\tau(T)$, and, in particular, since the genus of X is at least two, these “third” edges are not fixed by τ . Thus, a 2-cell has either zero, one, or two fixed edges. Note that the number of 2-cells that have two fixed edges is even.

We claim that there exist at least four 2-cells that each have two fixed edges. Indeed, if, on the contrary, there are at most two such 2-cells, then there are at least $4g - 4$ remaining 2-cells that each have at most one fixed edge. Thus, there are at most $2g - 2$ fixed Delaunay edges associated to these 2-cells, and at most 2 edges associated to the 2-cells that have two fixed edges. But, there are $2g + 1 > (2g - 2) + 2$ fixed edges, and so we have a contradiction.

The four 2-cells form two cylinders each bounded by two systolic edges. Thus, there are at most $6g - 5$ homotopy classes of systoles.

If there are exactly $6g - 5$ homotopy classes of cylinders, then there are two maximal cylinders each bounded by two systolic edges. The integral of ω over the middle curve of each cylinder is nonzero, and hence the middle curve is not null-homologous. The induced action of a hyperelliptic involution on $H_1(X)$ is the antipodal map, and so τ preserves each cylinder and has exactly two fixed points on the interior of each cylinder. It follows that there are exactly four Delaunay 2-cells each having two fixed edges.

$6g - 4$ systolic edges: Suppose that there are exactly $6g - 4$ systolic Delaunay edges. It follows that exactly $4g - 4$ Delaunay 2-cells are equilateral triangles. The complement, K , of the union of these equilateral triangles is (the interior of) a rhombus.

Since τ is an isometry, τ preserves K , and hence the center c of the rhombus is a fixed point of τ . The other Delaunay 2-cells are equilateral triangles and hence do not contain fixed points. Therefore, since τ has exactly $2g - 2$ fixed points and $\tau(z_0) = z_0$, exactly $2g$ systolic edges are fixed by τ .

If an edge e in ∂K is fixed by τ , then e is equal to the opposing edge and in particular $K \cup e$ is a cylinder. Indeed, if e were fixed by τ , then the segment in K joining the midpoint of e to c would be “rotated” by τ to a segment joining c to the midpoint of the edge e' opposite to e . Hence the midpoint of e would equal the midpoint of e' , and thus $e = e'$.

Since X is connected and of genus at least two, not all four edges of ∂K can be fixed by τ . Thus either \bar{K} is a rhombus with no fixed edges or a cylinder with no fixed boundary edges.

Suppose that \bar{K} is a rhombus. Among the remaining $4g - 4$ two-cells — equilateral triangles — there are exactly $2g$ fixed points. Hence there exist equilateral triangles that have at least two fixed edges. If there were $4g - 6$ equilateral triangles that each had at most one fixed edge, then there would be only $2g - 3 + 2 = 2g - 1$ fixed points among the $4g - 4$ equilateral triangles. It follows that there are at least four equilateral triangles that each have two fixed edges, and thus there exist two distinct maximal cylinders bounded by systoles.

Suppose that \bar{K} is a cylinder. In this case, neither of the two equilateral triangles that share edges with \bar{K} can have two fixed edges. Indeed, using an argument as above with a segment joining the center of the rhombus, we would see that the edges would be identified in such a way to form a torus.

Consider the two equilateral triangles T_+ , T_- that have an edge in \bar{K} . If an edge e of T_{\pm} is fixed then τ , then using the symmetry about c , one shows as above that e is identified with an edge of T_{\mp} . Because the X is connected and of genus at least two, T_{\mp} has at most one edge fixed by τ . It follows that among the remaining $4g - 6$ triangles there are at least $2g + 2 - 1 - 3 = 2g - 2$ fixed points. It follows that there exist at least one equilateral that has two fixed edges, and hence there exists another maximal cylinder bounded by systoles.

In either case, we have two maximal cylinders bounded by systoles, and therefore there are at most $6g - 6$ homotopy classes of systoles.

6g - 5 systolic edges: Suppose that there are exactly $6g - 5$ systolic edges. Then there are $4g - 6$ Delaunay 2-cells that are equilateral triangles. The complement, K , of the union of these equilateral triangles consists of either an equilateral hexagon or two disjoint rhombi.

Suppose that K is an equilateral hexagon. Then since τ preserves the Delaunay partition, we have $\tau(K) = K$. Hence K contains exactly one fixed point c and K is convex. Thus, arguing as above, we find that if a boundary edge of K is fixed by τ , then the edge equals an opposite edge. Since X is connected with genus at least two, all six edges can not be identified, and hence there are at most 3 fixed points in \bar{K} .

We claim that at least one pair of equilateral triangles each have exactly two fixed edges. If not, then each of the $4g - 6$ equilateral triangles contains at most one fixed edge. Thus, there are at most $2g - 3$ such edges, and hence $(2g - 3) + 3 + 1 = 2g + 1$ fixed points in total. But the total number of fixed points is $2g + 2$. Thus, we have a pair of equilateral triangles that share a pair of fixed edges. The union is a cylinder bounded by two systolic edges, and so we have at most $6g - 6$ homotopy classes of systoles in this case.

Finally suppose that K is the disjoint union of two rhombi R_+ and R_- . Since τ preserves the Delaunay partition, either $\tau(R_{\pm}) = R_{\pm}$ or $\tau(R_{\pm}) = R_{\mp}$.

If $\tau(R_{\pm}) = R_{\pm}$, then each rhombus contains a fixed point. If an edge of R_{\pm} is fixed, then $\overline{R_{\pm}}$ is a cylinder bounded by systolic edges and so there are at most $6g - 6$ homotopy classes of systoles. If neither rhombus has boundary edges fixed by τ , then \overline{K} contains exactly two fixed points. If there is not a pair of equilateral triangles that share fixed boundary edges, then each of the $4g - 6$ equilateral triangles would have at most one fixed edge, and so there would be at most $2g - 3 + 2 + 1 = 2g$ fixed points, a contradiction. Hence we have a systolic cylinder and at most $6g - 6$ homotopy classes of systoles.

If $\tau(R_{\pm}) = R_{\mp}$, then the rhombi do not contain fixed points. If an edge in ∂R_{\pm} is fixed by τ , then R_{\pm} shares this edge with R_{\mp} . It follows that there are at most three fixed points in \overline{K} , and one may argue as in the case of the hexagon, to find that there are at most $6g - 6$ homotopy classes of systoles.

If no edge in ∂R_{\pm} is fixed by τ , then among the $4g - 6$ equilateral triangles there are $2g + 1$ fixed points. It follows that there is an equilateral triangle that has two edges fixed by τ , and hence there is a maximal cylinder bounded by systoles. \square

Since each genus two surface is hyperelliptic, we have the following corollary.

Corollary 3.4. *Let X be a surface of genus two. If ω is a holomorphic 1-form on X that has exactly one zero, then the number of homotopy classes of systoles of (X, d_{ω}) is at most 7.*

3.2. Lengths of systoles. Although our main concern is the number of systoles, we observe in this section that it is quite straightforward to find a sharp upper bound on the length of systoles of translation surfaces provided they have a single cone point singularity. One of the ingredients is the Delaunay triangulation described in §2.5. The other ingredient is a result due to Fejes Tóth which we state in the form of the following lemma.

Lemma 3.5. *Let T be a Euclidean triangle embedded in the plane and let r be the maximal positive real number so that the open balls of radius r around the three vertices are disjoint. Then*

$$r^2 \leq \frac{\text{Area}(T)}{\sqrt{3}}$$

with equality if and only if T is equilateral.

This can be stated differently in terms of ratios of areas. Consider the area A_r of a triangle found at distance r from the vertices of T and so that the interior of the three sectors do not overlap. Then the ratio A_r/T never exceeds that of the equilateral triangle with r equal to half the length of a side.

Theorem 3.6. *If $(X, \omega) \in \mathcal{H}(2g - 2)$, then*

$$\frac{\text{sys}^2(X)}{\text{area}(X)} \leq \frac{4}{(4g - 2) \cdot \sqrt{3}} \quad (3.1)$$

with equality if and only if X is tiled by equilateral triangles.

Proof. Let z_0 denote the zero of ω . By Proposition 2.3, each systole that passes through z_0 lies in the 1-skeleton of the Delaunay cell decomposition of (X, d_ω) . Thus, if r_0 is the radius of the largest open Euclidean ball that can be embedded in (X, d_ω) with center z_0 , then $r_0 = \text{sys}(X)/2$. Therefore

$$\text{sys}^2(X) = 4 \cdot r_0^2. \quad (3.2)$$

Each open 2-cell P of the Delaunay cell-decomposition is isometric to a convex Euclidean polygon. We may further subdivide each 2-cell into Euclidean triangles to obtain a triangulation of X with one vertex, namely z_0 . By Proposition 2.4, there are at exactly $4g - 2$ such triangles. Thus, Lemma 3.5 implies that

$$\text{area}(X) = \sum_T \text{area}(T) \geq (4g - 2) \cdot \sqrt{3} \cdot r_0^2$$

where the sum is over triangles in the triangulation. By combining this estimate with (3.2), we obtain the desired inequality (3.1). Moreover, if equality holds in (3.1), then equality holds in Lemma 3.5, and so each triangle is an equilateral triangle. \square

We note that there is a unique surface (up to homothety) in $\mathcal{H}(2)$ tiled by equilateral triangles (illustrated previously in Figure 3). This surface realizes the maximum number of systoles and the maximum systolic ratio over $\mathcal{H}(2)$. In contrast, as indicated in the introduction, the maximum systolic ratio over $\mathcal{H}(1, 1)$ is not realized by the unique surface that realizes the maximum number of homotopy classes of systoles.

4. Geodesics on a surface in $\mathcal{H}(1, 1)$

In this section, X will denote a $\mathcal{H}(1, 1)$ surface of genus two equipped with a translation structure with two cone points c_+ and c_- each of angle 4π . The tangent bundle of a translation surface is parallelizable. In particular, each oriented segment has a direction. The hyperelliptic involution $\tau: X \rightarrow X$ is an isometry that reverses the direction of each oriented segment. The isometry τ has exactly six fixed points, the *Weierstrass points*.

Lemma 4.1. *The hyperelliptic involution τ interchanges cone points: $\tau(c_{\pm}) = c_{\mp}$*

Proof. Since τ is an isometry the set $\{c_+, c_-\}$ is permuted. If $\tau(c_+) = c_+$, then in a neighborhood of c_+ , the isometry τ acts as a rotation of π radians. But the cone angle is 4π , and hence it is impossible for τ^2 to be the identity. \square

By Lemma 4.1, the quotient $X/\langle\tau\rangle$ is a sphere with one cone point c^* with angle 4π and six cone points $\{c_1, \dots, c_6\}$ each of angle π . Let $p: X \rightarrow X/\langle\tau\rangle$ denote the degree 2 covering map branched at $\{c_1, \dots, c_6\}$. If γ is a simple geodesic loop, then either γ passes through two Weierstrass points in which case p maps γ onto a geodesic arc joining two distinct π cone points, or $p \circ \gamma$ is a simple geodesic loop that misses the π cone points.

A *flat torus* is a closed translation surface (necessarily of genus one). A *slit torus* is a flat torus with finitely many disjoint simple geodesic arcs removed. Each removed arc is called a *slit*. The completion of a slit torus (with respect to the natural length space structure) is obtained by adding exactly two geodesic segments for each removed arc. The interior angle between each pair of segments is 2π . This property characterizes slit tori.

Lemma 4.2. *Let Y be a topological torus with a closed disc removed. If Y is equipped with a translation structure such that the boundary³ consists of at most two geodesic segments, then Y is isometric to a slit torus.*




Figure 9. Identify the edges with the same labels via elements of $\text{Isom}(\mathbb{R}^2)$ to obtain a torus with a disc removed equipped with a flat structure such that the boundary consists of exactly two geodesics. The angles between the geodesics are not both π though they sum to 4π .

³By *boundary* we mean the set of points added by taking the metric completion of the length structure associated to the translation structure.

Remark 4.3. Figure 9 shows that Lemma 4.2 is false if one replaces the assumption of translation structure with the assumption of flat structure.

Proof of Lemma 4.2. Let Z be the boundary of Y . Let A be the intersection of the maximal geodesic segments in Z . By assumption A is either empty, contains one point, or contains two points. Let $\alpha: [0, 1] \rightarrow Z$ be a parameterization of Z such that if A is nonempty, then $\alpha(0) = \alpha(1) \in A$. Let $\bar{\alpha}$ be the development of α into the plane \mathbb{C} as discussed in §2.

Since $[\alpha] \in \pi_1(Y)$ is a commutator and \mathbb{C} is abelian, the holonomy of $[\alpha]$ equals 0. Hence by (2.3), we have $\bar{\alpha}(1) - \bar{\alpha}(0) = \text{dev}([\alpha] \cdot \tilde{\alpha}(0)) = 0$, and therefore $\bar{\alpha}(1) = \bar{\alpha}(0)$.

If A is empty or consists of one point, then $\bar{\alpha}$ is a line segment, but this is impossible as line segments in \mathbb{C} have distinct endpoints. If A consists of two points, then the curve $\bar{\alpha}$ consists of two line segments. Since $\bar{\alpha}(1) = \bar{\alpha}(0)$, the line segments coincide. Removing this segment and its translates by $\text{hol}(\pi_1(Y))$ and quotienting it by $\text{hol}(\pi_1(Y))$ gives a surface isometric to Y . \square

As a corollary, we have the following sharpening of Theorem 1.7 in [11].

Corollary 4.4. *If α is a separating simple closed geodesic on X , then $X - \alpha$ is the disjoint union of two slit tori. Moreover, each slit torus contains exactly three Weierstrass points, and the hyperelliptic involution τ preserves α .*

Proof. Since α is separating and X is closed of genus two, the complement of α consists of two one-holed tori Y_+ and Y_- . Since α is geodesic, the boundaries of Y_+ and Y_- are piecewise geodesic. Since α is simple and there are only two cone points, the number of geodesic pieces of Y_{\pm} is at most two. Lemma 4.2 implies that each component is a slit torus.

The restriction of τ to a slit torus component determines an elliptic involution $\underline{\tau}$ of the torus. The endpoints of each slit correspond to the cone points c_+ and c_- , and so they are preserved by the induced elliptic involution. Since τ preserves the cone points, the map $\underline{\tau}$ preserves the slit, and hence α is preserved by τ . In particular, the midpoint of the slit is fixed by $\underline{\tau}$ and the three other fixed points of $\underline{\tau}$ are fixed points of τ . \square

A cylinder of girth ℓ and width w is an isometrically embedded copy of $(\mathbb{R}/\ell\mathbb{Z}) \times [-w/2, w/2]$. Each cylinder is foliated by geodesics indexed by $t \in [-w/2, w/2]$. We will refer to the geodesic that corresponds to $t = 0$ as the *middle geodesic*. By Corollary 4.4, if a simple closed geodesic lies in a cylinder, then it is nonseparating.

A cylinder C is said to be *maximal* if it is not properly contained in another cylinder. If a closed translation surface has a cone point, then each geodesic that does not pass through a cone point lies in a unique maximal cylinder.

Because the hyperelliptic involution τ reverses the orientation of isotopy classes of simple curves, the map τ restricts to an orientation reversing isometry of each

maximal cylinder C , and thus it restricts to an orientation reversing isometry of the middle geodesic $\gamma \subset C$. In particular, it contains two Weierstrass points.

Proposition 4.5. *If γ is a nonseparating simple closed geodesic, then γ is homotopic to a unique geodesic γ' such that the restriction of τ to γ' is an isometric involution of γ' .*

Proof. If γ does not contain a cone point, then γ belongs to a maximal cylinder. If γ belongs to a maximal cylinder C , then it is homotopic to the middle geodesic $\gamma' \subset C$.

If γ does not belong to a cylinder, then γ is the unique geodesic in its homotopy class. Since τ reverses the orientation of the homotopy classes of simple loops, it acts like an orientation reversing isometry on γ . \square

Proposition 4.5 reduces the counting of homotopy classes of nonseparating systoles to a count of nonseparating systoles that pass through exactly two Weierstrass points. In the next two sections we analyse such geodesics.

5. Direct Weierstrass arcs

If γ is a simple closed geodesic on X that passes through two Weierstrass points, then the projection $p(\gamma)$ is an arc on $X/\langle\tau\rangle$ that joins one angle π cone point to another angle π cone point. We will call each such an arc a *Weierstrass arc*. Note that the p inverse image of a Weierstrass arc is a geodesic and so we obtain a one-to-one correspondence between homotopy classes of nonseparating simple geodesic loops on X and Weierstrass arcs on $X/\langle\tau\rangle$. A Weierstrass arc that is the image of a systole will be called a *systolic Weierstrass arc*. Note that each systolic Weierstrass arc has length equal to $\text{sys}(X)/2$.

The Weierstrass arcs come in two flavors. We will say that a Weierstrass arc is *indirect* if it passes through the angle 4π cone point, and otherwise we will call it *direct*.

Lemma 5.1. *There is at most one direct systolic Weierstrass arc joining two angle π cone points.*

Proof. Suppose to the contrary that there exist two distinct direct systolic Weierstrass arcs that both join the angle π cone point c to the angle π cone point $c' \neq c$. These arcs lift to closed systoles γ_+ and γ_- that intersect transversally at two Weierstrass points corresponding to c and c' . In particular, the Weierstrass points divide each geodesic into two arcs. By concatenating a shorter⁴ arc of γ_+ with a shorter arc of γ_- we construct a piecewise geodesic closed curve α that has length at most the systole. Since the angle between the arcs is strictly between 0 and π , we can perturb α to

⁴If the arcs have the same length, then choose either arc.

obtain a shorter curve whose length is strictly less than the systole. This contradicts the assumption that γ_+ and γ_- are both systoles. \square

Remark 5.2. The argument in the above lemma was that the concatenation of two geodesic arcs that meet with an angle strictly less than π cannot be of minimal length in their homotopy class. In particular, they can't form a systole. This argument will be used several times.

Proposition 5.3. *Let γ_+ and γ_- be distinct nonseparating systoles on X . If each contains two Weierstrass points and neither contains a 4π cone point, then the intersection $\gamma_+ \cap \gamma_-$ is either empty or consists of a single Weierstrass point. In particular, the geometric intersection number $i(\gamma_+, \gamma_-)$ equals either zero or one.*

Proof. Each projection $\alpha_{\pm} = p(\gamma_{\pm})$ is a direct systolic Weierstrass arc. By Lemma 5.1, at most one angle π cone point lies in the intersection $\alpha_+ \cap \alpha_-$, and hence $\gamma_+ \cap \gamma_-$ contains at most one Weierstrass point.

Suppose (to the contrary) that the intersection $\gamma_+ \cap \gamma_-$ were to contain a point on X that is not a Weierstrass point. Then $\alpha_+ \cap \alpha_-$ would contain a point p that is not an angle π cone point. Since, γ_{\pm} is a systole, there would exist a subarc, β_{\pm} , of α_{\pm} that joins p to an endpoint of α_{\pm} whose length is at most $\text{sys}(X)/4$. By concatenating β_+ and β_- and perturbing, we would obtain an arc joining two angle π cone points whose length would be strictly less than $\text{sys}(X)/2$. This arc would lift to a closed curve on X whose length is less than $\text{sys}(X)/2$, a contradiction. \square

The following result is central to the proof of Theorem 1.1.

Theorem 5.4. *If c is a cone point on $X/\langle\tau\rangle$ with angle π , then at most two direct systolic Weierstrass arcs have an endpoint at c . Thus, there are at most six direct systolic Weierstrass arcs.*

The remainder of this section is devoted to the proof of Theorem 5.4. The proof is a complicated proof by contradiction that involves many cases. We suppose that there exist three direct systolic Weierstrass arcs that end at c . We cut along these arcs and we cut along the two (necessarily direct) minimal arcs that join the remaining two angle π cone points to the angle 4π cone point on $X/\langle\tau\rangle$. The result of these cuts is an annulus with piecewise geodesic boundary that contains the remnants of the cone points. The various cases considered are based on the holonomy of the translation structure of the annulus as well as the relative positions of the cone points on the boundary of the annulus. To obtain a contradiction in each case, we use the fact that the distance between any two cone points can be no less than $\text{sys}(X)/2$.

We now begin the proof of Theorem 5.4.

Proof. Suppose to the contrary that there exist three direct systolic Weierstrass arcs each having c as an endpoint. Let $\theta_1 \leq \theta_2 \leq \theta_3$ denote the angles between the arcs at c . Since c is an angle π cone point, we have $\theta_1 + \theta_2 + \theta_3 = \pi$. Label the arcs α_i ,

$i \in \mathbb{Z}/3\mathbb{Z}$, so that the angle between α_{i-1} and α_i equals θ_i . By Lemma 5.1, the other endpoints of the α_i are all distinct. Label the other endpoint of α_i with c_i . Let c_4 and c_5 denote the two remaining angle π cone points.

The lift, $\tilde{\alpha}_i$, of each α_i to X is a non-separating direct simple closed geodesic on X . The involution preserves $G := \tilde{\alpha}_1 \cup \tilde{\alpha}_2 \cup \tilde{\alpha}_3$ and hence the complement $A := X - G$. We have $\chi(A) = \chi(X) - \chi(G) = 2 - 2 = 0$, and since A contains the fixed points c_4 and c_5 , it follows that A is connected and, moreover, is homeomorphic to an annulus.

Let γ be a shortest geodesic in X that represents the free homotopy class corresponding to a generator of $\pi_1(A) \subset \pi_1(X)$. Because $\theta_i < \pi$ and each $\tilde{\alpha}_i$ is a geodesic, the geometric intersection number of γ and each $\tilde{\alpha}_i$ is zero. In particular, γ can not coincide with some $\tilde{\alpha}_i$ as the intersection number $i(\tilde{\alpha}_i, \tilde{\alpha}_j) = 1$ for $i \neq j$ (see Proposition 5.3). Therefore, $\tilde{\alpha}_i$ and γ are disjoint for each $i \in \mathbb{Z}/3\mathbb{Z}$, and γ lies in A .

In the remainder of the proof, we will consider separately the two cases: (1) the closed geodesic γ is direct and (2) γ passes through an angle 4π cone point.

γ is direct: If γ is direct, then it belongs to a maximal cylinder C . Without loss of generality, γ is the middle geodesic of this cylinder. Since γ is nonseparating, τ preserves C and γ , and in particular, the fixed points c_4 and c_5 lie on γ . To obtain the desired contradiction in this case, it suffices to show that the length of γ is less than $\text{sys}(X)$.

Each component of ∂C consists of a direct geodesic segment β_{\pm} joining an angle 4π cone point c_{\pm}^* to itself. The geometric intersection number of β_{\pm} and each $\tilde{\alpha}_i$ equals zero, and hence β_{\pm} does not intersect any of the $\tilde{\alpha}_i$. Hence the complement $A - C$ consists of two topological annuli K_+ and K_- with $\beta_{\pm} \subset \partial K_{\pm}$. Because τ preserves each maximal cylinder as well as A , we have $\tau(K_{\pm}) = \tau(K_{\mp})$. Thus, we will now limit our attention to only one of the two annuli, $K := K_+$. One boundary component of K is the direct geodesic segment $\beta := \beta_+$ joining an angle 4π cone point, $c^* := c_+^*$, to itself. The other boundary component, β' , of K consists of three geodesic segments $\bar{\alpha}_1$, $\bar{\alpha}_2$, and $\bar{\alpha}_3$ corresponding respectively to $\tilde{\alpha}_1$, $\tilde{\alpha}_2$, and $\tilde{\alpha}_3$. Moreover, the interior angle between $\bar{\alpha}_{i-1}$ and $\bar{\alpha}_i$ is equal to θ_i . See the left hand side of Figure 10.

Since β and γ are parallel geodesics in the same cylinder C , it suffices to show that the length of β is less than $\text{sys}(X)$. Since β is a direct geodesic segment, the length of β equals the length of the holonomy vector associated to β . Since β and β' are homotopic, their holonomy vectors have the same length. Thus, it suffices to show that the length of the holonomy vector associated to β' is less than $\text{sys}(X)$.

Since, by assumption, each $\tilde{\alpha}_i$ is a systole, the length of β' is $b := 3 \cdot \text{sys}(X)$. Let $\beta': [0, b] \rightarrow \partial_{\pm}$ be a parameterization of β' so that $\beta'(0) = \bar{\alpha}_3 \cap \bar{\alpha}_1 = \beta'(1)$. The development, β' , consists of three line segments, each of length $\text{sys}(X)$, joined end to end with consecutive angles θ_2 and θ_3 . See the right hand side of Figure 10.




Figure 10. On the left is the topological annulus K case when the closed geodesic γ is direct. The right side shows the development of $\beta' = \bar{\alpha}_1 \cup \bar{\alpha}_2 \cup \bar{\alpha}_3$.

Since $2\pi/3 \leq \theta_2 + \theta_3 < \pi$ and the three sides of $\bar{\beta}'$ have the same length, an elementary fact from Euclidean geometry applies to give that the distance between $\text{dev}(\beta'(0))$ and $\text{dev}(\beta'(1))$ is less than $\text{sys}(X)$. Thus the holonomy vector of β' has length less than $\text{sys}(X)$ as desired.

γ is indirect: In the remainder of the proof we consider the case in which $\pi_1(A)$ is not generated by a direct simple closed geodesic. In this case, the shortest geodesic γ that generates $\pi_1(A)$ is unique in its homotopy class. In particular, since τ induces a nontrivial automorphism of $\pi_1(A) \cong \mathbb{Z}$, the isometry τ preserves γ and reverses its orientation. It follows that γ is a union of two geodesic segments each joining the two 4π angle cone points, and each segment contains as its midpoint one of the remaining two Weierstrass points. Let σ_+ denote the segment containing c_4 , and let σ_- denote the segment containing c_5 .

The complement of γ consists of two topological annuli K_+ and K_- that are isometric via τ . We limit our attention to one of the annuli, K . One boundary component of K consists of the geodesic segments $\bar{\alpha}_1, \bar{\alpha}_2,$ and $\bar{\alpha}_3$ with the interior angle between $\bar{\alpha}_{i-1}$ and $\bar{\alpha}_i$ equal to θ_i . The other boundary component consists of σ_+ and σ_- . See Figure 11.

Let c_+^* and c_-^* denote the angle 4π cone points. Let θ_{\pm} denote the interior angle between σ_+ and σ_- at c_{\pm}^* . Because τ interchanges the two components of $A - \gamma$, we have $\theta_+ + \theta_- = 4\pi$. Since γ is not direct, there is no direct geodesic segment joining c_4 and c_5 inside K . Indeed, if there were such a segment δ , then $\delta \cup \tau(\delta)$ would be a direct simple closed geodesic that generates $\pi_1(A)$ contradicting our assumption. It follows that $\theta_{\pm} \geq \pi$.




Figure 11. The topological annulus K .

We claim that $\theta_1 < \pi/3$. Indeed if not, then since $\theta_1 + \theta_2 + \theta_3 = \pi$ and $\theta_1 \leq \theta_2 \leq \theta_3$, we would have $\theta_i = \pi/3$ for each i and in particular, the holonomy of $\beta = \tilde{\alpha}_1 \cup \tilde{\alpha}_2 \cup \tilde{\alpha}_3$ would be zero. Thus, since $\sigma_+ \cup \sigma_-$ is homotopic to β , the holonomy of $\sigma_+ \cup \sigma_-$ would be trivial. Since σ_{\pm} is a geodesic segment, the angle at c_{\pm}^* would equal 2π and the lengths of σ_+ and σ_- would be equal. It would follow that the developing map would map \bar{K} onto the an equilateral triangle T having sidelength $\text{sys}(X)$. Moreover, $\text{dev}(\sigma_+) = \text{dev}(\sigma_-)$ would be a segment σ in the interior of T and the restriction of dev to $\bar{K} - (\sigma_+ \cup \sigma_-)$ would be injective. By elementary Euclidean geometry, the distance from each interior point of T to the set of midpoints of the sides of T is less than $\text{sys}(X)/2$. In particular, it would follow that there would be a direct geodesic segment in \bar{K} joining the set $\{c_4, c_5\}$ and $\{c_1, c_2, c_3\}$ having length less than $\text{sys}(X)/2$. This would contradict the definition of $\text{sys}(X)$.

Thus, in the remainder of the proof of Theorem 5.4, we may assume that $\theta_1 < \pi/3$. Our next goal is the show that this implies that there exists a direct geodesic that joins v_1 to one of the two 4π cone points, c_{\pm}^* .

Lemma 5.5. *There exists a (direct) geodesic segment $\delta \subset K$ that joins v_1 to either c_+^* or c_-^* .*

Proof of Lemma 5.5. Let V be the set of points $x \in K$ such that there exists a direct geodesic segment in K joining v_1 to x . By lifting to \tilde{X} and applying the developing map, the set V is mapped injectively onto a subset of the Euclidean sector S of angle θ_1 . In particular, v_1 is mapped to the vertex \bar{v}_1 of S . The bounding rays of S contain the respective images, \bar{c}_1 and \bar{c}_3 , of the points c_1 and c_3 .

Let T be the convex hull of $\{\bar{v}_1, \bar{c}_1, \bar{c}_3\}$ The set T is an isosceles triangle with $|\bar{v}_1\bar{c}_1| = \text{sys}(X)/2 = |\bar{v}_1\bar{c}_3|$, and the angle $\angle \bar{c}_1\bar{v}_1\bar{c}_3$ is less than $\pi/3$. In particular,

the side of T that joins \bar{c}_1 and \bar{c}_3 has length less than $\text{sys}(X)/2$, and the distance from \bar{v}_1 to any other point of T is at most $\text{sys}(X)/2$.

Let $x^* \in \overline{S - V}$ be a point such that $\text{dist}(x^*, v_1)$ equals the distance between \bar{v}_1 and the $\overline{S - V}$. We claim that x^* is the image of an angle 4π cone point, and hence that there exists a direct geodesic joining v_1 and this angle 4π cone point. See Figure 12.




Figure 12. The point x^* in the triangle T .

To verify the claim, we first note that x^* lies in the interior of T . Indeed if it did not, then since the developing map is injective on V , the side of T that joins \bar{c}_1 to \bar{c}_3 would be the image of a direct geodesic segment joining c_1 and c_3 having length less than $\text{sys}(X)/2$. This would contradict the definition of $\text{sys}(X)$.

Because $\theta_1 \leq \theta_2 \leq \theta_3$, the distance between v_1 and $\tilde{\alpha}_2$ is at least $\text{sys}(X)/2$, and hence the point x^* can not belong to $\text{dev}(\tilde{\alpha}_2)$. Thus, x^* is the image of a point in σ_+ or σ_- . Thus to verify the claim, it suffices to show that x^* is not the image of an interior point of σ_{\pm} .

Suppose to the contrary that x^* were the image of an interior point σ_{\pm} . Then the segment $\text{dev}(\sigma_{\pm})$ would lie in $\overline{S - V}$, and hence by the definition of x^* , the segment $\text{dev}(\sigma_{\pm})$ would be perpendicular to the segment joining v_1 and x^* , and hence parallel to the side of T that opposes v_1 . The segment $\text{dev}(\sigma_{\pm})$ does not intersect either $\text{dev}(\tilde{\alpha}_1)$ or $\text{dev}(\tilde{\alpha}_3)$, and hence the midpoint of $\text{dev}(\sigma_{\pm})$ would lie in T . The segment joining the midpoint and \bar{v}_1 corresponds to a direct geodesic segment joining v_1 to either c_4 or c_5 . Since this segment has length less than $\text{sys}(X)/2$, we would obtain a contradiction.

Thus, x^* is the image of either c_-^* or c_+^* . □

By relabeling if necessary, we may assume that $\text{dev}(c_+^*) = x^*$. Let δ denote the direct geodesic joining v_1 and c_+^* .

Let P denote the metric completion of $K - \delta$. The metric space P is a topological disk bounded by seven geodesic segments. The ‘‘polygon’’ P has seven vertices: the points v_2 and v_3 , two vertices, p_+ and p_- , corresponding to c_+^* , one vertex, q , corresponding to c_-^* , and two vertices, v_+ and v_- , corresponding to v_1 . See Figure 13.




Figure 13. The polygon P .

Continuing with our proof of Theorem 5.4, our next goal is to prove that P may be regarded as a polygon in the plane. In particular, we wish to show that the restriction of the developing map to P is injective.⁵

First, note that since the geodesics $\tilde{\alpha}_i$ all have the same length and the sum of the angles $\theta_2 + \theta_3$ is strictly larger than $\pi/3$, the set $\text{dev}(\tilde{\alpha}_1 \cup \tilde{\alpha}_2 \cup \tilde{\alpha}_3)$ is a simple piecewise linear arc in the plane with endpoints \bar{v}_+ and \bar{v}_- corresponding to v_+ and v_- respectively. In particular, the convex hull of $\{\bar{v}_+, \bar{v}_2, \bar{v}_3, \bar{v}_-\}$ is a quadrilateral Q , and the line segments $\bar{\alpha}_i := \text{dev}(\tilde{\alpha}_i)$ constitute three of the sides of Q .

Let $\delta_{\pm} \subset P$ be the segment that joins v_{\pm} and p_{\pm} , and let $\bar{\delta}_{\pm} := \text{dev}(\delta_{\pm})$. Since $\theta_1 < \pi/3$, the angle between $\tilde{\alpha}_1$ and δ_+ and the angle between $\tilde{\alpha}_3$ and δ_- are both less than $\pi/3$. It follows that the segment $\bar{\delta}_{\pm}$ lies in Q and that the point p_{\pm} lies in the interior of Q .

Let θ_z denote the interior angle at a vertex z of P .

Lemma 5.6. *We have $\pi < \theta_q < 2\pi$, $\theta_{p_{\pm}} < 2\pi$, and $\theta_{p_+} + \theta_q + \theta_{p_-} = 4\pi$.*

⁵One may regard P as a subset of \tilde{X} by lifting its interior and then taking the closure.

Proof. Suppose to the contrary that $\theta_q \leq \pi$. Since the angles $\theta_{v_+}, \theta_{v_-}, \theta_2, \theta_3$ are all less than π , the shortest path from p_+ to p_- and the shortest path from c_4 to c_5 are both direct. Because $\theta_2 + \theta_3 < \pi$, the Euclidean distance $|\bar{p}_+ \bar{p}_-| = |p_+ p_-|$ is strictly less than $|\bar{v}_2 \bar{v}_3| = \text{sys}(X)$. Since c_4 and c_5 are midpoints, it follows that $|c_4 c_5| = |\bar{c}_4 \bar{c}_5| < \text{sys}(X)/2$. This is a contradiction.

It follows that the point $\bar{q} := \text{dev}(q)$ lies in the closed half-plane bounded by the line through \bar{p}_+ and \bar{p}_- that does not contain \bar{v}_+ or \bar{v}_- . Hence, the angle θ_{p_\pm} is at most the angle between $\bar{\delta}_\pm$ and $\overline{p_+ p_-}$, and this is less than 2π . The angle θ_q equals $2\pi - \psi$ where ψ the angle opposite the segment $\bar{p}_+ \bar{p}_-$ in the (perhaps degenerate) triangle $\bar{p}_+ \bar{q} \bar{p}_-$.

As discussed in the analysis of Figure 11, we have $\theta_+ + \theta_- = 4\pi$. It follows that $\theta_{p_+} + \theta_q + \theta_{p_-} = 4\pi$. □

Proposition 5.7. *The metric space P is isometric to a simply connected polygon in the Euclidean plane.*

Proof. It suffices to show that the developing map is injective. Let $x, x' \in P$. Since P is path connected and compact, there exists a minimal geodesic arc η that joins x to x' . To prove the claim it suffices to show that the endpoints of $\text{dev} \circ \eta$ are distinct. If η is a direct geodesic segment, then $\text{dev} \circ \eta$ is a single Euclidean line segment and so $\text{dev}(x) \neq \text{dev}(x')$. If η is not direct, then η is a concatenation of a finite number direct geodesic segments, $\gamma_1, \dots, \gamma_n$, such that $\gamma_i \cap \gamma_{i+1}$ is a vertex v_i and the angle ψ_i between γ_i and γ_{i+1} satisfies $\pi \leq \psi_i \leq \theta_v$ where θ_v is the angle between boundary segments at v . Since the angles at v_\pm, v_2, v_3 are less than π , the minimal geodesic η can only pass through the vertices p_+, p_- , or q , and if η does pass through one of these vertices, then it passes through the vertex at most once.

Each of the angles θ_q, θ_{p_\pm} is positive, and so if η passes through exactly one of the points q, p_\pm , then the path $\text{dev} \circ \eta$ is a simple arc. In particular, the endpoints $\text{dev}(x)$ and $\text{dev}(x')$ are distinct.

Suppose that η passes through exactly two vertices say $v_1, v_2 \in \{p_+, p_-, q\}$. Lemma 5.6 implies that $\psi_1 + \psi_2 < 3\pi$. We also have $\psi_i \geq \pi$. An elementary argument in Euclidean geometry shows that $\text{dev}(\eta)$ is a simple arc.

Finally, suppose that η passes through each of p_+, p_-, q .

Hence $\psi_1 + \psi_2 + \psi_3 \leq 4\pi$. We also have $\psi_i \geq \pi$. An elementary Euclidean geometry argument shows that $\text{dev} \circ \eta$ is a simple arc. □

In what follows, we will identify the polygon P with its image in \mathbb{C} . See Figure 13.

Lemma 5.8. *The shortest geodesic joining c_1 (resp. c_3) to p_+ (resp. p_-) is direct.*

Proof. Recall the triangle T described in Figure 12. The point p_+ corresponds to $x^* = c_+^*$, and so if the shortest geodesic joining c_1 and p_+ were not direct, then the shortest geodesic in X joining c_1 to c_+^* would also pass through c_-^* . Hence c_-^* would also belong to the triangle T described above, and so either the image of σ_+

or the image of σ_- would lie in T . But then the midpoint c_4 of σ_+ or the midpoint c_5 of σ_- would belong to T . Hence $|v_1c_4|$ or $|v_1c_5|$ would be less than $\text{sys}(X)/2$, a contradiction.

A similar argument shows that the shortest geodesic from c_3 to p_- is direct. \square

Because x^* belongs to the interior of T , we have $\angle \bar{v}_1\bar{c}_1x^* < \angle \bar{v}_1\bar{c}_1\bar{c}_3$. Since T is isosceles, we have $2 \cdot \angle \bar{v}_1\bar{c}_1\bar{c}_3 + \theta_1 = \pi$. Thus, it follows that

$$\angle v_+c_1p_+ < \frac{\pi - \theta_1}{2}. \tag{5.1}$$

(A similar argument shows that $\angle v_-c_3p_- < (\pi - \theta_1)/2$.)

We will use (5.1) to prove the following

Lemma 5.9. *The minimal geodesic joining c_3 to c_5 is direct.*

Proof. Let ℓ_1 be the line parallel to $\overline{p_+p_-}$ that passes through c_3 , and let ℓ_2 be the line parallel to $\overline{v_-v_3}$ that passes through v_+ . Since $\theta_2 < \pi/2$ and $|v_+v_2| = |v_2v_3|$, the points v_2 and v_3 lie in distinct components of $\mathbb{C} - \ell_2$. Because p_- lies in the component of $\mathbb{C} - \overleftrightarrow{v_-v_3}$ that contains v_2 and $\overline{p_+p_-}$ is a translate of $\overline{v_+v_-}$, the point p_+ lies in the component H_2 of $\mathbb{C} - \ell_2$ that contains v_2 . See Figure 14.




Figure 14. The segment that joins c_3 to c_5 belongs to P .

Let x be the point of intersection of ℓ_1 and ℓ_2 , and let ℓ_3 be the line passing through c_1 and x . Since $|v_+x| = |v_-c_3| = \text{sys}(X)/2 = |v_+c_1|$, the triangle Δc_1xv_+ is

analogous to that used to derive (5.1) gives the inequality $\angle p_-c_3v_- < (\pi - \theta)/2$. Therefore, p_- lies in the component H_3 of $\mathbb{C} - \ell_3$ that contains v_- .

If we let ℓ'_3 denote the line parallel to ℓ_3 that passes through c_1 , then, since $\overline{p_+p_-}$ is a translate of $\overline{c_1x}$, the point p_+ lies in the component H'_3 of $\mathbb{C} - \ell'_3$ that contains v_- . Thus, to prove that there is a direct segment from c_1 to c_4 , it suffices to show that q lies in $\mathbb{C} - H'_3$ for then $\angle c_1p_+c_4 < \pi$.

Let m be the midpoint of $\overline{c_1x}$, and let ℓ''_3 be the line parallel to ℓ_3 that passes through m . To show that $q \in \mathbb{C} - H'_3$, it suffices to show that c_5 lies in the closure of the component H''_3 of $\mathbb{C} - \ell''_3$ that contains v_2 . Indeed, c_5 is the midpoint of $\overline{p_-q}$ and we know that p_- lies in H_3 .

Since there is a direct segment joining c_5 to c_3 , the point c_5 lies outside the ball B of radius $\text{sys}(X)/2$ with center at c_3 . We also know that c_5 lies in Q , the convex hull of $\{v_+, v_2, v_3, v_-\}$, and that c_5 belongs to $\mathbb{C} - H_1$. An elementary geometric argument shows that $(Q - B) \cap (\mathbb{C} - H_1)$ lies in H''_3 . Thus, $c_5 \in H''_3$ and there exists a direct segment joining c_1 to c_4 as desired. \square

Given that there are direct segments between c_1 and c_4 and between c_3 and c_5 , we will now derive a contradiction and thus complete the proof of Theorem 5.4 as follows.

Let ℓ_+ be the line that passes through v_+ and c_4 , let ℓ_- denote the line that passes through v_- and c_5 , and let x be the intersection of ℓ_+ and ℓ_- . See Figure 16. Because $\overline{v_+p_+}$ and $\overline{v_-p_-}$ are parallel and of the same length and the points c_4, c_5 are the respective midpoints of $\overline{p_+q}, \overline{p_-q}$, the points c_4, c_5 are also the respective midpoints of $\overline{v_+x}, \overline{v_-x}$. Since c_1 is the midpoint of $\overline{v_+v_2}$, we have $|xv_2| = 2 \cdot |c_1c_4|$. Since the geodesic from c_1 to c_4 is direct, we have $|c_1c_4| \geq \text{sys}(X)/2$ and hence $|xv_2| \geq \text{sys}(X)$. Similarly, since the geodesic from c_3 to c_5 is direct, we find that $|xv_3| \geq \text{sys}(X)$.

In other words, if we let B_+ (resp. B_-) be the ball of radius $\text{sys}(X)$ about v_2 (resp. v_3), then x lies outside $B_+ \cup B_-$. Since $\{v_+, v_2, v_3, v_-\}$ is contained $\overline{B_+} \cup \overline{B_-}$, the polygon P is contained in the convex hull of $\overline{B_+} \cup \overline{B_-}$.

Let ℓ_{23} denote the line passing through v_2 and v_3 , and let $y: \mathbb{C} \rightarrow \mathbb{R}$ denote the real affine 1-form such that $|y(z)| = \text{dist}(z, \ell_{23})$ and such that $y(v_+) > 0$. Because $\theta_2 \leq \theta_3 < \pi$, we have that $y(z) \geq 0$ for each $z \in P$.

Note that $y(x) < y(q)$. Indeed, since $\angle c_1v_+p_+ < \theta_1$ and $\theta_1 + \theta_2 < \pi$, it follows that $y(v_+) > y(p_+)$. The segment \overline{xq} is the reflection of $\overline{v_+p_+}$ about the point c_4 , and hence $y(x) < y(q)$.

Let x' be the intersection point of ℓ_{23} and the line passing through x and q . The point x' lies in the line segment $\overline{v_2v_3}$. Indeed, because $\theta_2 + \theta_3 < \pi$, the line through v_+ and v_2 and the line through v_- and v_3 intersect at a unique point z , and moreover, the polygon P lies in the convex hull T' of $\{z, v_2, v_3\}$. Because $\overline{p_-v_-}$ and $\overline{p_+v_+}$ are parallel, p_+ and p_- lie in T' , v_+ lies in $\overline{zv_2}$, and v_- lies in $\overline{zv_3}$, any line parallel




Figure 16. The points c_4 and c_5 are the respective midpoints of $\overline{v_+x}$ and $\overline{v_-x}$. Thus, $|v_2x| = 2|c_1c_4|$ and $|v_3x| = 2|c_3c_5|$.

to p_+v_+ that intersects T' must intersect ℓ_{23} at a point in the segment $\overline{v_2v_3}$. In particular, the point x' lies in $\overline{v_2v_3}$.

We claim that $y(x) > 0$. Indeed, suppose not. Then x' would lie in the segment \overline{xq} . Thus, $|x'x| \leq |xq| = |v_{\pm}p_{\pm}| < \text{sys}(X)/2$, and hence x would belong to the set, A , of points whose distance from $\overline{v_2v_3}$ is at most $\text{sys}(X)/2$. Elementary geometry shows that $A \subset B \cup B_+$, but x lies in the complement of $B_- \cup B_+$, a contradiction.

Let Q be the convex hull of $\{v_+, v_2, v_3, v_-\}$. We have $P \subset Q$ and hence $q \in Q$. Since $0 < y(x) < y(q)$ and the line through x and q meets $\ell_{23} = \ker(y)$ at $x' \in \overline{v_2v_3}$, the point x also belongs to Q . The set Q is contained in the convex hull of $\overline{B_+} \cup \overline{B_-}$. Therefore, x lies inside the convex hull of $\overline{B_+} \cup \overline{B_-}$ and outside $B_+ \cup B_-$. Since $x' \in \overline{v_2v_3}$ it follows that $\pi/4 \leq \angle v_2x'x \leq 3\pi/4$, and, therefore, since $y(q) > y(x)$, we find that q is also outside $B_+ \cup B_-$. See Figure 17.

Since x and q both lie inside the convex hull of $B_+ \cup B_-$ but outside $B_+ \cup B_-$, we have $y(q) - y(x) < (1 - \sqrt{3}/2) \cdot \text{sys}(X)$. Since $\pi/4 \leq \angle v_2x'x \leq 3\pi/4$, we have $|xq| \leq \sqrt{2} \cdot |y(q) - y(x)|$ and hence

$$|v_{\pm}p_{\pm}| \leq \sqrt{2} \cdot \left(1 - \frac{\sqrt{3}}{2}\right) \cdot \text{sys}(X) < \frac{\text{sys}(X)}{4}. \tag{5.2}$$




Figure 17. The distances $|xv_2|$ and $|xv_3|$ are at least $\text{sys}(X)$, and $y(x) > 0$.

Let ℓ_p be the line through p_+ and p_- and let ℓ_v be the line through v_+ and v_- . Let ℓ_{\pm}^* denote the line passing through v_{\pm} and p_{\pm} . By Lemma 5.6, the interior angle θ_- at $q \in P$ is greater than π , and hence the point q lies in the component of $\mathbb{C} - \ell_p$ that contains the segment $\overline{v_2v_3}$, and hence q lies in the component of $\mathbb{C} - \ell_v$ that contains $\overline{v_2v_3}$. Since q lies outside $B_+ \cup B_-$, it follows that q lies in the bounded component of $\mathbb{C} - (\ell_{23} \cup \ell_v \cup \ell_+^* \cup \ell_-^*)$.

Let ℓ_q be the line through q that is parallel to ℓ_v . Let A be the parallelogram that is the bounded component of $\mathbb{C} - (\ell_q \cup \ell_v \cup \ell_+ \cup \ell_-)$. Let b_{\pm} be the intersection of ℓ_{\pm} and ℓ_q . Then A is the convex hull of $\{b_+, b_-, v_+, v_-\}$. Because q lies in the component of $\mathbb{C} - \ell_p$ that contains $\overline{x_2x_3}$, the point p_{\pm} lies in $\overline{v_{\pm}b_{\pm}}$.

The line ℓ^* through x and q is parallel to the sides corresponding to ℓ_+ and ℓ_- . Let x'' be the intersection of ℓ^* with the side $\overline{v_+v_-}$ of A corresponding to ℓ_v . Since $v_{\pm} \in B_+ \cup B_-$, the point x'' lies in the convex hull of $B_+ \cup B_-$. By applying the argument that led to (5.2) to this situation, we find that $|x''q| < \text{sys}(X)/4$.

We have $|b_+b_-| = |v_+v_-| < \text{sys}(X)$ and hence either $|b_+q| < \text{sys}(X)/2$ or $|b_-q| < \text{sys}(X)/2$. Suppose that $|b_+q| < \text{sys}(X)/4$. The midpoint, c_4 , of $\overline{p_+q}$ lies in A . Let a_+ be the point of intersection of ℓ_+ and the line through c_4 that is parallel to ℓ_q . Then a_+ lies in the segment $\overline{p_+b_+}$.

By the triangle inequality, we have

$$|v_+p_+| + |p_+c_4| \leq |v_+a_+| + |a_+c_4| < \frac{\text{sys}(X)}{4} + \frac{\text{sys}(X)}{4} = \frac{\text{sys}(X)}{2}$$

But v_+ and c_4 are both Weierstrass points, and hence we would have a curve of length less than $\text{sys}(X)/2$. A similar contradiction is obtained in the case when $|b_-q| < \text{sys}(X)/2$. \square

The following is immediate.

Corollary 5.11. *There are at most six homotopy classes of nonseparating systoles.*

6. Indirect Weierstrass arcs

The angle 4π cone point c^* divides each systolic indirect Weierstrass arc on $X/\langle\tau\rangle$ into two subarcs. We will call each such subarc a *prong*.

Let C_ϵ be the set of points at distance ϵ from c^* . For ϵ sufficiently small, the set C_ϵ is a topological circle, and each prong intersects C_ϵ exactly once. Thus, the prongs divide the circle C_ϵ into disjoint arcs. Two prongs are said to be *adjacent* if they are joined by one of these arcs, and the *angle* between two adjacent arcs is the arclength divided by ϵ .

If a systolic indirect Weierstrass arc is the union of two adjacent prongs then the angle between the two prongs must be at least π . Indeed, otherwise one can shorten the arc by perturbing it near c^* .

The sum of the lengths of any two prongs is at least $\text{sys}(X)/2$. Indeed, otherwise the concatenation of the two prongs would lift to a geodesic loop on X that would have length less than $\text{sys}(X)$. On the other hand, for each prong, there is another prong so that the sum of the lengths of the two prongs equals $\text{sys}(X)/2$.

In particular, the minimum, ℓ , of the lengths of the prongs is at most $\text{sys}(X)/4$. If $\ell < \text{sys}(X)/4$, then there is a unique shortest prong and the remaining prongs have length $\text{sys}(X)/2 - \ell$.

If $\ell = \text{sys}(X)/4$, then each prong has length $\text{sys}(X)/4$, and each pair of adjacent prongs determines a systolic Weierstrass arc. Since the angle between each adjacent pair is at least π and c^* has total angle 4π , there are at most four adjacent pairs and if there are exactly four pairs, then each angle equals π . In sum, we have

Proposition 6.1. *If all of the prongs have the same length, then the number of prongs is at most four. If there are exactly four such prongs, then the angle between each pair of adjacent prongs is exactly π .*

We will show below that if one of the prongs is shorter than the others then there are at most five prongs. To do this we will use the following lemma.

Lemma 6.2. *Two distinct prongs can not end at the same angle π cone point, c' .*

Proof. Suppose not. Then the concatenation, α , of the two prongs would be a closed curve that divides the sphere $X/\langle\tau\rangle$ into two discs. Since there are five other cone points, one of the discs, D , would contain at most two cone points. There are no Euclidean bigons and so D would have to contain at least one cone point.

If D were to contain two angle π cone points, then α would be homotopic to the concatenation of the two oriented minimal arcs joining the two cone points. The length of the unoriented minimal arc is at least $\text{sys}(X)/2$, and hence, since the length of each prong is less than $\text{sys}(X)/2$, we would have a contradiction.

If D were to contain one angle π cone point c , then α would be homotopic to the concatenation of the two oriented minimal arcs joining c and c' . We would then arrive at a contradiction as in the case of two cone points. \square

Since there are exactly six Weierstrass points, Lemma 6.2 implies that there are at most six prongs. In fact, we have the following.

Proposition 6.3. *There are at most five prongs.*

Proof. Suppose to the contrary that there are six prongs. Let e_1 denote the unique shortest prong, let ℓ be its length, and let c_1 denote its endpoint. Let e_1, \dots, e_6 be a cyclic ordering of the remaining prongs, let $L = \text{sys}(X)/2 - \ell$ denote their common length, and let c_2, \dots, c_6 denote their respective endpoints.

Since $\ell(e_1 + e_2) = \text{sys}(X)/2 = \ell(e_1 + e_6)$, the angles $\angle c_1 c^* c_2$ and $\angle c_1 c^* c_6$ are each at least π . (Otherwise, by perturbation near the 4π cone point we could construct a direct Weierstrass arc with length less than $\text{sys}(X)/2$.) Each of the other four angles between adjacent prongs is greater than $\pi/3$. Indeed, otherwise, since $L < \text{sys}(X)/2$, we would have a segment joining two angle π cone points having length less than $\text{sys}(X)/2$ which contradicts the definition of systole. Since $\angle c_1 c^* c_2 + \angle c_1 c^* c_6 \geq 2\pi$ it follows that each of these four angles is less than π . Moreover, since the angle at c^* equals 4π , the sum $\angle c_1 c^* c_2 + \angle c_1 c^* c_6 < 8\pi/3 < 3\pi$ and individually $\angle c_1 c^* c_2 < 5\pi/3$ and $\angle c_1 c^* c_6 < 5\pi/3$.

By cutting along the prongs and taking the length space completion, we obtain a closed topological disc D whose boundary consists of a topological disc bounded by six geodesic segments. The midpoint of each segment corresponds to an end point of a prong. The developing map provides an immersion of D into the Euclidean plane. Since $\angle c_k c^* c_{k+1} + \angle c_1 c^* c_2 < 3\pi$ and $\angle c_i c^* c_{i+1} < \pi$ for $i = 2, \dots, 5$, this immersion is an embedding. In other words, we may regard D as Euclidean hexagon.

Let v_i denote the vertex of D corresponding to c^* that lies between c_{i-1} and c_i . The length of the side $\overline{v_1 v_2}$ is 2ℓ , and the common length of the other sides is $2L$.

From above, the interior angles at v_1 and v_6 are between π and $5\pi/3$, and the angles at the other four vertices lie between $\pi/3$ and π . Without loss of generality, $c_1 = (0, 0)$, $v_1 = (\ell, 0)$, $v_6 = (-\ell, 0)$ and an H -neighborhood of c_1 lies in the upper half plane (see Figure 18).




Figure 18. The points $v_i, i = 1, \dots, 6$.

Since the angle at v_2 (resp. v_5) is greater than $\pi/3$, and the edges $\overline{v_1v_2}$ and $\overline{v_2v_3}$ (resp. $\overline{v_4v_5}$ and $\overline{v_5v_6}$) have length $2L$, the vertex v_3 (resp. v_4) lies outside the ball of radius $2L$ centered at v_1 (resp. v_6). It follows that if both v_4 and v_3 both lie in the lower half plane then the shortest arc in H that joins v_4 to v_3 has distance at least $2L + 2\ell$. This contradicts the equality $|v_3v_4| = 2L$.

Since the angle at v_1 (resp. v_6) is at least π and the angle at v_2 (resp. v_5) is greater than $\pi/3$, if v_3 (resp. v_4) lies in the upper half plane, then v_3 (resp. v_4) lies in the half plane $V_+ = \{(x_1, x_2) \mid x_1 > \ell + L\}$ (resp. $V_- = \{(x_1, x_2) \mid x_1 < -\ell - L\}$). Since the distance between U_+ and U_- equals $2L + 2\ell$, if v_3 and v_4 both lie in U , then we contradict $|v_3v_4| = 2L$.

If v_3 lies in the upper half plane and that v_4 lies in the lower half plane but not in U_- , then v_4 lies in the half plane that is bounded by the line through v_3 and v_6 and contains v_1 . In particular, the shortest path in D between v_3 and v_4 passes through v_6 . But the distance from v_6 to U_+ is equal to $2L + \ell$, and the distance from v_6 to v_4 is greater than $2L$. Thus, we contradict $|v_3v_4| = 2L$.

A symmetric argument rules out the remaining case in which the rôles of v_3 and v_4 are reversed. \square

Theorem 6.4. *There are at most six systolic indirect Weierstrass arcs. Equality occurs if and only if there are exactly four prongs and these four prongs have the same length.*

Proof. If the prongs are not all of the same length, then one prong has length less than $\text{sys}(X)/4$ and hence the others have length greater than $\text{sys}(X)/4$. Therefore, concatenations of none of the others constitute a systolic Weierstrass arcs. By Proposition 6.3, there are at most five prongs and hence at most five systolic indirect Weierstrass arcs.

If the prongs all have the same length — namely $\text{sys}(X)/4$ — then by Proposition 6.1 there are at $n \leq 4$ prongs. Each concatenation of a pair prongs constitutes a systolic Weierstrass arc, and so there are exactly $n \cdot (n - 1)/2$ prongs and hence at most six. Six occurs if and only if $n = 4$. \square

7. A separating systole

In this section we wish to prove the following:

Theorem 7.1. *If X has a separating systole α , then X has at most nine homotopy classes of closed curves with systolic representatives.*

We will use the argument explained in Remark 5.2 in the three lemmas that follow. We first observe:

Lemma 7.2. *X has at most one separating systole.*

Proof. Suppose there are two separating systoles. Each angle of intersection between the two curves must be at least π , otherwise one can find a shorter non-homotopically trivial curve by a cut-and-paste argument. Hence intersection points between the systoles occur at the 4π cone points. But as any two separating curves intersect at least 4 times, this is impossible because there are only two angle 4π cone points. \square

Lemma 7.3. *If α is a separating systole and γ is a direct systolic Weierstrass arc, then γ does not intersect the projection of α to $X/\langle\tau\rangle$.*

Proof. Suppose not. The lift, $\tilde{\gamma}$, of γ to X is a systole that does not pass through an angle 4π cone point. Since α is separating, the curve $\tilde{\gamma}$ intersects α at least twice. Let p_- and p_+ be two of the intersection points. The points p_+ and p_- divides α (resp. $\tilde{\gamma}$) into a pair of arcs. One of the arcs, α_- (resp. $\tilde{\gamma}_-$), has length at most $\text{sys}(X)/2$. By concatenating α_- and $\tilde{\gamma}_-$, we obtain a non null homotopic closed curve β of length at most $\text{sys}(X)$. Since each intersection point is not a cone point and the geodesics are distinct, the angle at each intersection point $\tilde{\gamma}_-$ is less than π . Thus, a perturbation of β near an intersection point produces a curve homotopic to β that has shorter length, a contradiction. \square

Lemma 7.4. *If X has a separating systole α , then each prong of X has length equal to $\text{sys}(X)/4$. Moreover, the angle between the projection $p(\alpha)$ and each prong is at least π .*

Proof. If not, then by the discussion at the beginning of §6, there would exist a prong of length strictly less than $\text{sys}(X)/4$. The preimage of a prong under p is an arc γ of length $\text{sys}(X)/2$ that joins one angle 4π cone point c_-^* to the other angle 4π cone point c_+^* . By Corollary 4.4, the separating systole α passes through both c_-^* and c_+^* , and the complement $\alpha \setminus \{c_-^*, c_+^*\}$ consists of two arcs α_+ and α_- each of

length $\text{sys}(X)/2$. By concatenating α_{\pm} with γ we would obtain a non-null homotopic closed curve having length less than $\text{sys}(X)$, a contradiction.

If the angle between the prong and $p(\alpha)$ were less than π , then one could perturb the concatenation of α_{\pm} and γ to obtain a non-null homotopic closed curve whose length would be less than $\text{sys}(X)/2$, a contradiction. \square

Proof of Theorem 7.1. Let α denote the separating systole to $X/\langle\tau\rangle$ which is unique by Lemma 7.2. By Lemma 7.4, each prong has length equal to $\text{sys}(X)/4$ and the angle between $p(\alpha)$ and each prong is at least π . Thus, since the total angle at c^* is 4π , there are at most two prongs. Hence there are at most two indirect systolic Weierstrass arcs.

By Theorem 5.4, there are at most six direct systolic Weierstrass arcs. Thus, by Proposition 4.5 and the discussion at the beginning of §5, there are at most eight homotopy classes of non-separating closed curves that have systolic representatives. Since α is the unique separating systole, the claim is proven. \square

As a corollary of the proof of Theorem 7.1 and Lemma 7.4, we have the following.

Corollary 7.5. *If X has a separating systole, then X has either no prongs or exactly two prongs of the same length.*

8. Crossing systoles

In this section we prove the following:

Theorem 8.1. *Suppose that $X/\langle\tau\rangle$ has exactly four prongs and each of these prongs has length equal to $\text{sys}(X)/4$. Then at most ten homotopy classes of closed curves are represented by systoles. Moreover, if X has exactly ten homotopy classes of systoles, then X is homothetic to the surface described in Figure 1, and otherwise X has at most eight homotopy classes of systoles.*

Proof. By Corollary 7.5, the surface X has no separating systole. By Theorem 6.4, there are exactly six indirect systolic Weierstrass arcs. Thus, by Proposition 4.5 and the discussion at the beginning of §5, to prove the first claim it suffices to show that there are at most four direct systolic Weierstrass arcs.

By Proposition 6.1, the angle between adjacent prongs equals π . Thus, by cutting along the four prongs we obtain a topological disc D bounded by a geodesic β with no corners. The geodesic β has length $8 \cdot (\text{sys}(X)/4) = 2 \cdot \text{sys}(X)$ and contains one point corresponding to each of the angle π cone points that are endpoints of the four prongs. Label those cone points in cyclic order c_1, c_2, c_3 , and c_4 . For each i , there is a unique point c_i^* on β lying between c_i and c_{i+1} that corresponds to c^* . The distances satisfy $\text{dist}(c_i, c_i^*) = \text{sys}(X)/4 = \text{dist}(c_i^*, c_{i+1})$. The interior angle at each c_i, c_i^* is π .

The two remaining angle π cone points, c_5 and c_6 , lie in the interior of the disc D . Because β is a geodesic (without corners), the disk is geodesically convex, and there exists a direct Weierstrass arc γ joining c_5 and c_6 . By cutting along γ we obtain a topological annulus A with geodesic boundary components β and β' . Since X is a translation surface, A is a Euclidean cylinder isometric to $[0, h] \times (\mathbb{R}/\ell \cdot \mathbb{Z})$ where $\ell = 2 \cdot \text{sys}(X)$ is the common length of β and β' .

The length of γ equals $(1/2) \cdot \ell$, and hence γ is not systolic. The distance between c_5 (resp. c_6) and $\{c_1, c_2, c_3, c_4\}$ is at least $\text{sys}(X)/2$. It follows that the height h of the cylinder A is at least $(\sqrt{3}/4) \cdot \text{sys}(X)$. As a consequence, there does not exist a direct systolic Weierstrass arc joining two distinct points in $\{c_1, c_2, c_3, c_4\}$.

In sum, if δ is a direct systolic Weierstrass arc, then δ joins a point in $\{c_5, c_6\}$ to a point in $\{c_1, c_2, c_3, c_4\}$. Since A is a Euclidean annulus, there are at most two direct systolic Weierstrass arcs joining c_5 (resp. c_6) to $\{c_1, c_2, c_3, c_4\}$, and hence at most ten systolic Weierstrass arcs in total.

Moreover, since the points $\{c_1, c_2, c_3, c_4\}$ are evenly spaced around β , and the points $\{c_5, c_6\}$ are evenly spaced about β' , there are exactly four systolic arcs only if the respective shortest segments, σ_5 and σ_6 , joining c_5 and c_6 to β bisect arcs joining successive points in $\{c_1, c_2, c_3, c_4\}$, that is, only if σ_5 and σ_6 have endpoints in $\{c_1^*, c_2^*, c_3^*, c_4^*\}$. In this case, $h = (\sqrt{3}/4) \cdot \text{sys}(X)$. It follows that X is homothetic to the surface described in Figure 1.

Finally, if there is only one direct systolic Weierstrass arc joining c_5 (resp. c_6) to $\{c_1, c_2, c_3, c_4\}$, then there is only one direct systolic Weierstrass arc joining c_6 (resp. c_5). Hence, if X is not homothetic to the surface described in Figure 1, then X has at most eight homotopy classes of simple closed curves with systolic representatives. \square

9. One short prong

In this section we prove the following:

Theorem 9.1. *If $X/\langle\tau\rangle$ has one short prong, then X has at most nine homotopy classes of closed curves that are represented by systoles.*

Proof. By Corollary 7.5, the surface X has no separating systole. By Proposition 6.3, there are at most five prongs, and so by assumption there is one prong of length $\ell < \text{sys}(X)/4$ and four prongs of length $L = \text{sys}(X)/2 - \ell$. Thus, there are at most four indirect systolic Weierstrass arcs. Thus, it suffices to show that X has at most five direct systolic Weierstrass arcs.

By cutting $X/\langle\tau\rangle$ along the five prongs, we obtain a topological disc D with one angle π cone point in the interior. The boundary consists of five geodesic arcs each of whose endpoints — vertices — corresponds to the angle 4π cone point. The midpoint of each arc corresponds to an angle π cone point on $X/\langle\tau\rangle$. Choose an

orientation of the boundary, and let c_1^* and c_2^* denote the endpoints of the oriented arc that corresponding to the short prong. Label the other vertices c_3^* , c_4^* , and c_5^* according to the orientation. Denote by c_i the midpoint of the arc with endpoints c_i^* and c_{i+1}^* . There remains one angle π cone point, c_6 , that belongs to the interior of D .

By Theorem 5.4, for each angle π cone point c_i , there are at most two direct systolic Weierstrass arcs ending at c_i . Thus, to prove the claim, it suffices to show that c_1 is the endpoint of at most one direct systolic Weierstrass arc. We will show that if c_1 is the endpoint of a direct systolic Weierstrass arc, then the other endpoint must be c_6 .

Since systolic Weierstrass arcs can not intersect except at a cone point, a direct Weierstrass arc joining c_1 to another angle π cone point can not pass through the boundary of D . In particular, if α is a direct Weierstrass arc joining c_1 to either c_2 , c_3 , c_4 , or c_5 , then the complement of α consists of two disks, one that contains c_6 and one that does not.

Suppose that α is a direct geodesic segment that joins c_1 and c_2 . Consider the component, D' , of $D \setminus \alpha$, containing c_1^* . If D' does not contains c_6 , then D' is a flat surface bounded by three geodesic segments. Since the angle at c_1^* is at least π , the Gauss–Bonnet formula implies that the angles at c_1 and c_2 are both zero, and hence α is not direct.

If D' contains c_6 , then by cutting D' along the geodesic segment joining c_6 and c_1^* we obtain a quadrilateral Q with a side corresponding to α . The endpoints of α correspond to c_1 and c_2 . Let x_- and x_+ denote the vertices of Q distinguished by $|x_-c_1| = \ell$ and $|x_+c_2| = L$. If α is systolic, then, by the triangle inequality, $|c_1x_+| \leq L + \text{sys}(X)/2$ with equality if and only if c_1 , c_2 and x_+ are colinear. The midpoint of $\overline{x_-x_+}$ is c_6 , and thus by the triangle inequality

$$|c_1c_6| \leq \frac{|c_1x_-|}{2} + \frac{|c_1x_+|}{2} \leq \ell + L = \frac{\text{sys}(X)}{2}$$

with equality c_1 , c_6 , and c_2 are colinear. Thus, either $|c_1c_6| < \text{sys}(X)/2$ or $|c_2c_6| < \text{sys}(X)/2$, a contradiction. Therefore, there is no direct systolic Weierstrass arc joining c_1 and c_2 . Similarly, there is no direct systolic Weierstrass arc joining c_1 and c_5 .

Suppose that α is a direct geodesic segment that joins c_1 to c_3 . Let D' denote the component of $X \setminus \alpha$ that contains c_2 . If D' does not contain c_6 , then D' is a quadrilateral with vertices c_1 , c_1^* , c_2^* , and c_3 . Since $|c_2c_2^*| = L = |c_2^*c_3|$, the angle $\angle c_3c_2c_2^*$ is less than $\pi/2$, and thus $\angle c_1^*c_2c_3 > \pi/2$. Therefore $|c_1^*c_3| > |c_2c_3| \geq \text{sys}(X)/2$. Because $|c_1^*c_2^*| = 2L$ and $|c_2^*c_3| = L$, the angle $\angle c_2c_1^*c_3$ is acute. Thus, since the interior angle at c_1^* is at least π , the angle $\angle c_1c_1^*c_3$ greater than π . In particular, $|c_1c_3| > |c_1^*c_3|$, and so, in sum, the length of α is greater than $\text{sys}(X)/2$.

If D' contains c_6 , then the other component of $D \setminus \alpha$, is a pentagon with vertices c_1, c_3, c_3^*, c_4^* , and c_5^* . Using the triangle inequality, we have

$$L + |c_3c_5^*| \geq |c_3c_3^*| + |c_3^*c_5^*| \geq |c_3^*c_5^*| = 2|c_4c_5| \geq \text{sys}(X) = 2\ell + 2L,$$

and therefore $|c_3c_5^*| \geq 2\ell + L > \ell + L = \text{sys}(X)/2$.

Since $|c_3^*c_5^*| \geq \text{sys}(X) > 2L = |c_3^*c_4^*| = |c_4^*c_3^*|$, the angle $\angle c_3^*c_5^*c_3$ is less than $\pi/3$. Because $|c_3^*c_5^*| > 2L = |c_3^*c_3|$, we have $\angle c_3^*c_5^*c_3 < \pi/6$. Thus, since the interior angle at c_5^* is at least π , the angle $\angle c_1c_5^*c_3$ is greater than $\pi/2$. Therefore, $|c_1c_3| > |c_3c_5^*|$. In sum, $|c_1c_3| > \text{sys}(X)/2$, and hence α is not systolic. Therefore, there is no direct systolic Weierstrass arc joining c_1 to c_3 . A similar argument shows that there is no direct systolic Weierstrass arc joining c_1 to c_4 . \square

References

- [1] C. Boissy and S. Geninska, Systoles in translation surfaces, *preprint*. arXiv:1707.05060
- [2] S. A. Broughton and C. Judge, Ellipses in translation surfaces, *Geom. Dedicata*, **157** (2012), 111–151. Zbl 1241.57023 MR 2893481
- [3] M. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften, 319, Springer-Verlag, Berlin, 1999. Zbl 0988.53001 MR 1744486
- [4] L. Fejes Tóth, *Regular figures*, A Pergamon Press Book, The Macmillan Co., New York, 1964. Zbl 0134.15705 MR 165423
- [5] G. Forni and C. Matheus, Introduction to Teichmüller theory and its applications to dynamics of interval exchange transformations, flows on surfaces and billiards, *J. Mod. Dyn.*, **8** (2014), no. 3–4, 271–436. Zbl 1377.37057 MR 3345837
- [6] E. Gutkin and C. Judge, Affine mappings of translation surfaces: geometry and arithmetic, *Duke Math. J.*, **103** (2000), no. 2, 191–213. Zbl 0965.30019 MR 1760625
- [7] F. Jenni, Über den ersten Eigenwert des Laplace-Operators auf ausgewählten Beispielen kompakter Riemannscher Flächen, *Comment. Math. Helv.*, **59** (1984), no. 2, 193–203. Zbl 0541.30034 MR 749104
- [8] M. G. Katz and S. Sabourau, An optimal systolic inequality for CAT(0) metrics in genus two, *Pacific J. Math.*, **227** (2006), no. 1, 95–107. Zbl 1156.53019 MR 2247874
- [9] J. Malestein, I. Rivin, and L. Theran, Topological designs, *Geom. Dedicata*, **168** (2014), 221–233. Zbl 1284.57020 MR 3158040
- [10] H. Masur and J. Smillie, Hausdorff dimension of sets of nonergodic measured foliations, *Ann. of Math. (2)*, **134** (1991), no. 3, 455–543. Zbl 0774.58024 MR 1135877
- [11] C. T. McMullen, Dynamics of $SL_2(\mathbb{R})$ over moduli space in genus two, *Annals of Mathematics*, **165** (2007), no. 2, 397–456. Zbl 1143.37033 MR 2299738
- [12] P. M. Pu, Some inequalities in certain nonorientable Riemannian manifolds, *Pacific J. Math.*, **2** (1952), 55–71. Zbl 0046.39902 MR 48886

- [13] S. Sabourau, Systoles des surfaces plates singulières de genre deux, *Math. Z.*, **247** (2004), no. 4, 693–709. Zbl 1074.53034 MR 2077416
- [14] J. Smillie and B. Weiss, Characterizations of lattice surfaces, *Invent. Math.*, **180** (2010), no. 3, 535–557. Zbl 1195.57041 MR 2609249
- [15] W. P. Thurston, Shapes of polyhedra and triangulations of the sphere, in *The Epstein birthday schrift*, 511–549, Geom. Topol. Monogr., 1, Geom. Topol. Publ., Coventry, 1998. Zbl 0931.57010 MR 1668340
- [16] A. Wright, From rational billiards to dynamics on moduli spaces, *Bull. Amer. Math. Soc. (N.S.)*, **53** (2016), no. 1, 41–56. Zbl 1353.37076 MR 3403080

Received May 23, 2017

C. Judge, Department of Mathematics, Indiana University,
Bloomington, IN 47405-7106, USA

E-mail: cjudge@indiana.edu

H. Parlier, Mathematics Research Unit, University of Luxembourg,
L-4365 Esch-sur-Alzette, Luxembourg

E-mail: hugo.parlier@uni.lu