

Zeitschrift: Commentarii Mathematici Helvetici
Herausgeber: Schweizerische Mathematische Gesellschaft
Band: 94 (2019)
Heft: 3

Artikel: Counterexamples to the complement problem
Autor: Poloni, Pierre-Marie
DOI: <https://doi.org/10.5169/seals-846786>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.10.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Counterexamples to the complement problem

Pierre-Marie Poloni

Abstract. We provide explicit counterexamples to the so-called Complement Problem in every dimension $n \geq 3$, i.e. pairs of nonisomorphic irreducible algebraic hypersurfaces $H_1, H_2 \subset \mathbb{C}^n$ whose complements $\mathbb{C}^n \setminus H_1$ and $\mathbb{C}^n \setminus H_2$ are isomorphic. Since we can arrange that one of the hypersurfaces is singular whereas the other is smooth, we also have counterexamples in the analytic setting.

Mathematics Subject Classification (2010). 14R10; 14J26, 32M17.

Keywords. Affine algebraic geometry, complements of hypersurfaces, Danielewski surfaces.

1. Introduction

The Complement Problem (in the affine n -space) is one of the “challenging problems” considered by Hanspeter Kraft in his survey on affine algebraic geometry at the Bourbaki seminar [5]. It is formulated as follows.

Given two irreducible hypersurfaces $H_1, H_2 \subset \mathbb{C}^n$ and an isomorphism of their complements, does it follow that H_1 and H_2 are isomorphic?

Let us specify that we work here in the context of algebraic geometry. In particular, the hypersurfaces considered above are algebraic, i.e. defined as the zero sets of some polynomials $f_1, f_2 \in \mathbb{C}[x_1, \dots, x_n]$, and the isomorphisms are isomorphisms of algebraic varieties. Moreover, we recall that the complement $\mathbb{C}^n \setminus H$ of an hypersurface $H \subset \mathbb{C}^n$ is also an affine algebraic variety.

The Complement Problem is a very natural question: We want to retrieve some information about a subvariety $X \subset M$ from its complement $M \setminus X$. Such questions make of course sense in various contexts (as e.g. in knot theory, see [3] and [4]). Closer to our immediate interests, Jérémy Blanc [1] gave counterexamples to the Complement Problem for curves in the projective plane \mathbb{P}^2 . Actually, his main motivation was to disprove another conjecture, due to Hisao Yoshihara [8], which stated that if two irreducible curves $\Gamma_1, \Gamma_2 \subset \mathbb{P}^2$ have isomorphic complements $\mathbb{P}^2 \setminus \Gamma_1 \simeq \mathbb{P}^2 \setminus \Gamma_2$, then they should be equivalent, i.e. there should exist an automorphism of \mathbb{P}^2 sending Γ_1 onto Γ_2 .

The purpose of this note is to answer Kraft's Complement Problem in the negative for every $n \geq 3$. More precisely, we will give explicit counterexamples of several different types, as described in the main theorem below.

Theorem 1.1. *For every integer $n \geq 3$, there exist examples of:*

- (1) *irreducible hypersurfaces $H_1, H_2 \subset \mathbb{C}^n$ with isomorphic complements $\mathbb{C}^n \setminus H_1 \simeq \mathbb{C}^n \setminus H_2$ such that H_1 and H_2 are smooth and nonisomorphic;*
- (2) *irreducible hypersurfaces $H_1, H_2 \subset \mathbb{C}^n$ with isomorphic complements $\mathbb{C}^n \setminus H_1 \simeq \mathbb{C}^n \setminus H_2$ such that H_1 is smooth but H_2 is singular;*
- (3) *irreducible hypersurfaces $H_1, H_2 \subset \mathbb{C}^n$ with isomorphic complements $\mathbb{C}^n \setminus H_1 \simeq \mathbb{C}^n \setminus H_2$ such that H_1 and H_2 are isomorphic, although there is no automorphism of \mathbb{C}^n mapping H_1 onto H_2 .*

At the time that the first version of the present paper was being written, the case of irreducible curves on \mathbb{C}^2 was still wide open. But since then it has been solved, again in the negative, by Blanc, Furter and Hemmig in a remarkable paper [2] in which they make use of totally different methods to find counterexamples to the Complement Problem in the case where $n = 2$.

We remark that the second kind of examples in above Theorem 1.1 provide counterexamples to the Complement Problem in the analytic setting too. On the other hand, the nonisomorphic algebraic varieties that we will produce in case (1) are biholomorphic, and we do not get any examples of smooth affine algebraic varieties $V_1, V_2 \subset \mathbb{C}^n$ (i.e. of affine algebraic manifolds in \mathbb{C}^n), which are not biholomorphic, although their complements $\mathbb{C}^n \setminus V_1$ and $\mathbb{C}^n \setminus V_2$ are. By contrast, it is shown in [2] that if two nonisomorphic irreducible affine curves have isomorphic complements in \mathbb{C}^2 , then they are necessarily smooth and biholomorphic.

All our examples will be realized as hypersurfaces of \mathbb{C}^{m+2} defined by an equation of the form $x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m (z^2 - \alpha)^k = \alpha$ for some integer $k \geq 0$ and some constant $\alpha \in \mathbb{C}$. These varieties were first studied by Lucy Moser-Jauslin and the author in [6] for the case where $m = 1$ and then in [7] for the general case. In particular, it was observed that there exist such polynomials, say P and Q , whose zero sets $\{P = 0\}$ and $\{Q = 0\}$ are not isomorphic, whereas their other fibers $\{P = c\}$ and $\{Q = c\}$ are isomorphic for all $c \in \mathbb{C}^*$. The main ingredient of the present paper will be to use the isomorphisms $\{P = c\} \simeq \{Q = c\}$ to construct an isomorphism between the complements $\mathbb{C}^{m+2} \setminus \{P = 0\}$ and $\mathbb{C}^{m+2} \setminus \{Q = 0\}$.

Acknowledgements. The author thanks the referee for helpful comments and suggestions, and for pointing out an error in a previous version of Proposition 3.4.

2. Preliminaries

Let us start by recalling some notations and results from [7] that we will use in the sequel.

Throughout this paper, we fix an integer $m \geq 1$ and a coordinate system x_1, \dots, x_m, y, z on the complex affine space \mathbb{C}^{m+2} . If $P \in \mathbb{C}[x_1, \dots, x_m, y, z]$, then $V(P)$ denotes the zero set of P in \mathbb{C}^{m+2} .

Notation 2.1. Given a polynomial $q(t) \in \mathbb{C}[t]$, we denote by P_q the polynomial of $\mathbb{C}[x_1, \dots, x_m, y, z]$ defined by

$$P_q = x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m q(z^2).$$

It was shown in [7] that the algebraic varieties $V(P_0)$, $V(P_0 - 1)$, $V(P_1)$, and $V(P_1 - 1)$ are pairwise nonisomorphic. Moreover, every fiber $V(P_q - c) = P_q^{-1}(c) \subset \mathbb{C}^{m+2}$ is isomorphic to one of these four and we have the following classification result.

Proposition 2.2 ([7, Lemma 2.2 and Proposition 2.5]). *Let $q(t) \in \mathbb{C}[t]$ and $c \in \mathbb{C}$. Then, the variety $V(P_q - c)$ is isomorphic to $V(P_{q(c)} - c)$. Moreover, the latter is isomorphic to:*

- $V(P_0)$ if and only if $c = 0$ and $q(c) = 0$;
- $V(P_0 - 1)$ if and only if $c \neq 0$ and $q(c) = 0$;
- $V(P_1)$ if and only if $c = 0$ and $q(c) \neq 0$;
- $V(P_1 - 1)$ if and only if $c \neq 0$ and $q(c) \neq 0$.

Finally, we recall the classification of the hypersurfaces $V(P_q - c) \subset \mathbb{C}^{m+2}$ up to equivalence, i.e. up to automorphisms of the ambient space.

Proposition 2.3 ([7, Proposition 3.2]). *Let $q_1(t), q_2(t) \in \mathbb{C}[t]$ be two polynomials and $c_1, c_2 \in \mathbb{C}$ be two constants. Then, the following are equivalent.*

- (1) *There exists an algebraic automorphism of \mathbb{C}^{m+2} which maps the hypersurface $V(P_{q_1} - c_1)$ onto $V(P_{q_2} - c_2)$.*
- (2) *There exist $\lambda, \mu \in \mathbb{C}^*$ such that $c_2 = \mu^{-1}c_1$ and $q_2(t) = \lambda q_1(\mu t)$.*

3. Explicit examples

All our examples will consist in hypersurfaces $H_{\alpha,k}$ in \mathbb{C}^{m+2} defined by an equation of the form

$$x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m (z^2 - \alpha)^k = \alpha$$

for some integer $k \geq 0$ and some constant $\alpha \in \mathbb{C}$. By Proposition 2.2, the variety $H_{\alpha,k} = V(P_{(t-\alpha)^k} - \alpha)$ is isomorphic to:

$$\begin{cases} V(P_0) & \text{if } \alpha = 0 \text{ and } k \geq 1, \\ V(P_0 - 1) & \text{if } \alpha \neq 0 \text{ and } k \geq 1, \\ V(P_1) & \text{if } \alpha = 0 \text{ and } k = 0, \\ V(P_1 - 1) & \text{if } \alpha \neq 0 \text{ and } k = 0. \end{cases}$$

In particular, since $V(P_0)$, $V(P_0 - 1)$, $V(P_1)$ and $V(P_1 - 1)$ are pairwise nonisomorphic, we observe that $H_{\alpha,k} \not\simeq H_{\alpha,0}$ if $k \neq 0$.

Lemma 3.1. *The hypersurfaces $H_{\alpha,k}$ and $H_{\alpha,k'}$ have isomorphic complements, i.e. $\mathbb{C}^{m+2} \setminus H_{\alpha,k} \simeq \mathbb{C}^{m+2} \setminus H_{\alpha,k'}$ for all $\alpha \in \mathbb{C}$ and all $k, k' \geq 0$.*

Proof. Following the notation of the previous section, we have that $H_{\alpha,k} = V(P_{q_k} - \alpha)$, where $q_k(t) = (t - \alpha)^k \in \mathbb{C}[t]$. To prove that $H_{\alpha,k}$ and $H_{\alpha,k'}$ have isomorphic complements, it suffices to prove that $\mathbb{C}^{m+2} \setminus H_{\alpha,k} \simeq \mathbb{C}^{m+2} \setminus H_{\alpha,0}$ for all $\alpha \in \mathbb{C}$ and all $k \geq 1$. We do this by giving an explicit isomorphism.

We set $P = P_{q_k} - \alpha$ and $Q = P_{q_0} - \alpha$, so that the coordinate rings of $\mathbb{C}^{m+2} \setminus H_{\alpha,k}$ and $\mathbb{C}^{m+2} \setminus H_{\alpha,0}$ are isomorphic to the rings $\mathbb{C}[x_1, \dots, x_m, y, z, \frac{1}{P}]$ and $\mathbb{C}[x_1, \dots, x_m, y, z, \frac{1}{Q}]$, respectively.

Next, we consider the morphisms

$$\begin{aligned} \Phi : \quad \mathbb{C}^{m+2} \setminus H_{\alpha,0} &\rightarrow \mathbb{C}^{m+2} \\ (x_1, \dots, x_m, y, z) &\mapsto \left(\frac{x_1}{Q^k}, x_2, \dots, x_m, yQ^{2k} + Q^k \frac{Q^k - (z^2 - \alpha)^k}{x_1 \cdots x_m}, z \right) \end{aligned}$$

and

$$\begin{aligned} \Psi : \quad \mathbb{C}^{m+2} \setminus H_{\alpha,k} &\rightarrow \mathbb{C}^{m+2} \\ (x_1, \dots, x_m, y, z) &\mapsto \left(P^k x_1, x_2, \dots, x_m, \frac{1}{P^{2k}} \left(y - \frac{P^k - (z^2 - \alpha)^k}{x_1 \cdots x_m} \right), z \right) \end{aligned}$$

We remark that the above morphisms are well defined, since

$$\frac{Q^k - (z^2 - \alpha)^k}{x_1 \cdots x_m} \quad \text{and} \quad \frac{P^k - (z^2 - \alpha)^k}{x_1 \cdots x_m}$$

are both elements of $\mathbb{C}[x_1, \dots, x_m, y, z]$.

One checks by a straightforward calculation that $P \circ \Phi = Q$ and $Q \circ \Psi = P$. This shows that

$$\Phi(\mathbb{C}^{m+2} \setminus H_{\alpha,0}) \subset \mathbb{C}^{m+2} \setminus H_{\alpha,k} \quad \text{and} \quad \Psi(\mathbb{C}^{m+2} \setminus H_{\alpha,k}) \subset \mathbb{C}^{m+2} \setminus H_{\alpha,0}.$$

Finally, one easily checks that

$$\Phi \circ \Psi = \text{id}_{\mathbb{C}^{m+2} \setminus H_{\alpha,k}} \quad \text{and} \quad \Psi \circ \Phi = \text{id}_{\mathbb{C}^{m+2} \setminus H_{\alpha,0}}. \quad \square$$

Combining Proposition 2.2 with Lemma 3.1, we obtain the following counterexamples to the Complement Problem.

Proposition 3.2. *Let $m \geq 1$ and let H_1 and H_2 be the irreducible hypersurfaces of \mathbb{C}^{m+2} that are defined by the equations*

$$x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m (z^2 - 1) = 1$$

and

$$x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m = 1,$$

respectively. Then, H_1 and H_2 are smooth and not isomorphic, although they have isomorphic complements $\mathbb{C}^{m+2} \setminus H_1 \simeq \mathbb{C}^{m+2} \setminus H_2$.

Proof. On the one hand, Proposition 2.2 implies that the hypersurfaces $H_1 \simeq V(P_0 - 1)$ and $H_2 = V(P_1 - 1)$ are not isomorphic. On the other hand, their complements are isomorphic by Lemma 3.1. \square

Remark 3.3. Even if they are not isomorphic as algebraic varieties, the above hypersurfaces $H_1 \simeq V(P_0 - 1)$ and $H_2 = V(P_1 - 1)$ are biholomorphic [7, Remark 2.6].

We now give counterexamples in the analytic category. For this, we remark that the hypersurface $H_{0,0}$ is smooth in the case where $m = 1$. Nevertheless, by Lemma 3.1, its complement $\mathbb{C}^3 \setminus H_{0,0}$ in \mathbb{C}^3 is isomorphic to that of the singular hypersurface $H_{0,1}$. Considering the cylinders over these two hypersurfaces, we obtain nonbiholomorphic counterexamples to the Complement Problem in any dimension $n \geq 3$.

Proposition 3.4. *Let $m = 1$ and denote by S_1 and S_2 the irreducible hypersurfaces of $\mathbb{C}^{m+2} = \mathbb{C}^3$ that are defined by the equations*

$$x_1^2 y + z^2 + x_1 z^2 = 0$$

and

$$x_1^2 y + z^2 + x_1 = 0,$$

respectively. Let $m' \geq 0$ be any nonnegative integer and consider the hypersurfaces $H'_1 = S_1 \times \mathbb{C}^{m'}$ and $H'_2 = S_2 \times \mathbb{C}^{m'}$ in $\mathbb{C}^{m'+3}$.

Then, the complements $\mathbb{C}^{m'+3} \setminus H'_1$ and $\mathbb{C}^{m'+3} \setminus H'_2$ are isomorphic. However, since H'_1 is singular and H'_2 is smooth, H'_1 and H'_2 are not biholomorphic.

Proof. It is straightforward to check that S_1 is singular and that S_2 is smooth. Hence, H'_1 is singular and H'_2 is smooth. Since $S_1 = H_{0,1}$ and $S_2 = H_{0,0}$, their complements $\mathbb{C}^3 \setminus S_1$ and $\mathbb{C}^3 \setminus S_2$ are isomorphic by Lemma 3.1. This implies that H'_1 and H'_2 have isomorphic complements in $\mathbb{C}^{m'+3}$. \square

Let us conclude by giving, thanks to Proposition 2.3, an example of two smooth nonequivalent hypersurfaces which are isomorphic and have isomorphic complements.

Proposition 3.5. *Let $m \geq 1$ and let H_1'' and H_2'' be the hypersurfaces of \mathbb{C}^{m+2} that are defined by the equations*

$$x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m (z^2 - 1) = 1$$

and
$$x_1^2 \cdots x_m^2 y + z^2 + x_1 \cdots x_m (z^2 - 1)^2 = 1,$$

respectively. Then, H_1'' and H_2'' are smooth irreducible varieties which are isomorphic and have isomorphic complements in \mathbb{C}^{m+2} . Nevertheless, no automorphisms of \mathbb{C}^{m+2} map H_1'' onto H_2'' .

Proof. Proposition 2.3 shows that the hypersurfaces $H_1'' = V(P_{(t-1)} - 1) = H_{1,1}$ and $H_2'' = V(P_{(t-1)^2} - 1) = H_{1,2}$ are not equivalent. Nevertheless, their complements are isomorphic by Lemma 3.1. \square

References

- [1] J. Blanc, The correspondence between a plane curve and its complement, *J. Reine Angew. Math.*, **633** (2009), 1–10. Zbl 1203.14035 MR 2561193
- [2] J. Blanc, J.-P. Furter, and M. Hemmig, Exceptional isomorphisms between complements of affine plane curves, *Duke Math. J.*, **168** (2019), no. 12, 2235–2297. MR 3999446
- [3] S. E. Cappell and J. L. Shaneson, There exist inequivalent knots with the same complement, *Ann. of Math. (2)*, **103** (1976), no. 2, 349–353. Zbl 0338.57008 MR 413117
- [4] C. McA. Gordon and J. Luecke, Knots are determined by their complements, *J. Amer. Math. Soc.*, **2** (1989), no. 2, 371–415. Zbl 0678.57005 MR 965210
- [5] H. Kraft, Challenging problems on affine n -space, in *Séminaire Bourbaki, Vol. 1994/95*, Exp. No. 802, 5, 295–317, Astérisque, 237, 1996. Zbl 0892.14003 MR 1423629
- [6] L. Moser-Jauslin and P.-M. Poloni, Embeddings of a family of Danielewski hypersurfaces and certain \mathbb{C}^+ -actions on \mathbb{C}^3 , *Ann. Inst. Fourier (Grenoble)*, **56** (2006), no. 5, 1567–1581. Zbl 1120.14056 MR 2273864
- [7] P.-M. Poloni, A note on the stable equivalence problem, *J. Math. Soc. Japan*, **67** (2015), no. 2, 753–761. Zbl 1338.14058 MR 3340194
- [8] H. Yoshihara, On open algebraic surfaces $\mathbb{P}^2 - C$, *Math. Ann.*, **268** (1984), no. 1, 43–57. Zbl 0523.14013 MR 744327

Received December 15, 2016

P.-M. Poloni, Mathematisches Institut, Universität Bern,
 Sidlerstrasse 5, CH-3012 Bern, Switzerland
 E-mail: pierre.poloni@math.unibe.ch